

POSTER PRESENTATION

Open Access

Adsorption of amino acids on MFI-type zeolite: DFT calculations and experimental results

Kai Stueckenschneider^{1*}, Achim Zielesny², Gerhard Schembecker¹

From 7th German Conference on Chemoinformatics: 25 CIC-Workshop
Goslar, Germany. 6-8 November 2011

Adsorption is a common unit operation in separation and purification of biotechnological products where chromatography steps can make up more than half the amount of the total purification costs [1]. The molecular mechanisms of adsorption are still not understood in detail. Further understanding of interactions between adsorbent surfaces and adsorptives could help to facilitate process design in a more cost efficient manner.

In this work the interaction of MFI-type zeolite MFI-27 (Al/Si=13) with Alanine and Phenylalanine is investigated by quantum chemical (QM) calculations which are compared to experimental adsorption data.

For the QM calculations T3-clusters are used as MFI-27 surface models which were shown to be successful e. g. in the case of protolytic cracking of alkanes [2,3]. In order to model different pH-values Alanine and Phenylalanine are applied in their protonated, zwitterionic and deprotonated state. Geometry optimisations and frequency analysis of all molecular structures are performed with Density Functional Theory (DFT) using the B3LYP functional with different basis sets. Calculated complex energies are corrected for BSSE and ZPE. Adsorption isotherms are derived from corresponding experiments.

Regarding the adsorption isotherms it is shown that a high adsorption of Alanine and Phenylalanine on MFI-27 takes place at low pH values (near the pK_a of the amino acids). Less adsorption occurs with an increased pH equalling the amino acids' isoelectric points. At the pH of the amino acids' pK_b values adsorption is no longer observed.

These trends can be correlated with the corresponding QM calculations. High binding energies are calculated

for the protonated amino acids. Zwitterionic states lead to lower binding energies. The deprotonated amino acids do not show any binding affinity to MFI-27.

These first results indicate the reliability of the applied methods for our model system.

Author details

¹Department of Biochemical and Chemical Engineering, TU Dortmund, Dortmund, 44227, Germany. ²Institute for Bioinformatics and Chemoinformatics, University of Applied Sciences Gelsenkirchen, Recklinghausen, 45665, Germany.

Published: 1 May 2012

References

1. Ladisch MR: *Bioseparations Engineering*. Wiley and Sons; 2001, 1-5.
2. Zheng X, Blowers P: *Reactivity of Alkanes on Zeolites: A Computational Study of Propane Conversion Reactions*. *J Phys Chem A* 2005, **109**:10734-10741.
3. Zheng X, Blowers P: *Reactivity of Isobutane on Zeolites: A First Principles Study*. *J Phys Chem A* 2006, **110**:2455-2460.

doi:10.1186/1758-2946-4-S1-P38

Cite this article as: Stueckenschneider et al.: Adsorption of amino acids on MFI-type zeolite: DFT calculations and experimental results. *Journal of Cheminformatics* 2012 **4**(Suppl 1):P38.

Publish with **ChemistryCentral** and every scientist can read your work free of charge

"Open access provides opportunities to our colleagues in other parts of the globe, by allowing anyone to view the content free of charge."

W. Jeffery Hurst, The Hershey Company.

- available free of charge to the entire scientific community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
<http://www.chemistrycentral.com/manuscript/>


ChemistryCentral

* Correspondence: Kai.Stueckenschneider@bci.tu-dortmund.de

¹Department of Biochemical and Chemical Engineering, TU Dortmund, Dortmund, 44227, Germany

Full list of author information is available at the end of the article