

Article

Application of the RDoC Framework to Predict Alcohol Use and Suicidal Thoughts and Behaviors among Early Adolescents in the Adolescent Brain and Cognitive Development (ABCD) Study

Laika D. Aguinaldo , Clarisa Coronado, Diego A. Gomes, Kelly E. Courtney  and Joanna Jacobus *

Department of Psychiatry, University of California, La Jolla, CA 92093, USA;
laaguinaldo@health.ucsd.edu (L.D.A.); clari.coronado@gmail.com (C.C.); gomesadiego@gmail.com (D.A.G.);
kecourtney@health.ucsd.edu (K.E.C.)

* Correspondence: jjacobus@health.ucsd.edu

Abstract: Alcohol use confers risk for suicidal thoughts and behaviors (ideation, attempt) in early adolescents. The Research Domain Criteria provides a framework for examination of multidimensional and modifiable risk factors. We examined distinct latent profiles based on patterns of positive valence (reward responsivity) and cognitive systems (neurocognition) from the ABCD Study (age 9–10, $N = 10,414$) at baseline enrollment. Longitudinal associations were determined between baseline positive valence and cognitive profiles and group classification (alcohol use, suicidal thoughts and behaviors, or their co-occurrence) two-years after initial assessment (ages 11–12). Three unique profiles of positive valence, cognition, alcohol use, and suicidal thoughts and behaviors were identified. Two baseline profiles predicted alcohol use and suicidal thoughts and behaviors, two-years after initial assessment. Low positive valence with high cognition (but low impulsivity) predicted alcohol use ($OR = 1.414, p \leq 0.001$), while high positive valence with low cognition (but high impulsivity) predicted suicidal thoughts and behaviors ($OR = 1.25, p = 0.038$), compared to average positive valence and cognition. Unique profiles of positive valence and cognitive systems among 9–12-year-olds may be predictive of alcohol use and suicidal thoughts and behaviors over a two-year period. Findings underscore the potential for trajectory research on positive valence and cognitive profiles to enhance prevention for early-adolescents.

Keywords: RDoC; suicide prevention; suicide intervention; pre-teen youth; alcohol use



Citation: Aguinaldo, L.D.; Coronado, C.; Gomes, D.A.; Courtney, K.E.; Jacobus, J. Application of the RDoC Framework to Predict Alcohol Use and Suicidal Thoughts and Behaviors among Early Adolescents in the Adolescent Brain and Cognitive Development (ABCD) Study. *Brain Sci.* **2022**, *12*, 935. <https://doi.org/10.3390/brainsci12070935>

Academic Editor: Matteo Marti

Received: 21 May 2022

Accepted: 14 July 2022

Published: 17 July 2022

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Alcohol use is linked to suicidal thoughts and behaviors during early adolescence (pubescence) [1]. In our prior research, early adolescents who reported lifetime alcohol use (i.e., ≥ 1 sip) showed a nearly two-fold increase in their odds of lifetime suicidal ideation and attempts, compared to early adolescents with no previous alcohol use [1]. Given that the alcohol use (ALC)–suicidal thoughts and behaviors (STB) association can be detected in early adolescence (as young as age 9), it is important to determine the mechanisms that may increase vulnerability to the ALC–STB association, given that the comorbidity is particularly deadly [2]. However, the extant literature on the ALC–STB association examines the same set of risk factors (e.g., internalizing, and externalizing symptoms, STB history, and demographic and culture/environment characteristics) studied in isolation often using cross-sectional designs with older adolescents, adult populations, and/or small sample sizes [3–8]. There is a need for research on the ALC–STB association that characterizes novel risk factors, examines the relationships that exist between them, and utilizes prospective designs with populations underrepresented in the comorbid ALC and STB literature (e.g., early-adolescents).

The National Institute of Mental Health has developed a research framework for investigating mental disorders, the Research Domain Criteria (RDoC). The goal of RDoC is to overcome one-dimensional limitations, and promote novel approaches that will lead to better diagnosis, prevention, and intervention. RDoC is a conceptual research framework that prioritizes neurobiological and transdiagnostic risk factors that are present across mental illness (e.g., STBs) [8–10], with a focus on risk factor integration across human functioning domains. The framework currently includes six major functional domains (arousal/regulatory, positive valence, negative valence, sensorimotor, cognitive, and social processes). Different aspects of each domain are represented by three to six psychological/biological dimensions, or constructs, which are studied along the full range of functioning from normal to abnormal. Both behavioral and biological aspects of functioning change and mature throughout pre- to post-pubescence; thus, research during these critical development periods is important. The RDoC framework encourages researchers to measure and integrate many classes of variables (units of analysis, e.g., behavioral, physiological, and self-report data) in order to seek a comprehensive understanding of the construct(s) under study. Advancing the ALC-STB association literature by using this framework, coupled with advanced statistical methods, allows for the exploration of brain–behavior relationships using complex computational approaches to explore increasing risk for STBs [8,11].

Associations between the RDoC cognitive and positive valence system domains are understudied. Positive valence systems (PVS) are responsible for responses to positive motivational situations or contexts, such as reward-seeking, consummatory behavior, and reward/habit learning. A recent review by Glenn et al. (2018) used RDoC to examine relationships between cognitive and positive valence risk factors and STBs in adult samples [8]. Poorer working memory increased the risk for STBs, while findings for the association between cognitive control (e.g., impulsiveness, problem-solving (approach avoidance, personal control)) and STBs were mixed [8]. In the domain of positive valence, adolescents who self-reported STBs demonstrated blunted responses to reward responsiveness [7,12–18]. In a recent study that used an RDoC approach to test blunted positive valence in 57 adult men and women heavy binge drinkers compared to healthy controls during functional magnetic resonance imaging, binge alcohol drinkers showed abnormally blunted activity in PVS regions, such as the striatum, compared with healthy controls [19]. PVS brain regions are well established in the adult literature (e.g., striatum, dopamine, and opioid system) [19–21]. However, research is needed to further examine these understudied neuroscience-based risk factors using statistical models that consider complex relationships among several variables simultaneously and in pre–post pubescent samples [7,8,12–14,22,23].

Latent Profile Analysis (LPA) is a statistical approach that considers complex relationships among several predictors simultaneously. LPA has been used in adult populations to identify profiles (i.e., subgroups) of people who engage in STBs. Prior research has focused on patterns among self-reports of thoughts of death and suicide and healthy lifestyle patterns [24–31]. To our knowledge, only one study exists examining suicidality using the RDoC framework and LPA [10]. Podlogar and colleagues (2018) used LPA to assess overlapping and distinct features of depression and anxiety in relation to suicide risk among 616 adult outpatients [10]. Results demonstrated that those identified as showing a higher suicide risk profile reported significantly higher levels of negative affect and anxious distress and reported significantly lower levels of positive affect than other classes [10]. To date, the application of LPA models to studies of child and adolescent STBs within the RDoC framework is an understudied area of research.

Early life (prepubescent) ALC, STBs, reward responsivity, and cognitive control are critical risk factors that share similar characteristics and predict addictive and suicidal behaviors [1,3,7,8,12–18,22,23]. Driven by prior research, theory, and parsimony, a priori construct selection drove our hypothesis and aims [1,7,8,12–18,22,23,32]. The primary aims of this study were to examine latent profiles of (1) positive valence (PVS) and cognitive (CS) systems, ALC, and STBs among 9- to 10-year-olds in the ABCD baseline cohort and (2) examine if PVS, CS, ALC, and STB profiles prospectively predict ALC, STBs, and their

co-occurrence two years after initial assessment. We hypothesized that profiles of high PVS and low CS would predict ALC, STBs, and their co-occurrence and that these patterns in profiles could predict future ALC, STBs, and their co-occurrence.

2. Materials and Methods

2.1. Protocol

ABCD is a 10-year longitudinal study conducted across 21 sites in the United States that recruited 11,878 participants and is funded by the National Institutes of Health [33,34]. The Institutional Review Board at the University of California, San Diego, approved the study, and each study site has a detailed protocol to address reports of STBs. A detailed account of the recruitment strategy has been previously published [35–37]. ABCD primarily utilized a probability sample recruited through schools, with school selection based on sex, race and ethnicity, socioeconomic status, and urbanicity.

Participants and their parents or guardians attended study session(s) at their local research site to complete the baseline and follow-up visits. Parents/guardians provided written consent, with each early adolescent providing written assent. Participants and their parent or guardian were in separate private rooms during study participation to maintain confidentiality. The baseline and year two measures included self- and parent/guardian-report questionnaires, neurocognitive testing, biological samples, and an MRI scan [38–40]. Study assessments were completed over an 8-h research session (or two 4-h sessions), and parents and youths were compensated financially. The cross-sectional and longitudinal analyses presented herein drew from self- and parent-report questionnaires and neurocognitive assessments from the baseline and year two follow-up data of the ABCD data (Release 4.0).

2.2. Measures

2.2.1. Positive Valence Systems (PVS; Trait-Based Measures)

Sensitivity to Reward and Punishment: The 24 item BIS/BAS survey was administered to all youth to assess for avoidance (BIS) and approach (BAS) sensitivities reflective of motivational traits [41–44]. The cross-sectional LPA included the BIS summary and BAS reward responsiveness subscale scores. Scoring is based on youth self-reports per subscale (BIS, BAS reward responsiveness), such that high values on BIS correspond to high avoidance sensitivities, low values on BIS correspond to low avoidance sensitivities, and high values on BAS reward responsiveness correspond to high approach sensitivities to reward responsiveness, and low values on BAS reward responsiveness correspond to low approach sensitivities to reward responsiveness.

2.2.2. NIH Neurocognitive Toolbox (Cognitive Systems; CS)

Inhibitory Control and Attention-Flanker (uncorrected standard score): The NIH Toolbox Flanker Inhibition Control and Attention Task, a variant of the Eriksen Flanker task, was used to measure attention and the ability to inhibit automatic responses that interfere with achieving goals [45]. Participants were presented with five arrows on the iPad screen with four flanking stimuli (two on the outer left and two on the outer right) all facing the same way, either left or right. The middle arrow either faced the same way (congruent trial) or a different way (incongruent trial) as the flanking stimuli. Participants pressed a button to indicate whether the middle stimuli faced left or right. The word MIDDLE also appeared on the screen for all participants. For participants ages 8 to 11 years, an audio recording stated “MIDDLE” to remind participants where to focus. Scoring is based on speed and accuracy (e.g., higher normative scores indicate better ability to attend to relevant stimuli and ignore irrelevant stimuli), with high values corresponding to higher cognitive control and attention, and low values corresponding to lower cognitive control and attention.

Processing Speed and Information Processing (uncorrected standard score): The NIH Toolbox Pattern Comparison Processing Speed Test[®] [46] was used as the measure of rapid visual processing. Participants were shown two pictures and asked to determine whether

the pictures were the same or not. High scores can be interpreted as a better and faster processing speed and low scores can be interpreted as a slower processing speed.

Episodic Learning and Memory-Picture Sequence Memory Task (uncorrected standard score; PSMT): The PSMT is modeled from memory tests asking children to imitate a sequence of actions using props and measures episodic memory for a sequence of pictured events [47,48]. Participants were presented with a series of pictures, not in any intrinsic order, depicting activities or events that could occur in a particular setting (i.e., going to the fair) [47,49]. After the last item was presented, the pictures were randomly displayed in the center of the screen, and the participants were asked to place the pictures in the same order they were presented. There were three different sets of test items for this measure: (1) Play in the Park (Form A), (2) Go to the Fair (Form B), and (3) Work on the Farm (Form C); each form consists of different test items that yield equivalent scores and can be used in a repeated-measures research design to minimize practice effects [47,48]. Form A (Play in the Park) was administered to all participants at baseline and Form B (Go to the Fair) was administered to all participants at the two-year follow-up. Scoring is based on the total number of adjacent pairs of pictures placed correctly across two learning trials (high values are interpreted as a better working memory, while low values are interpreted as a poorer episodic memory).

Verbal Learning. Rey Auditory Verbal Learning Test (short and long delay total correct raw score): The Rey Auditory Verbal Learning Test (RAVLT) was used as a measure of learning and memory across the lifespan that is sensitive to various influences, including psychopathology across development [50–52]. A customized automated version of the RAVLT, created with Pearson's Q-interactive platform, was used [53]. Participants listened to and recalled a list of 15 unrelated words over five learning trials. Following the fifth initially learned list, participants are asked to listen and recall a distractor list, then they are asked to recall the initially learned list. To assess longer-term retention, recall is reassessed following a 30-min delay during which other non-verbal tasks from the neurocognitive battery were administered. Alternate forms were used at baseline and two-year follow-up to facilitate longitudinal testing [38]. Q-interactive automatically calculated the number of correctly recalled words for each trial and the number of perseverations and intrusions. A higher uncorrected raw score represents better verbal learning and recall performance, while lower uncorrected raw score represents poorer verbal learning and recall performance.

Impulsivity: The shortened 20-item youth version of the Urgency, Premeditation, Perseverance, Sensation Seeking, and Positive Urgency (UPPS-P) [54,55] scale was administered to youth at baseline and two-year follow-up. Recent research has demonstrated no significant differences in psychometric performance of this measure across gender, race/ethnicity, and socioeconomic status amongst youths [55]. High scores on each subscale can be interpreted as higher impulsivity, and low scores on subscales can be interpreted as lower impulsivity.

2.2.3. Measures of Alcohol Use, Suicidal Thoughts and Behaviors, and Demographic Covariates

Lifetime Low-Level Alcohol Use (ALC): Youth completed the iSay Sip Inventory to characterize participants' endorsement of low-level alcohol use (any alcohol drink and/or sip) [52].

Suicidal Thoughts and Behaviors (STBs): Youths' reports of lifetime SI and SA were gathered from a computerized version of the Kiddie-Schedule for Affective Disorders and Schizophrenia (K-SADS) [56]. Participants at baseline and year two follow-up of the ABCD cohort were separated into three groups: (1) lifetime low-level alcohol use and no lifetime STBs were categorized as ALC, (2) no lifetime low-level alcohol use and lifetime STBs were categorized as STBs, and (3) the co-occurrence of lifetime low-level alcohol use and lifetime STBs were categorized as ALC + STBs.

Demographic Covariates: Demographic covariates were chosen based on prior evidence of associations with the ALC and STB variables [53,57–66] (Table 1). The parent-reported demographic variables of age, sex (assigned at birth), parental education, and ethno-racial identity, are items extracted from the PhenX toolkit [62].

Table 1. Descriptive statistics of the positive valence systems (PVS) and neurocognitive systems (CS) profiles of suicidal thoughts and behaviors (STBs) at baseline enrollment ($n = 10,414$) with children (age 9 to 10 years) in the Adolescent Brain and Cognitive Development (ABCD) study.

Characteristics (<i>n</i> , %/Mean, Std)	Total Sample <i>N</i> = 10,414	Profile 1 Average PVS and CS <i>n</i> = 4940 (Reference Group)	Profile 2 High PVS and Low CS (But High Impulsivity) <i>n</i> = 1470	Profile 3 Low PVS and High CS (But Low Impulsivity) <i>n</i> = 4004	Tests for Group Differences [†]	
					Profile 1 vs. 2 <i>p</i> Value	Profile 1 vs. 3 <i>p</i> Value
Age (years)	9.9 (0.6)	9.9 (0.6)	9.8 (0.6)	10.0 (0.6)	≤0.001	≤0.001
Sex (Biological)						
Female	4961 (47.6%)	2245 (45.4%)	568 (38.6%)	2148 (53.6%)	≤0.001	≤0.001
Male	5453 (52.4%)	2695 (54.6%)	902 (61.4%)	1856 (46.4%)		
Parental Education						
>High School Graduate	7106 (68.3%)	3210 (65%)	761 (51.8%)	3138 (78.4%)	≤0.001	≤0.001
<High School Graduate	3308 (31.7%)	1730 (35%)	709 (48.2%)	866 (21.6%)		
Ethno-Racial Identity						
White	5602 (53.8%)	2508 (50.8%)	525 (35.7%)	2569 (64.2%)	≤0.001	≤0.001
Hispanic/Latino	2086 (20.0%)	1088 (22%)	332 (22.6%)	666 (16.6%)		
Black	1422 (13.7%)	758 (15.3%)	432 (29.4%)	232 (5.8%)		
Asian/Pacific Islander	216 (2.1%)	85 (1.7%)	17 (1.2%)	114 (2.8%)		
Other	1088 (10.4%)	501 (10.1%)	164 (11.2%)	423 (10.6%)		
Lifetime Alcohol Use (ALC)						
Yes (>1 Sip of Alcohol)	985 (9.5%)	366 (7.4%)	83 (5.6%)	536 (13.4%)	≤0.001	≤0.001
Lifetime Suicidal Thoughts and Behaviors (STBs)						
Yes	739 (7.1%)	336 (6.8%)	143 (9.7%)	263 (6.6%)	≤0.001	≤0.001
Lifetime Co-Occurring ALC + STBs						
Yes	177 (1.7%)	83 (1.7%)	22 (1.5%)	71 (1.8%)	0.779	<0.001

Table 1. Cont.

Characteristics (<i>n</i> , %/Mean, Std)	Total Sample <i>N</i> = 10,414	Profile 1 Average PVS and CS <i>n</i> = 4940 (Reference Group)	Profile 2 High PVS and Low CS (But High Impulsivity) <i>n</i> = 1470	Profile 3 Low PVS and High CS (But Low Impulsivity) <i>n</i> = 4004	Tests for Group Differences [†]	
					Profile 1 vs. 2 <i>p</i> Value	Profile 1 vs. 3 <i>p</i> Value
Positive Valence Systems (PVS)						
Reward Sensitivity						
BIS Sum	9.5 (3.7)	9.4 (3.7)	9.8 (4.0)	9.6 (3.6)	0.007 **	0.278
BAS Reward Responsiveness	11.0 (2.9)	11.1 (2.9)	11.1 (3.2)	10.8 (2.8)	0.407	<0.001 ***
Cognitive Systems (CS; Neurocognition)						
Inhibitory Control/ Attention-Flanker	94.2 (8.9)	94.2 (8.4)	87.9 (11.3)	96.5 (7.4)	≤0.001 ***	≤0.001 ***
Processing Speed/ Information Processing— Pattern Comparison	88.2 (14.5)	87.2 (14.1)	80.6 (14.2)	92.1 (13.8)	≤0.001 ***	≤0.001 ***
Episodic Memory—Picture Sequence	103.1 (12.1)	100.2 (10.3)	93.2 (9.3)	110.2 (10.9)	≤0.001 ***	≤0.001 ***
Verbal Learning (RAVLT)						
Short Delay	58.8 (13.3)	55.3 (6.9)	37.9 (9.0)	70.9 (6.8)	≤0.001 ***	≤0.001 ***
Long Delay	9.2 (3.2)	8.4 (1.8)	4.3 (2.1)	12.1 (1.6)	≤0.001 ***	≤0.001 ***
Impulsivity						
Negative Urgency	8.5 (2.6)	8.5 (2.7)	9.1 (2.9)	8.3 (2.5)	≤0.001 ***	≤0.001 ***
Positive Urgency	8.0 (2.9)	8.0 (3.0)	8.9 (3.2)	7.5 (2.7)	≤0.001 ***	≤0.001 ***
Lack of Planning	7.7 (2.4)	7.7 (2.4)	8.0 (2.7)	7.7 (2.2)	≤0.001 ***	0.546
Sensation Seeking	9.8 (2.7)	9.8 (2.7)	9.6 (2.9)	9.8 (2.6)	0.014 *	0.919
Lack of Perseverance	7.0 (2.2)	7.1 (2.2)	7.6 (2.7)	6.8 (2.0)	≤0.001 ***	≤0.001 ***

Note: *p* value “*” ≤ 0.05, “***” ≤ 0.01, “****” ≤ 0.001. [†] Derived from t-test for continuous variables, and chi-square tests for categorical variables with adjusted *p*-values from multiple comparisons.

2.3. Statistical Methods

Missing data, demographics, and covariates: ABCD data harmonization, multiple imputation, and data analyses were conducted in R (Version 1.2.5033, Auckland, New Zealand). Descriptive statistics were used to summarize baseline characteristics of the total sample ($n = 10,414$) and by ALC/STBs group (ALC, STBs, and ALC + STBs) prior to multiple imputation. Between-group differences among the different latent profiles (Profile 1: average PVS and CS vs. Profile 2: high PVS and low CS (high impulsivity), Profile 1: average PVS and CS vs. Profile 3: low PVS and high CS) were assessed using t-tests for PVS, CS, and age (treated as continuous variables), and chi-square tests for categorical variables (sex, parental education, and ethno-racial identity), with adjusted p -values due presented due to Bonferroni corrections from multiple comparisons; Table 1). Demographic covariates were included in the longitudinal analyses and included age, sex (biological), ethno-racial identity, and parent education [64–72].

Multiple imputation with chained equations using the mice package in R were used to obtain more complete data sets and to better protect against bias due to data missing at random mechanisms [72]. This process produced a total of 25 new data sets with the observed and imputed scores [73].

Cross-Sectional Baseline Latent Profile Analysis. To test for the existence of discrete profiles based on PVS, CS, and ALC/STBs patterns, we conducted an LPA at baseline enrollment (Table 2) using an imputed data set that was the most similar to the means and standard deviations in the total sample [74]. LPA is a model-based data reduction method for identifying latent profiles (i.e., subgroups) within a population based on response patterns among a set of variables (PVS, CS, and ALC/STBs) [75]. We employed LPA using all continuous positive valence variables (i.e., reward sensitivity), all continuous neurocognitive variables (i.e., language/verbal intellect, inhibitory control and attention, processing speed/information processing, episodic memory, reading ability/language/academic achievement, verbal learning, and impulsivity), and lifetime ALC/STBs variables (dichotomous).

Table 2. Fit statistics for latent profile model specification at baseline study enrollment ($N = 10,414$).

Model Specification	AIC	BIC	Entropy *	Number of Individuals per Profile						p Value †
				1	2	3	4	5	6	
ABCD Participants who Completed Study Enrollment at Baseline and Year 2 Follow-Up ($N = 10,414$)										
Two-Profile	434,252	434,586	0.75	3972	6442					≤0.001
Three-Profile	431,091	431,541	0.77	4940	1470	4004				0.036
Four-Profile	428,304	429,823	0.739	1492	1896	3424	3602			0.085
Five-Profile	426,956	427,638	0.75	3557	763	1228	1904	2962		0.111

Note: AIC, * Entropy is an index of how well the latent profiles are separated: it ranges from zero to one with higher values to be a sign of a useful model. † Derived from the Vuong Lo Mendell Rubin adjusted test that assesses whether the number of profiles provides improved model fit compared to the model using one fewer profiles.

In LPA, fit indices are compared across models to determine the optimal number of profiles, firstly evaluating a two-profile model fit and incrementally adding latent profiles up to a five-profile model until the best profile solution is found. The final model selection uses several fit indices, including *information criteria*, *likelihood ratios*, and *entropy*. For the information criteria, we used the Akaike Information Criterion (AIC) [75], and the Bayesian Information Criterion (BIC) [76], in which lower values in information criterion statistics indicate a better fit. We relied on the Lo-Mendell-Rubin adjusted likelihood ratio test (LRT) [77] of statistically significant difference ($p < 0.05$) to suggest the number of profiles that are the best fit. Lastly, we relied on a standardized measure of entropy which is an index of model-based classification accuracy. Entropy indicates how accurately the model defines the participant's classification to a latent profile. Relying on the standardized entropy values range from 0 to 1, higher values indicate more precise assignment of participants to latent profiles, and better separation of identified profiles (e.g., entropy value close to 1 is ideal) [78,79].

Longitudinal Multinomial Logistic Regression Analysis. We used multinomial logistic regression models to test PVS, CS, and ALC/STBs baseline profiles as predictors of ALC, STBs, and ALC + STBs at year two follow-up. Specifically, our multinomial logistic regression model tested a nominal variable for profile class as the predictor of interest, with the reference group being the average PVS and CS group. Next, this nominal variable for each LPA profile was examined as a potential predictor of four of the outcome groups (NO ALC/STBs, ALC, STBs, and ALC + STBs; nominal outcomes) at year two follow-up. The longitudinal multinomial model controlled for demographic covariates of sex, ethno-racial identity, and study site.

3. Results

3.1. Descriptive Analysis

A total of 10,414 participants (average age 9.9 years, 52.2% male, 52% non-Hispanic White) who completed baseline enrollment and year two follow-up were included in the analyses (Table 1).

3.2. Cross-Sectional, Latent Profile Analysis

Our data best supported a three-profile model based on recommended decision criteria for evaluating latent patterns (Table 2) [74]. Specifically, the model fit indices for the three-profile model indicated lower AIC and BIC values than the four-profile solution, a significant Vuong Lo Mendell Rubin adjusted p -value ($p = 0.036$), and an entropy value of 0.77). Each of the three profiles were adequately populated (profile 1, $n = 4940$; profile 2, $n = 1470$; profile 3, $n = 4004$; Table 1), which was not true for the other profile solutions. Furthermore, the three-profile model yielded conceptually meaningful configurations of participant profiles: (1) average PVS and CS, (2) high PVS with low CS (high impulsivity), and low PVS with high CS. The three profiles were statistically different ($p = 0.036$) in PVS, CS, ALC, and STBs, but not co-occurring ALC + STBs (Table 1, Figure 1).




Figure 1. Positive valence and cognitive system domains (X Axis) and mean score (Y Axis) by profile at baseline enrollment in the Adolescent Brain and Cognitive Development study (N = 10,414).

3.3. Prospective, Multinomial Logistic Regression Analyses

A multinomial logistic regression analysis identified whether baseline PVS, CS, ALC, STBs, profiles significantly predicted ALC, STBs, and ALC + STBs two years after initial assessment. All three of the PVS, CS, and ALC STB profiles from the cross-sectional LPA were included as a nominal predictor (e.g., Profile 1: average PVS and CS = REFERENCE GROUP, compared to Profile 2: high PVS with low CS (but high impulsivity), and Profile 3: low PVS with high CS (but low impulsivity) in the multinomial logistic regression predicting classification to NO ALC/STBs (reference group), ALC, STBs, and ALC + STBs (Table 3). Two of the baseline PVS, CS, ALC, and STB profiles predicted ALC and STBs, assessed two years later. The low PVS with high CS (but low impulsivity) predicted ALC (OR = 1.41, 95% CI 1.17–1.71, $p \leq 0.001$), while high PVS with low CS (high impulsivity) predicted STBs (OR = 1.25, $p = 1.01$ –1.56). The multinomial logistic regression controlled for baseline demographics of age ($p \leq 0.001$ ALC, $p = 0.024$ STB, $p = 0.322$ ALC + STBs), sex ($p = 0.001$ ALC, $p \leq 0.001$ STB, $p = 0.002$ ALC + STBs), ethno-racial identity ($p \leq 0.001$ ALC, $p = 0.05$ STBs, $p = 0.07$ ALC + STBs), and parental education ($p \leq 0.001$ ALC, STBs, and ALC + STBs).

Table 3. Longitudinal multinomial logistic regression standardized models for baseline profiles (predictors) showing associations with year two follow-up classification of lifetime alcohol use (ALC), lifetime suicidal thoughts and behaviors (STBs), and lifetime co-occurring ALC + STBs (outcomes).

Profile (Predictor) (REF = Profile 1: Average PVS and CS)	Odds Ratio	95% Confidence Interval	<i>p</i> Value
Outcome: Lifetime Alcohol Use (ALC)			
Profile 2: High PVS with Low CS (but high impulsivity)	0.67	0.47–0.96	0.028 *
Profile 3: Low PVS with High CS (but low impulsivity)	1.41	1.17–1.71	≤ 0.001 ***
Outcome: Lifetime Suicidal Thoughts and Behaviors (STBs)			
Profile 2: High PVS with Low CS (but high impulsivity)	1.25	1.01–1.56	0.038 *
Profile 3: Low PVS with High CS (but low impulsivity)	0.95	0.80–1.13	0.593
Outcome: Lifetime Co-occurring Alcohol Use and Suicidal Thoughts and Behaviors (ALC + STBs)			
Profile 2: High PVS with Low CS (but high impulsivity)	0.55	0.22–1.42	0.217
Profile 3: Low PVS with High CS (but low impulsivity)	1.38	0.88–2.17	0.158

Note: *p*-value "*" ≤ 0.05 , "***" ≤ 0.001 . Covariates included in longitudinal multinomial logistic regression standardized models included age ($p \leq 0.001$ ALC, $p = 0.024$ STB, $p = 0.322$ ALC + STBs), sex (biological; $p = 0.001$ ALC, $p \leq 0.001$ STB, $p = 0.002$ ALC + STBs), ethno-racial identity ($p \leq 0.001$ ALC, $p = 0.05$ STBs, $p = 0.07$ ALC + STBs), and parental education ($p \leq 0.001$ ALC, STBs, and ALC + STBs).

4. Discussion

This study demonstrated that the combination of the RDoC framework and data-driven advanced statistical modeling provides a novel and promising approach to research on the ALC-STBs association in early adolescence. We found two conceptually meaningful participant profiles of PVS and CS among youth ages 9–10 at the baseline time point. We found evidence of clinical meaning as the unique profile of low PVS with high CS (but low impulsivity) was prospectively associated with ALC at two-year follow-up; while the profile of high PVS with low CS (but high impulsivity) prospectively predicted STBs independently at two-year follow-up, implying that early adolescents who endorse STBs have high PVS and low CS (but high impulsivity) at ages 9–10 and are at increased risk

for suicide at ages 11–12. This finding has not been well researched in the early adolescent literature, and while the simultaneous modeling of the ALC-STB association using PVS and CS constructs is sparse, research on adult populations showing the high PVS-STB link has been replicated [4] while the low CS-STB findings have been less consistent [4,6].

To our knowledge, this is the first study to examine unique profiles of PVS, CS, ALC, and STB patterns simultaneously using data-driven, advanced statistical modeling in children as young as 9–10-years. Notably, the most common ($n = 4940$, ~47%) PV/CS/ALC/STB profile among children in the ABCD baseline cohort can be described as the average PVS and CS group, followed by the low PVS and high CS (but low impulsivity; $n = 4004$, ~38%), and the least common profile was the high PVS with low CS (but high impulsivity; $n = 1470$, ~14%). Few studies have explored PVS and CS profiles and their ability to predict alcohol use, suicidal thoughts and behaviors, and their co-occurrence at a later time point, thus it is difficult to make comparisons with other research. However, the characterization of unique PVS and CS profiles might facilitate a deeper understanding of the ALC-STB association and inform future research and targets for intervention.

These findings of neurocognitive and behavioral profiles have notable clinical relevance in both ALC and STB prevention and intervention. First, these findings may inform future research on the screening and detection of ALC, STBs, their co-occurrence, and risk factors, in youth as young as 9 years of age. A tiered approach to screening with validated (population- and setting-specific) instruments that identify increased ALC and STBs risk may benefit from considering combined neurocognitive (language/vocabulary comprehension, cognitive control, episodic memory, and impulsivity), and positive valence (reward sensitivity) assessment [80,81]. Second, patterns in positive valence and cognitive system deficits may influence treatment outcomes, thereby making patients less amenable or responsive to psychotherapeutic approaches that have demonstrated effectiveness at treating ALC, STBs, or their co-occurrence. Thus, thoughtful consideration in tailoring interventions to address the positive valence and neurocognitive heterogeneity is important. It may be that patients who present with deficits in these domains, could benefit from adjunctive treatments directed at ameliorating deficits in positive valence and cognitive systems, such as cognitive training. Third, given the co-occurrence of ALC and STBs in youth as young as 9 years, personalized treatments to meet this high-risk group are needed. To date, there is currently no standard approach to treating co-occurring ALC and STBs, but motivational enhancement therapy or cognitive behavioral therapy approaches have demonstrated efficacy [82,83]. Unfortunately, ALC and STBs are often treated in two separate systems of care, and usually with older adolescents, highlighting an important and unmet need [82,83]. However, one intervention to highlight is a two-site randomized control trial that recruited 95 participants aged 13–21, with depression and substance use, led by John Curry, PhD, MD, and Yifrah Kaminer, MD, MBA at the University of Connecticut (J. Curry, personal communication, 14 June 2022) [83]. The trial includes an adaptive method, such that initial substance use treatment (CBT) is supplemented with one of two depression treatments (CBT-for-depression [CBT-D], or treatment-as-usual [TAU] only when needed. Study findings included that 35 of the 95 adolescents had positive depression response to CBT for substance use within one month; and that for the remaining adolescents, there were no differences in depression or substance use outcomes between CBT-D and TAU at the end of treatment. All groups showed reduced depression and reduced substance use. Although this trial excluded youth who were at high risk for suicide (past-month SA or current SI with intent or plan), the investigators did include measures that assess STBs, that they plan to examine in subsequent work. This research is both timely and novel given the increase in STBs, and early adolescent youth as young as age 9 incur additional risk when ALC co-occurs. Interventions that can treat the co-occurrence of ALC and STBs in youth as young as age 9 is an important and understudied area of research.

While this study's findings are critical additions to the literature, limitations exist. As these analyses include two time points (baseline study enrollment, year two follow-up), directionality and causal inferences cannot be conclusively derived. Patterns of

PVS and CS may be due to other mechanisms (e.g., family history of STBs, psychiatric diagnosis). In follow-up prospective ABCD analyses, we will extrapolate the directionality of the relationship between decrements in PVS, CS, ALC, and STBs longitudinally while examining potential confounds within-group differences (children and adolescents for whom cognitive PVS and CS may be linked to STBs changes) using a structural equation modeling framework (e.g., mediation and moderation analyses) with more time points. These analyses present small effect sizes: however, these findings are clinically meaningful, as early identification of behaviors that increase suicide risk even slightly can prevent devastating outcomes and provide clues to better understanding STBs. One limitation of all self-report data is self-reporting biases arising from social desirability, recall period, or selective recall. It is possible that participants under-reported STBs experiences due to fears of parents and authority figures being informed of their behavior. Although, ABCD is designed to ensure youths feel comfortable providing honest answers; research assistants underwent extensive training, including motivational interviewing techniques, to alleviate concerns about disclosing responses to parents. Youths and parents were informed that only specific responses (e.g., posing an immediate and real danger to themselves or others) would be shared with parents.

5. Conclusions

Few studies have examined the interplay between PVS and CS in relation to ALC and STBs through an RDoC framework. Our manuscript is the first to examine PVS, CS, and STB profiles in early adolescents (ages 9–12 years) and suggests that unique profiles related to these constructs prospectively predict ALC, STBs, and their co-occurrence two years after initial assessment. A better understanding of the relationship between PVS, CS, and STBs will help to identify neurobehavioral markers of increased risk, an important first step in moving beyond clinicians' reliance on self-report and subjective impressions to predict suicide. Further, this research can inform early intervention, precision medicine strategies, and suggest avenues for neuroscience-informed STB interventions.

Author Contributions: Conceptualization, L.D.A. and J.J.; methodology, L.D.A. and J.J.; formal analysis, L.D.A. and J.J.; investigation, L.D.A., C.C., and J.J.; data curation, D.A.G. and L.D.A.; writing—original draft preparation, L.D.A., C.C. and J.J.; writing—review and editing, L.D.A., C.C., K.E.C. and J.J.; visualization, L.D.A. and D.A.G.; supervision, J.J. and K.E.C. All authors have read and agreed to the published version of the manuscript.

Funding: Work on this paper was supported by the National Institute on Drug Abuse grants U01 DA041089 (L.D.A., D.G. and J.J.), U01 DA041147 (C.C.), R21 DA047953 (J.J.), R01 DA054106 (J.J.); and the California Tobacco-Related Disease Research Grants Program Office of the University of California grant 580264 (J.J.). Data used in the preparation of this article were obtained from the Adolescent Brain Cognitive DevelopmentSM Study (ABCD Study[®]) (<https://abcdstudy.org>, access date 7 January 2022), held in the NIMH Data Archive (NDA). This is a multisite, longitudinal study designed to recruit more than 10,000 children aged 9–10 and follow them over 10 years into early adulthood. The ABCD Study is supported by the National Institutes of Health and additional federal partners under award numbers: U01DA041048, U01DA050989, U01DA051016, U01DA041022, U01DA051018, U01DA051037, U01DA050987, U01DA041174, U01DA041106, U01DA041117, U01DA041028, U01DA041134, U01DA050988, U01DA051039, U01DA041156, U01DA041025, U01DA041120, U01DA051038, U01DA041148, U01DA041093, U01DA041089, U24DA041147, and U24DA041123. A full list of supporters is available at <https://abcdstudy.org/federal-partners/>, accessed on 20 May 2022. A listing of participating sites and a complete listing of the study investigators can be found at <https://abcdstudy.org/principal-investigators.html>, accessed on 20 May 2022. ABCD Study consortium investigators designed and implemented the study and/or provided data but did not necessarily participate in analysis or writing of this report. This manuscript reflects the views of the authors and may not reflect the opinions or views of the NIH or ABCD Study consortium investigators.

Institutional Review Board Statement: UCSD is the IRB site (160091).

Informed Consent Statement: Written informed consent was obtained from the participants to publish this paper.

Data Availability Statement: The ABCD data repository grows and changes over time. The ABCD data used in this report came from Annual Release 4.0, doi:10.15154/1519007. DOIs can be found at <https://ndar.nih.gov/study.html?id=721> (accessed on 20 May 2022).

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Aguinaldo, L.D.; Goldstone, A.; Hasler, B.P.; Brent, D.A.; Coronado, C.; Jacobus, J. Preliminary analysis of low-level alcohol use and suicidality with children in the adolescent brain and cognitive development (ABCD) baseline cohort. *Psychiatry Res.* **2021**, *299*, 113825. [[CrossRef](#)] [[PubMed](#)]
2. Esang, M.; Ahmed, S. A Closer Look at Substance Use and Suicide. *Am. J. Psychiatry Resid. J.* **2018**, *13*, 6–8. [[CrossRef](#)]
3. Franklin, J.C.; Ribeiro, J.D.; Fox, K.R.; Bentley, K.H.; Kleiman, E.M.; Huang, X.; Musacchio, K.M.; Jaroszewski, A.C.; Chang, B.P.; Nock, M.K. Risk factors for suicidal thoughts and behaviors: A meta-analysis of 50 years of research. *Psychol. Bull.* **2017**, *143*, 187–232. [[CrossRef](#)]
4. Glenn, C.R.; Nock, M.K. Improving the Short-Term Prediction of Suicidal Behavior. *Am. J. Prev. Med.* **2014**, *47*, S176–S180. [[CrossRef](#)] [[PubMed](#)]
5. O'Connor, R.C.; Nock, M.K. The psychology of suicidal behaviour. *Lancet Psychiatry* **2014**, *1*, 73–85. [[CrossRef](#)]
6. Stewart, J.G.; Polanco-Roman, L.; Duarte, C.S.; Auerbach, R.P. Neurocognitive Processes Implicated in Adolescent Suicidal Thoughts and Behaviors: Applying an RDoC Framework for Conceptualizing Risk. *Curr. Behav. Neurosci. Rep.* **2019**, *6*, 188–196. [[CrossRef](#)] [[PubMed](#)]
7. Zelazny, J.; Melhem, N.; Porta, G.; Biernesser, C.; Keilp, J.G.; Mann, J.J.; Oquendo, M.A.; Stanley, B.; Brent, D.A. Childhood maltreatment, neuropsychological function and suicidal behavior. *J. Child Psychol. Psychiatry* **2019**, *60*, 1085–1093. [[CrossRef](#)]
8. Glenn, C.R.; Kleiman, E.M.; Cha, C.B.; Am, C.A.D.; Franklin, J.C.; Nock, M.K. Understanding suicide risk within the Research Domain Criteria (RDoC) framework: A meta-analytic review. *Depress. Anxiety* **2018**, *35*, 65–88. [[CrossRef](#)]
9. Nock, M.K.; Millner, A.J.; Joiner, T.E.; Gutierrez, P.M.; Han, G.; Hwang, I.; King, A.; Naifeh, J.A.; Sampson, N.A.; Zaslavsky, A.M.; et al. Risk factors for the transition from suicide ideation to suicide attempt: Results from the Army Study to Assess Risk and Resilience in Servicemembers (Army STARRS). *J. Abnorm. Psychol.* **2018**, *127*, 139–149. [[CrossRef](#)]
10. Podlogar, M.C.; Rogers, M.L.; Stanley, I.H.; Hom, M.A.; Chiurliza, B.; Joiner, T.E. Anxiety, depression, and the suicidal spectrum: A latent class analysis of overlapping and distinctive features. *Cogn. Emot.* **2018**, *32*, 1464–1477. [[CrossRef](#)]
11. Insel, T.; Cuthbert, B.; Garvey, M.; Heinssen, R.; Pine, D.S.; Quinn, K.; Sanislow, C.; Wang, P. Research Domain Criteria (RDoC): Toward a New Classification Framework for Research on Mental Disorders. *Am. J. Psychiatry* **2010**, *167*, 748–751. [[CrossRef](#)] [[PubMed](#)]
12. Dougherty, D.M.; Mathias, C.W.; Marsh-Richard, D.M.; Prevette, K.N.; Dawes, M.A.; Hatzis, E.S.; Palmes, G.; Nouvion, S.O. Impulsivity and clinical symptoms among adolescents with non-suicidal self-injury with or without attempted suicide. *Psychiatry Res.* **2009**, *169*, 22–27. [[CrossRef](#)] [[PubMed](#)]
13. Mathias, C.W.; Dougherty, N.M.; James, L.M.; Richard, D.M.; Dawes, M.A.; Acheson, A.; Hill-Kapturczak, N. Intolerance to Delayed Reward in Girls with Multiple Suicide Attempts. *Suicide Life-Threat. Behav.* **2011**, *41*, 277–286. [[CrossRef](#)]
14. Pan, L.A.; Batezati-Alves, S.C.; Almeida, J.; Segreti, A.; Akkal, D.; Hassel, S.; Lakdawala, S.; Brent, D.A.; Phillips, M.L. Dissociable Patterns of Neural Activity During Response Inhibition in Depressed Adolescents with and Without Suicidal Behavior. *J. Am. Acad. Child Adolesc. Psychiatry* **2011**, *50*, 602–611. [[CrossRef](#)] [[PubMed](#)]
15. Auerbach, R.P.; Millner, A.J.; Stewart, J.G.; Esposito, E.C. Identifying differences between depressed adolescent suicide ideators and attempters. *J. Affect. Disord.* **2015**, *186*, 127–133. [[CrossRef](#)] [[PubMed](#)]
16. Tsypes, A.; Owens, M.; Gibb, B.E. Blunted Neural Reward Responsiveness in Children with Recent Suicidal Ideation. *Clin. Psychol. Sci.* **2019**, *7*, 958–968. [[CrossRef](#)] [[PubMed](#)]
17. Tsypes, A.; Owens, M.; Hajcak, G.; Gibb, B.E. Neural responses to gains and losses in children of suicide attempters. *J. Abnorm. Psychol.* **2017**, *126*, 237–243. [[CrossRef](#)]
18. Sauder, C.L.; Derbidge, C.M.; Beauchaine, T.P. Neural responses to monetary incentives among self-injuring adolescent girls. *Dev. Psychopathol.* **2016**, *28*, 277–291. [[CrossRef](#)]
19. Tolomeo, S.; Macfarlane, J.A.; Baldacchino, A.; Koob, G.F.; Steele, J.D. Alcohol Binge Drinking: Negative and Positive Valence System Abnormalities. *Biol. Psychiatry Cogn. Neurosci. Neuroimaging* **2021**, *6*, 126–134. [[CrossRef](#)]
20. Koob, G.F.; Volkow, N.D. Neurocircuitry of Addiction. *Neuropsychopharmacology* **2010**, *35*, 217–238. [[CrossRef](#)]
21. Koob, G.F.; Volkow, N.D. Neurobiology of addiction: A neurocircuitry analysis. *Lancet Psychiatry* **2016**, *3*, 760–773. [[CrossRef](#)]
22. Bridge, J.A.; McBee-Strayer, S.M.; Cannon, E.A.; Sheftall, A.H.; Reynolds, B.; Campo, J.V.; Pajer, K.A.; Barbe, R.P.; Brent, D.A. Impaired Decision Making in Adolescent Suicide Attempters. *J. Am. Acad. Child Adolesc. Psychiatry* **2012**, *51*, 394–403. [[CrossRef](#)] [[PubMed](#)]
23. Ackerman, J.P.; McBee-Strayer, S.M.; Mendoza, K.; Stevens, J.; Sheftall, A.H.; Campo, J.V.; A Bridge, J. Risk-Sensitive Decision-Making Deficit in Adolescent Suicide Attempters. *J. Child Adolesc. Psychopharmacol.* **2015**, *25*, 109–113. [[CrossRef](#)] [[PubMed](#)]
24. Hamza, C.A.; Willoughby, T. Nonsuicidal Self-Injury and Suicidal Behavior: A Latent Class Analysis among Young Adults. *PLoS ONE* **2013**, *8*, e59955. [[CrossRef](#)] [[PubMed](#)]

25. Klonsky, E.D.; Olino, T.M. Identifying clinically distinct subgroups of self-injurers among young adults: A latent class analysis. *J. Consult. Clin. Psychol.* **2008**, *76*, 22–27. [[CrossRef](#)]
26. Dhingra, K.; Boduszek, D.; Klonsky, E.D. Empirically Derived Subgroups of Self-Injurious Thoughts and Behavior: Application of Latent Class Analysis. *Suicide Life-Threat. Behav.* **2016**, *46*, 486–499. [[CrossRef](#)]
27. Peterson, A.L.; Chen, J.I.; Karver, M.S.; Labouliere, C.D. Frustration with feeling: Latent classes of non-suicidal self-injury and emotion regulation difficulties. *Psychiatry Res.* **2019**, *275*, 61–70. [[CrossRef](#)]
28. Whitlock, J.; Muehlenkamp, J.; Eckenrode, J. Variation in Nonsuicidal Self-Injury: Identification and Features of Latent Classes in a College Population of Emerging Adults. *J. Clin. Child Adolesc. Psychol.* **2008**, *37*, 725–735. [[CrossRef](#)]
29. Stoep, A.V.; McCauley, E.; Flynn, C.; Stone, A. Thoughts of Death and Suicide in Early Adolescence. *Suicide Life-Threat. Behav.* **2009**, *39*, 599–613. [[CrossRef](#)]
30. Xiao, Y.; Romanelli, M.; Lindsey, M.A. A latent class analysis of health lifestyles and suicidal behaviors among US adolescents. *J. Affect. Disord.* **2019**, *255*, 116–126. [[CrossRef](#)]
31. Thompson, M.; Kuruwita, C.; Foster, E.M. Transitions in Suicide Risk in a Nationally Representative Sample of Adolescents. *J. Adolesc. Health* **2009**, *44*, 458–463. [[CrossRef](#)] [[PubMed](#)]
32. Mann, J.J.; Waternaux, C.; Haas, G.L.; Malone, K.M. Toward a clinical model of suicidal behavior in psychiatric patients. *Am. J. Psychiatry* **1999**, *156*, 181–189. [[CrossRef](#)] [[PubMed](#)]
33. Heeringa, S.G.; Berglund, P.A. A Guide for Population-based Analysis of the Adolescent Brain Cognitive Development (ABCD) Study Baseline Data. *bioRxiv* **2020**. [[CrossRef](#)]
34. Garavan, H.; Bartsch, H.; Conway, K.; Decastro, A.; Goldstein, R.Z.; Heeringa, S.; Jernigan, T.; Potter, A.; Thompson, W.; Zahs, D. Recruiting the ABCD sample: Design considerations and procedures. *Dev. Cogn. Neurosci.* **2018**, *32*, 16–22. [[CrossRef](#)]
35. Barch, D.M.; Albaugh, M.D.; Avenevoli, S.; Chang, L.; Clark, D.B.; Glantz, M.D.; Hudziak, J.J.; Jernigan, T.L.; Tapert, S.F.; Yurgelun-Todd, D.; et al. Demographic, physical and mental health assessments in the adolescent brain and cognitive development study: Rationale and description. *Dev. Cogn. Neurosci.* **2018**, *32*, 55–66. [[CrossRef](#)]
36. Volkow, N.D.; Koob, G.F.; Croyle, R.T.; Bianchi, D.W.; Gordon, J.A.; Koroshetz, W.J.; Pérez-Stable, E.J.; Riley, W.T.; Bloch, M.H.; Conway, K.; et al. The conception of the ABCD study: From substance use to a broad NIH collaboration. *Dev. Cogn. Neurosci.* **2018**, *32*, 4–7. [[CrossRef](#)]
37. Luciana, M.; Bjork, J.M.; Nagel, B.J.; Barch, D.M.; Gonzalez, R.; Nixon, S.J.; Banich, M.T. Adolescent neurocognitive development and impacts of substance use: Overview of the adolescent brain cognitive development (ABCD) baseline neurocognition battery. *Dev. Cogn. Neurosci.* **2018**, *32*, 67–79. [[CrossRef](#)]
38. Lisdahl, K.M.; Sher, K.J.; Conway, K.P.; Gonzalez, R.; Ewing, S.W.F.; Nixon, S.J.; Tapert, S.; Bartsch, H.; Goldstein, R.Z.; Heitzeg, M. Adolescent brain cognitive development (ABCD) study: Overview of substance use assessment methods. *Dev. Cogn. Neurosci.* **2018**, *32*, 80–96. [[CrossRef](#)]
39. Uban, K.A.; Horton, M.K.; Jacobus, J.; Heyser, C.; Thompson, W.K.; Tapert, S.F.; Madden, P.A.; Sowell, E.R. Biospecimens and the ABCD study: Rationale, methods of collection, measurement and early data. *Dev. Cogn. Neurosci.* **2018**, *32*, 97–106. [[CrossRef](#)]
40. Watts, A.L.; Smith, G.T.; Barch, D.M.; Sher, K.J. Factor structure, measurement and structural invariance, and external validity of an abbreviated youth version of the UPPS-P Impulsive Behavior Scale. *Psychol. Assess.* **2020**, *32*, 336–347. [[CrossRef](#)]
41. O'Connor, R.C.; Forgan, G. Suicidal Thinking and Perfectionism: The Role of Goal Adjustment and Behavioral Inhibition/Activation Systems (BIS/BAS). *J. Ration. Cogn. Ther.* **2007**, *25*, 321–341. [[CrossRef](#)]
42. Carver, C.S.; White, T.L. Behavioral inhibition, behavioral activation, and affective responses to impending reward and punishment: The BIS/BAS Scales. *J. Pers. Soc. Psychol.* **1994**, *67*, 319–333. [[CrossRef](#)]
43. Gershon, R.C.; Cook, K.F.; Mungas, D.; Manly, J.J.; Slotkin, J.; Beaumont, J.L.; Weintraub, S. Language Measures of the NIH Toolbox Cognition Battery. *J. Int. Neuropsychol. Soc.* **2014**, *20*, 642–651. [[CrossRef](#)] [[PubMed](#)]
44. Eriksen, B.A.; Eriksen, C.W. Effects of noise letters upon the identification of a target letter in a nonsearch task. *Percept. Psychophys.* **1974**, *16*, 143–149. [[CrossRef](#)]
45. Weintraub, S.; Dikmen, S.S.; Heaton, R.K.; Tulsky, D.S.; Zelazo, P.D.; Slotkin, J.; Carlozzi, N.E.; Bauer, P.J.; Wallner-Allen, K.; Fox, N.; et al. The Cognition Battery of the NIH Toolbox for Assessment of Neurological and Behavioral Function: Validation in an Adult Sample. *J. Int. Neuropsychol. Soc.* **2014**, *20*, 567–578. [[CrossRef](#)]
46. Carlozzi, N.E.; Tulsky, D.S.; Kail, R.V.; Beaumont, J. VI.cNIH TOOLBOX COGNITION BATTERY (CB): MEASURING PROCESSING SPEED. *Monogr. Soc. Res. Child Dev.* **2013**, *78*, 88–102. [[CrossRef](#)]
47. Taylor, E.M. *Psychological Appraisal of Children with Cerebral Defects*; Harvard University Press: Cambridge, MA, USA, 1959.
48. Strauss, E.; Sherman, E.M.; Spreen, O. *A Compendium of Neuropsychological Tests: Administration, Norms, and Commentary*; American Chemical Society: Washington, WA, USA, 2006.
49. Bauer, P.J.; Dikmen, S.S.; Heaton, R.K.; Mungas, D.; Slotkin, J.; Beaumont, J. III. NIH TOOLBOX COGNITION BATTERY (CB): MEASURING EPISODIC MEMORY. *Monogr. Soc. Res. Child Dev.* **2013**, *78*, 34–48. [[CrossRef](#)]
50. Günther, T.; Holtkamp, K.; Jolles, J.; Herpertz-Dahlmann, B.; Konrad, K. Verbal memory and aspects of attentional control in children and adolescents with anxiety disorders or depressive disorders. *J. Affect. Disord.* **2004**, *82*, 265–269. [[CrossRef](#)]
51. Daniel, M.H.; Wahlstrom, D.; Zhang, O. Equivalence of Q-interactive™ and Paper Administrations of Cognitive Tasks: WISC®–V. *Q-Interact. Tech. Rep.* **2014**, *8*.

52. Jackson, K.M.; Barnett, N.P.; Colby, S.M.; Rogers, M.L. The Prospective Association between Sipping Alcohol by the Sixth Grade and Later Substance Use. *J. Stud. Alcohol Drugs* **2015**, *76*, 212–221. [[CrossRef](#)]
53. Beautrais, A.L.; Joyce, P.R.; Mulder, R.T. Risk Factors for Serious Suicide Attempts among Youths Aged 13 through 24 Years. *J. Am. Acad. Child Adolesc. Psychiatry* **1996**, *35*, 1174–1182. [[CrossRef](#)] [[PubMed](#)]
54. Ammerman, B.A.; Kleiman, E.M.; Jenkins, A.L.; Berman, M.; McCloskey, M. Using Propensity Scores to Examine the Association Between Behavioral Inhibition/Activation and Nonsuicidal and Suicidal Self-Injury. *Crisis* **2017**, *38*, 227–236. [[CrossRef](#)] [[PubMed](#)]
55. Khosravani, V.; Baseri, A.; Kamali, Z.; Mohammadzadeh, A.; Amirinezhad, A. Direct and Indirect Effects of Behavioral Inhibition/Activation Systems on Depression and Current Suicidal Ideation Through Rumination and Self-Reflection. *Arch. Suicide Res.* **2020**, *24*, 568–588. [[CrossRef](#)] [[PubMed](#)]
56. Kaufman, J.; Birmaher, B.; Brent, D.A.; Ryan, N.D.; Rao, U. K-SADS-PL. *J. Am. Acad. Child Adolesc. Psychiatry* **2000**, *39*, 1208. [[CrossRef](#)]
57. Brent, D.A.; Baugher, M.; Bridge, J.; Chen, T.; Chiappetta, L. Age- and Sex-Related Risk Factors for Adolescent Suicide. *J. Am. Acad. Child Adolesc. Psychiatry* **1999**, *38*, 1497–1505. [[CrossRef](#)]
58. Bridge, J.A.; Asti, L.; Horowitz, L.M.; Greenhouse, J.B.; Fontanella, C.A.; Sheftall, A.H.; Kelleher, K.J.; Campo, J.V. Suicide Trends Among Elementary School-Aged Children in the United States From 1993 to 2012. *JAMA Pediatr.* **2015**, *169*, 673–677. [[CrossRef](#)]
59. Cash, S.J.; A Bridge, J. Epidemiology of youth suicide and suicidal behavior. *Curr. Opin. Pediatr.* **2009**, *21*, 613–619. [[CrossRef](#)]
60. Gould, M.S.; Fisher, P.; Parides, M.; Flory, M.; Shaffer, D. Psychosocial Risk Factors of Child and Adolescent Completed Suicide. *Arch. Gen. Psychiatry* **1996**, *53*, 1155–1162. [[CrossRef](#)]
61. Gould, M.S.; Greenberg, T.; Velting, D.M.; Shaffer, D. Youth Suicide Risk and Preventive Interventions: A Review of the Past 10 Years. *J. Am. Acad. Child Adolesc. Psychiatry* **2003**, *42*, 386–405. [[CrossRef](#)]
62. Stover, P.J.; Harlan, W.R.; Hammond, J.A.; Hendershot, T.; Hamilton, C.M. PhenX: A toolkit for interdisciplinary genetics research. *Curr. Opin. Lipidol.* **2010**, *21*, 136–140. [[CrossRef](#)]
63. Miranda-Mendizabal, A.; Castellví, P.; Parés-Badell, O.; Alayo, I.; Almenara, J.; Alonso, I.; Blasco, M.J.; Cebrià, A.; Gabilondo, A.; Gili, M.; et al. Gender differences in suicidal behavior in adolescents and young adults: Systematic review and meta-analysis of longitudinal studies. *Int. J. Public Health* **2019**, *64*, 265–283. [[CrossRef](#)] [[PubMed](#)]
64. Bilsen, J. Suicide and Youth: Risk Factors. *Front. Psychiatry* **2018**, *9*, 540. [[CrossRef](#)] [[PubMed](#)]
65. Baiden, P.; LaBrenz, C.A.; Asiedua-Baiden, G.; Muehlenkamp, J.J. Examining the intersection of race/ethnicity and sexual orientation on suicidal ideation and suicide attempt among adolescents: Findings from the 2017 Youth Risk Behavior Survey. *J. Psychiatr. Res.* **2020**, *125*, 13–20. [[CrossRef](#)] [[PubMed](#)]
66. Arshanapally, S.; Werner, K.B.; Sartor, C.E.; Bucholz, K.K. The Association between Racial Discrimination and Suicidality among African-American Adolescents and Young Adults. *Arch. Suicide Res.* **2017**, *22*, 584–595. [[CrossRef](#)]
67. Holtmann, M.; Buchmann, A.F.; Esser, G.; Schmidt, M.H.; Banaschewski, T.; Laucht, M. The Child Behavior Checklist-Dysregulation Profile predicts substance use, suicidality, and functional impairment: A longitudinal analysis. *J. Child Psychol. Psychiatry* **2011**, *52*, 139–147. [[CrossRef](#)]
68. Achenbach, T.M.; Ruffle, T.M. The Child Behavior Checklist and Related Forms for Assessing Behavioral/Emotional Problems and Competencies. *Pediatr. Rev.* **2000**, *21*, 265–271. [[CrossRef](#)]
69. Van Meter, A.R.; Algorta, G.P.; Youngstrom, E.; Lechtman, Y.; Youngstrom, J.K.; Feeny, N.C.; Findling, R.L. Assessing for suicidal behavior in youth using the Achenbach System of Empirically Based Assessment. *Eur. Child Adolesc. Psychiatry* **2018**, *27*, 159–169. [[CrossRef](#)]
70. Johnson, D.; McLennan, J.D.; Heron, J.; Colman, I. The relationship between profiles and transitions of internalizing and externalizing symptoms in children and suicidal thoughts in early adolescence. *Psychol. Med.* **2020**, *50*, 2566–2574. [[CrossRef](#)]
71. Lecloux, M.; Culp, S.; O'Brien, K.; Wolff, J.; Andrade, L. The relationship between internalizing and externalizing symptomatology and parent/adolescent suicide report concordance: Implications for prevention and clinical practice. *Suicidol. Online* **2017**, *8*, 1–8.
72. Van Buuren, S.; Groothuis-Oudshoorn, K. mice: Multivariate Imputation by Chained Equations in R. *J. Stat. Softw.* **2011**, *45*, 1–67. [[CrossRef](#)]
73. Rubin, D.B. *Multiple Imputation for Nonresponse in Surveys*; John Wiley & Sons: Hoboken, NJ, USA, 2004; Volume 81.
74. Lanza, S.T.; Bray, B.C.; Collins, L.M. An Introduction to Latent Class and Latent Transition Analysis. *Handb. Psychol.* **2012**, *2*, 691–716. [[CrossRef](#)]
75. Akaike, H. Factor analysis and AIC. *Psychometrika* **1987**, *52*, 317–332. [[CrossRef](#)]
76. Schwarz, G. Estimating the Dimension of a Model. *Ann. Stat.* **1978**, *6*, 461–464. [[CrossRef](#)]
77. Lo, Y.; Mendell, N.R.; Rubin, D.B. Testing the number of components in a normal mixture. *Biometrika* **2001**, *88*, 767–778. [[CrossRef](#)]
78. Lubke, G.H.; Muthén, B. Investigating population heterogeneity with factor mixture models. *Psychol. Methods* **2005**, *10*, 21–39. [[CrossRef](#)] [[PubMed](#)]
79. Wang, M.-C.; Deng, Q.; Bi, X.; Ye, H.; Yang, W. Performance of the entropy as an index of classification accuracy in latent profile analysis: A Monte Carlo simulation study. *Acta Psychol. Sin.* **2017**, *49*, 1473–1482. [[CrossRef](#)]
80. Aguinaldo, L.D.; Sullivant, S.; Lanzillo, E.C.; Ross, A.; He, J.-P.; Bradley-Ewing, A.; Bridge, J.A.; Horowitz, L.M.; Wharff, E.A. Validation of the ask suicide-screening questions (ASQ) with youth in outpatient specialty and primary care clinics. *Gen. Hosp. Psychiatry* **2021**, *68*, 52–58. [[CrossRef](#)]
81. Fleming, M.F. Screening and Brief Intervention in Primary Care Settings. *Alcohol Res. Health* **2008**, *28*, 57–62.

82. O'Brien, K.H.M.; Sellers, C.M.; Battalen, A.W.; Ryan, C.A.; Maneta, E.K.; Aguinaldo, L.; White, E.; Spirito, A. Feasibility, acceptability, and preliminary effects of a brief alcohol intervention for suicidal adolescents in inpatient psychiatric treatment. *J. Subst. Abus. Treat.* **2018**, *94*, 105–112. [[CrossRef](#)]
83. Curry, J.F.; Kaminer, Y.; Goldston, D.B.; Chan, G.; Wells, K.C.; Burke, R.H.; Inscoc, A.B.; Meyer, A.E.; Cheek, S.M. Adaptive Treatment for Youth with Substance Use and Depression: Early Depression Response and Short-term Outcomes. *J. Am. Acad. Child Adolesc. Psychiatry* **2021**, *61*, 508–519. [[CrossRef](#)]