

17. Ravi V, Sharma PS, Patel NR, Dommaraju S, Zalavadia DV, Garg V *et al.* New-onset atrial fibrillation in left bundle branch area pacing compared with right ventricular pacing. *Circ Arrhythm Electrophysiol* 2022;**15**:e010710.
18. Barbier P, Solomon SB, Schiller NB, Glantz SA. Left atrial relaxation and left ventricular systolic function determine left atrial reservoir function. *Circulation* 1999;**100**: 427–36.
19. Liu Q, Yang J, Bolun Z, Pei M, Ma B, Tong Q *et al.* Comparison of cardiac function between left bundle branch pacing and right ventricular outflow tract septal pacing in the short-term: a registered controlled clinical trial. *Int J Cardiol* 2021;**322**:70–6.
20. Wang Z, Zhu H, Li X, Yao Y, Liu Z, Fan X. Left bundle branch area pacing versus right ventricular pacing in patients with persistent atrial fibrillation requiring ventricular pacing. *Pacing Clin Electrophysiol* 2021;**44**:2024–30.

EP CASE EXPRESS

<https://doi.org/10.1093/europace/euac166>

Stridor and dyspnoea after ablation for atrial fibrillation

Khalid Bin Waleed ¹, Pavandeep Toor ¹, Zaki Akhtar ¹, Jonathan Aron², Paul Govewalla³, and Mark M. Gallagher ^{1*}

¹Department of Cardiology, St George's University Hospitals NHS Foundation Trust, St George's University of London, Blackshaw Road, SW17 0QT London, UK;


²Department of Critical Care, St George's University Hospitals NHS Foundation Trust, St George's University of London, Blackshaw Road, SW17 0QT London, UK; and

³Department of Cardiac Surgery, St George's University Hospitals NHS Foundation Trust, St George's University of London, Blackshaw Road, SW17 0QT London, UK

*Corresponding author. E-mail address: mark_m_gallagher@hotmail.com

A 74-year-old female underwent redo catheter ablation for symptomatic atrial fibrillation. Transeptal puncture was performed without difficulty. The pulmonary veins had remained isolated, so radiofrequency energy was applied at the left atrial roof and posterior wall, and cavotricuspid isthmus by QDOT® catheter (Biosense Webster) using the QMODE plus protocol without immediate complication. Approximately 10 h post-ablation, the patient reported a foreign-body sensation in the throat and then progressive dyspnoea with stridor. Chest X-ray showed widening of the upper mediastinum (*Panel A*). Computed tomography (CT) at 24 h post-ablation showed a 6 × 3.5 cm haematoma extending 12 cm from the mediastinum into the neck and compressing surrounding structures including displacement of the trachea (*Panel B*). The patient was intubated and ventilated without difficulty. Repeated CT on day 3 showed shrinkage of the haematoma and reduced tracheal compression. The patient was extubated; her subsequent recovery was uneventful despite the resumption of anticoagulation at 1 week.

To our knowledge, this is the first report of superior mediastinal haematoma following ablation. Management by intubation, positive pressure ventilation and cessation of anticoagulation was followed by slow resolution. The favourable outcome of this conservative approach may be a guide to management of similar cases.



The full-length version of this report can be viewed at: <https://www.escardio.org/Education/E-Learning/Clinical-cases/Electrophysiology>.

© The Author(s) 2023. Published by Oxford University Press on behalf of the European Society of Cardiology.

This is an Open Access article distributed under the terms of the Creative Commons Attribution-NonCommercial License (<https://creativecommons.org/licenses/by-nc/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited. For commercial re-use, please contact journals.permissions@oup.com