

Review



Cite this article: Millins C, Gilbert L, Medlock J, Hansford K, Thompson DBA, Biek R. 2017 Effects of conservation management of landscapes and vertebrate communities on Lyme borreliosis risk in the United Kingdom. *Phil. Trans. R. Soc. B* **372**: 20160123. <http://dx.doi.org/10.1098/rstb.2016.0123>

Accepted: 17 October 2016

One contribution of 13 to a theme issue 'Conservation, biodiversity and infectious disease: scientific evidence and policy implications'.

Subject Areas:

ecology, environmental science, health and disease and epidemiology, microbiology

Keywords:

Lyme borreliosis, *Ixodes*, conservation management, biodiversity

Author for correspondence:

Caroline Millins
e-mail: caroline.millins@glasgow.ac.uk

Effects of conservation management of landscapes and vertebrate communities on Lyme borreliosis risk in the United Kingdom

Caroline Millins^{1,2,3}, Lucy Gilbert⁴, Jolyon Medlock^{5,6}, Kayleigh Hansford⁵, Des BA Thompson⁷ and Roman Biek^{1,2}

¹Institute of Biodiversity, Animal Health and Comparative Medicine, and ²The Boyd Orr Centre for Population and Ecosystem Health, University of Glasgow, Glasgow G12 8QQ, UK

³School of Veterinary Medicine, University of Glasgow, Glasgow G61 1QH, UK

⁴The James Hutton Institute, Craigiebuckler, Aberdeen AB15 8QH, UK

⁵Medical Entomology Group, Emergency Response Department, Public Health England, Salisbury, SP4 0JG, UK

⁶Health Protection Research Unit in Environment and Health, Porton Down, Salisbury SP4 0JG, UK

⁷Scottish Natural Heritage, 231 Corstorphine Road, Edinburgh, EH12 7AT, UK

CM, 0000-0002-2006-092X

Landscape change and altered host abundance are major drivers of zoonotic pathogen emergence. Conservation and biodiversity management of landscapes and vertebrate communities can have secondary effects on vector-borne pathogen transmission that are important to assess. Here we review the potential implications of these activities on the risk of Lyme borreliosis in the United Kingdom. Conservation management activities include woodland expansion, management and restoration, deer management, urban greening and the release and culling of non-native species. Available evidence suggests that increasing woodland extent, implementing biodiversity policies that encourage ecotonal habitat and urban greening can increase the risk of Lyme borreliosis by increasing suitable habitat for hosts and the tick vectors. However, this can depend on whether deer population management is carried out as part of these conservation activities. Exclusion fencing or culling deer to low densities can decrease tick abundance and Lyme borreliosis risk. As management actions often constitute large-scale perturbation experiments, these hold great potential to understand underlying drivers of tick and pathogen dynamics. We recommend integrating monitoring of ticks and the risk of tick-borne pathogens with conservation management activities. This would help fill knowledge gaps and the production of best practice guidelines to reduce risks.

This article is part of the themed issue 'Conservation, biodiversity and infectious disease: scientific evidence and policy implications'.

1. Introduction

The management of landscapes and habitats for conservation is often driven by policies aiming to enhance biodiversity, to improve ecosystem services or to manage invasive species. These policy-driven land management changes include native woodland regeneration and restoration to optimize biodiversity, vegetation management, urban greening and the management of invasive or pest species. However, there may be unintended consequences of these management actions, such as effects on infectious-disease risk due to changes in wild vertebrate and vector population distribution, abundance and movement patterns [1,2]. As most significant human and livestock pathogens can infect many host species, changes in the host community composition can affect the




Figure 1. Overview figure of selected conservation management activities including invasive species management, woodland regeneration, urban greening and deer management that can affect vertebrate host communities, tick populations, pathogen transmission and the risk of Lyme borreliosis (© Diogo Guerra).

persistence and prevalence of these pathogens [3–6]. This is because species within a host community contribute differently to pathogen dynamics due to differences in their abundance, infection prevalence and infectiousness [7].

In recent decades, there has been a global increase in the incidence of many emerging and endemic vector-borne diseases in humans [8,9]. Although some important vector-borne diseases such as malaria and dengue fever are maintained in a human-vector cycle, most vector-borne zoonotic pathogens are maintained by vectors and multiple wild vertebrate hosts, with humans acting as dead-end hosts [10]. Shifts in land use, alterations to host populations and climate change have been identified as the main drivers of endemic vector-borne pathogen emergence, which can result in invasion of the vector-borne pathogen into new areas, or cause increased intensity of transmission within enzootic areas [1,8,11]. Therefore, conservation management activities that alter the land use or control populations of invasive or pest species are likely to result in changes to the distribution and transmission dynamics of vector-borne pathogens. This is due to effects on vertebrate host communities, which provide blood meals for vectors, and effects on the abiotic conditions for vector development and off-host survival. For example, increased deer populations in Europe have been linked to rising numbers of *Ixodes ricinus* ticks, which transmit a number of pathogens important for human health and livestock [11]. Management of deer populations in areas where they are highly abundant is often necessary as part of woodland regeneration and biodiversity projects, with consequences for ecological communities, tick populations and pathogen transmission [12,13] (see §2).

Lyme borreliosis is among the most important vector-borne zoonoses in the Northern hemisphere and has increased in distribution and incidence across large parts of North America and the higher altitudes and latitudes of Europe, including the United Kingdom [14,15]. Transmitted by Ixodid ticks, the causative agents of Lyme borreliosis are spirochaete bacteria belonging to the *Borrelia burgdorferi sensu*

lato species complex. There are at least 19 genospecies of *B. burgdorferi* s.l. in this group, some of which are pathogenic to humans [16,17]. In Western Europe, *I. ricinus* is the primary vector and is widely distributed across a range of ecoregions and habitats, and feeds on many species of birds, mammals and reptiles [18]. In the United Kingdom, four genospecies of *B. burgdorferi* s.l. have been detected in questing *I. ricinus* ticks [19–21]. These are small-mammal-associated *B. afzelii*, bird-associated *B. garinii*, *B. valaisiana* and the generalist genospecies *B. burgdorferi sensu stricto* which can be transmitted by Ixodid vectors feeding on competent reservoir hosts such as birds and small mammals [22–26]. The main route of transmission of *B. burgdorferi* s.l. is considered to be via transtadial transmission by ticks feeding on infected hosts that maintain infection through to the next life stage. Co-feeding and transovarial transmission can contribute to transmission in some circumstances [27–29]. The environmental risk to humans from Lyme borreliosis is defined as the density of infected tick vectors in the environment. The density of infected nymphs is usually focused on as the most abundant tick life-stage carrying the pathogen and is referred to as Lyme borreliosis risk throughout the rest of this paper. The probability of human exposure to these infected ticks will depend on human behaviour and how people interact with the environment. Therefore, awareness of tick-borne disease risk as well as mitigation strategies such as animal and habitat management can help to reduce human exposure to ticks.

Here, we consider the effect of common conservation management practices on the environmental risk of Lyme borreliosis in the UK. We identify four relevant areas of conservation activities: deer management, woodland management and regeneration, control of invasive species, and urban greening (figure 1). We review the evidence that these management activities have on vector populations, host communities, pathogen transmission and the environmental risk of *B. burgdorferi* s.l. and discuss knowledge gaps and policy implications (table 1). While we focus on data and examples from the UK and continental Europe, the ecological mechanisms we discuss

Table 1. Summary of the potential effects of different conservation management actions on vertebrate communities, *Ixodes ricinus* abundance and the risk of Lyme borreliosis.

type of conservation management	effect on vertebrate community?	effect on tick abundance?	effect on risk of Lyme borreliosis? (density of infected nymphs)
deer management (fencing or culling)	deer management can result in reduced deer abundance. Subsequent increases in vegetation from reduced browsing by deer can result in increased densities of competent small mammal hosts [12,30]	tick abundance would be expected to be reduced in the absence of alternate hosts for adult ticks [12]	increases in competent small mammal hosts may lead to an increased prevalence of infection, but reduced deer will reduce overall tick density, therefore the density of infected ticks (risk) will likely fall (assuming there are few alternative hosts for adult ticks)
woodland regeneration	an increase in populations of small mammals and birds (competent hosts) is predicted based on habitat-specific densities [31–33]. Deer populations (incompetent hosts) may increase if not controlled or excluded by fencing [12,34]	if deer are controlled to aid tree growth, ticks may be reduced or, if not, ticks may increase due to more favourable abiotic conditions for tick survival	the prevalence of infection is likely to increase, but the density of infected ticks may be increased or decreased depending on whether deer management is carried out
invasive species management	culling of invasive grey squirrels can result in decreased populations of this host. Populations of red squirrels may increase where present [35], populations of other competent hosts such as small mammals and birds may also increase in response	invasive species management is likely to have a limited effect in areas with tick reproduction hosts (e.g. deer)	unknown, but may lead to a decreased risk in the short term. Longer-term changes are dependent on the response of other competent small-mammal and bird populations to the removal of grey squirrels
urban greening	increased urban greenspace and connectivity will increase the abundance of both competent and incompetent vertebrate hosts in urban environments	a range of vertebrates support ticks and if larger animals (e.g. deer) are able to access urban greenspace then ticks can establish. The role of cats and dogs as tick hosts should be investigated [36]	although tick abundance may be lower in urban greenspace compared to rural areas, there is evidence that pathogen prevalence may be higher in those ticks (given that there is likely to be less of a dilution effect from large mammals) [37]. Also human exposure is likely to be higher in urban areas

are likely to be relevant to any geographical area affected by Lyme borreliosis.

2. Deer management

Conservation objectives often require the close management of large herbivores in order to improve habitat quality. Prime examples include woodland regeneration and improvement projects for biodiversity enhancement, which require management of deer to avoid damage of young trees and vegetation. Reducing grazing or browsing by deer can be achieved by

exclusion fencing or culling [12], figure 2. As deer are important hosts of Ixodid ticks in many areas, changes in deer density can affect tick abundance with implications for the transmission of tick-borne pathogens [12,38].

Deer can feed large numbers of adult female ticks, which then lay eggs and produce the next generation of immature ticks, and deer are thus termed ‘tick reproduction hosts’ [39]. A great many studies have shown that deer can be instrumental in maintaining tick populations, such that areas with more deer also have more ticks [12,19,40–51] although there is some uncertainty in the precise relationship between deer density and tick density [52]. Some of these studies specifically tested



Figure 2. Woodland regeneration projects often incorporate exclusion fences for deer to reduce browsing (© Caroline Millins).

the impact of deer management methods and, when deer numbers were reduced through culling or fencing, there were dramatic declines in the tick population. For example, a study examining *I. ricinus* tick density in response to deer management methods reported reductions of 73% on heather moorland and 94% in woodlands due to culling, and 86–88% due to fencing moorlands and 96% reductions due to fencing woodlands [12].

However, as deer are considered incompetent reservoir hosts for *B. burgdorferi* s.l. and do not infect feeding ticks [53–55], but see [56,57], increasing deer densities might not necessarily result in a higher risk of Lyme borreliosis. Mathematical models of tick-borne pathogens have predicted a non-linear relationship between deer density and the prevalence of tick-borne pathogens [58–60]. According to these models, initial increases in deer density cause increased pathogen prevalence, because more adult ticks feed successfully. Ticks in their early life stages tend to feed preferentially on small hosts that are often competent to transmit the pathogen. However, at very high deer densities, a reduction in prevalence is predicted because more and more immature ticks feed on deer that are not competent to transmit, known as a ‘dilution effect’ [38,61]. A dilution effect is defined as occurring when the addition of one or more host species to a community reduces the prevalence of a pathogen and decreases the likelihood of pathogen persistence [62].

Consistent with these predictions, a large empirical study in Italy found an increase in the prevalence of *B. burgdorferi* s.l. in ticks with increasing deer density up to a threshold of 15 deer/100 hectares (ha), after which prevalence decreased [43]. Due to positive effects of deer density on tick density, the risk of Lyme borreliosis (density of infected nymphs) continued to increase up to 60 deer/100 ha before decreasing slightly [43]. Other empirical studies have reported variable associations between deer density and *B. burgdorferi* s.l. prevalence, from positive [19] to negative [41,63] or neutral [51,64–66]. These inconsistent effects might be due to sampling that usually only covers part of the range of deer and competent reservoir host densities, which limits the chance of detecting non-linear relationships, as well as local variation in climatic factors affecting vector populations and host–vector interactions. Despite the variable

effects of deer density on *B. burgdorferi* s.l. prevalence, it remains possible that deer density may be more consistently linked to the risk of Lyme borreliosis, apart from at exceptionally high deer densities when a dilution effect may occur. Some studies have reported positive effects of deer density on Lyme borreliosis risk [43,51,67], while other studies have found no significant effect [65,66,68]. Reported differences among studies probably relate to variation in the density of competent reservoir hosts between studies.

While reducing deer densities by fencing or culling will almost certainly result in dramatically decreased tick populations when there are no suitable alternative hosts, and may decrease the risk of Lyme, there are several important issues concerning both fencing and intensive culling. These include expense, ethics, public opinion and conflicting land management objectives. For example roe deer (*Capreolus capreolus*) are increasingly present in urban green space and the peri-urban fringe and act as important tick hosts. Public opinion and practical concerns may make it extremely difficult to manage urban deer populations by culling or fencing. Furthermore, deer move within urban areas by moving along green corridors, which can include peri-domestic habitats such as gardens. Habituation of urban deer populations to humans and attractive feeding areas within these areas can increase tick densities close to human dwellings and the risk of human exposure to ticks. Culling of deer to reduce the population density is only likely to be effective when conducted at the landscape scale, which may require cooperation between private and public land managers [69]. As deer are iconic animals with cultural value for tourism and hunting, intensive culls are not always desirable [12].

Managing deer populations where alternate hosts are present is likely to be less successful in reducing tick numbers. Livestock can act as hosts for all life stages of *I. ricinus*, and other species such as mountain hares (*Lepus timidus*) can maintain *I. ricinus* populations by feeding all three tick life-stages in the absence of larger vertebrates [27,70–72]. It may be difficult to maintain fenced enclosures to prevent all deer from entering, and may not be feasible to fence large areas. Fencing can be unsightly and unpopular with countryside users and can pose risks to birds of conservation importance such as capercaillie (*Tetrao urogallus*) and black grouse (*Tetrao tetrix*) [73].

There is also evidence that fencing can lead to shifts in host communities, as seen with mountain hare (*Lepus timidus*) and small mammal densities increasing on the inside of fenced areas [12,30]. Such increases in competent host densities in response to changes in vegetation could, in principle, result in a higher prevalence of *B. burgdorferi* s.l. inside fenced areas. Studies that monitor changes in small mammal communities and tick infestations as well as questing tick abundance and *B. burgdorferi* s.l. prevalence in response to deer density control are therefore needed to assess the effect of fencing on Lyme borreliosis risk.

3. Woodland regeneration and management

Many parts of Europe are currently experiencing major land use change due to woodland expansion. For Europe as a whole, there was an increase in forested land cover of more than 600 000 km² between 1993 and 2006 [74]. While some woodland expansion is unintended due to reduced pasture management, increasing woodland cover is currently being encouraged under international and national policies [74–77]. Implementation of changes to land services is through existing policies including the European Union (EU) Biodiversity Action Plan, Article 10 of the Habitats Directive and the Strategic Environmental Assessment and Environmental Impact Assessment Directives [74]. Though, as the UK voted to leave the EU in 2016, EU policies may soon not apply.

The primary aim of these policies is to improve ecosystem services for climate change mitigation and to enhance biodiversity, water quality and human well-being. In the UK, forested land cover is currently much reduced from historic levels and is low in comparison with other countries globally and across Europe [75,78]. Government policy in Scotland is to increase woodland cover from 18% to 25% by 2050, requiring the creation of 10–15 000 hectares of woodland a year, while in England a target has been set to increase woodland cover from 9% to 12% by 2060 [76,78,79]. A proportion of this new woodland is planned to be close to urban areas to facilitate people's access and enjoyment of the outdoors [75], see §5 of this paper.

In addition to increased forest cover, there is a drive to improve woodland quality in terms of biodiversity potential and aesthetics for recreational purposes [80,81]. Existing semi-natural broadleaf woodlands, which are important for biodiversity and conservation, are often highly fragmented and embedded in a landscape of agriculture, commercial coniferous plantations and moorland [80,82]. To aid biodiversity, targets have been set to increase the area of semi-natural mixed/broadleaf woodland and to reduce fragmentation by developing ecologically functional forest habitat networks that facilitate the colonization and movement of animals and plants [77,83]. Re-wilding initiatives also aim to restore habitats, and native broadleaf woodland regeneration can form part of these projects. As a consequence, semi-natural mixed/broadleaf woodland is now the most commonly planted woodland type. These changes in woodland cover, type and connectivity are predicted to result in changes to host communities, tick populations and Lyme borreliosis risk.

As well as ecosystem-service benefits from increased woodland, there can be disservices such as increased numbers of pests, such as insects, which are damaging for agriculture or disease-transmitting vectors [78]. Although *I. ricinus* ticks can

be found in meadows, open hillside and heather moorland, the highest densities of *I. ricinus* are typically found in woodland. This is due to more favourable abiotic conditions that promote increased tick activity and survival, and due to increased densities of hosts [12,13,27,48,84]. Desiccation is a significant risk to *I. ricinus* survival as the majority of the life cycle is spent in the environment. The saturation deficit, a product of temperature and humidity and a measure of the drying power of the environment, affects the likelihood of *I. ricinus* host seeking or questing [85]. Decaying leaf litter, ground vegetation and canopy cover in woodlands tend to provide more favourable microclimatic conditions than surrounding grassland, moorland or farmland, with reduced saturation deficit allowing more frequent and prolonged questing activity [13]. Importantly, there are also generally higher densities of hosts (especially birds, rodents and roe deer) in woodlands in comparison to moorland and grassland habitats, which increases the probability of ticks obtaining a blood meal.

Based on reported habitat-specific densities, reservoir host densities for *B. burgdorferi* s.l. such as rodents, shrews and birds are predicted to increase as open habitats are converted into woodland [31–33]. This could lead to an increased prevalence of *B. burgdorferi* s.l. in questing nymphs, and, unless tick control measures are put in place, increased densities of infected ticks and an increased risk of Lyme borreliosis. Indeed, a study from central France found a higher prevalence of *B. burgdorferi* s.l. in *I. ricinus* collected from woodland habitat compared to adjacent pasture [86]. Similarly, a study from Scotland found densities of infected *I. ricinus* nymphs (a measure of the risk of Lyme borreliosis) to be five times higher in woodland compared to adjacent open habitats [34].

There is some evidence that the type of woodland may be important for the risk of Lyme borreliosis. In New York State a positive association between *B. burgdorferi* s.s. infection in small mammals and woodlands with lower canopy height and increased amounts of denser ground vegetation was detected [87]. An association between a higher prevalence of *B. burgdorferi* s.l. in questing ticks from semi-natural mixed/broadleaved woodlands compared to coniferous plantation was found in surveys from northern England and Scotland [19,20]. This was suggested to be due to higher densities of bird and rodent transmission hosts in semi-natural woodlands compared to plantations. However, a third study from Scotland did not find a difference in prevalence of *B. burgdorferi* s.l. or risk of Lyme borreliosis in semi-natural mixed/broadleaf woodland compared to coniferous plantations, indicating the difficulty in generalizing between broad habitat types [66]. The distribution of woodlands across the landscape, including levels of connectivity, fragmentation and patch size, will affect the movement of animals, tick abundance and persistence and risk from tick-borne pathogens. Increasing woodland fragmentation has been found by some researchers to be associated with increased *B. burgdorferi* s.l. prevalence and risk [88]. This may be caused by 'edge effects' and an increased proportion of suitable ecotonal habitat leading to higher densities of small mammals [86,89], increased densities of roe deer, which can maintain tick populations, and suitable humid microclimatic conditions, which are important for tick survival. Similar ecotonal habitat is found in woodland rides (linear non-wooded herbaceous habitat alongside tracks in woodlands) [13]. Management of woodland rides to maximize biodiversity by reducing scrub encroachment and encouraging wide 'sunny' rides increases browsing by deer and provides




Figure 3. Maps of red (*Sciurus vulgaris*) (a) and grey squirrel (*Sciurus carolinensis*) (b) distribution in the United Kingdom in 1945 and 2010. (Distribution maps © Red Squirrel Survival Trust, red squirrel photograph © Steve Ransome, grey squirrel photograph © Aileen Adam).

habitat suitable for small mammal reservoir hosts. This combination of ecological factors could result in increased densities of infected ticks [13]. Therefore, the creation of additional small woodlands and management for biodiversity may lead to higher densities of infected ticks and an increased risk of Lyme borreliosis particularly within ecotonal habitat at woodland edges and along woodland rides. Incorporation of tick-targeted management such as seasonal mowing and management of mulch/mat could be considered to mitigate this increase [13].

The spatial arrangement and connectivity of woodland patches is likely to affect the risk of Lyme borreliosis due to effects on host movement patterns. Isolated and fragmented small forest patches, such as those surrounded by agricultural land in France, or wooded islands within a large water body in Scotland, had a lower risk of Lyme borreliosis compared to nearby continuous forest [90,91]. This is in contrast to reports of increased Lyme borreliosis risk with increased forest fragmentation and smaller forest patch size in North America [88]. In North America, increased Lyme borreliosis risk was associated with decreased mammal-species diversity in smaller forest patches and increased densities of a competent reservoir host, the white footed mouse (*Peromyscus leucopus*). Inclusion of the surrounding landscape in a study of the effect of habitat fragmentation on *I. ricinus* abundance in northern Spain found that ‘stepping stone’ patches of habitat that facilitate host movement had the highest tick density, while patches of suitable but isolated habitat away from main movement networks had lower tick density [92]. Landscape structure and fragmentation of woodland may also affect the genospecies composition of *B. burgdorferi* s.l. This is of public health significance as different genospecies can have different clinical presentations in humans [93,94]. In a study of small wooded islands in a large lake in Scotland, only bird-associated and generalist genospecies were present in fragmented woodland habitat on islands, while both mammal- and bird-associated genospecies were present nearby on the mainland [91]. Lack of persistence of mammal-associated *B. afzelii* on islands may be associated with rodent populations falling below a critical community size during troughs in population cycles, and restricted host movements between islands [91,95].

Natural regeneration of woodlands is often limited by grazing herbivores, particularly deer [77]. It is therefore generally essential to cull deer or exclude them by fencing in order to

protect new saplings for the creation of woodlands, as described in §2 of this paper (figure 2). If large herbivores are successfully excluded from regenerating woodlands, this should mitigate the otherwise predicted increase in tick abundance and the risk of Lyme borreliosis. Increasing woodland cover without long-term strategies for deer management could lead to an increased usage of woodlands by deer, increased tick populations and elevated Lyme borreliosis risk.

4. Invasive-species management

Introduced and invasive species are a widespread challenge in natural-resource management and can affect biodiversity, ecosystem function and human health [96]. Negative effects of these species on native communities can involve predation, direct competition and the introduction of novel parasites [2,97]. In addition, introduced species can act as hosts for endemic pathogens and change the transmission dynamics and infection risk for native species and humans [2]. For example, Siberian chipmunks (*Tamias sibiricus barberi*) introduced to France have been found to be infected with multiple species of *B. burgdorferi* s.l., to harbour more ticks and to contribute more to the local risk of Lyme borreliosis than native rodents [98–100]. The North American grey squirrel (*Sciurus carolinensis*) is an invasive species of major conservation concern in the UK. Following introduction to a small number of sites in the late 19th and early 20th century, this species has invaded large parts of the UK, and now has an estimated population of over two million [31,101] (figure 3). Introduced grey squirrels in the UK thrive in urban parks and gardens as well as rural woodlands [101]. Their introduction has coincided with the decline of the formerly widespread native red squirrel (*S. vulgaris*; see figure 3)—probably the result of direct and/or apparent competition; grey squirrels are asymptomatic carriers of the squirrelpox virus, which is highly pathogenic to red squirrels [97,102–104]. As a consequence, plans for red squirrel conservation rely on halting the spread and reducing the range of grey squirrels through culling. These control efforts are concentrated on areas where the red squirrel is still present, such as large parts of Scotland (figure 3), and have led to recovery of red squirrels in some areas [35]. Targeted culls are also carried out in many other parts of the UK to protect woodlands, since grey squirrels cause significant damage to woodlands by stripping bark

from young trees [101]. This can affect broadleaf/semi-natural woodlands planted for conservation and biodiversity as well as economically important conifer plantations.

The effect of removal of grey squirrels on Lyme borreliosis risk will depend on their relative contribution to pathogen transmission and the effect on the host community competence following their removal, including changes in other reservoir host species densities. Grey squirrels in the UK have been found to carry relatively high numbers of immature *I. ricinus*, they are commonly infected with *B. burgdorferi* s.l., and infected individuals are able to transmit the pathogen to feeding larvae [105–107]. In Scotland, grey squirrels have been found to be infected with all genospecies found in questing ticks, and most commonly with *B. garinii*, a genospecies normally associated with bird hosts [107]. In addition, this study found that *B. burgdorferi* s.l. prevalence in grey squirrels rose and fell seasonally with changes in tick activity, indicating that infection in grey squirrels might be relatively short-lived [107]. Overall, infection patterns in invasive grey squirrels suggest that the species might be a spill-over rather than a maintenance host for *B. burgdorferi* s.l. [107].

While our current understanding of the role of grey squirrels in Lyme borreliosis epidemiology is still incomplete, the information available indicates that the removal of grey squirrels by itself could result in a reduction in Lyme borreliosis risk, as they host immature ticks and can transmit locally circulating strains of *B. burgdorferi* s.l. [105–107]. However, further studies are needed in particular focusing on the survival of ticks fed on grey squirrels compared to native hosts, and to measure the transmission competence of grey squirrels for different *B. burgdorferi* genospecies. In the long term, effects of grey squirrel removal will depend on the response of the host community, particularly other reservoir hosts, including red squirrels, small mammals and songbirds. Several studies have associated red squirrels with *B. burgdorferi* s.s., and it is possible that the prevalence of this genospecies could rise in local areas where populations of red squirrels increase [108–110]. There is some evidence that grey squirrels may affect populations of songbirds and small mammals [111], which are competent hosts for *B. garinii* and *B. afzelii*, respectively, so the transmission dynamics of these genospecies could also change. Dedicated studies to determine the effects of grey squirrel removal on tick abundance and *B. burgdorferi* s.l. prevalence are clearly needed to better understand and predict the consequences of grey squirrel culling management decisions. As *B. burgdorferi* s.l. strains detected in grey squirrels may reflect the locally circulating strains in questing ticks, this species may be useful as a sentinel for public health surveillance purposes [107]. Studies comparing strains in grey squirrels and in ticks at a finer scale within local woodland areas would be of great interest to investigate their usefulness as sentinels further.

While grey squirrels represent the most obvious example for a potential link between invasive species management and the risk of Lyme borreliosis in the UK, such a link might also exist for other non-native species. For some, the effect could be negligible. Sika (*Cervus nippon*) and fallow deer (*Dama dama*) for example are thought to be equally non-competent to transmit *B. burgdorferi* s.l. and occur at similar densities as native deer species, so their role in Lyme borreliosis epidemiology is likely to be similar. Wild boars (*Sus scrofa*) have been accidentally re-introduced to a few places in England and can host *I. ricinus* [112], though their role as reservoir

hosts is uncertain. In contrast, in another introduced species the ring-necked pheasant (*Phasianus colchicus*), high tick burdens, *B. burgdorferi* s.l. infection and transmission of *B. garinii* and *B. valaisiana* to feeding immature *I. ricinus* have been demonstrated [22,25,106,113]. Over 20 million pheasants are released for hunting each year in the UK [114]. This raises questions about the effect of management of host densities on the risk of Lyme borreliosis, but with pheasants being a popular and economically important game bird rather than a designated pest species as for grey squirrels.

5. Urban greening

There are several drivers for urban greening, including climate change mitigation to keep cities cooler, and peri-urban woodland regeneration and restoration projects for human well-being and biodiversity. Encroachment of towns and cities into woodland areas and increasing urban and peri-urban greenspace also increase the likelihood of human–tick contact and exposure to tick-borne pathogens. A recent review of the likely impacts of climate change on vector-borne disease in the UK has raised concerns over the possible indirect effects on vector-borne disease systems via climate change mitigation [9]. In line with these predictions, a tick surveillance scheme run by Public Health England has found increased reports of ticks acquired in urban areas (Hansford K, Medlock J. 2016 personal communication). This could suggest that like other urban areas in Europe, similar UK habitats may also be suitable for ticks and may therefore pose a hazard to humans from tick-borne diseases.

For *I. ricinus* to survive in urban areas, it needs the same ecological and environmental requirements that it has in the countryside; having access to suitable vegetation and animal hosts. The requirement for low saturation deficit (including a high relative humidity of over 80%) to reduce desiccation-associated mortality [85] restricts *I. ricinus* to urban parks, forest patches and gardens. Urban areas can provide a mosaic of ecotonal habitats, with areas of woodland, hedges, grasslands managed as meadows for flowers and insects, and parks all in close proximity. In some cases, expansion of towns has brought ancient woodland, and more traditional tick habitat, closer to the urban environment, with some cities engulfing woodland within its limits. Urban areas are home to many wildlife tick hosts including increasing numbers of urban deer and red foxes (*Vulpes vulpes*) [115,116], but also dogs and cats which can provide an abundant host for the nymphal and adult tick stages [36]. The proportion of ticks feeding on dogs and cats which are able to return to habitat that could support their development is unknown; however, it is likely that some engorged ticks will find suitable habitat to develop a subsequent generation. Dogs and cats may also carry ticks into human homes and increase human–tick exposure. Higher temperatures in urban areas compared to surrounding rural areas may also affect tick development rates [117,118] and tick–host interactions.

Like many organisms, ticks also benefit from habitat connectivity, and urban areas with well-connected habitats via green corridors would probably support more ticks. Also, urban greenspace on the margins of towns and cities, with direct connectivity to the countryside, may also be significantly more suitable for tick survival than fragmented habitats. These connected spaces are also used by members

of the public (walking or walking dogs) to navigate through urban areas, providing opportunities for exposure to ticks and potentially the movement of ticks between fragmented habitat on canine hosts. Urban tick densities are likely to be lower than those in the countryside, due to lower host density, but human exposure is likely to be much higher. The latter factor is difficult to quantify, but it is conceivable that low tick densities in a publicly accessible site will present a greater risk to human health than high tick densities in a remote woodland [37,119–121]. Behavioural research directed towards understanding the probabilities of human–tick contact in different environments is needed; human disease risk cannot be estimated with any accuracy with data relating to the environmental risk alone (i.e. density of infected nymphs).

The first report indicating a risk of Lyme borreliosis in urban green space in the UK identified a clinical case of *B. burgdorferi* s.l. infection in a dog that had visited Richmond and Bushy Park in London in 1988, and testing of questing ticks from the parks found a prevalence of *B. burgdorferi* s.l. of 8% [122]. Serological evidence of *B. burgdorferi* s.l. exposure was later detected in park workers [123]. Many further studies on *B. burgdorferi* s.l. prevalence in ticks have included Richmond Park as a site of interest [21,124–126], but this does not represent a typical urban area and its large population of red deer contribute significantly to tick survival. More recent research in cities in Southern England have investigated tick density and the prevalence of *B. burgdorferi* s.l. in more typical urban greenspace such as small parks, vegetated pathways, housing estates bordering woodland, urban grasslands managed as meadows, as well as woodlands within urban settings. Higher tick densities were found in urban woodlands or localities (e.g. parks, grasslands) adjacent to woodland, particularly on the urban fringe. Seasonal variation in *B. burgdorferi* s.l. prevalence and tick densities was found and high *B. burgdorferi* s.l. prevalence rates of up to 30% were detected in some of these habitats [37] compared to a mean of 7.5% in woodland habitat from outside urban areas in England [20]. Early findings suggest that the presence and prevalence of *B. burgdorferi* s.l. infected ticks across tick-infested habitats are heterogeneous, lending strength to the importance of connectivity in determining *B. burgdorferi* s.l. prevalence as well as tick density. Differences in habitat connectivity could impact the ability for larger animals (e.g. deer) to access urban habitats, thus increasing tick densities, but possibly reducing *B. burgdorferi* s.l. prevalence. There is also evidence from across Europe that urban habitats support ticks and associated tick-borne pathogens [127–129], with various studies comparing tick abundance and *B. burgdorferi* s.l. prevalence in both rural and urban areas [130–135], and others looking at variation across different urban green spaces [119,127,132,135–137].

For increasing parts of southern England at least, ticks appear to be emerging as an urban issue (Medlock J, Hansford K. 2016 personal communication), and owing to the complex ecology of *B. burgdorferi* s.l., the hazard of Lyme borreliosis in such areas might be comparable to more typical, rural habitats. Similar mitigation strategies can be employed in both rural and urban areas to help reduce exposure to ticks but the challenge now is how we use these control measures alongside managing greenspace for nature and the health and well-being of the public.

6. Conclusion

There are many positive effects of conservation and biodiversity management, including benefits to human well-being from spending time in nature [138]. As well as these benefits, any strategy that increases biodiversity may also increase the abundance of some organisms considered to be pests, such as vectors, and could result in a risk to public health. This review synthesizes current knowledge on the effects of conservation management of landscapes and host communities on the risk of Lyme borreliosis in the UK. This is useful to identify knowledge gaps and future research directions, and similar approaches could be made with other infectious-disease systems. Each of the conservation management actions, including deer management, woodland regeneration, invasive species management and urban greening, is predicted to result in changes to host communities and movements, tick abundance and the risk of Lyme borreliosis (table 1). Understanding and predicting the effects of particular management changes on the risk posed by a vector-borne pathogen can be complex [139]. For example, from this review, removal of a large herbivore such as deer can result in cascading effects on vegetation, small mammals and vector populations resulting in altered pathogen transmission [12].

There is significant uncertainty in the effects of some of these types of management considered in this review on Lyme borreliosis risk due to a lack of empirical studies. Land use management actions are essentially large-scale ecological experiments, which are otherwise rare and difficult to conduct. As such they provide unique opportunities to gain mechanistic insights about ecological interactions relevant to disease transmission [140]. Research studies to answer knowledge gaps outlined in each of the main sections could be designed as part of existing or new conservation projects in order to answer these questions. For example, where biodiversity monitoring is conducted to measure the success of conservation projects such as woodland regeneration or restoration, tick surveys and pathogen testing could be incorporated as part of these studies.

As the cycles of many vector-borne pathogens occur in nature, interdisciplinary collaboration between veterinary and human public health scientists with stakeholders in conservation and forest management is needed to assess and minimize these environmental hazards [10,141]. Although conservation management decisions will necessarily consider many factors, we suggest that inclusion of vector-borne pathogen dynamics and mitigation should be part of environmental impact assessments [142]. It will be important for habitat creation projects to demonstrate that in addition to measuring increases in biodiversity, such projects do not create pest and vector issues. Indeed such a lack of assessment and monitoring has blighted a number of wetland creation projects in relation to mosquito risk [143]. Ideally, the review of information and assessment of whether the risk of a vector-borne pathogen will increase or decrease with a conservation management action would be part of an interdisciplinary framework, with involvement of stakeholders in medical and veterinary public health, environmental management and biodiversity enhancement to provide advice to government [10,141].

In line with efforts in relation to managing habitats for mosquitoes [142], further research on how conservation management impacts Lyme borreliosis risk would inform best-practice guidelines for practitioners in how to manage woodlands accessible by the public in order to minimize

the exposure of people to infected ticks. This could include guidelines on environmental management to minimize the hazard from ticks, for example by managing the height of ground vegetation close to paths [13]. Another important measure would be to provide appropriate risk messaging about ticks and tick-borne diseases, with surveys to assess the effectiveness of these measures [79].

Authors' contributions. C.M. and R.B. designed the review; C.M., L.G., J.M., K.H., D.B.A.T. and R.B. contributed to the initial drafts and revisions of the manuscript. All authors reviewed and approved the final manuscript.

Competing interests. The authors declare that they have no competing interests.

Funding. C.M. received support from the School of Veterinary Medicine, University of Glasgow and the Biotechnology and Biological Sciences Research Council (BBSRC) Doctoral Training grant (BB/

F016786/1). J.M. was partly funded by the National Institute for Health Research Health Protection Research Unit (NIHR HPRU) in Environmental Change and Health at the London School of Hygiene & Tropical Medicine in partnership with Public Health England (PHE), and in collaboration with the University of Exeter, University College London, and the Met Office; and partly funded by the NIHR HPRU on Emerging Infections and Zoonoses at the University of Liverpool in partnership with PHE and Liverpool School of Tropical Medicine. The views expressed are those of the authors and not necessarily those of the National Health Service, the NIHR, the Department of Health, or PHE. L.G. was supported by the Scottish Government's Rural and Environment Science and Analytical Services Division (RESAS).

Acknowledgements. Thank you to the Red Squirrel Survival Trust for giving permission to use maps of grey and red squirrel distribution in the UK. The illustration in figure 1 was created by Diogo Guerra, www.diogoguerra.com. Thank you to anonymous reviewers who provided valuable comments on an earlier version of the manuscript.

References

- Lambin EF, Tran A, Vanwambeke SO, Linard C, Soti V. 2010 Pathogenic landscapes: interactions between land, people, disease vectors, and their animal hosts. *Int. J. Health Geogr.* **9**, 54–67. (doi:10.1186/1476-072X-9-54)
- Telfer S, Bown K. 2012 The effects of invasion on parasite dynamics and communities. *Funct. Ecol.* **26**, 1288–1299. (doi:10.1111/j.1365-2435.2012.02049.x)
- Woolhouse MEJ, Taylor L, Haydon DT. 2001 Population biology of multihost pathogens. *Science* **292**, 1109–1112. (doi:10.1126/science.1059026)
- Haydon DT, Cleaveland S, Taylor LH, Laurenson MK. 2002 Identifying reservoirs of infection: a conceptual and practical challenge. *Emerg. Infect. Dis.* **8**, 1468–1473. (doi:10.3201/eid0812.010317)
- LoGiudice K, Ostfeld RS, Schmidt KA, Keesing F. 2003 The ecology of infectious disease: effects of host diversity and community composition on Lyme disease risk. *Proc. Natl Acad. Sci. USA* **100**, 567–571. (doi:10.1073/pnas.0233733100)
- LoGiudice K, Duerr STK, Newhouse MJ, Schmidt KA, Killilea ME, Ostfeld RS. 2008 Impact of host community composition on Lyme disease risk. *Ecology* **89**, 2841–2849. (doi:10.1890/07-1047.1)
- Streicker DG, Fenton A, Pedersen AB. 2013 Differential sources of host species heterogeneity influence the transmission and control of multihost parasites. *Ecol. Lett.* **16**, 975–984. (doi:10.1111/ele.12122)
- Kilpatrick AM, Randolph SE. 2012 Drivers, dynamics, and control of emerging vector-borne zoonotic diseases. *Lancet* **380**, 1946–1955. (doi:10.1016/S0140-6736(12)61151-9)
- Medlock JM, Leach SA. 2015 Effect of climate change on vector-borne disease risk in the UK. *Lancet Infect. Dis.* **15**, 721–730. (doi:10.1016/S1473-3099(15)70091-5)
- Braks M *et al.* 2014 Vector-borne disease intelligence: strategies to deal with disease burden and threats. *Front. Public Health* **2**, 280. (doi:10.3389/fpubh.2014.00280)
- Medlock JM *et al.* 2013 Driving forces for changes in geographical distribution of *Ixodes ricinus* ticks in Europe. *Parasit. Vectors* **6**, 1. (doi:10.1186/1756-3305-6-1)
- Gilbert L, Maffey GL, Ramsay SL, Hester A. 2012 The effect of deer management on the abundance of *Ixodes ricinus* in Scotland. *Ecol. Appl.* **22**, 658–667. (doi:10.1890/11-0458.1)
- Medlock JM, Shuttleworth H, Copley V, Hansford K, Leach S. 2012 Woodland biodiversity management as a tool for reducing human exposure to *Ixodes ricinus* ticks: a preliminary study in an English woodland. *J. Vector Ecol.* **37**, 307–315. (doi:10.1111/j.1948-7134.2012.00232.x)
- Linard C, Lamarque P, Heyman P, Ducoffre G, Luyasu V, Tersago K, Vanwambeke SO, Lambin EF. 2007 Determinants of the geographic distribution of Puumala virus and Lyme borreliosis infections in Belgium. *Int. J. Health Geogr.* **6**, 1–14. (doi:10.1186/1476-072X-6-15)
- Rizzoli A, Hauffe HC, Carpi G, Vourc'h GI, Neteler M, Rosà R. 2011 Lyme borreliosis in Europe. *Eurosurveillance* **16**, 19906.
- Margos G, Vollmer SA, Ogden NH, Fish D. 2011 Population genetics, taxonomy, phylogeny and evolution of *Borrelia burgdorferi* sensu lato. *Infect. Genet. Evol.* **11**, 1545–1563. (doi:10.1016/j.meegid.2011.07.022)
- Ivanova LB *et al.* 2014 *Borrelia chilensis*, a new member of the *Borrelia burgdorferi* sensu lato complex that extends the range of this genospecies in the Southern Hemisphere. *Environ. Microbiol.* **16**, 1069–1080. (doi:10.1111/1462-2920.12310)
- Gray JS. 1991 The development and seasonal activity of the tick *Ixodes ricinus*: a vector of Lyme borreliosis. *Rev. Med. Vet. Entomol.* **79**, 323–333.
- James MC, Bowman AS, Forbes KJ, Lewis F, McLeod JE, Gilbert L. 2013 Environmental determinants of *Ixodes ricinus* ticks and the incidence of *Borrelia burgdorferi* sensu lato, the agent of Lyme borreliosis, in Scotland. *Parasitology* **140**, 237–246. (doi:10.1017/S003118201200145X)
- Bettridge J, Renard M, Zhao F, Bown KJ, Birtles RJ. 2013 Distribution of *Borrelia burgdorferi* sensu lato in *Ixodes ricinus* populations across central Britain. *Vector Borne Zoonotic Dis.* **13**, 139–146. (doi:10.1089/vbz.2012.1075)
- Hansford KM, Fonville M, Jahfari S, Sprong H, Medlock JM. 2014 *Borrelia miyamotoi* in host-seeking *Ixodes ricinus* ticks in England. *Epidemiol. Infect.* **143**, 1079–1087. (doi:10.1017/S0950268814001691)
- Kurtenbach K, Carey D, Hoodless AN, Nuttall PA, Randolph SE. 1998 Competence of pheasants as reservoirs for Lyme disease spirochetes. *J. Med. Entomol.* **35**, 77–81. (doi:10.1093/jmedent/35.1.77)
- Comstedt P, Jakobsson T, Bergström S. 2011 Global ecology and epidemiology of *Borrelia garinii* spirochetes. *Infect. Ecol. Epidemiol.* **1**, 9545. (doi:10.3402/iee.v1i0.9545)
- Gern L, Siegenthaler M, Hu CM, Humair PF, Moret J. 1994 *Borrelia burgdorferi* in rodents (*Apodemus flavicollis* and *A. sylvaticus*): duration and enhancement of infectivity for *Ixodes ricinus*. *Eur. J. Epidemiol.* **10**, 75–80. (doi:10.1007/BF01717456)
- Kurtenbach K, Peacey M, Rijpkema SGT, Hoodless AN, Nuttall PA, Randolph SE. 1998 Differential transmission of the genospecies of *Borrelia burgdorferi* sensu lato by game birds and small rodents in England. *Appl. Environ. Microbiol.* **64**, 1169–1174.
- Hanincová K, Schäfer SM, Etti S, Sewell HS, Taragelová V, Ziak D, Labuda M, Kurtenbach K. 2003 Association of *Borrelia afzelii* with rodents in Europe. *Parasitology* **126**, 11–20. (doi:10.1017/S0031182002002548)
- Ogden NH, Nuttall PA, Randolph SE. 1997 Natural Lyme disease cycles maintained via sheep by co-feeding ticks. *Parasitology* **115**, 591–600. (doi:10.1017/S0031182097001868)
- Voordouw MJ. 2015 Co-feeding transmission in Lyme disease pathogens. *Parasitology* **142**, 290–302. (doi:10.1017/S0031182014001486)
- van Duijvendijk G *et al.* 2016 Larvae of *Ixodes ricinus* transmit *Borrelia afzelii* and *B. miyamotoi* to vertebrate hosts. *Parasit. Vectors* **9**, 97. (doi:10.1186/s13071-016-1389-5)

30. Perkins SE, Cattadori IM, Tagliapietra V, Rizzoli AP, Hudson PJ. 2006 Localized deer absence leads to tick amplification. *Ecology* **87**, 1981–1986. (doi:10.1890/0012-9658)
31. Harris S, Morris P, Wray S, Yalden D. 1995 *A review of British mammals: population estimates and conservation status of British mammals other than cetaceans*. Peterborough, UK: Joint Nature Conservation Committee.
32. Gilbert L, Jones LD, Hudson PJ, Gould EA, Reid HW. 2000 Role of small mammals in the persistence of Louping-ill virus: field survey and tick co-feeding studies. *Med. Vet. Entomol.* **14**, 277–282. (doi:10.1046/j.1365-2915.2000.00236.x)
33. British Trust for Ornithology. 2009 BBS Maps of population density and trends. See <http://www.bto.org/volunteer-surveys/bbs/latest-results/maps-population-density-and-trends> (accessed 17 June 2016).
34. Gilbert L. 2016 How landscapes shape Lyme borreliosis risk. In *Ecology and control of vector borne diseases* (eds MAH Braks, SE van Wieren, W Takken, H Sprong), pp. 161–171. Wageningen, Netherlands: Wageningen Academic Publishers (WAP).
35. Shuttleworth CM, Lurz PWW, Halliwell EC (eds). 2015 *Shared experience of red squirrel conservation practice*. Warwickshire, UK: European Squirrel Initiative.
36. Abdullah S, Helps C, Tasker S, Newbury H, Wall R. 2016 Ticks infesting domestic dogs in the UK: a large-scale surveillance programme. *Parasit. Vectors* **9**, 391. (doi:10.1186/s13071-016-1673-4)
37. Hansford KM *et al.* 2017 Lyme borreliosis risk posed by ticks found in greenspace within an urban/peri-urban area in southern England. *Ticks Tick Borne Dis.* **8**, 353–361. (doi:10.1016/j.ttbdis.2016.12.009)
38. Rosà R, Pugliese A, Norman R, Hudson PJ. 2003 Thresholds for disease persistence in models for tick-borne infections including non-viraemic transmission, extended feeding and tick aggregation. *J. Theor. Biol.* **224**, 359–376. (doi:10.1016/S0022-5193(03)00173-5)
39. Gray JS. 1998 The ecology of ticks transmitting Lyme borreliosis. *Exp. Appl. Acarol.* **22**, 249–258. (doi:10.1023/A:1006070416135)
40. Wilson ML, Ducey AM, Litwin TS, Gavin TA, Spielman A. 1990 Microgeographic distribution of immature *Ixodes dammini* ticks correlated with that of deer. *Med. Vet. Entomol.* **4**, 151–159. (doi:10.1111/j.1365-2915.1990.tb00273.x)
41. Gray JS, Kahl O, Janetzki C, Stein J. 1992 Studies on the ecology of Lyme disease in a deer forest in County Galway, Ireland. *J. Med. Entomol.* **29**, 915–920. (doi:10.1093/jmedent/29.6.915)
42. Deblinger RD, Wilson ML, Rimmer DW, Spielman A. 1993 Reduced abundance of immature *Ixodes dammini* (Acari: Ixodidae) following incremental removal of deer. *J. Med. Entomol.* **30**, 144–150. (doi:10.1093/jmedent/30.1.144)
43. Rizzoli A, Merler S, Furlanello C, Genchi C. 2002 Geographical information systems and bootstrap aggregation (bagging) of tree-based classifiers for Lyme disease risk prediction in Trentino, Italian Alps. *J. Med. Entomol.* **39**, 485–492. (doi:10.1603/0022-2585-39.3.485)
44. Stafford KC, DeNicola AJ, Kilpatrick HJ. 2003 Reduced abundances of *Ixodes scapularis* (Acari: Ixodidae) and the tick parasitoid *Ixodiphagus hookeri* (Hymenoptera: Encyrtidae) with reduction of white-tailed deer. *J. Med. Entomol.* **40**, 642–652. (doi:10.1603/0022-2585-40.5.642)
45. Rand PW, Lubelczyk C, Lavigne GR, Elias S, Holman MS, Lacombe EH, Smith RP. 2003 Deer density and the abundance of *Ixodes scapularis* (Acari: Ixodidae). *J. Med. Entomol.* **40**, 179–184. (doi:10.1603/0022-2585-40.2.179)
46. Rand PW, Lubelczyk C, Holman MS, Lacombe EH, Smith RP. 2004 Abundance of *Ixodes scapularis* (Acari: Ixodidae) after the complete removal of deer from an isolated offshore island, endemic for Lyme Disease. *J. Med. Entomol.* **41**, 779–784. (doi:10.1603/0022-2585-41.4.779)
47. Gilbert L. 2010 Altitudinal patterns of tick and host abundance: a potential role for climate change in regulating tick-borne diseases? *Oecologia (Berlin)* **162**, 217–225. (doi:10.1007/s00442-009-1430-x)
48. Ruiz-Fons F, Gilbert L. 2010 The role of deer as vehicles to move ticks, *Ixodes ricinus*, between contrasting habitats. *Int. J. Parasitol.* **40**, 1013–1020. (doi:10.1016/j.ijpara.2010.02.006)
49. Vor T, Kiffler C, Hagedorn P, Niedrig M, Rühle F. 2010 Tick burdens on European roe deer (*Capreolus capreolus*). *Exp. Appl. Acarol.* **51**, 405–417. (doi:10.1007/s10493-010-9337-0)
50. Vourc'h G, Abrial D, Bord S, Jacquot M, Masseglia S, Poux V, Pisanu B, Bailly X, Chapuis JL. 2016 Mapping human risk of infection with *Borrelia burgdorferi* sensu lato, the agent of Lyme borreliosis, in a periurban forest in France. *Ticks Tick Borne Dis.* **7**, 644–652. (doi:10.1016/j.ttbdis.2016.02.008)
51. Werden L, Barker IK, Bowman J, Gonzales EK, Leighton PA, Lindsay LR, Jardine CM. 2014 Geography, deer, and host biodiversity shape the pattern of Lyme disease emergence in the Thousand Islands archipelago of Ontario, Canada. *PLoS ONE* **9**, 85640. (doi:10.1371/journal.pone.0085640)
52. Kilpatrick AM *et al.* 2017 Lyme disease ecology in a changing world: consensus, uncertainty and critical gaps for improving control. *Phil. Trans. R. Soc. B* **372**, 20160117. (doi:10.1098/rstb.2016.0117)
53. Telford SR, Mather TN, Moore SI, Wilson ML, Spielman A. 1988 Incompetence of deer as reservoirs of the Lyme disease spirochete. *Am. J. Trop. Med. Hyg.* **39**, 105–109.
54. Jaenson TGT, Tälleklint L. 1992 Incompetence of roe deer (*Capreolus capreolus*) as reservoirs of the Lyme disease spirochete. *J. Med. Entomol.* **29**, 813–817. (doi:10.1093/jmedent/29.5.813)
55. Kurtenbach K, De Michelis S, Etti S, Schafer SM, Sewell H-S, Brade V, Kraiczky P. 2002 Host association of *Borrelia burgdorferi* sensu lato—the key role of host complement. *Trends Microbiol.* **10**, 74–79. (doi:10.1016/S0966-842X(01)02298-3)
56. Oliver J, Stallknecht D, Chandler F, James A, McGuire B, Howerth E. 1992 Detection of *Borrelia burgdorferi* in laboratory-reared *Ixodes dammini* (Acari: Ixodidae) fed on experimentally inoculated white-tailed deer. *J. Med. Entomol.* **29**, 980–984. (doi:10.1093/jmedent/29.6.980)
57. Kimura K, Isogai E, Isogai H, Kamewaka Y, Nishikawa T, Ishii N, Fujii N. 1995 Detection of Lyme disease spirochaetes in the skin of naturally infected wild Sika deer (*Cervus nippon yezoensis*). *Appl. Environ. Microbiol.* **61**, 1641–1642.
58. Gilbert L, Norman R, Laurenson KM, Reid HW, Hudson PJ. 2001 Disease persistence and apparent competition in a three-host community: an empirical and analytical study of large-scale, wild populations. *J. Anim. Ecol.* **70**, 1053–1061. (doi:10.1046/j.0021-8790.2001.00558.x)
59. Hartemink NA, Randolph SE, Davis SA, Heesterbeek JAP. 2008 The basic reproduction number for complex disease systems: defining R_0 for tick-borne infections. *Am. Nat.* **171**, 743–754. (doi:10.1086/587530)
60. Mannelli A, Bertolotti L, Gern L, Gray J. 2012 Ecology of *Borrelia burgdorferi* sensu lato in Europe: transmission dynamics in multi-host systems, influence of molecular processes and effects of climate change. *FEMS Microbiol. Rev.* **36**, 837–861. (doi:10.1111/j.1574-6976.2011.00312.x)
61. Ostfeld R, Keesing F. 2000 Biodiversity and disease risk: the case of Lyme disease. *Conserv. Biol.* **14**, 722–728. (doi:10.1046/j.1523-1739.2000.99014.x)
62. Begon M. 2008 Effects of host diversity on disease dynamics. In *Infectious disease ecology: effects of ecosystems on disease and of disease on ecosystems* (eds R Ostfeld, F Keesing, V Eviner), pp. 12–29. Princeton, NJ: Princeton University Press.
63. Mysterud A, Easterday WR, Quiller L, Viljugrein H, Yttrup B. 2013 Spatial and seasonal variation in the prevalence of *Anaplasma phagocytophilum* and *Borrelia burgdorferi* sensu lato in questing *Ixodes ricinus* ticks in Norway. *Parasit. Vectors* **6**, 187. (doi:10.1186/1756-3305-6-187)
64. Pichon B, Mousson L, Figureau C, Rodhain F, Perez-Eid C. 1999 Density of deer in relation to the prevalence of *Borrelia burgdorferi* s.l. in *Ixodes ricinus* nymphs in Rambouillet forest, France. *Exp. Appl. Acarol.* **23**, 267–275. (doi:10.1023/A:1006023115617)
65. Ostfeld RS, Canham CD, Oggenfuss K, Winchcombe RJ, Keesing F. 2006 Climate, deer, rodents, and acorns as determinants of variation in Lyme-disease risk. *PLoS Biol.* **4**, 1058–1068. (doi:10.1371/journal.pbio.0040145)
66. Millins C, Gilbert L, Johnson P, James M, Kilbride E, Birtles R, Biek R. 2016 Spatial and temporal heterogeneity in the abundance and distribution of *Ixodes ricinus* and *Borrelia burgdorferi* (sensu lato) in Scotland: implications for risk prediction. *Parasit. Vectors* **9**, 595. (doi:10.1186/s13071-016-1875-9)
67. Kilpatrick HJ, Labonte AM, Stafford KC. 2014 The relationship between deer density, tick abundance, and human cases of Lyme disease in a residential community. *J. Med. Entomol.* **51**, 777–784. (doi:10.1603/ME12322)
68. Jordan RA, Schulze TL, Jahn MB. 2007 Effects of reduced deer density on the abundance of *Ixodes*

- scapularis* (Acari: Ixodidae) and Lyme disease incidence in a northern New Jersey endemic area. *J. Med. Entomol.* **44**, 752–757. (doi:10.1093/jmedent/44.5.752)
69. Wäber K, Spencer J, Dolman PM. 2013 Achieving landscape-scale deer management for biodiversity conservation: the need to consider sources and sinks. *J. Wildl. Manage.* **77**, 726–736. (doi:10.1002/jwmg.530)
70. Jaenson TGT, Tälleklint L. 1996 Lyme borreliosis spirochetes in *Ixodes ricinus* (Acari: Ixodidae) and the varying hare on isolated islands in the Baltic Sea. *Exp. Appl. Acarol.* **33**, 339–343. (doi:10.1007/BF00053310)
71. Laurenson AMK, Norman RA, Gilbert L, Reid HW, Hudson PJ, Journal S, Jan N. 2003 Identifying disease reservoirs in complex systems: mountain hares as reservoirs of ticks and louping-ill virus, pathogens of red grouse. *J. Anim. Ecol.* **72**, 177–185. (doi:10.1046/j.1365-2656.2003.00688.x)
72. Medlock J *et al.* 2008 Investigation of ecological and environmental determinants for the presence of questing *Ixodes ricinus* (Acari: Ixodidae) on Gower, South Wales. *J. Med. Entomol.* **45**, 314–325. (doi:10.1093/jmedent/45.2.314)
73. Baines D, Summers RW. 1997 Assessment of bird collisions with deer fences in Scottish forests. *J. Appl. Ecol.* **34**, 941–948. (doi:10.2307/2405284)
74. IEEP. 2010 Reflecting environmental land use needs into EU policy: preserving and enhancing the environmental benefits of land services: soil sealing, biodiversity corridors, intensification/marginalisation of land use and permanent grassland. See http://www.ieep.eu/assets/637/Land_services_-_Executive_summary.pdf. (accessed 5 May 2016).
75. Forestry Commission. 2009 The Scottish Government's rationale for woodland expansion. See <http://scotland.forestry.gov.uk/images/corporate/pdf/> (accessed 7 May 2016).
76. Woodland Expansion Advisory Group. 2012 Report of the Woodland Expansion Advisory Group to the cabinet secretary for rural affairs and environment, Richard Lochhead MSP. See <http://scotland.forestry.gov.uk/supporting/management/annual-review/woodland-expansion> (accessed 5 May 2016).
77. Quine CP, Cahalan C, Hester A, Humphrey J, Kirby K, Moffat A, Valatin G. 2011 Woodlands. In *UK National Ecosystem Assessment: Technical Report*, pp. 241–295. Cambridge, UK: UNEP-WCMC.
78. Thomas HJD, Paterson JS, Metzger MJ, Sing L. 2015 Towards a research agenda for woodland expansion in Scotland. *For. Ecol. Manage.* **349**, 149–161. (doi:10.1016/j.foreco.2015.04.003)
79. Quine CP *et al.* 2011 Frameworks for risk communication and disease management: the case of Lyme disease and countryside users. *Phil. Trans. R. Soc. B* **366**, 2010–2022. (doi:10.1098/rstb.2010.0397)
80. Peterken GFF. 1977 Habitat conservation priorities in British and European woodland. *Biol. Conserv.* **11**, 223–236. (doi:10.1016/0006-3207(77)90006-4)
81. O'Brien L, Morris J. 2009 Active England: the woodland projects. Available online [http://www.forestry.gov.uk/pdf/active_england_final_report.pdf/\\$FILE/active_england_final_report.pdf](http://www.forestry.gov.uk/pdf/active_england_final_report.pdf/$FILE/active_england_final_report.pdf) (accessed 6 May 2016).
82. Gimona A, Poggio L, Brown I, Castellazzi M. 2012 Woodland networks in a changing climate: threats from land use change. *Biol. Conserv.* **149**, 93–102. (doi:10.1016/j.biocon.2012.01.060)
83. Bailey SA. 2007 Increasing connectivity in fragmented landscapes: an investigation of evidence for biodiversity gain in woodlands. *For. Ecol. Manage.* **338**, 7–23. (doi:10.1016/j.foreco.2006.09.049)
84. Walker AR, Alberdi MP, Urquhart KA, Rose H. 2001 Risk factors in habitats of the tick *Ixodes ricinus* influencing human exposure to *Ehrlichia phagocytophila* bacteria. *Med. Vet. Entomol.* **15**, 40–49. (doi:10.1046/j.1365-2915.2001.00271.x)
85. Randolph SE. 2004 Tick ecology: processes and patterns behind the epidemiological risk posed by ixodid ticks as vectors. *Parasitology* **129**, S37–S65. (doi:10.1017/S0031182004004925)
86. Halos L, Bord S, Cotté V, Gasqui P, Abrial D, Barnouin J, Boulouis H-J, Vayssier-Taussat M, Vourc'h G. 2010 Ecological factors characterizing the prevalence of bacterial tick-borne pathogens in *Ixodes ricinus* ticks in pastures and woodlands. *Appl. Environ. Microbiol.* **76**, 4413–4420. (doi:10.1128/AEM.00610-10)
87. Prusinski MA *et al.* 2006 Habitat structure associated with *Borrelia burgdorferi* prevalence in small mammals in New York State. *Environ. Entomol.* **35**, 308–319. (doi:10.1603/0046-225X-35.2.308)
88. Allan B, Keesing F, Ostfeld R. 2003 Effect of forest fragmentation on Lyme disease risk. *Conserv. Biol.* **17**, 267. (doi:10.1046/j.1523-1739.2003.01260.x)
89. Simon JA *et al.* 2014 Climate change and habitat fragmentation drive the occurrence of *Borrelia burgdorferi*, the agent of Lyme disease, at the northeastern limit of its distribution. *Evol. Appl.* **7**, 750–764. (doi:10.1111/eva.12165)
90. Ferquel E, Garnier M, Marie J, Berne C, Baranton G, Pe C. 2006 Prevalence of *Borrelia burgdorferi* sensu lato and Anaplasmataceae members in *Ixodes ricinus* ticks in Alsace, a focus of Lyme borreliosis endemicity in France. *Appl. Environ. Microbiol.* **72**, 3074–3078. (doi:10.1128/AEM.72.4.3074)
91. Millins C. 2016 Ecological drivers of a vector borne pathogen: Distribution and abundance of *Borrelia burgdorferi* sensu lato and its vector *Ixodes ricinus* in Scotland. PhD thesis, University of Glasgow, UK.
92. Estrada-Peña A. 2003 The relationships between habitat topology, critical scales of connectivity and tick abundance *Ixodes ricinus* in a heterogeneous landscape in northern Spain. *Ecography* **26**, 661–671. (doi:10.1034/j.1600-0587.2003.03530.x)
93. van Dam AP, Kuiper H, Vos K, Widjokusumo A, de Jongh BM, Spanjaard L, Ramselaar AC, Kramer MD, Dankert J. 1993 Different genospecies of *Borrelia burgdorferi* are associated with distinct clinical manifestations of Lyme borreliosis. *Clin. Infect. Dis.* **17**, 708–717. (doi:10.1093/clinids/17.4.708)
94. Stanek G, Reiter M. 2011 The expanding Lyme *Borrelia* complex-clinical significance of genomic species? *Clin. Microbiol. Infect.* **17**, 487–493. (doi:10.1111/j.1469-0691.2011.03492.x)
95. Lloyd-Smith JO, Cross PC, Briggs CJ, Daugherty M, Getz WM, Latto J, Sanchez MS, Smith AB, Swei A. 2005 Should we expect population thresholds for wildlife disease? *Trends Ecol. Evol.* **20**, 511–519. (doi:10.1016/j.tree.2005.07.004)
96. Crowl TA, Crist TO, Parmenter RR, Belovsky G, Lugo AE. 2008 The spread of invasive species and infectious disease as drivers of ecosystem change. *Front. Ecol. Environ.* **6**, 238–246. (doi:10.1890/070151)
97. Tompkins DM, Sainsbury AW, Nettleton P, Buxton D, Gurnell J. 2002 Parapoxvirus causes a deleterious disease in red squirrels associated with UK population declines. *Proc. R. Soc. Lond. B* **269**, 529–533. (doi:10.1098/rspb.2001.1897)
98. Pisanu B, Marsot M, Marmet J, Chapuis J-L, Réale D, Vourc'h G. 2010 Introduced Siberian chipmunks are more heavily infested by Ixodid ticks than are native bank voles in a suburban forest in France. *Int. J. Parasitol.* **40**, 1277–1283. (doi:10.1016/j.ijpara.2010.03.012)
99. Marsot M, Sigaud M, Chapuis J-L, Ferquel E, Cornet M, Vourc'h G. 2011 Introduced Siberian chipmunks (*Tamias sibiricus barberi*) harbor more-diverse *Borrelia burgdorferi* sensu lato genospecies than native bank voles (*Myodes glareolus*). *Appl. Environ. Microbiol.* **77**, 5716–5721. (doi:10.1128/AEM.01846-10)
100. Marsot M, Chapuis J-L, Gasqui P, Dozières A, Masségla S, Pisanu B, Ferquel E, Vourc'h G. 2013 Introduced Siberian chipmunks (*Tamias sibiricus barberi*) contribute more to Lyme borreliosis risk than native reservoir rodents. *PLoS ONE* **8**, e55377. (doi:10.1371/journal.pone.0055377)
101. Mayle BA, Broome AC. 2013 Changes in the impact and control of an invasive alien: the grey squirrel (*Sciurus carolinensis*) in Great Britain, as determined from regional surveys. *Pest Manag. Sci.* **69**, 323–333. (doi:10.1002/ps.3458)
102. Rushton SP, Lurz PWW, Gurnell J, Nettleton P, Bruemmer C, Shirley MDF, Sainsbury AW. 2006 Disease threats posed by alien species: the role of a poxvirus in the decline of the native red squirrel in Britain. *Epidemiol. Infect.* **134**, 521–533. (doi:10.1017/S0950268805005303)
103. Sainsbury AW *et al.* 2008 Poxviral disease in red squirrels *Sciurus vulgaris* in the UK: spatial and temporal trends of an emerging threat. *Ecohealth* **5**, 305–316. (doi:10.1007/s10393-008-0191-z)
104. Chantrey J, Dale TD, Read JM, White S, Whitfield F, Jones D, McInnes CJ, Begon M. 2014 European red squirrel population dynamics driven by squirrelpox at a grey squirrel invasion interface. *Ecol. Evol.* **4**, 3788–3799. (doi:10.1002/ece3.1216)
105. Craine NG, Randolph SE, Nuttall PA. 1995 Seasonal variation in the role of grey squirrels as hosts of *Ixodes ricinus*, the tick vector of the Lyme disease spirochaete, in a British woodland. *Folia Parasitol. (Praha)*. **42**, 73–80.
106. Craine NG, Nuttall PA, Marriott AC, Randolph SE. 1997 Role of grey squirrels and pheasants in the

- transmission of *Borrelia burgdorferi sensu lato*, the Lyme disease spirochaete, in the U.K. *Folia Parasitol. (Praha)*. **44**, 155–160.
107. Millins C, Magierecka A, Gilbert L, Eloff A, Brereton A, Kilbride E, Denwood M, Birtles R, Biek R. 2015 An invasive mammal (gray squirrel, *Sciurus carolinensis*) commonly hosts diverse and atypical genotypes of the zoonotic pathogen *Borrelia burgdorferi sensu lato*. *Appl. Environ. Microbiol.* **81**, 109–115. (doi:10.1128/AEM.00109-15)
 108. Humair P-F, Gern L. 1998 Relationship between *Borrelia burgdorferi sensu lato* species, red squirrels (*Sciurus vulgaris*) and *Ixodes ricinus* in enzootic areas in Switzerland. *Acta Trop.* **69**, 213–227. (doi:10.1016/S0001-706X(97)00126-5)
 109. Pisanu B, Chapuis J-L, Dozières A, Basset F, Poux V, Vourc'h G. 2014 High prevalence of *Borrelia burgdorferi* s.l. in the European red squirrel *Sciurus vulgaris* in France. *Ticks Tick Borne Dis.* **5**, 1–6. (doi:10.1016/j.ttbdis.2013.07.007)
 110. James MC, Gilbert L, Bowman AS, Forbes KJ. 2014 The heterogeneity, distribution, and environmental associations of *Borrelia burgdorferi sensu lato*, the agent of Lyme borreliosis, in Scotland. *Front. Public Health* **2**, 129–139. (doi:10.3389/fpubh.2014.00129)
 111. Hewson CM, Fuller RJ. 2003 Impacts of grey squirrels on woodland birds: an important predator of eggs and young? *Br. Trust Ornithol. Res. Rep. No.* 328.
 112. Ruiz-Fons F, Fernandez-de-Mera IG, Acevedo P, Hofle U, Vicente J, de la Fuente J, Gortazar C. 2006 Ixodid ticks parasitizing Iberian red deer (*Cervus elaphus hispanicus*) and European wild boar (*Sus scrofa*) from Spain: geographical and temporal distribution. *Vet. Parasitol.* **140**, 133–142. (doi:10.1016/j.vetpar.2006.03.033)
 113. Hoodless AN, Kurtenbach K, Nuttall PA, Randolph SE. 2002 The impact of ticks on pheasant territoriality. *Oikos* **96**, 245–250. (doi:10.1034/j.1600-0706.2002.960206.x)
 114. Tapper SC. 1999 *A question of balance: game animals and their role in the British countryside*. Fordingbridge, UK: The Game Conservancy Trust.
 115. Macdonald DW, Newdick MT. 1982 The distribution and ecology of foxes, *Vulpes vulpes* (L.) in urban areas. In *Urban ecology* (eds JA Lee, R Bornkamm, MRD Seaward), pp. 123–135. Oxford, UK: Blackwell Scientific Publications.
 116. Rotherham I, Derbyshire M. 2012 Deer in the Peak District and its urban fringe. *Br. Wildl.* **23**, 256–264.
 117. Macleod J. 1935 *Ixodes ricinus* in relation to its physical environment III. Climate and reproduction. *Parasitology* **27**, 489–500.
 118. Randolph S, Green RM, Hoodless A, Peacey MF. 2002 An empirical quantitative framework for the seasonal population dynamics of the tick *Ixodes ricinus*. *Int. J. Parasitol.* **32**, 979–989. (doi:10.1016/S0020-7519(02)00030-9)
 119. Junttila J, Peltomaa M, Soini H, Marjamäki M, Viljanen MK. 1999 Prevalence of *Borrelia burgdorferi* in *Ixodes ricinus* ticks in urban recreational areas of Helsinki. *J. Clin. Microbiol.* **37**, 1361–1365.
 120. Jaenson TGT, Eisen L, Comstedt P, Mejlon HA, Lindgren E, Bergström S, Olsen B. 2009 Risk indicators for the tick *Ixodes ricinus* and *Borrelia burgdorferi sensu lato* in Sweden. *Med. Vet. Entomol.* **23**, 226–237. (doi:10.1111/j.1365-2915.2009.00813.x)
 121. Soleng A, Kjelland V. 2013 *Borrelia burgdorferi sensu lato* and *Anaplasma phagocytophilum* in *Ixodes ricinus* ticks in Brønnøysund in northern Norway. *Ticks Tick Borne Dis.* **4**, 218–221. (doi:10.1016/j.ttbdis.2012.11.006)
 122. Guy EC, Farquhar RG. 1991 *Borrelia burgdorferi* in urban parks. *Lancet* **338**, 253. (doi:10.1016/0140-6736(91)90392-3)
 123. Rees DHE, Axford JS. 1994 Evidence for Lyme disease in urban park workers: a potential new health hazard for city inhabitants. *Rheumatology* **33**, 123–128. (doi:10.1093/rheumatology/33.2.123)
 124. Vollmer SA *et al.* 2011 Host migration impacts on the phylogeography of Lyme Borreliosis spirochaete species in Europe. *Environ. Microbiol.* **13**, 184–192. (doi:10.1111/j.1462-2920.2010.02319.x)
 125. Nelson C, Banks S, Jeffries CL, Walker T, Logan JG. 2015 Tick abundances in South London parks and the potential risk for Lyme borreliosis to the general public. *Med. Vet. Entomol.* **29**, 448–452. (doi:10.1111/mve.12137)
 126. Sorouri R, Ramazani A, Karami A, Ranjbar R, Guy EC. 2015 Isolation and characterization of *Borrelia burgdorferi* strains from *Ixodes ricinus* ticks in the southern England. *BiolImpacts* **5**, 71–78. (doi:10.15171/bi.2015.08)
 127. Rizzoli A *et al.* 2014 *Ixodes ricinus* and its transmitted pathogens in urban and peri-urban areas in Europe: new hazards and relevance for public health. *Front. Public Health* **2**, 1–26. (doi:10.3389/fpubh.2014.00251)
 128. Uspensky I. 2014 Tick pests and vectors (Acari: Ixodoidea) in European towns: introduction, persistence and management. *Ticks Tick Borne Dis.* **5**, 41–47. (doi:10.1016/j.ttbdis.2013.07.011)
 129. LaDeau SL, Allan BF, Leisnham PT, Levy MZ. 2015 The ecological foundations of transmission potential and vector-borne disease in urban landscapes. *Funct. Ecol.* **29**, 889–901. (doi:10.1111/1365-2435.12487)
 130. Maetzel D, Maier WA, Kampen H. 2005 *Borrelia burgdorferi* infection prevalences in questing *Ixodes ricinus* ticks (Acari: Ixodidae) in urban and suburban Bonn, western Germany. *Parasitol. Res.* **95**, 5–12. (doi:10.1007/s00436-004-1240-3)
 131. Wielinga PR *et al.* 2006 Longitudinal analysis of tick densities and *Borrelia*, *Anaplasma*, and *Ehrlichia* infections of *Ixodes ricinus* ticks in different habitat areas in the Netherlands. *Appl. Environ. Microbiol.* **72**, 7594–7601. (doi:10.1128/AEM.01851-06)
 132. Cekanac R, Pavlovic N, Gledovic Z, Grgurevic A, Stajkovic N, Lepsanovic Z, Ristanovic E. 2010 Prevalence of *Borrelia burgdorferi* in *Ixodes ricinus* ticks in Belgrade area. *Vector Borne Zoonotic Dis.* **10**, 447–452. (doi:10.1089/vbz.2009.0139)
 133. Reye AL, Hübschen JM, Sausy A, Muller CP. 2010 Prevalence and seasonality of tick-borne pathogens in questing *Ixodes ricinus* ticks from Luxembourg. *Appl. Environ. Microbiol.* **76**, 2923–2931. (doi:10.1128/AEM.03061-09)
 134. Reis C, Cote M, Paul REL, Bonnet S. 2011 Questing ticks in suburban forest are infected by at least six tick-borne pathogens. *Vector Borne Zoonotic Dis.* **11**, 907–916. (doi:10.1089/vbz.2010.0103)
 135. Corrain R, Drigo M, Fenati M, Menandro ML, Mondini A, Pasotto D, Martini M. 2012 Study on ticks and tick-borne zoonoses in public parks in Italy. *Zoonoses Public Health* **59**, 468–476. (doi:10.1111/j.1863-2378.2012.01490.x)
 136. Plch J, Basta J. 1999 Incidence of spirochetes (*Borrelia* sp.) in the tick *Ixodes ricinus* in the urban environment (capital of Prague) between 1994–1997. *Zentralbl. Bakteriol.* **289**, 79–88. (doi:10.1016/S0934-8840(99)80127-3)
 137. Pangráčová L, Derdákova M, Pekárik L, Hviščová I, Víchová B, Stanko M, Hlavatá H, Peško B. 2013 *Ixodes ricinus* abundance and its infection with the tick-borne pathogens in urban and suburban areas of Eastern Slovakia. *Parasit. Vectors* **6**, 238. (doi:10.1186/1756-3305-6-238)
 138. Kilpatrick AM, Salkeld DJ, Titcomb G, Hahn MB. 2017 Conservation of biodiversity as a strategy for improving human health and well-being. *Phil. Trans. R. Soc. B* **372**, 20160131. (doi:10.1098/rstb.2016.0131)
 139. Gottdenker NL, Streicker DG, Faust CL, Carroll CR. 2014 Anthropogenic land use change and infectious diseases: a review of the evidence. *EcoHealth* **11**, 619–632. (doi:10.1007/s10393-014-0941-z)
 140. Viana M, Mancy R, Biek R, Cleveland S, Cross PC, Lloyd-Smith JO, Haydon DT. 2014 Assembling evidence for identifying reservoirs of infection. *Trends Ecol. Evol.* **29**, 270–279. (doi:10.1016/j.tree.2014.03.002)
 141. Medlock JM, Jameson LJ. 2010 Ecological approaches to informing public-health policy and risk assessments on emerging vector-borne zoonoses. *Emerg. Health Threats J.* **33**: e1. (doi:10.3402/ehtj.v3i0.7095)
 142. Medlock JM, Vaux AGC. 2015 Impacts of the creation, expansion and management of English wetlands on mosquito presence and abundance—developing strategies for future disease mitigation. *Parasit. Vectors* **8**, 142. (doi:10.1186/s13071-015-0751-3)
 143. Medlock AJM, Vaux AGC. 2013 Colonization of UK coastal realignment sites by mosquitoes: implications for design, management, and public health. *J. Vector Ecol.* **38**, 53–62. (doi:10.1111/j.1948-7134.2013.12008.x)