



Research Article

Impacts of the invasive species *Caulerpa cylindracea* Sonder, 1845 on the algae flora of the west coast of Algeria

Benabdallah Bachir Bouiadjra[‡], Malika Ghellai[§], Mohamed Daoudi[‡], Ibrahim Elkhailil Behmene[‡], Mohammed El Amine Bachir Bouiadjra[§]

[‡] Laboratory of Animal Production Sciences and Techniques (LSTPA), University of Mostaganem, Mostaganem, Algeria
[§] University of Relizane, Relizane, Algeria

Corresponding author: Mohamed Daoudi (mohamed.daoudi@univ-mosta.dz)

Academic editor: Anne Thessen

Received: 17 Feb 2021 | Accepted: 10 May 2021 | Published: 26 May 2021

Citation: Bachir Bouiadjra B, Ghellai M, Daoudi M, Behmene IE, Bachir Bouiadjra MEA (2021) Impacts of the invasive species *Caulerpa cylindracea* Sonder, 1845 on the algae flora of the west coast of Algeria. Biodiversity Data Journal 9: e64535. <https://doi.org/10.3897/BDJ.9.e64535>

Abstract

The assessment of the impacts of the expansion of the invasive species on taxonomic diversity, the abundance and dominance of groups of algae, the presence and/or absence of species of ecological interest that may or may not be indicative of water quality well mentioned, through the installation of a 20 × 20 cm quadrat representing the minimum area. The observation stations were visited monthly, during a repetitive three-year cycle, during the spring, summer and autumn seasons, periods of maximum growth and development of the algal flora and the results suggest the following facts. The invasive alga *Caulerpa cylindracea* Sonder, 1845 tends to colonise disturbed ecosystems reflecting a reduction in native algal diversity; in fact, we note a drastic impoverishment of the invaded algal community, represented by a limited number of Macrophyte algae accompanying the invasive taxon in phytosociological surveys and a Shannon-Weaver Diversity Index (H') and Equitability reduced by 4.49 and 0.77 n the heavily affected station. The number of macroalgal species accompanying the invasive species has dropped by 52% in Salamandre. In addition, the multidimensional analysis, represented by the Hierarchical Ascendant Clustering applied to this case, confirms our results.

Keywords

algae, invasive, macrophytes, diversity, index, phytosociological, taxon

Introduction

Studies on ENI date back to the 1970s (Ojaveer et al. 2018). Since then, global research in this area has developed very rapidly (Giakoumi et al. 2019).

The Mediterranean Sea is today considered to be one of the most affected of the regional seas, due to the various anthropogenic pressures on the different ecosystems of the Mediterranean basin (Mannino et al. 2017). This favors it to be an important place for non-native species, which can become invasive leading to a loss of biodiversity and changes in ecosystem services (Brunel et al. 2013, Giakoumi 2014).

The presence of mud-flats, estuaries and coral reefs along lagoons and rocky beaches provides ideal habitat for sustainable algae growth (Premarathna et al. 2019).

The concept of invasive species introduction is applied when the naturalisation of a species outside its usual geographic range is directly or indirectly linked to a human activity (Carlton 1985). The impact of introduced species has, thus, become the second most important cause of biodiversity loss in the world (Wilson 1997), after the outright disappearance of habitats. Bright (1998) defines the impact of introduced species on the planet's biodiversity as "evolution in reverse".

Amongst these bioinvasions, we are interested in *Caulerpa cylindracea* (Sonder 1845), recently reported on the Algerian west coast and which presents a strongly invasive behaviour during its colonisation (Manconi et al. 2020). It is considered to be one of the most threatening invasions of the Mediterranean Sea (Piazzi et al. 2016) and its impacts on algal biodiversity and seagrass beds are well reported in literature (Gribben et al. 2018, Montefalcone et al. 2015).

Indeed, the algal flora of the Algerian west coast remains largely unexplored. Direct observation of the algal settlement constituting the phytobenthos and its evolution is essential for monitoring possible changes in algal biodiversity. Therefore, the purpose of this research is to:

- establish a quantitative and qualitative inventory, located in time and space, of macrophyte species inhabiting two sites located on the coast of Mostaganem (west coast of Algeria Fig. 1), one affected by *C. cylindracea*, the other spare.
- assess the impact of invasive species expansion on taxonomic and specific diversity, abundance and dominance of algal groups, presence and/or absence of species of ecological interest that may or may not be indicative of water quality.

Material and methods

The choice of harvesting stations was primarily based on the importance of the algal marine flora, the location of the stations studied shown in Fig. 1.



Figure 1. [doi](#)

The geographical location of the studied sites (Google Earth modified).

The area studied extends over 50 km of coastline, from Salamandre each to Sidi Lakhdar each (small port). Both stations were visited monthly during a repetitive cycle of three years, in spring, summer and autumn seasons, periods of maximum growth and development of the algal flora. In each station, we prospected a 50 to 100 m long and 10 to 15 m wide coastal line, with a depth of 5 to 10 m.

Salamandre Station: The Salamandre site (Fig. 1) is 5 km from the town of Mostaganem, housing a new fishing port, its length around 4000 m. Surveys conducted in sheltered, quiet areas within a rocky beach with the following coordinates: 35°55' N / 0°03' E.

Small Port Station: This zone is located 35 km east of Mostaganem (Fig. 1), the sampling stations located in places with sandy and rocky bottoms on a beach 3500 m long sheltering a clear-seeded *Posidonia oceanica* meadow, associated with macrophyte algae with geographical coordinates, 36°12' N / 0°23' E.

Sampling, sorting and conservation of material.

Sampling was performed on both soft and hard variable substrates. Sampling random, raking the rocks at each survey, three surveys per observation station and per year, according to climatological conditions and without biotope delimitation so that the sample is as representative as possible of the zone (Fig. 2).



Figure 2. [doi](#)

Quadrat used for sampling (20 × 20 cm).

Species identification

Different keys for the determination of macrophyte algal flora have been consulted see Fischer et al. 1987, Mojetta and Ghisotti 1996, Gayral 1958, Gayral 1966, Carrillo Marta Sanson 1999, Coppejans 1983, Hamel 1931, Hamel 1939, Verlaque 1990, Ballesteros et al. 2007, Boudouresque and Boudouresque 1969, Feldmann 1931, Feldmann 1933, Bliding 1968, Belhissoun 1995, allow us to draw up a catalogue which is by no means exhaustive of the species of algae inhabiting the two stations considered.

Minimum Area

The minimum area is the area where one has the best chance of finding all the species of the settlement studied to within 10% according to Gounout (1969). For our case, the minimum area adopted is 20 × 20 cm a surface of 400 cm² Fig. 2.

Analytical Parameters

For the quantitative and qualitative analysis of vegetation, we used the terrestrial phytosociological methods that several authors have adopted in the marine environment (Boudouresque 1971a, Boudouresque 1971b, Cormaci et al. 1997, Scammacca et al.

1993). The results are expressed by their average, with data analysis carried out by the Statistica software (version 7.0) for all the parameters retained.

Recovery

The recovery (R_i) is the approximate percentage of the substrate surface covered in projection by species i . Given the stratification of vegetation, the total coverage of a survey $\sum R_i$ is generally greater than 100%. The importance of recovery is expressed in class according to the following scale:

+ = negligible recovery

1 = less than 5% of the surface is covered

2 = between 5 and 25% of the surface area

3 = between 25 and 50% of the surface area

4 = between 50% and 75% of the surface area

5 = more than 75% of the surface area

Overall average recovery

Each class of the recovery coefficient R_i is assigned a conventional monthly average value (class centre) called average recovery.

Absence = 0; + = 01%; 1 = 2.5%; 2 = 15%; 3 = 37.5%; 4 = 62.5%; 5 = 87.5%

The RMG (overall average recovery) of species i in a set of N recorded is the average of these successive average recoveries.

$$RMG = \sum R_i / N$$

The RMG of a subset E of n species (ecological group, systematic unit or phytogeographic elements) is the sum of the constituent species.

Species richness (Q).

The Q coefficient is the specific population size of any subset in a survey (ecological group, systematic unit or biogeographic features) and the Q (highlighted) of a group of species representing its average species size.

Quantitative dominance.

The dominance based on recovery ($\sum DR_i$) of a group of species in a survey (in a survey table), is the ratio, expressed as a percentage of the sum of their recovery (or their RMG) to the total recovery of species in the survey (or to the total average recovery of species in the survey table).

Qualitative dominance (DQ)

This is the ratio expressed as a percentage of Q, (where Q highlighted) to T (where T highlighted).

Index of Specific Diversity (Shannon-Wiener)

This index (H') measures, in a settlement, the amount of information resulting from species differentiation. The H' value reflects the degree of structural evolution, maturity and stability of the ecosystem under consideration. It was calculated from the dominance of each species (R_i/R_t) according to Shannon's formula (in Boudouresque and Cinelli 1971).

$$H' = - \sum P_i * \text{Log}_2 P_i \text{ with } P_i = R_i/R_t \text{ } R_t = \text{total recovery}$$

The calculations were made from Suppl. materials 1, 2 under R software version 3.0.3 R Core Team 2014

Evenness or Regularity E

This is the ratio between the actual community diversity index and its maximum value for the number of species present $H'_{\max} = \text{Log}_2 T$ with T = total number of taxa

$$E = H'/H'_{\max}$$

The calculations were made from Suppl. materials 1, 2 under R software version 3.0.3 R Core Team 2014

Taxonomic diversity

Three taxonomic indices have also been adopted for the study of taxonomic diversity (Δ , Δ^* and Δ^+). These were calculated from a taxonomic tree constructed on the basis of phylogenetic classification (Clarke and Warwick 1998), considering eight taxonomic levels: species, genus, family, order, class, phylum, under-reign and reign. From this tree, a taxonomic distance w_{ij} is quantified between each pair of species, assuming that this distance is equal to 100 for two species related to the highest taxonomic level considered in the study (Clarke and Warwick 1999). This calculated distance (w_{ij}) will then be integrated into the calculation of the various taxonomic indices. The calculations were performed using Suppl. materials 3, 1, 2 software R version 3.0.3 (R Core Team 2014). The phylogenetic classification is based on the Worms Editorial Board (Guiry and Guiry 2020).

With x_i ($i = 1, \dots, S$): the abundance of the i^{th} species, N ($= \sum x_i$): the total number of individuals in the sample and w_{ij} : the distance to be covered between species i and the first common node with species j in the hierarchical classification:

- Δ "Taxonomic diversity" (Warwick and Clarke 1995):

Δ represents the average taxonomic distance between each pair of randomly selected individuals in the sample.

- Δ^* "**Taxonomic Distinctness**" (Warwick and Clarke 1995):

Δ^* is the average taxonomic distance between two randomly selected individuals belonging only to different species.

- Δ^+ "**Average Taxonomic Distinctness**", on data in presence/absence, Δ and Δ^* are simplified by Δ^+ (Clarke and Warwick 1998):

Δ^+ measures the average taxonomic distance between two randomly selected species.

- **Funnel Test:**

In order to detect assemblages whose taxonomic diversity would be influenced by disturbances, Warwick and Clarke (1998) proposed the "Funnel" test. This test consists comparing the taxonomic variety of a site to the area to which it belongs, from 1000 random draws of size m species made on the global list of S species listed in the area under consideration.

Hierarchical Ascendant Clustering (HAC)

The hierarchical classification has been used in a wide variety of disciplines. It has been described by several authors including Daget 1979, Dagnelie 1986, Legendre and Legendre 1984, Legendre and Legendre 1998, Herrera and Legac 2002 and Bouroche and Saporta 2005. Our study was confirmed by the use of this multidimensional analysis which consists in partitioning the objects (or descriptors) of the study into groups and subgroups, passing through the condensation of the information provided through the data matrix into a similarity or distance matrix. The latter will then be classified according to different classification algorithms (Ward, single, complete, average), which will each provide a corresponding dendrogram. The calculation procedure for this analysis was performed using Suppl. materials 1, 2 under software R version 3.0.3 (R Core Team 2014).

Results

Results are reported in Table 1 and Table 3 for algal stand composition at the Salamandre and Small Port Stations respectively and in Tables 2, 4 for analytical parameters (RMG, DRi, Q DQ) at the same stations. Table 5 represents the set of taxonomic indices at the two study stations and Fig. 3 illustrates the Funnel Test. The Hierarchical Ascendant Classification of the study stations based on the algae stands surveyed is illustrated in Fig. 4.

Table 1.

Algal settlement sheet: Salamandre Station.

Statement number	1	2	3	RMG	Pr
Month	Mar.	Nov.	June		
Surface in cm ²	400	400	400		
Cover	100%	90%	90%		
Exposition	N	NE	NW		
<i>Erythrotrichia carnea</i>	+	+	+	0.10	3
<i>Acrochaetium cheminii</i>	0	+	+	0.07	2
<i>Colaconema daviesii</i>	+	+	+	0.10	3
<i>Asparagopsis armata</i>	2	1	1	6.67	3
<i>Falkenbergia rufolanosa</i>	1	1	1	2.50	3
<i>Dasya rigidula</i>	+	+	+	0.10	3
<i>Taenio mananum</i>	1	1	0	1.67	2
<i>Chondria coerulescens</i>	0	+	+	0.07	2
<i>Chondriamairei</i> + <i>C. capillaris</i>	+	+	+	0.10	3
<i>Digenea simplex</i>	1	1	+	1.70	3
<i>Halopithyin curvus</i>	0	1	1	1.67	2
<i>Herposiphonia secunda</i> f. <i>secunda</i>	0	0	+	0.03	1
<i>Herposiphonia secunda</i> f. <i>tenella</i>	0	+	0	0.03	1
<i>Lophocladia lallemandii</i>	+	+	0	0.07	2
<i>Vertebrata fruticulosa</i>	+	+	+	0.10	3
<i>Vertebrata furcellata</i>	0	+	+	0.07	2
<i>Polysiphonia spinulosa</i>	0	0	+	0.03	1
<i>Amphiroa rigida</i>	2	1	+	5.87	3
<i>Ellisolandia elongata</i>	2	2	2	15.00	3
<i>Corallina officinalis</i>	0	2	1	5.83	2
<i>Jania rubens</i>	+	1	1	1.70	3
<i>Hypnea musciformis</i>	1	1	1	2.50	3
<i>Grateloupia filicina</i>	0	+	+	0.07	2
<i>Gastroclonium clavatum</i>	0	+	+	0.07	2
<i>Dictyopteris polypodioides</i>	+	+	1	0.90	3
<i>Dictyopteris divaricata</i>	+	1	1	1.70	3
<i>Asperococcus bullosus</i>	0	0	+	0.03	1
<i>Feldmannia globifera</i>	0	+	+	0.07	2
<i>Colpomenia peregrina</i>	+	+	+	0.10	3

Statement number	1	2	3	RMG	Pr
Month	Mar.	Nov.	June		
Surface in cm ²	400	400	400		
Cover	100%	90%	90%		
Exposition	N	NE	NW		
<i>Colpomenia sinuosa</i>	+	+	0	0.07	2
<i>Bryopsis hypnoides</i>	0	0	+	0.03	1
<i>Bryopsis muscosa</i>	0	+	0	0.03	1
<i>Bryopsis plumosa</i>	+	+	0	0.07	2
<i>Bryopsis secunda</i>	+	+	0	0.07	2
<i>Caulerpa prolifera</i>	1	1	1	2.50	3
<i>Caulerpa racemosa</i>	2	2	1	10.83	3
<i>Codium bursa</i>	0	1	0	0.83	1
<i>Codium effusum</i>	1	1	0	1.67	2
<i>Codium fragile</i>	2	1	+	5.87	3
<i>Codium tomentosum</i>	0	+	1	0.87	2
<i>Chaetomorpha aerea</i>	1	1	1	2.50	3
<i>Chaetomorpha mediterranea</i>	0	1	1	1.67	2
<i>Chaetomorpha linum</i>	0	1	1	1.67	2
<i>Cladophora coelothrix</i>	0	1	0	0.83	1
<i>Cladophora laetevirens</i>	1	0	0	0.83	1
<i>Cladophora prolifera</i>	1	1	1	2.50	3
<i>Cladophora coelothrix</i>	+	+	+	0.10	3
<i>Cladophora rupestris</i>	0	0	1	0.83	1
<i>Bldingia marginata</i>	+	+	+	0.10	3
<i>Ulva compressa</i>	1	1	1	2.50	3
<i>Ulva intestinalis</i>	2	1	1	6.67	3
<i>Ulva linza</i>	+	+	+	0.10	3
<i>Ulva prolifera</i>	+	+	+	0.10	3
<i>Ulva clathrata</i>	0	0	+	0.03	1
<i>Ulva fasciata</i>	+	+	+	0.10	3
<i>Ulva lactuca</i>	1	1	1	2.50	3
<i>Ulva rigida</i>	2	1	1	6.67	3
Number of species per statement	34	49	46	101.79	
Average number of species per statement	43			Pr: the presence of the species	
R/P per statement	3.5	4.4	4.2		
R/P average	4.03				

Statement number	1	2	3	RMG	Pr
Month	Mar.	Nov.	June		
Surface in cm ²	400	400	400		
Cover	100%	90%	90%		
Exposition	N	NE	NW		
Diversity index H'	4.49				
Evenness E	0.77				

Table 2.
Analytical parameters at the Salamandre Station.

	ΣRMG	Dri	Q	DQ
Chlorophyceae	53.91%	52.96%	19.66	46.08%
Phaeophyceae	2.85%	2.79%	4	9.37%
Rhodophyceae	45.11%	44.31%	19	44.53%
Total	101.87%			

Table 3.
Algal settlement sheet: Small Port Station.

Statement number	1	2	3	RMG	Pr	Statement number	1	2	3	RMG	Pr
Month	Mar.	Nov.	June			Date	Mar.	Nov.	June		
Surface in cm ²	400	400	400			Surface in cm ²	400	400	400		
Cover	100%	90%	90%			Cover	100%	90%	90%		
Exposition	N	NE	NW			Exposition	N	NE	NW		
<i>Porphyra umbilicalis</i>	+	0	0			0.03	1	<i>Cladosiphon mediterraneu</i>	+		
<i>Pyropia leucosticta</i>	+	0	+	0.07	2	<i>Myriactula graciliariae</i>	0	0	+	0.03	1
<i>Sahlingia subintegra</i>	+	+	0	0.07	2	<i>Myriactula rigida</i>	0	0	+	0.03	1
<i>Antithamnion amphigeneum</i> A. J. K. Millar	0	0	+	0.03	1	<i>Myriactula rivulariae</i>	0	0	+	0.03	1
<i>Asparagopsis armata</i>	2	1	0	5.83	2	<i>Ectocarpus fasciculatus</i> var. <i>fasciculatus</i>	+	0	+	0.07	2
<i>Falkenbergia rufolanosa</i>	1	1	0	1.67	2	<i>Ectocarpus commensalis</i>	0	0	+	0.03	1
<i>Anotrichium tenue</i>	+	+	0	0.07	2	<i>Ectocarpus siliculosus</i>	+	0	0	0.03	1
<i>Centrocerca clavulatum</i>	+	0	+	0.07	2	<i>Feldmannia globifera</i>	+	+	0	0.07	2

Statement number	1	2	3	RMG	Pr	Statement number	1	2	3	RMG	Pr
Month	Mar.	Nov.	June			Date	Mar.	Nov.	June		
Surface in cm ²	400	400	400			Surface in cm ²	400	400	400		
Cover	100%	90%	90%			Cover	100%	90%	90%		
Exposition	N	NE	NW			Exposition	N	NE	NW		
<i>Corallophila cinnabarina</i>	+	+	+	0.10	3	<i>Feldmannia simplex</i>	+	+	+	0.10	3
<i>Ceramium diaphanum</i>	2	1	+	5.87	3	<i>Feldmannia mitchelliae</i>	1	+	0	0.87	2
<i>Spyridia filamentosa</i>	0	+	+	0.07	2	<i>Hinckia sandriana</i>	1	+	0	0.87	2
<i>Dasya rigidula</i>	+	+	0	0.07	2	<i>Ralfsia verrucosa</i>	1	1	0	1.67	2
<i>Taenio mananum</i>	+	+	0	0.07	2	<i>Cystoseira algeriensis</i>	1	+	1	1.70	3
<i>Chondria coerulescens</i>	+	+	+	0.10	3	<i>Cystoseira barbata</i>	1	+	+	0.90	3
<i>Chondria dasyphylla</i>	+	+	0	0.07	2	<i>Cystoseira compressa</i>	1	+	1	1.70	3
<i>Chondria mairei</i>	+	+	+	0.10	3	<i>Cystoseira brachycarpa</i> var. <i>balearica</i>	1	1	1	2.50	3
<i>Digenea simplex</i>	1	1	0	1.67	2	<i>Cystoseira crinita</i>	0	0	2	5.00	1
<i>Halopithys incurva</i>	1	1	1	2.50	3	<i>Cystoseira sedoides</i>	1	1	1	2.50	3
<i>Herposiphonia secunda</i> f. <i>secunda</i>	+	0	+	0.07	2	<i>Cystoseira mediterranea</i>	1	+	1	1.70	3
<i>Herposiphonia secunda</i> f. <i>tenella</i>	+	0	+	0.07	2	<i>Cystoseira amentacea</i> var. <i>stricta</i>	1	+	1	1.70	3
<i>Laurencia microcladia</i>	1	+	1	1.70	3	<i>Cystoseira tamariscifolia</i>	1	1	1	2.50	3
<i>Laurencia obtusa</i>	0	1	1	1.67	2	<i>Sargassum acinarium</i>	0	0	1	0.83	1
<i>Palisada perforata</i>	+	1	1	1.70	3	<i>Sargassum vulgare</i>	1	0	+	0.87	2
<i>Lophocladia lallemandii</i>	+	0	0	0.03	1	<i>Colpomenia peregrine</i>	1	1	1	2.50	3
<i>Ellisolandia elongata</i>	2	1	+	5.87	3	<i>Colpomenia sinuosa</i>	1	1	1	2.50	3
<i>Corallina officinalis</i>	1	1	1	2.50	3	<i>Cladostephus spongiosus</i>	1	+	0	0.87	2
<i>Jania longifurca</i>	1	+	+	0.90	3	<i>Sphacelaria cirrosa</i>	+	0	+	0.07	2
<i>Jania rubens</i>	1	+	+	0.90	3	<i>Sphacelaria plumula</i>	+	0	+	0.07	2
<i>Gelidiella ramellosa</i>	+	0	1	0.87	2	<i>Halopteris scoparia</i>	0	1	1	1.67	2
<i>Gelidium crinale</i>	+	0	1	0.87	2	<i>Bryopsis duplex</i>	+	+	0	0.07	2
<i>Gelidium spinosum</i>	0	0	1	0.83	1	<i>Bryopsis hypnoides</i>	+	+	0	0.07	2
<i>Gelidium corneum</i>	+	0	0	0.03	1	<i>Bryopsis muscosa</i>	+	0	0	0.03	1
<i>Pterocladia capillacea</i>	+	0	0	0.03	1	<i>Bryopsis plumosa</i>	+	0	0	0.03	1

Statement number	1	2	3	RMG	Pr	Statement number	1	2	3	RMG	Pr
Month	Mar.	Nov.	June			Date	Mar.	Nov.	June		
Surface in cm ²	400	400	400			Surface in cm ²	400	400	400		
Cover	100%	90%	90%			Cover	100%	90%	90%		
Exposition	N	NE	NW			Exposition	N	NE	NW		
<i>Chondracanthus acicularis</i>	1	1	+	1.70	3	<i>Bryopsis secunda</i>	+	+	0	0.07	2
<i>Chondracanthus teedei</i>	0	1	+	0.87	2	<i>Caulerpa prolifera</i>	1	+	1	1.70	3
<i>Rissoella verruculosa</i>	+	+	0	0.07	2	<i>Codium bursa</i>	0	1	1	1.67	2
<i>Gracilaria bursa-pasoris</i>	1	+	+	0.90	3	<i>Codium effusum</i>	1	+	1	1.70	3
<i>Gracilariopsis longissima</i>	+	+	0	0.07	2	<i>Codium fragile</i>	1	0	1	1.67	2
<i>Hypnea musciformis</i>	2	2	1	10.83	3	<i>Codium tomentosum</i>	0	1	1	1.67	2
<i>Grateloupia filicina</i>	+	0	+	0.07	2	<i>Chaetomorpha aerea</i>	1	1	2	6.67	3
<i>Peyssonnelia polymorpha</i>	+	0	+	0.07	2	<i>Chaetomorpha mediterranea</i>	1	1	1	2.50	3
<i>Peyssonnelia rubra</i>	0	0	+	0.03	1	<i>Cladophora coelothrix</i>	0	1	+	0.87	2
<i>Peyssonnelia squamaria</i>	+	0	0	0.03	1	<i>Cladophora laetevirens</i>	1	1	1	2.50	3
<i>Sphaerococcus coronopifolius</i>	0	+	0	0.03	1	<i>Cladophora prolifera</i>	+	0	0	0.03	1
<i>Nemalion helminthoides</i>	+	+	0	0.07	2	<i>Acetabularia acetabulum</i>	+	0	0	0.03	1
<i>Liagora viscida</i>	+	0	1	0.87	2	<i>Bldingia marginata</i>	+	+	+	0.10	3
<i>Gastroclonium clavatum</i>	+	0	0	0.03	1	<i>Ulva compressa</i>	1	1	1	2.50	3
<i>Irvinea boergesenii</i>	+	0	0	0.03	1	<i>Ulva intestinalis</i>	+	0	0	0.03	1
<i>Cutleria adspersa</i>	0	0	+	0.03	1	<i>Ulva linza</i>	+	+	0	0.07	2
<i>Dictyota dichotoma</i>	1	1	0	1.67	2	<i>Ulva prolifera</i>	+	1	1	1.70	3
<i>Dictyota fasciola</i>	1	+	0	0.87	2	<i>Ulva clathrata</i>	2	1	1	6.67	3
<i>Dictyopteris polypodioides</i>	2	0	1	5.83	2	<i>Ulva fasciata</i>	+	+	+	0.10	3
<i>Dictyota spiralis</i>	0	+	1	0.87	2	<i>Ulva lactuca</i>	1	1	1	2.50	3
<i>Padina pavonica</i>	0	0	+	0.03	1	<i>Ulva rigida</i>	2	1	1	6.67	3
<i>Asperococcus bullosus</i>	+	0	+	0.07	2						
						Number of species per statement	89	68	72	130.4	

Statement number	1	2	3	RMG	Pr	Statement number	1	2	3	RMG	Pr
Month	Mar.	Nov.	June			Date	Mar.	Nov.	June		
Surface in cm ²	400	400	400	Surface in cm ²	400	400	400				
Cover	100%	90%	90%	Cover	100%	90%	90%				
Exposition	N	NE	NW	Exposition	N	NE	NW				
				Average number of species per statement	76.33			Pr: the presence of the species			
				R/P per statement	1.61	1.45	1.07				
				R/P Average	1.37						
				Diversity index H'	5.56						
				Evenness E	0.82						




Figure 3. [doi](#)

Funnel test.

Table 4.
Analytical parameters at the Small Port Station.

	ΣRMG	Dri	Q	DQ
Chlorophyceae	31.69%	24.30%	19	25.11%
Phaeophyceae	42.61%	32.67%	24	31.72%
Rhodophyceae	56.10%	43.02%	32.66	43.17%
Total	130.40%			

Table 5.
Taxonomic diversity indices.

Region	S	delta	delta*	delta+
Small port	109	79.98	81.80	83.99
Salamandre	57	72.163	76.26	77.84
Expected		160.61	78.59	83.37

The results shown in Table 1 and Table 3, with illustrations and descriptions of sample specimens for the two observation stations are contained in a doctoral thesis (Bachir Bouiadjra 2012).




Figure 4. [doi](#)

Classification of phytosociological surveys carried out at the Small Port and the Salamandre Station by the average link.

Discussion

Salamandre Station

The reading of the sheet on the algal settlement of the Salamandre site (Table 1) and the result of the structural parameters shown in Table 2, indicate a quantitative and qualitative dominance, respectively of 52% and 46% with an overall average recovery exceeding 50% of the species belonging to Chlorophyceae. Within this systematic unit, we note an overall average recovery rate of five invasive taxa, *C. cylindracea*, *C. fragile*, *A. armata*, *F. rufolanosa* *L. lallemandii*, which is of the order of 24.5%, 10.5% of which is reserved for the *C. cylindracea*, which is progressing in some surveys by joining its small colonies (Figs 5, 6) to which are added the eutrophication indicator species of the environment and nitrophilic strongly accumulating nitrogen are represented by the order of Ulvales in the number of 8 species (Table 1) with a global average cover rate of 18% on a total global

cover of 50% of Chlorophyceae in the Salamandre Station. This high presence rate is an indicator of a polluted environment which is confirmed by domestic and urban discharges (Khiari et al. 2017, Damak et al. 2019) duly observed at Salamandre Station, in addition the presence rate of the invasive species *C. cylindracea* amount to 10.83% (Table 1) which confirms an increasingly advanced colonisation.



Figure 5. [doi](#)

Central colonies of *C. cylindracea* at the Salamandre Station.



Figure 6. [doi](#)

Dense colonies of *Caulerpa cylindracea* in the heavily affected station of Salamandre.

The algal population of Phaeophyceae occupies a negligible average cover rate of 2.85% at the Salamandre Station and reveals the total absence of macrophyte species sensitive to pollution by detergents and water turbidity such as cystoseires (*Cystoseira stricta*, *C. algeriensis*, *C. crinita* ... etc) (Gamulin-Brida et al. 1967, Bellan-Santini 1966, Verlaque and Fritayre 1994a) which confirms that this site is strongly impacted by *C. cylindracea* and disturbed, given the strong presence of the invasive species. On the other hand the site of the small harbor displays a rate of recovery of Phaeophyceae macrophytes of 42.61% (Table 4) with a presence of nine species of Cystoseires suggesting a good water quality and an average algal specific richness of 76 species (Table 3) per survey contrary to the

previous site, which displays an average number of species of only 43 per survey (Table 1).

As for Rhodophyceae the overall cover rate is respectively 45.11% and 56.10% at the Salamandre and small port Station with a qualitative dominance of 44% and 43% (Tables 2, 4) which suggests protection and valorisation of some taxa rich in mineral elements and chemical components (Sambusiti et al. 2015, Cardoso et al. 2014, Shobana et al. 2017) see *Ellisolandia elongata*, *Hypnea musciformis*, *Gelidium crinale*, *Gelidium spinosum*, *Gelidium corneum*, *Jania longifurca*, *Jania rubens* and *Gracilaria bursa-pasoris*.

This is easily verified in the number of species per phytosociological survey, which is much lower than those reported in the Small Port site and the reduction in the Diversity and Evenness Indices, respectively by 4.49 and 0.77 at the Salamandre Station, in relation to the results obtained at the Small Port. In this station invaded by *C. cylindracea*, a native species, *Caulerpa prolifera* (Forsskal) Lamouroux are recorded in surveys, with a low recovery rate of 2.5% compared to that of the invasive taxon (10.5%) suggesting a tendency for *C. prolifera* to be replaced by the invasive species (Piazzi and Ceccherelli 2006). Over time, according to Ceccherelli et al. 2001, Jaubert et al. 2003), the homogenisation of the underwater landscape observed in some Mediterranean regions heavily affected by the invasive species has resulted. It is important to remember that for the four other invasive species of Rhodophyceae inventoried and subservient to the photophilic infralittoral of hard substrates, seem to find favorable conditions for their development, without any threat to the ecosystem. Nevertheless, the *Bonnemaisonnea* of the genus *Asparagopsis* found in Salamandre, has major assets its avoidance by herbivorous fish because secreting secondary metabolites and its ability to multiply vegetatively and by apomeiotic spores (Feldmann 1954) hence its expansion to the site of Salamandre with a recovery rate of 1%, thus forming a settlement in *Asparagopsis*.

In addition, the beach of Salamandre with coastal developments operating in recent years is a vulnerable site to closely monitor to avoid erosion of its algal richness due to an expansion of invasive taxon *C. cylindracea*.

Small Port Station

It is a site rich in species, 109 taxa listed (Table 3), 80% of the total number recorded on the coast of Mostaganem (Bachir Bouiadjra 2012). The recovery rate of the entire algal flora exceeds 100%, with a Diversity Index (H') of 5.56 which is higher than that of the previous station (Salamandre), as well as that of Bordj El kifan (1.62) east of Algiers reported by Serridi (2003). The Evenness Index is about 0.82, which suggests, according to Daget (1979), a balanced algal stand.

The overall average overlap among Rhodophyceae is clearly dominant at 56%, the same observation is made for their quantitative and qualitative dominance estimated at 43%, followed by the Phaeophyceae group with an RMG of 42% and dominance (Dri) and (DQ) of 32% and finally the Chlorophyceae group with an RMG of 31% (Table 4). This represents a well-structured settlement with species constituting the crusty, grassy, erect

and epiphytic strata according to the algal settlement sheet (Table 3). Note the absence of the invasive taxon *C. cylindracea* however, the fishing port adjacent to the observation sites, will constitute a potential vector for caulerpe transmission in the area. Hence the need for increased vigilance and follow-up with awareness campaigns among fishing professionals to avoid contamination of the site.

Taxonomic Diversity Indices

Analysis of the different taxonomic Diversity Indices (Table 5) indicates a higher taxonomic diversity at the Small Port than at the Salamandre Station. This means that the species recorded at the Small Port are more taxonomically distant than those found in Salamandre. In other words, the average taxonomic distance between each pair of randomly selected individuals is greater at the Small Port than at Salamandre.

The Funnel Test (Fig. 3) based on the values of $\Delta+$ obtained for each Station, in comparison with the mean expected value in the study area, shows that the Salamandre Station is outside the 95% confidence interval of $\Delta+m$, indicating a low taxonomic diversity compared to that expected in the area. This suggests that this site is subject to anthropogenic disturbances, influencing the taxonomic composition of the settlement by generating a poor assemblage of species, in addition to the presence of invasive species in this plant, associated with untreated wastewater discharges, confirm this reduction in algal diversity (Verlaque and Fritayre 1994b, Ceccherelli et al. 2001, Piazzzi et al. 2001, Piazzzi and Cinelli 2003, Piazzzi and Ceccherelli 2006).

Hierarchical Ascendant Clustering

Our study was confirmed by a multidimensional analysis which is the Hierarchical Ascendant Clustering (HAC). The latter (Fig. 4) shows a clear separation between the two sites considered by indicating four distinct groups of phytosociological surveys. The first two, represented by the summer and spring surveys at the Small Port Station, are the most dissimilar of the other groups and represent the greatest distance. The two surveys, therefore, represent the most species-rich removal recorded, indicating 89 and 72 species recorded respectively during the spring and summer seasons at the Small Port. However, we note a decrease in the number of species in the surveys conducted during the winter season at the same site with 68 species recorded, which results in a smaller distance in the classification tree for this survey. The last group consists of the surveys carried out in the Salamandre Station and represents the smallest distance in the dendrogram due to the low number of species recorded due to the dominance of the settlement by species belonging to Chlorophytes, neutrophils associated with five other invasive species including *C. cylindracea*, *C. fragile*, *A. armata*, *F. rufolanosa* and *L. lallemandii*, which testifies to the disturbance and degradation of the environment in the Salamandre Station.

Conclusions

The invasive alga *Caulerpa cylindracea* Sonder, 1845 tends to colonise disturbed ecosystems reflecting a reduction in indigenous algal diversity (Ceccherelli et al. 2001, Piazzzi et al. 2001, Piazzzi and Cinelli 2003, Piazzzi and Ceccherelli 2006). This is confirmed in the heavily colonised observation site of Salamandre Beach, through the statistical analysis tools used to see the parameters of abundance recovery, qualitative and quantitative dominance of algal groups, Equitability Index and taxonomic diversity and where there is a drastic depletion of the invaded algal community, represented by a limited number of Macrophyte algae accompanying the invasive taxon in phytosociological surveys and a low Shannon-Weaver Diversity Index (H) of 4.49. On the other hand, the Station of the Small Port not affected by *C. cylindracea* hosts a population exceeding in some surveys about twenty taxa or the Diversity Index is 5.56 suggesting a well-structured and stable algal population. The number of macroalgal species accompanying the invasive species has dropped by 52% in Salamandre. Verlaque and Fritayre (1994b) attribute this decrease to a rate ranging from 25 to 55%. This is consistent with our results. Furthermore, the analysis of the floral procession mentioned in the surveys of the Salamandre Station indicates a decrease in the cover of erect shrub algae (-70 to -80%) and epiphytic and filamentous algae in the autumn season, while encrusting algae of the genus *Corallina* dominate and best resist the occupation of the ecological niche by the invasive species *C. cylindracea*. This study provides an inventory of the algal population associated with the recently introduced *C. cylindracea* at the Mostaganem coast, which needs to be monitored at different levels of the affected coastline in order to effectively limit its expansion to the detriment of existing algal diversity.

Acknowledgements

The authors would like to thank the Directorate General of Scientific Research and Technological Development for its accompaniment of the Projects for University Research and Formation.

References

- Bachir Bouiadjra B (2012) Étude de la flore algale benthique et les impacts de ses espèces invasives devant la côte Mostaganemoise. Thèse de doctorat. Université Abdelhamid Ibn Badis de Mostaganem, Algérie, 143 pp. URL: <http://e-biblio.univ-mosta.dz/handle/123456789/1092>
- Ballesteros E, Toras X, Pinedo S, Garcia M, Mangialajo L, De Tores M (2007) A new methodology based on littoral community cartography dominated by macroalgae for the implementation of the European Water framework directive. Marine Pollution Bulletin 55 (2007): 172-180. <https://doi.org/10.1016/j.marpolbul.2006.08.038>
- Belhissoun S (1995) Contribution à l'étude Phyto-écologique du Phytobenthos Marin de la région de Rabat- Mohammedia (Atlantique Marocain). Thèse du diplôme d'études

supérieures de 3ème Cycle. Université Mohammed V. Faculté des Sciences.
Rabat, Maroc.

- Bellan-Santini D (1966) influences des eaux usées sur la faune et la flore marine benthique dans la région Marseillaise. [Techniques et sciences municipales](#) 61 (7): 285-292.
- Bliding C (1968) A critical survey of European taxa in Ulvales, II: Ulva, Ulvaria, Monostroma, Kormania. *Botaniska Notiser* 121: 535-629.
- Boudouresque CF, Boudouresque E (1969) Contribution à la flore des Algues de l'Algérie. *Bulletin du Musée d'histoire naturelle de Marseille*. Fr 29: 19-36.
- Boudouresque CF (1971a) Méthodes d'étude qualitative et quantitative du benthos (en particulier le phytobenthos). *Téthys*, Fr 3 (1): 79-104.
- Boudouresque CF (1971b) Recherche de bionomie analytique, structurale et expérimentale sur les peuplements benthiques sciaphiles de Méditerranée occidentale (fraction algale). La sous-stratesciaphile des peuplements de grande *Cystoseira* de mode battu. *Bulletin du Musée d'Histoire Naturelle de Marseille*, Fr 31: 141-151.
- Boudouresque CF, Cinelli F (1971) The algae population of superficial biotopes sciaphiles of the kind found in the island of Ischia (Naples Gulf, Italy). *Pubblicazioni della Stazione Zoologica Napoli, Italy* 39: 1-43.
- Bourouche JM, Saporta G (2005) L'analyse des données. 9ème edition. Presses Universitaires de France, Paris, 127 pp.
- Bright C (1998) Life out of bounds. Bioinvasion in a Borderless World. *Worlworth Institute: Norton Publication*.
- Brunel S, Fernandez-Galiano E, Genovesi P, Heywood VH, Kueffer C, Richardson DM (2013) Invasive alien species: A growing but neglected reat? In late lessons from early warning: Science, precaution, innovation. *essons for preventing harm*. EEA Report 1/2013, Copenhagen 518-540.
- Cardoso S, Carvalho L, Silva P, Rodrigues M, Pereira O, Pereira L (2014) Bioproducts from Seaweeds: A Review with Special Focus on the Iberian Peninsula. *Current Organic Chemistry* 18 (7): 896-917. <https://doi.org/10.2174/138527281807140515154116>
- Carlton J, (1985) transoceanic and antarctic dispersal of coastal marine organisms: the biology of ballast water. *Oceanography and marine biology* 3: 265-273.
- Carrillo Marta Sanson JA (1999) Algas, Hongros y fanerogamas marinas de Las Islas Canarias. Clave analítica. Service de publicaciones. Universidad de la Laguna, 143 pp.
- Ceccherelli G, Piazzini I, Cinelli F, Gravez V, Ruitton S, Boudouresque CF, Le Direac'h L, Meinesz A, Scabbia G, Verlaque M (2001) The development of *Caulerpa racemosa* at the margin of *Posidonia oceanica*. Fourth international workshop on *Caulerpa taxifolia* edition, GIS Posidonie Publication. 376-384.
- Clarke KR, Warwick RM (1998) A taxonomic distinctness index and its statistical properties. *Journal of Applied Ecology* 35: 523-531. <https://doi.org/10.1046/j.1365-2664.1998.3540523.x>
- Clarke KR, Warwick RM (1999) The taxonomic distinctness measure of biodiversity: weighting of step lengths between hierarchical levels. *Marine Ecology Progress Series* 184: 21-29. <https://doi.org/10.3354/meps184021>

- Coppejans E (1983) Iconographie d'Algues Méditerranéennes. Chlorophyta, Phaeophyta, Rhodophyta. Bibliothecaphycologica, Herausgegeben von J. Cramer, 317 pp.
- Cormaci M, Lanfranco E, Borg JA, Buttigieg S, Furnari G, Micallef SA, Mifsud C, Pizzuto F, Scammacca B, Serio D (1997) Contribution to the Knowledge of benthic marine algae on rocky substrata of the Maltese Islands (Mediterranean Sea). *Botanica Marina*. 40: 203-215.
- Daget J (1979) Les modèles mathématiques en écologie. 8ème ed. Collection d'écologie. Masson, 172 pp.
- Dagnelie P (1986) Analyse statistique à plusieurs variables. 3 ième réimpression. Les presses agronomiques de Gembloux, Gembloux, 362 pp.
- Damak M, Frontalini F, Elleuch B, Kallel M (2019) Benthic foraminiferal assemblages as pollution proxies along the coastal fringe of the Monastir Bay (Tunisia). *Journal of African Earth Sciences* 150: 379-388. <https://doi.org/10.1016/j.jafrearsci.2018.11.013>
- Feldmann J (1931) Contribution à la flore Algologique Marine de l'Algérie. Les Algues de Cherchell. *Bulletin de la Société d'Histoire Naturelle de l'Afrique du Nord* Alger 22: 229-254.
- Feldmann J (1933) Contribution à la flore Algologique Marine de l'Algérie. *Bulletin de la Société d'Histoire Naturelle de l'Afrique du Nord* Alger 24 (Fascicule 2): 360-366.
- Feldmann J (1954) Inventaire de la flore marine de Roscoff (algues, champignons, lichens et spermatophytes). *Travaux de la Station Biologique de Roscoff*. Fr 6 (suppl.): 1-252.
- Fischer W, Schneider M, Bauchot ML (1987) Méditerranée et Mer Noire, Zone de pêche 37 Révision 1, volume 1, Végétaux et Invertébrés. Fiches d'identification, 760 pp.
- Gamulin-Brida H, Giaccone G, Golubić S (1967) Contribution aux études des biocoenoses subtidales. *Helgoländer Wissenschaftliche Meeresuntersuchungen* 15: 429-444. <https://doi.org/10.1007/bf01618639>
- Gayral P (1958) Algues de la côte Atlantique Marocaine. La nature au Maroc II, Casablanca, 523 pp.
- Gayral P (1966) Les Algues des Côtes Françaises (Manche et Atlantique). *Notions Fondamentales sur l'écologie, la biologie et la systématique des Algues Marines*. Edition Doin- Deren & Cie, 632 pp.
- Giakoumi S (2014) Distribution patterns of the invasive herbivore *Siganus luridus* (Ruppell, 1829) and its relation to native benthic communities in the Central Aegean Sea, Northeastern Mediterranean. *Marine Ecology* 35 (12059): 96-105. <https://doi.org/10.1111/maec.12059>
- Giakoumi S, Katsanevakis S, Albano PG, Azzurro E, Cardoso AC, Cebrian E (2019) Management priorities for marine invasive species. *Science of the Total Environment* 688: 976-982. <https://doi.org/10.1016/j.scitotenv.2019.06.282>
- Gounout M (1969) Méthodes d'étude quantitative de la végétation. Masson édit, Paris, 320 pp.
- Gribben P, Thomas T, Pusceddu A, Bonechi L, Bianchelli S, Buschi E, Nielsen S, Ravaglioli C, Bulleri F (2018) Below-ground processes control the success of an invasive seaweed. *Journal of Ecology* 106 (5): 2082-2095. <https://doi.org/10.1111/1365-2745.12966>
- Guiry MD, Guiry GM (2020) *Algae-Base*. World-wide electronic publication, National University of Ireland, Galway (taxonomic information republished from *Algae-Base* with

- permission of M.D Guiry). Accessed through: World Register of Marine Species at: <http://www.marinespecies.org/aphia.php?p=taxdetails&id=382517> on 2020-03-28.
- Hamel G (1931) Chlorophycées des côtes Françaises. Revue algologique T. 5-6 33 pp.
 - Hamel G (1939) Phaeophycées de France. F. I-V, Paris, 432 pp.
 - Herrera RR, Legac DS (2002) Initiation à l'analyse factorielle des données: fondements mathématiques et interprétations. Ellipses, Paris, 315 pp.
 - Jaubert JM, Chisholm JR, Minghelli-Roman A, Marchioretto M, Morrow JH, Ripley HT (2003) Re-evaluation of the extent of *Caulerpa taxifolia* development in the northern Mediterranean using airborne spectrographic sensing. Marine Ecology Progress Series 263: 75-82. <https://doi.org/10.3354/meps263075>
 - Khiari N, Atoui A, Khalil N, Charef A, Aleya L (2017) Dynamics of sediments along with their core properties in the Monastir-Bekalta coastline (Tunisia, Central Mediterranean). Journal of African Earth Sciences 134: 320-331. <https://doi.org/10.1016/j.jafrearsci.2017.06.028>
 - Legendre L, Legendre P (1984) Ecologie numérique. Tome 1. Le traitement multiple des données écologiques: 260 p. Tome 2. La structure des données écologiques: 247p. 2^{ème} édition. Collection d'écologie 12. Masson, Paris.
 - Legendre P, Legendre L (1998) Numerical Ecology. Elsevier science, Amsterdam, 853 pp. [https://doi.org/10.1016/S0167-8892\(98\)80052-3](https://doi.org/10.1016/S0167-8892(98)80052-3)
 - Manconi R, Padiglia A, Padedda BM, Pronzato R (2020) Invasive green algae in a western Mediterranean Marine Protected Area: interaction of photophilous sponges with *Caulerpa cylindracea*. Journal of the Marine Biological Association of the United Kingdom 100 (3): 361-373. <https://doi.org/10.1017/s0025315420000193>
 - Mannino A, Balistreri P, Deidun A (2017) The marine biodiversity of the Mediterranean Sea in a changing climate: The impact of biological invasions. Mediterranean Identities - Environment, Society, Culture <https://doi.org/10.5772/intechopen.69214>
 - Mojetta A, Ghisotti A (1996) Flore et Faune de la Méditerranée. Edition Solar, Paris, 317 pp.
 - Montefalcone M, Morri C, Parravicini V, Bianchi CN (2015) A tale of two invaders: divergent spreading kinetics of the alien green algae *Caulerpa taxifolia* and *Caulerpa cylindracea*. Biological Invasions 17 (9): 2717-2728. <https://doi.org/10.1007/s10530-015-0908-1>
 - Ojaveer H, Galil BS, Carlton JT, Alleway H, Gouletquer P, Lehtiniemi M, et al. (2018) Historical baselines in marine bioinvasions: implications for policy and management. PLOS One 13 (e0202383). <https://doi.org/10.1371/journal.pone.0202383>.
 - Piazzi L, Ceccherelli G, Cinelli F, Gravez V, Ruitton S, Boudouresque CF, Le Direac'h L, Meinesz A, Scabbia G, Verlaque M (2001) Effet de *Caulerpa racemosa* sur la structure des communautés algales benthiques. Fourth international workshop on *Caulerpa taxifolia* edition, GIS Posidonie ublication. Fr: 371-37.
 - Piazzi L, Cinelli F (2003) Evaluation of benthic macroalgal invasion in a harbour area of the western Mediterranean Sea. European Journal of Phycology 38: 223-231. <https://doi.org/10.1080/1364253031000136358>
 - Piazzi L, Ceccherelli G (2006) Persistence of biological invasion effects: Recovery of macroalgal assemblages after removal of *Caulerpa racemosa* var. *cylindracea*. Estuarine Coastal and Shelf Science 68: 455-461. <https://doi.org/10.1016/j.ecss.2006.02.011>

- Piazzi L, Balata D, Bulleri F, Gennaro P, Ceccherelli G (2016) The invasion of *Caulerpa cylindracea* in the Mediterranean: the known, the unknown and the knowable. *Marine Biology* 163 (7). <https://doi.org/10.1007/s00227-016-2937-4>
- Premarathna AD, Sarvananda L, Jayasooriya AP, Amarakoon SA (2019) Review on pathogenic diseases on corals associated risk factors and possible devastations in future in the globe. *Journal of Marine Science: Research & Development* 9 (269).
- R Core Team (2014) R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria.
- Sambusiti C, Bellucci M, Zabaniotou A, Beneduce L, Monlau F (2015) Algae as promising feedstocks for fermentative biohydrogen production according to a biorefinery approach: A comprehensive review. *Renewable and Sustainable Energy Reviews* 44: 20-36. <https://doi.org/10.1016/j.rser.2014.12.013>
- Scammacca B, Giaccone G, Pizzuto F, Alongi G (1993) La vegetazione marina di substratoduro del l'isola di Lampedusa (Isole Pelagie). *Bollettino Accademia Gioenia di Scienze Naturali* 26 (341): 85-126.
- Serridi H (2003) *Algues Macrophytes, biodiversité Marine et littorale Algérienne*. ISBN N°9961- 9547-O-X79-94.
- Shobana S, Kumar G, Bakonyi P, Saratale G, Al-Muhtaseb AH, Nemestóthy N, Bélafi-Bakó K, Xia A, Chang J (2017) A review on the biomass pretreatment and inhibitor removal methods as key-steps towards efficient macroalgae-based biohydrogen production. *Bioresource Technology* 244: 1341-1348. <https://doi.org/10.1016/j.biortech.2017.05.172>
- Verlaque M (1990) *Végétation Marine de la Corse (Méditerranée) VIII. Documents pour la flore des Algues*. *Vie et Milieu* 40 (1): 79-92.
- Verlaque M, Fritayre P (1994a) Incidence de l'algue introduite *Caulerpa taxifolia* sur le phytobenthos de Méditerranée occidentale. - 2. Les peuplements d'algues photophiles de l'infralittoral. GIS Posidonie publication 349-353.
- Verlaque M, Fritayre P (1994b) Mediterranean algal communities are changing in face of the invasive alga *Caulerpa taxifolia* (Vahl) C. Agardh. *Oceanologia Acta* 17: 659-672.
- Warwick RM, Clarke KR (1995) New biodiversity measures reveal a decrease in taxonomic distinctness with increasing stress. *Marine Ecology Progress Series* 129: 301-305. <https://doi.org/10.3354/meps129301>
- Warwick RM, Clarke KR (1998) Taxonomic distinctness and environmental assessment. *Journal of Applied Ecology* 35: 532-543. <https://doi.org/10.1046/j.1365-2664.1998.3540532.x>
- Wilson O (1997) Foreword. In *Strangers in Paradis. Impact and Management of nonindigenous species in Florida*. Washington DC Island Press

Supplementary materials

Suppl. material 1: average recovery: Salamandre Station [doi](#)

Authors: Bachir Bouiadjra B., Ghellai M., Daoudi M., Behmene I.E., Bachir Bouiadjra M.A.

Data type: Table.CSV

Brief description: average recovery by species and by season, Salamandre Station

[Download file](#) (2.44 kb)

Suppl. material 2: average recovery: Small Port Station [doi](#)

Authors: Bachir Bouiadjra B., Ghellai M., Daoudi M., Behmene I.E., Bachir Bouiadjra M.A.

Data type: Table.CSV

Brief description: average recovery by species and by season, Small Port Station

[Download file](#) (4.74 kb)

Suppl. material 3: taxonomic classification of the species recorded [doi](#)

Authors: Bachir Bouiadjra B., Ghellai M., Daoudi M., Behmene I.E., Bachir Bouiadjra M.A.

Data type: table.CSV

Brief description: the taxonomic classification table, based on the phylogenetic classification, used for the calculation of taxonomic indices

[Download file](#) (10.86 kb)