

OPEN ACCESS

EDITED AND REVIEWED BY
Celia J. A. Morgan,
University of Exeter, United Kingdom

*CORRESPONDENCE Shannon Dames shannon.dames@viu.ca

SPECIALTY SECTION

This article was submitted to Psychopharmacology, a section of the journal Frontiers in Psychiatry

RECEIVED 06 June 2022 ACCEPTED 01 July 2022 PUBLISHED 19 July 2022

CITATION

Dames S, Kryskow P and Watler C (2022) Corrigendum: A cohort-based case report: The impact of ketamine-assisted therapy embedded in a community of practice framework for healthcare providers with PTSD and depression.

Front. Psychiatry 13:962882.
doi: 10.3389/fpsyt.2022.962882

COPYRIGHT

© 2022 Dames, Kryskow and Watler. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.

Corrigendum: A cohort-based case report: The impact of ketamine-assisted therapy embedded in a community of practice framework for healthcare providers with PTSD and depression

Shannon Dames^{1*}, Pamela Kryskow¹ and Crosbie Watler²

¹Health and Human Services, Vancouver Island University, Nanaimo, BC, Canada, ²Private Practice, Duncan, BC, Canada

KEYWORDS

ketamine-assisted psychotherapy, healthcare providers (HCP), resilience, psychedelic therapy, group therapy, community of practice, post-traumatic stress disorder (PTSD), depression

A corrigendum on

A Cohort-Based Case Report: The Impact of Ketamine-Assisted Therapy Embedded in a Community of Practice Framework for Healthcare Providers With PTSD and Depression

by Dames, S., Kryskow, P., and Watler, C. (2022). Front. Psychiatry 12:803279. doi: 10.3389/fpsyt.2021.803279


In the published article, there was an error in Figure 2 as published. The PTSD labels in the pre- and post- rows of the figure were incorrectly positioned underneath the Anxiety score, and vice versa. The corrected version of Figure 2 and its caption is published below.

The authors apologize for this error and state that this does not change the scientific conclusions of the article in any way. The original article has been updated.

Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

Dames et al. 10.3389/fpsyt.2022.962882



Cohort 1, 2, and 3 aggregate: mean scores. The "n" represents the number of participants who screened positive for PTSD, depression, or generalized anxiety disorder upon entry into the RTT-KaT program. The pre-results were taken within 1 month of the program beginning and post results were taken directly after the 12-week program was completed. If data was missing, it was not included.