

SYSTEMATIC REVIEW

A narrative synthesis scoping review of life course domains within health service utilisation frameworks [version 1; peer review: 2 approved]

Mary-Ann O'Donovan¹, Phillip McCallion¹, Mary McCarron¹, Louise Lynch ¹, Hasheem Mannan ², Elaine Byrne ³

v1

First published: 29 Mar 2019, 2:6 (

https://doi.org/10.12688/hrbopenres.12900.1)

Latest published: 29 Mar 2019, 2:6 (

https://doi.org/10.12688/hrbopenres.12900.1)

Abstract

Background: Current thinking in health recognises the influence of early life experiences (health and otherwise) on later life outcomes. The life course approach has been embedded in the work of the World Health Organisation since the Ageing and Health programme was established in 1995. Yet there has been limited debate on the relevancy of a life course lens to understanding health service utilisation.

Aim: The aim of the review was twofold. Firstly, identify existing healthcare utilisation frameworks other than the dominant Andersen's behavioural model currently in use. Secondly, to identify if current frameworks incorporate the advocated life course perspective in understanding health service utilisation.


Methods: A scoping review of PubMed, Cinahl Plus, Emerald, PsycINFO, Web of Knowledge and Scopus was conducted. Data extraction used a framework approach with meta-synthesis guided by the four domains of the life course proposed by Elder (1979): human agency, location, temporality and relationships, and interdependencies.

Results: A total of 551 papers were identified, with 70 unique frameworks (other than Andersen's Behavioural Model) meeting the inclusion criteria and included in the review.

Conclusion: To date there has been limited explicit discussion of health service utilisation from a life course perspective. The current review highlights a range of frameworks that draw on aspects of the life course, but have been used with this perspective in mind. The life course approach highlights important gaps in understanding and assessing health service utilisation (HSU), such as utilisation over time. HSU is a complex phenomenon and applying a structured framework from a life course perspective would be of benefit to researchers, practitioners and policy makers.

Keywords

health services utilisation, life course, framework, model



¹School of Nursing and Midwifery, Trinity College Dublin, Dublin, Ireland

²School of Nursing and Midwifery, University College Dublin, Dublin, Ireland

³Department of Epidemiology and Public Health Medicine, Royal College of Surgeons in Ireland, Dublin, Ireland



This article is included in the Ageing Populations collection.

Corresponding author: Mary-Ann O'Donovan (odonovm3@tcd.ie)

Author roles: O'Donovan MA: Conceptualization, Data Curation, Formal Analysis, Funding Acquisition, Investigation, Methodology, Validation, Writing – Original Draft Preparation, Writing – Review & Editing; McCallion P: Writing – Review & Editing; McCarron M: Writing – Review & Editing; Lynch L: Writing – Review & Editing; Mannan H: Writing – Review & Editing; Byrne E: Supervision, Writing – Review & Editing

Competing interests: No competing interests were disclosed.

Grant information: Health Research Board Ireland [PHD/2007/16].

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Copyright: © 2019 O'Donovan MA et al. This is an open access article distributed under the terms of the Creative Commons Attribution Licence, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: O'Donovan MA, McCallion P, McCarron M *et al.* A narrative synthesis scoping review of life course domains within health service utilisation frameworks [version 1; peer review: 2 approved] HRB Open Research 2019, 2:6 (https://doi.org/10.12688/hrbopenres.12900.1)

First published: 29 Mar 2019, 2:6 (https://doi.org/10.12688/hrbopenres.12900.1)

Introduction

The Global Health Strategy of 'Health for All in the 21st century' strives for equality, quality and security of health access, while extending the number of healthy life years lived by all (WHO, 1997). Within this strategy is a recognition that individual health experiences differ across and within populations, which ultimately results in divergent and unequal health outcomes that are linked to both medical and non-medical determinants of health (WHO, 1997). The preceding Ageing and Health Programme 1995 laid the foundations for the prominence of a life course perspective in health, and as a central thesis of World Health Organization (WHO) activities, recognises the importance of understanding the impact of the life course on health and how health systems are developed (WHO, 2000). This was further emphasized in the 2015 World Report on Ageing and Health (WHO, 2015).

Aligned to this is the belief that health outcomes in later life are impacted by early life circumstances (Braveman, 2014). This may be early-life health experiences, early life exposure to environmental, social, political and economic factors, or a combination of all of these early-life experiences in line with the life course view. The value of a life course approach in general is that it enables an understanding of the individual in relation to their environment (Elder Jr & Rockwell, 1979), past and present.

Life and the individual as they exist at present is a product of when, where, how an individual has lived previously, the choices made throughout life and the people and relationships that formed part of that life. The individual is shaped by their historical, social, cultural and familial context, as well as the extent of personal control and choice experienced throughout his/her life. Elder (1994) proposes that a life course approach is one which incorporates four elements: location, agency or personal control,

temporality and social relations and interdependencies (Braveman, 2014; Elder, 1994; Harrison, 2003; Slota & Martin, 2003).

In this approach, the importance of early life events on later life outcomes and behaviours (Braveman, 2014; Robison & Moen, 2000) is highlighted, as well as the relationships with family and dependencies and interdependencies with others over time. In addition, the extent of 'agency' or control a person has over their life is shaped by their environment (social, cultural, historical, locational and structural) and influences the direction and timing of life events and transitions.

It is all of these factors combined which form a life course perspective and this approach provides an opportunity to examine and address health, health inequities (Braveman, 2014) and health service utilisation (HSU) across groups and across time.

The changing demographic profile globally, and within individual countries, impacts directly on the delivery of health services (Goodyear-Smith & Janes, 2008), with the majority of developed countries facing the demands of an ageing population. Health service systems and the structure of service delivery needs to be informed by where and how the population is ageing to, as well as where and how the population has aged from. The supports and services that will be newly required or will continue to be used into the next stage of the life course will also inform these health service systems and delivery.

The study of HSU is commonly explored using the Andersen Behavioural Model of Healthcare Use (Andersen, 1995), depicted in Figure 1. The original Andersen model was proposed in 1968 with the focus on the use of formal personal health services, and the family as the unit of analysis. The purpose of the model was to help explain and predict use of health services. By the 1970s




Figure 1. Andersen's Behavioural Model of healthcare use. This figure has been reproduced with permission from (Andersen, 1995).

the unit of analysis had shifted from the family to the individual. Reference to the health system, to account for the role of health policy and its impact on health service use, was incorporated (Andersen, 1995). There have been many adaptations of the core Andersen model across the years (Goldsmith, 2007) and a number of reviews (Babitsch *et al.*, 2012; Guilcher *et al.*, 2012; Phillips *et al.*, 1998).

Although Andersen is the most frequently used model of HSU, it may not necessarily be the most appropriate framework for understanding HSU from a life course perspective. The Andersen model examines HSU at one point in time and there is no explicit recognition of the life course within this model. However, the generally accepted belief that health should be examined across the life course (Elder Jr & Rockwell (1979)), would imply that to understand HSU a life course approach may be beneficial.

Thus, the purpose of this paper is that if we are to understand HSU more fully in light of transitions in living and choice, we need to identify:

- 1. What are the range of frameworks currently applied to the study of HSU?
- 2. Whether these frameworks are informed (or could be adapted to be informed) by a life course approach.

Methods

For this scoping review, a methodical approach involving five steps of the systematic review process was followed to corroborate the paper objectives (Briner & Denyer, 2012):

- 1. Planning the review
- 2. Identifying studies
- 3. Evaluating study contribution
- 4. Analysing the information
- 5. Reporting accurate results

Planning the review

Approach

A narrative synthesis approach was used to identify and amalgamate the relevant framework information from appropriate

studies for this paper. A narrative synthesis uses a textual rather than a statistical approach for analysing results and drawing conclusions. This words-based approach is more suitable to investigating the life course domains within HSU.

A scoping review is a descriptive review, which is a broader and less defined search than a systematic review. It is suitable for literature that is large, complex or heterogeneous. A scoping review can summarise the extent, variety and characteristics of findings on a specific topic, in this case the use of life course domains within HSU, and identify gaps in knowledge (Tricco *et al.*, 2018).

Identifying studies

Search strategy

The search spanned a 20-year time frame (1995–2015); 1995 was selected as the starting point for the review as this was the year the major revision of the Andersen model was published in the academic literature. Email alerts were set up in the databases to ensure the review was current. A summary of the review criteria is presented in Table 1.

Databases

PubMed, Cinahl Plus, Emerald, PsycINFO, Web of Knowledge, and Scopus were used for the search. The key search terms used were health services utilization OR health services utilisation OR health services accessibility AND (framework OR model); and related MESH terms in PubMed. The searches in Scopus and Web of Knowledge were restricted to journal title and included titles relevant to health services, health policy, health management, value in health, health planning.

An example of a search string that was used in PubMed for a custom year span of 1995–2015 is:

((((health service* utilization) OR health service* utilisation) OR health service* Access*) AND Framework) OR model

Search procedure

Application of the broad search string indicated above returned 12,126 articles. See Figure 2 for the flow diagram of results.

At stage one and two of the screening process, many articles were excluded based on the title, abstract or full text not meeting key inclusion criteria, leaving 1698 articles for further review.

Table 1. Summary criteria for review.

Aim	Identify frameworks used to understand/study health service utilisation framework from 1995–2015.
Participants	General adult population including vulnerable groups (minority, disease specific, disabled, intellectual disability, low income)
Interventions	Use of health services; accessibility of health services where accessibility refers to or leads to use of services
Outcome measure	Application of specific health services utilisation/health services accessibility model or framework
Study design	Any
Inclusion criteria	English language articles only; Full article availability only from 1995 to 2013; Specific named model or framework applied or developed to understand/examine health services utilisation.
Exclusion criteria	General health systems (strengthening) frameworks/models; Social determinants of health; Health inequalities/health disparity; Health delivery frameworks/models; Middle and low income countries; Health care workers




Figure 2. Flow chart for systematic review search of models or frameworks to study health service utilisation.

The full text of the 1,698 articles were accessed and reviewed for the framework or model used in the study. Articles were excluded at this stage (stage 2) if statistical modelling alone as opposed to a conceptual or theoretical model for understanding health services utilisation was referenced, resulting in 551 articles remaining. If a framework was named or a new framework was proposed this was also recorded. As noted earlier only one article purposively selected with the occurrence of a

framework was included. A total of 164 frameworks with an associated primary article remained after this third stage of screening.

To ensure that the identified frameworks were unique and separate frameworks, and were applied in an HSU context, each full text article (n=164), whether quantitative or qualitative, was read and confirmed for inclusion by MOD (and confirmed by EB). Further duplicates or adaptations of the Andersen model

were excluded, resulting in 64 frameworks remaining. Six additional frameworks were subsequently identified through email alerts, and once reviewed against the same criteria, the final number of frameworks included in the review was 70. This list of frameworks was reviewed and consensus agreed for inclusion by the entire review team.

See Table 4 for a sample of the frameworks identified. The complete list of the 70 frameworks is available on Zenodo: https://doi.org/10.5281/zenodo.2590439 (O'Donovan *et al.*, 2019).

Evaluating study contribution

Frameworks

For the purposes of this review, the Behavioural model and any iteration will be referred to as the Andersen model/framework. The terms framework and model were used interchangeably and taken to be equal. Of note is that the context for use of the framework was to understand health services utilisation. Thus, the frameworks were identified as 'used by researchers', regardless of whether the original genesis or intent of the framework was to understand or examine this particular phenomenon.

The frequency with which the identified models were used was not recorded, as the purpose of the review was to identify the range rather than prevalence/utilization of specific frameworks. Hence one paper using the identified framework was purposively selected for further analysis. These selected articles met the criteria for the systematic review most clearly and incorporated the most complete use of the framework.

The unit of analysis was the individual's use of services. The terms health services, utilisation and population are defined in Table 2. Inpatient, surgery and diagnostic testing were not included in this review.

Screening

There were three levels of screening, led by the first author (MOD) with consultation meetings and review at each stage involving the entire review team (the remaining authors). The review team covered expertise across the areas of HSU (PMcC), health systems and qualitative research methods (EB), health and disability statistics (HM, PMcC) and intellectual disability and health (MMcC, PMcC). Each of these experts had previous experience in conducting systematic and scoping reviews. The review team systematically discussed the merit of each framework in relation to the aim of the review as well as the context of its application to understanding HSU. This process was on-going until a consensus was reached on the final list of frameworks for inclusion.

Analysing the information

Mapping

Once the frameworks were identified they were mapped against a matrix of HSU categories and variables. The Andersen model was the basis for 'a-priori' categories and any variables not easily aligned with the Andersen model were categorised as 'other specified'. The a-priori categories were the predisposing, enabling and need variables of the model. The rationale for this approach was to explore whether the Anderson model incorporated all the categories and variables of the other HSU models and if not, what these 'missing' variables were.

The Anderson model and the 'other specified' categories were then mapped to the life course domains of human agency, location, relationships and interdependencies, and temporality (Elder Jr & Rockwell (1979)). Table 3 explains each of the life course domains. This mapping exercise explored whether the frameworks incorporated a life course approach and could therefore be used in exploring HSU from a life course perspective. See Table 5 for details of the mapping.

Table 2. Definition of health services, utilisation data and population.

Health Services	Community based and primary care services, hospital-based therapy care (physiotherapy, Speech and language therapy, cancer care, chemotherapy; non anti-retroviral therapy care for people with HIV/Aids, dental services, emergency department, ambulance use, screening, preventative care, vaccinations and sexual health check screening)
Utilisation data	Use of the listed health services and accessing health services where it corresponds with use, visits, consultations, attendance, service consumption.
Population	People aged 18 years and over living in high income countries (based on World Bank listings)

Table 3. Explanation of life course domains.

Life Course Domain	Meaning
Human Agency	Refers to an individual's control, autonomy and choice in directing his/her own life path.
Location	Exposure to risks, whether the living environment is health-promoting and access to services can be determined by where a person lives
Relationships & interdependencies	Dependence (or not) on others for support to engage with a range of activities, functional, social, sexual, as well as support in asserting one's own identity and place in the world,
Temporality	Historical effects on life/the timing of lives

Table 4. Sample of frameworks identified and purposive sample of articles.

No.	Framework/model identified	Description	Sample article from current review
1	Access Triangle	This framework specifically focuses on access to dental care for underserved populations.	Guay, 2004; 135(11):1599–605
2	Cooper model of access to care	Copper model explores barriers and mediators to health service use across personal, structural and financial aspects	Messer et al., 2013
3	Frenk's domains of Access	"Frenk reserves the term access to denote the ability of the population to seek and obtain care"	Levesque et al., 2013
4	Gullifords model of healthcare access 2001	Gullifords model of access depicts relationship between factors and how they impact on access. This model was adapted by Alborz <i>et al.</i> for people with ID.	Alborz <i>et al.</i> , 2005;
5	Hispanic Farmworker Health Model (Ward, 2007)	The HFH model incorporates 4 determinants of health and 2 outcomes specific to farmworker of Hispanic ethnicity. Highlights importance of living and working conditions and social and cultural factors	López-Cevallos et al., 2013;.
6	ICF framework	The authors used the ICF "was chosen because it allows examination of personal, environmental, and contextual factors, as well as impairments, activity limitations, and participation".	Iversen <i>et al.</i> , 2011; WHO, 2001.
7	Modified model of access to dental care	This model is combination of elements of Penchansky & Thomas (1981) and Maxwell (1984) models to understand access to dental services by PwID.	Owens et al., 2010;
8	Penchansky & Thomas 1981 Concept of Access: definition and relationship to consumer satisfaction	This model introduces the concept of 'fit' between the individual and the service. Five dimensions considered important – availability, accessibility, accommodation, affordability, and acceptability.	Wallace & Macentee, 2012.
9	Complementary and Alternative Medicine (CAM) use – a conceptual model	Draws on Parson's sick role theory, Suchman's stages of illness theory and Andersen's Sociobehavioural Healthcare utilisation Model to examine use of complementary and alternative medicine.	Davis <i>et al.</i> , 2011
10	Non-urgent Emergency Department use – a conceptual model	Conceptual model of pathway to use emergency department	Uscher-Pines et al., 2013;

Note: N=2200 approximately if included all 70 models

This process of meta-synthesis was conducted by two of the authors (MOD and EB) and reviewed by the entire review team. It involved a process of sense-making and developing a cohesive structure with which to present the findings of the extraction.

Results

In total 29 of the 70 frameworks identified were specific HSU frameworks, with the remainder drawing on models from within psychology, sociology, disability studies and human rights, and applied to a HSU context. The frameworks are now discussed across the four life course domains. See Table 5.

1. Human agency

Human agency refers to the individual as an actor who plans and makes choices in their lives (Elder, 1994). In the context of this review, agency was identified in terms of choice and control. Fifteen frameworks indicate the role of choice in HSU through the inclusion of variables of lifestyle choice (Alborz *et al.*, 2005), individual decisions and decision points, and the

environment (Stoller *et al.*, 2011). This is described well in the work of Stoller *et al.* (2011), which is framed by Leventhal's concept of the individual as an active problem solver in relation to their health, and Cockerham's healthy lifestyle paradigm which includes agency and life chances (structure), and the interaction between both of these (Leventhal *et al.*, 2003; Stoller *et al.*, 2011).

The individual as self-directing is also captured by McCormack & Mackintosh (2001) in the conceptualisation of pathways to health. The concept of decision-points in relation to use or selection of services (Engelman *et al.*, 2011; Shaw *et al.*, 2013; Uscher-Pines *et al.*, 2013), informed choice (Essink-Bot *et al.*, 2013), ability to seek care (Levesque *et al.*, 2013) and individual involvement in decisions (Lee *et al.*, 2009) were also identified.

Control features in a more limited way than choice in the reviewed frameworks with six instances of control specifically mentioned: decisional control (Witucki & Wallace, 1998), low job control and how this is related to stress (Azagba & Sharaf, 2011), health

Table 5. Meta-synthesis of variables.

Andersen	Variables shared with other frameworks	Variables unique to other frameworks	Alignment to life course themes
Environment			
Health care system	Equity of services; Decreased disparity; Healthcare reaching; Health consequence; Accommodation; Accessibility; Quality of providers; Experience dealing with People with Intellectual disability (PwID); Willingness to serve; Liaison; Referral by medical practitioner; Reinforcing factors		Location Relationships & interdependencies
External environment	External environment Local advertising; Cues to action; Alleviating stigma	Iving conditions - physical, biological, social, cultural, material; traffic congestion/courtesy,	
Population			
Predisposing	Personal attachment style: Patience; Intrinsic motivation; Cognitive appraisal; Affective response, Self-confidence, Level of intellectual disability; ability factors at individual and family level; cognitive level; Cultural Beliefs and matching; Feminism; Readiness to change; Assumption of sick role; Equity in health, Disposition to act; Health competence; Health knowledge; Awareness and information factors; Perceived threat; Planning; Ethnic inequity;		Relationships and interdependencies
Enabling	Mother - age, race, marital status, education, employment status, if child is in or out of house; others - functional ability, burden; future worry); availability of carer/advocate; Relationships; Family separation, Advocacy; Cultural distance, socialization experience; Folk arena; Social inclusion; Social interconnectedness; Social capital – individual and community level; Sick person requires assistance to get better; Human rights; Person, behaviour, environment interaction; Resources (monetary, incentives);		Relationships and interdependencies
Need			
Health behaviour			
Personal health practices	Avoidance;	Life/lifestyle choices; Adherence;	Human agency
Use of health services	Multiple types of use options;	intention to use service; utilisation propensity; unmet need; delay or no treatment; ability to get care (perceive, seek, access);	Human agency Temporality

Andersen	Variables shared with other frameworks	Variables unique to other frameworks	Alignment to life course themes
Outcomes			
		employment outcome;	
Perceived health status	Outcome expectancy/efficacy; Inaccurate symptom interpretation;		
Evaluated health status	Disease activity index (RADI); Multi-dimensional health assessment questionnaire (MDHAQ); Perceived susceptibility; Days hospitalised; Disability days; Efficacy of treatment; Health outcomes at individual and community level;	Care appropriateness,	
Consumer satisfaction			
		Life events; Life changes/structure; Permanent change or modification of life situation; Life disclosure; Time;	Temporality
		Perceived control; High or low control of services; Agency; Human agency Decisional control; Informed choice and decision-making	Human agency
		Choosing/appraising; Decision to seek help; Process component - danger control/fear control; Fear Cost/benefit analysis; Competing priorities	Human agency
		Message – acceptance/rejection;	Human agency

locus of control (Davis *et al.*, 2011) and perceived control (Jo *et al.*, 2008; Wong *et al.*, 2006). Hubbell (2006) incorporates reference to danger control and fear control. Concepts of cost/benefit analysis were identified in the work of Fonseca-Becker *et al.* (2010) and Iversen *et al.* (2011) and imply a specific type of decision around competing priorities that an individual may make.

Choice and control were not identified in the Andersen model. The individual as an active participant in their lives, could be said to incorporate attitudes, beliefs, values and knowledge (Davis et al., 2011; Hubbell, 2006; Jo et al., 2008; McCormack & Macintosh, 2001; Wong et al., 2006), internal processing (Davis et al., 2011; Essink-Bot et al., 2013; Shaw et al., 2013), and disposition to act (Essink-Bot et al., 2013; Fonseca-Becker et al., 2010; Leventhal et al., 2003; Levesque et al., 2013; Stoller et al., 2011; Wong et al., 2006), which inform the decisions made and actions taken. These could be captured under the predisposing element of the Andersen model and were identified in 14 frameworks.

2. Location

Six frameworks incorporated a specific place variable, which included urban/rural distinction Hubbell, 2006; Seccombe, 1995; Stoller *et al.*, 2011), geographic location (Levesque *et al.*, 2013), living and housing conditions (Koopmans & Lamers, 2007; Lopez-Cevallos & Garside, 2013; Ståhlnacke *et al.*, 2005), residential settings (Essink-Bot *et al.*, 2013) and distance to travel and neighbourhood factors (Birkin *et al.*, 2008; Graves, 2009; Meade *et al.*, 2015; Mobley *et al.*, 2006). Essink-Bot *et al.* (2013) specifically refer to living in a residential setting but this is the only occurrence of such a place variable in the frameworks.

Transportation (Messer et al., 2013; Shaw et al., 2013) and service setting (Alborz et al., 2005) are other location variables identified as influencing HSU. Toloo et al. (2013) includes living status as a socio-demographic variable to consider in HSU. In this context, it refers to if the person lives alone or with someone else (partner, children, family etc.), thus highlighting the link between place and people (Toloo et al., 2013) and the role of relationships in HSU.

Andersen does not explicitly mention location in terms of geographic, physical and social make-up of the space within which an individual lives. Travel to health service location is incorporated within Andersen as an enabling factor.

3. Relationships and interdependencies

The relationships and interdependency domain, or as otherwise referred to by Elder (1994) as linked lives was identified in 20 of the reviewed frameworks. The importance of social interconnectedness (Levy *et al.*, 2014), psychosocial (Nauenberg *et al.*, 2011) and affective supports (Witucki & Wallace, 1998) to HSU were highlighted.

Communication and rapport with the healthcare provider was identified in the context of immigrants who were non-English speaking (Jo et al., 2008). Broader issues of human rights

(Gibson & Mykitiuk, 2012), feminism (Bergeron & Senn, 2003), stigma (Yalon-Chamovitz, 2009), and legal status (Hvinden, 2003) are included within this domain. This is because they impact on the status and roles an individual has within a given society and how others in society interact with the individual. Cultural beliefs (Barrett *et al.*, 2003) and folk values (Thurston *et al.*, 2014) also represent a type of interdependency that was recognised in some frameworks as impacting on HSU.

Relationships and interdependencies manifest in the Andersen model under the enabling factor of personal and family means to access services and availability of health personnel, as well as the social networks and interactions under the predisposing variable.

4. Temporality

Temporality within life course theory recognises the impact of when people are born, and societal and historical context within which the person is born, on their life course trajectory (Elder, 1994). Implicit within the concept of temporality of the life course, is that of change over time and the differing experiences and timings of change and transitions. With this, the current review identified temporality as any time-related variable or variables linked to change occurring, that have time implicit in the concept.

Changes in the life course were identified in a small number of the reviewed frameworks (n=4). Stoller *et al.* (2011) include life changes/structure, while Petersen (1990), as referenced in Ståhlnacke *et al.* (2005), includes permanent change or modification of life situation in his conceptualisation of health-related behaviour.

Life events were identified in two frameworks, with Setia *et al.* (2011) referring to Berry's (1997) life events and Kosteniuk & D'Arcy (2006) referring to early life as an important consideration for understanding utilisation of health services. Time as a factor was included in the conceptual model by Xiao *et al.* (2014). This was the only explicit occurrence of time in the frameworks.

The Andersen model presents a snapshot of health service utilisation and does not incorporate a change or time variable.

The PRIMSA-ScR checklist, PRIMSA flow diagram and Table S6 which contains all 70 frameworks identified in this review are available on Zenodo: https://doi.org/10.5281/zenodo.2590439 (O'Donovan *et al.*, 2019).

Discussion

Andersen and the life course

Andersen's framework (Andersen, 1973; Andersen 1995) provides a useful basis for understanding and exploring predictors of health services utilisation. This is illustrated by the frequent application and adaptation of the framework to different health service contexts. In addition to the application of the Andersen model, which is the dominant guiding framework in the study of HSU, this review indicates that HSU in some contexts, with some populations and specific service settings is not sufficiently

addressed through the Andersen model. This is evidenced by the emergence of such a range of alternative frameworks.

When assessing HSU in the context of the life course theory, Andersen is limited in its incorporation of the domains of the life course. Most notably, through the absence of issues related to human agency and temporality, limited reference to location, and some engagement with relationships and interdependencies. Though the traditional HSU model does not include core domains, recent reviews of the model indicate that the application of the model in this way has begun. For example, the domains of choice, decision-making and control are not explicitly evident in the Andersen framework, but the health locus of control concept was noted by Babitsch *et al.* (2012) as used in one study as a predisposing factor. This shows the potential to adapt the model to take on aspects of a life course perspective.

The particular strength of the Anderson model is very useful for considering the individual's interaction with the health system at one point in time for one condition/scenario. Temporality as envisaged within the life course is not addressed within the model, although 'time interval' does feature within phase two of the model. This time interval refers to time within the health system, such as waiting time. However, the alternative frameworks propose ways to capture the temporality of health and temporality of health care need and use over the life course.

Strong across both Anderson and non-Anderson models, is the recognition of the importance of systems of support around the individual to engage with and access the health system. Relationships and interdependencies are how this is relayed in the life course approach and it is apparent in both.

Andersen identifies family and community level variables as important enabling resources to HSU with social and emotional supports recorded by Babitsch *et al.* (2012). Guilcher *et al.* (2012) recommend the inclusion of quality of relationships/emotional support with informal and formal carers to future adaptations of Andersen. Andersen introduces the concept of 'site' of the health service in version two and 'external environment' in version three of his model, again related to the health system, and are location-related variables (Andersen, 1995).

Applications of the Andersen model have used broad place variables at the level of geographic location and community characteristics such as distance travelled, urban/rural distinction and region. There is an absence of a more refined place variable such as type of living setting, condition of housing and ease of navigation between home and wider neighbourhood/community space. Place and home have been shown to be important non-determinants of health (Field & Briggs, 2001; Macintyre *et al.*, 2003) and yet have limited expression in Andersen's model. The model is primed to further develop these domains to reflect the life course more fully.

In the review of Andersen's model by Guilcher et al. (2012), it is recommended that 'home layout' be included in future access studies for people with spinal cord injury. Applications of Andersen's model have incorporated the idea of distance

travelled to health services, but Andersen does not specify access to transport as a factor in influencing HSU.

Culture is another aspect of the life course incorporated into HSU frameworks. Babitsch *et al.* (2012) include culture as an additional domain when mapping previous applications of Andersen. Culture is not a specific domain in the Andersen model but can manifest under the social structure aspect of the pre-disposing factors. In addition, cultural factors have been identified as enabling in application of the Andersen model. It is important to consider that the role of other people and cultural and social norms, may in fact be prohibitive to using health services and engaging with the health system.

This scoping review has identified a number of frameworks that illustrate how elements of the life course can be incorporated into the study of health service utilisation, although no framework included all aspects. The additional frameworks identified in this review, demonstrate commonality in variables with the Andersen model. These commonalities are illustrated in Table 5 (column 2).

The unique variables identified in these frameworks have a strong propensity towards the human agency aspect of the life course and position the individual as a more active participant in their health care and life situation. There is also added focus on the timing of care and temporality overall which is not easily identified in Andersen and which the current analysis considers absent.

The life course and HSU

The WHO (2002) perspective of the individual as an active participant in their health and healthcare, is evidenced in the findings of this scoping review. The identified frameworks positioning the individual at the centre of their own lives and in particular their health experience. Choice, decision making and the individual demonstrating agency over their health experience and HSU experience is explicit.

Selecting among competing priorities is a judgement that individuals face in everyday life and in the healthcare context. Such competing decisions also exist at a health system and health policy level. This cost/benefit judgement will implicitly impact on if, when, how and the type of health services accessed by individuals and decisions to forgo required health services. Influencing factors at these decision points include the social network and connections that the individual has, as well as family, community and broader cultural and societal beliefs and values.

In particular, the role that environmental factors play in the HSU experience of minority or more vulnerable populations, such as health equity, healthcare reaching and cultural and social capital (Nauenberg *et al.*, 2011) is highlighted, and includes issues of discrimination and stigma as well as legal status of minority groups within the community (Lee *et al.*, 2009).

Social interconnectedness (Levy *et al.*, 2014), psychosocial (Nauenberg *et al.*, 2011) and affective supports (Witucki & Wallace, 1998) are also important influences on HSU. Availability

of supports, in addition to the availability of services, is another strong theme within the reviewed frameworks and align with the interdependency and relationship domain of the life course. Andersen's predisposing domain could incorporate social relations, but the context or nature of the role of social relations in HSU is unclear within the model.

The importance of the provider-patient relationship and ability to communicate and establish rapport in the HSU process was noted and highlighted, particularly for immigrants who were non-English speaking (Jo *et al.*, 2008). Broader issues of human rights, feminism, and stigma are included within the interdependency domain as they impact on the status and roles an individual has within a given society and how others in society interact with the individual.

Cultural beliefs and folk values also present a type of relationship and interdependency that informs HSU for some people. The influence of the individual's support networks and the interrelationship between the individual and the broader cultural belief system within which they live or have lived has been shown to influence type of health care, such as decision to use traditional or complementary health services compared with mainstream health services. The heightened focus on cultural factors and distinguishing culture from community is an important addition.

Andersen does not address cultural factors as distinct from community, however reviews of the Andersen model have identified the application of culture in the use of the model (Babitsch *et al.*, 2012) under the rubric of contextual factors. Similarly, social relationships are not explicitly mentioned in the Anderson model, but are implied through its application.

Adopting a life course perspective to HSU enables a disaggregation of the influence of the family unit, which changes over the trajectory of the life course. This impacts on the use of health services through availability of support from other people to access services. Others knowledge, attitudes and experience of the health system, predisposition to mainstream or alternative, complementary or culturally specific values, and beliefs of appropriate health system to engage with, can also affect HSU. The inclusion of living status by Toloo *et al.* (2013) in examining HSU, that is whether the person lives alone or with someone else (partner, children, family etc.), highlights the link between place and people.

Under location, the need to include a more refined and 'local' place variable to better understand HSU is highlighted. Though the location of the health care interaction needs consideration, so does the type of housing, structural issues, living situation, neighbourhood and community factors. These place variables, which were identified and differ to that of site of health service and distance to travel, which are included in Andersen, are shown to be useful additions. Transportation (Messer et al., 2013; Shaw et al., 2013) highlights the relevance of how an individual can navigate the space they live in to access services. Alborz et al. (2005) also indicate service setting as an important contributor to HSU, specifying the need to set characteristics of opening times and waiting times in understanding use. In

addition, the home setting and social connections within the home are also relevant locational factors. Where an individual lives and who an individual lives with, as well as how geographically close to networks of support they are, acknowledges the role of the social determinants of health (Dahlgreen & Whitehead, 1991; Emerson *et al.*, 2011).

Temporality is the fourth domain of Elder's life course perspective but there is limited focus on time and the impact of life events over time in the frameworks. The absence of time illustrates the continued tendency to view health access at one point in time. Of the four models with some reference to temporality, only one explicitly named time. However, life changes and life events were identified in others as potential trigger points for health service use. The inclusion of a time variable acknowledges that health is impacted by early life events as is current access and use of appropriate services (Norris & Aiken, 2006). HSU is not static. It is a complex process, involving multiple service interactions at any one time. Interaction with one element of the health system may be a trigger for later interactions with the same or other aspects of the system. Hser et al. (2007) introduces the concept of duration of use in addition to frequency and highlights the importance of capturing HSU data over multiple time points.

As noted above, not all of the frameworks identified are traditional HSU models but have been applied to understand, explain or predict service utilisation in the health sector. Of note, the health belief model was widely used when the service focus was screening and health checks, with studies of emergency care and ambulance service use more likely not to use a framework. It could be implied that the move beyond HSU frameworks is an indication of the limits of current HSU frameworks, including Andersen.

The identified models of HSU add to the understanding of HSU by including the individual as an active participant in health and utilization of services. This is demonstrated through the focus on decision-making and choice elements. The individual utilization of healthcare does not exist in isolation. HSU is demonstrated as informed by the geographical, social, temporal and cultural context within the person's life, and is more deeply explored in these models.

In addition to the differences between these models and the Andersen model, some similarities were identified, with many of the reviewed frameworks including variables that map to the traditional domains of the Andersen model; predisposing, enabling, need, health system characteristics and outcome variables. Most notable was the prevalence of predisposing and enabling variables. However nearly 40% (n=27) of the frameworks included variables that did not easily map to Andersen's framework. See Table 5, column 3 to identify the variables that were unique to other frameworks.

Conclusion

The dominant HSU model, developed by Andersen, does not offer a life course approach to the examination of HSU. The current review highlights that researchers are exploring aspects of the life course to understand how and why people choose to use health services. The role of personal choice and autonomy; the application of choice, and to a lesser extent control, in HSU frameworks have been highlighted.

This application of a life course framework to HSU not only helps to categorise and understand the additional variables/domains included within the reviewed frameworks, but also sheds light on some limitations of the Andersen framework. In particular, extending the understanding of HSU beyond the health system, the individual and community at one point in time, to encompass HSU over changing time and space is a particularly important learning.

Health and HSU are framed by where one lives and who one lives with and vice versa; choice of location or home is influenced by one's need for service. This systematic review of frameworks highlights how the individual is positioned at the centre of their own lives and in particular their health experience. The position of the individual in this and the extent of choice and autonomy is impacted by and impacts on both health and home.

None of the identified models took an explicit life course approach or incorporated all four elements of the life course. Thus, the reviewed frameworks provide evidence for how elements of the life course can be included in frameworks to understand HSU but further work is needed to develop a model that includes all life course domains explicitly and comprehensively. A limitation of HSU models overall is the failure to recognise the impact of earlier life events on health and HSU. Previous health system interactions and knowledge of HSU impact on current and future HSU as well as other people's previous HSU experience.

Important considerations in the utilisation of services by specific sub-groups of the populations should inform the content and structure of any future framework applied. The comprehensive examination of domains of the HSU frameworks has highlighted a range of factors that can potentially enhance the ability to plan the delivery and financing of future health care. Overall the issues and experience of the individual across the life course will likely impact on the ease with which required services are accessed at the appropriate time and place. Greater application of these domains within general HSU frameworks could inform

future HSU models for people currently more susceptible to health inequalities across the life course.

Limitations of the review

Due to the inclusion criteria for this review, it is likely that not all frameworks that have been used for the study of HSU were identified. Frameworks used for the specific HSU experience of children, low-middle income countries and people with mental health difficulties are not covered in this review. Some of the identified frameworks may be applied in these contexts but there may also be additional frameworks which have not been identified in this review.

The quality of the studies using frameworks was not assessed. The aim of this review was to identify the range of frameworks, other than Andersen's Behavioural Model, that have been applied to the understanding of HSU. The study design and methodological issues were not of relevance to this question as the focus was the presence and use of a framework.

Data availability

Zenodo: A Narrative synthesis Analysis of Life Course Domains with Health Service Utilisation Frameworks. https://doi.org/10.5281/zenodo.2613425 (O'Donovan *et al.*, 2019).

This project contains the following data:

Underlying data

 HSU_Narsyn_70FWs: The complete list of the 70 frameworks

Reporting guidelines

- PRISMA-ScR checklist
- PRSIMA flow diagram

Data are available under the terms of the Creative Commons Attribution 4.0 International license (CC-BY 4.0).

Grant information

Health Research Board Ireland [PHD/2007/16].

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

References

Alborz A, McNally R, Glendinning C: Access to health care for people with learning disabilities in the UK: mapping the issues and reviewing the evidence. *J Health Serv Res Policy*. 2005; **10**(3): 173–82.

PubMed Abstract | Publisher Full Text | Free Full Text

Andersen JG: Health services utilization: framework and review. Health Serv Res. 1973; 8(3): 184–99.

PubMed Abstract | Free Full Text

Andersen RM: Revisiting the behavioral model and access to medical care: does it matter? *J Health Soc Behav.* 1995; **36**(1): 1–10.

PubMed Abstract | Publisher Full Text

Azagba S, Sharaf MF: Psychosocial working conditions and the utilization of

health care services. BMC Public Health. 2011; 11: 642. PubMed Abstract | Publisher Full Text | Free Full Text

Babitsch B, Gohl D, von Lengerke T: Re-revisiting Andersen's Behavioral Model of Health Services Use: a systematic review of studies from 1998-2011. Psychosoc Med. 2012; 9: Doc11.

PubMed Abstract | Publisher Full Text | Free Full Text

Barrett B, Marchand L, Scheder J, et al.: Themes of holism, empowerment, access, and legitimacy define complementary, alternative, and integrative medicine in relation to conventional biomedicine. J Altern Complement Med. 2003; 9(6): 937–47.

PubMed Abstract | Publisher Full Text

Bergeron S, Senn CY: Health care utilization in a sample of Canadian lesbian women: predictors of risk and resilience. Women Health. 2003; 37(3): 19–35. PubMed Abstract | Publisher Full Text

Berry JW: Immigration, acculturation, and adaptation. Appl Psychol. 1997; 46(1): 5-34

Publisher Full Text

Birkin C, Anderson A, Seymour F, et al.: A parent-focused early intervention program for autism: who gets access? J Intellect Dev Disabil. 2008; 33(2):

PubMed A ostract | Publisher Full Text

Braveman P: What is health equity: and how does a life-course approach take us further toward it? Matern Child Health J. 2014; 18(2): 366-72. PubMed Abstract | Publisher Full Text

Briner R, Denyer D: Systematic review and evidence synthesis as a practice and scholarship tool. In Rousseau, D. (ed.), The Oxford Handbook of Evidencebased Management. Oxford: Oxford University Press, 2012; 112-129. **Reference Source**

Dahlgreen G, Whitehead M: Policies and Strategies to Promote Social Equity in Health, Stockholm, Sweden: Institute for Future Studies, 1991.

Reference Source

Davis MA, Weeks WB, Coulter ID: A proposed conceptual model for studying the use of complementary and alternative medicine. Altern Ther Health Med 2011: 17(5): 32-6.

PubMed Abstract

Elder GH Jr, Rockwell RC: The Life-Course and Human Development: An Ecological Perspective. Int J Behav Dev. 1979; 2(1): 1-21.

Publisher Full Text

Elder GH Jr: Time, Human Agency, and Social Change: Perspectives on the Life Course. Soc Psychol Q. 1994; 57(1): 4–15.

Emerson E, Madden R, Graham H, et al.: The health of disabled people and the social determinants of health. *Public Health*. 2011; **I25**(3): 145–7. PubMed Abstract | Publisher Full Text

Engelman KK, Cupertino AP, Daley CM: Engaging diverse underserved communities to bridge the mammography divide. BMC Public Health. 2011; 11:

PubMed Abstract | Publisher Full Text | Free Full Text

Essink-Bot ML, Lamkaddem M, Jellema P, et al.: Interpreting ethnic inequalities in healthcare consumption: a conceptual framework for research. Eur J Public Health. 2013: 23(6): 922-6.

PubMed Abstract | Publisher Full Text

Field KS, Briggs DJ: Socio-economic and locational determinants of accessibility and utilization of primary health-care. Health Soc Care Community. 2001; 9(5): 294-308.

PubMed Abstract | Publisher Full Text

Fonseca-Becker F, Perez-Patron MJ, Munoz B, et al.: Health competence as predictor of access to care among Latinos in Baltimore. J Immigr Minor Health. . 2010; **12**(3): 354–60.

PubMed Abstract | Publisher Full Text

Gibson BE, Mykitiuk R: Health care access and support for disabled women in Canada: falling short of the UN Convention on the Rights of Persons with Disabilities: a qualitative study. Womens Health Issues. 2012; 22(1): e111–8. PubMed Abstract | Publisher Full Text

Goldsmith LJ: Access to health care for disadvantaged individuals: A qualitative inquiry. Diss. The University of North Carolina at Chapel Hill, 2007. Reference Source

Goodyear-Smith F, Janes R: New Zealand rural primary health care workforce in 2005: more than just a doctor shortage. Aust J Rural Health. 2008; 16(1): 40–46. PubMed Abstract | Publisher Full Text

Graves A: A model for assessment of potential geographical accessibility: a case for GIS. Online J Rural Nurs Health Care. 2009; 9(1): 46-55.

Guay AH: Access to dental care: solving the problem for underserved populations. *J Am Dent Assoc.* 2004; **135**(11): 1599–1605; quiz 1623. PubMed Abstract | Publisher Full Text

Guilcher SJ, Craven CB, McColl MA, et al.: Application of the Andersen's health care utilization framework to secondary complications of spinal cord injury: a scoping review. Disabil Rehabil. 2012; 34(7): 531–41.

PubMed Abstract | Publisher Full Text

Harrison T: Women aging with childhood onset disability. A holistic approach using the life course paradigm. J Holist Nurs. 2003; 21(3): 242–59. PubMed Abstract | Publisher Full Text

Hser YI, Longshore D, Anglin MD: The life course perspective on drug use: a conceptual framework for understanding drug use trajectories. Eval Rev. 2007; 31(6): 515-47.

PubMed Abstract | Publisher Full Text

Hubbell AP: Mexican American women in a rural area and barriers to their ability to enact protective behaviors against breast cancer. Health Commun. 2006; 20(1): 35-44

PubMed Abstract | Publisher Full Text

Hvinden B: The Uncertain Convergence of Disability Policies in Western

Europe. Soc Policy Adm. 2003; 37(6): 609-24.

Publisher Full Text

Iversen MD, Chhabriya RK, Shadick N: Predictors of the use of physical therapy services among patients with rheumatoid arthritis. Phys Ther. 2011; 91(1):

PubMed Abstract | Publisher Full Text

Jo AM, Maxwell AE, Wong WK, et al.: Colorectal cancer screening among underserved Korean Americans in Los Angeles County. J Immigr Minor Health. 2008; 10(2): 119-26.

PubMed Abstract | Publisher Full Text | Free Full Text

Koopmans GT, Lamers LM: Gender and health care utilization: the role of mental distress and help-seeking propensity. Soc Sci Med. 2007; 64(6):

PubMed Abstract | Publisher Full Text

Kosteniuk J, D' Arcy C: Dental service use and its correlates in a dentate population: an analysis of the Saskatchewan population health and dynamics survey, 1999-2000. J Can Dent Assoc. 2006; 72(8): 731.

PubMed Abstract

Lee C, Ayers SL, Kronenfeld JJ: The association between perceived provider discrimination, healthcare utilization and health status in racial and ethnic minorities. Ethn Dis. 2009; 19(3): 330-7.

PubMed Abstract | Free Full Text

Leventhal H, Brissette I, Leventhal EA: The common-sense model of selfregulation of health and illness. In: Cameron LD, Leventhal H, The Self-Regulation of Health and Illness Behaviour, London Routledge, 2003; 42-65.

Levesque JF, Harris MF, Russell G: Patient-centred access to health care: conceptualising access at the interface of health systems and populations. Int J Equity Health. 2013; 12: 18.
PubMed Abstract | Publisher Full Text | Free Full Text

Levy ME, Wilton L, Philips G 2nd, et al.: Understanding structural barriers to accessing HIV testing and prevention services among black men who have sex with men (BMSM) in the United States. AIDS Behav. 2014; 18(5): 972–96. Med Abstract | Publisher Full Text | Free Full Text

López-Cevallos DF, Garside LI: Employment and family conditions are related to health care utilization among foreign-born farmworker men. Hisp Health Care Int. 2013; 11(4): 181-6.

PubMed Abstract | Publisher Full Text

Macintyre S, Ellaway A, Hiscock R, et al.: What features of the home and the area might help to explain observed relationships between housing tenure and health? Evidence from the west of Scotland. Health Place. 2003; 9(3): 207–18. PubMed Abstract | Publisher Full Text

Maxwell RJ: Quality assessment in health. Br Med J (Clin Res Ed). 1984; 288(6428): 1470-2

PubMed Abstract | Publisher Full Text | Free Full Text

McCormack D, Macintosh J: Research with homeless people uncovers a model of health. West J Nurs Res. 2001; **23**(7): 679–97.

PubMed Abstract | Publisher Full Text

Meade MA, Mahmoudi E, Lee SY: The intersection of disability and healthcare disparities: a conceptual framework. Disabil Rehabil. 2015; 37(7): 632–41. Med Abstract | Publisher Full Text

Messer LC, Quinlivan EB, Parnell H, et al.; Barriers and facilitators to testing.

treatment entry, and engagement in care by HIV-positive women of color. AIDS Patient Care STDS. 2013; 27(7): 398-407.

PubMed Abstract | Publisher Full Text | Free Full Text

Mobley LR, Root E, Anselin L, et al.: Spatial analysis of elderly access to primary care services. Int J Health Geogr. 2006; 5: 19.

PubMed Abstract | Publisher Full Text | Free Full Text Nauenberg E, Laporte A, Shen L: Social capital, community size and utilization

of health services: a lagged analysis. Health Policy. 2011; 103(1): 38-46. PubMed Abstract | Publisher Full Text

Norris TL, Aiken M: Personal access to health care: a concept analysis. Public Health Nurs. 2006; 23(1): 59-66.

PubMed Abstract | Publisher Full Text

O'donovan MA, McCallion P, McCarron M, et al.: A narrative synthesis analysis of lifecourse domains within health service utilisation frameworks [Data set]. Zenodo. 2019.

http://www.doi.org/10.5281/zenodo.2613425

Owens J, Dyer TA, Mistry K: People with learning disabilities and specialist services. Br Dent J. 2010; 208(5): 203-5.

PubMed Abstract | Publisher Full Text

Petersen PE: Social inequalities in dental health. Towards a theoretical explanation. Community Dent Oral Epidemiol. 1990; 18(3): 153-158.

PubMed Abstract | Publisher Full Text

Penchansky R, Thomas JW: The concept of access: definition and relationship to consumer satisfaction. Med Care. 1981; 19(2): 127–140.

PubMed Abstract | Publisher Full Text

Phillips KA, Morrison KR, Andersen R, et al.: Understanding the context of healthcare utilization: assessing environmental and provider-related variables in the behavioral model of utilization. Health Serv Res. 1998; 33(3 Pt 1): 571-96. PubMed Abstract | Free Full Text

Robison JT, Moen P: A life-course perspective on housing expectations and shifts in late midlife. Research on Aging. 2000; 22(5): 499-532

Publisher Full Text

Seccombe K: Health insurance coverage and use of services among low-income elders: does residence influence the relationship? J Rural Health. 1995: 11(2): 86-97

PubMed Abstract | Publisher Full Text

Setia MS, Quesnel-Vallee A, Abrahamowicz M, et al.: Access to health-care in Canadian immigrants: a longitudinal study of the National Population Health Survey. Health Soc Care Community. 2011; 19(1): 70–9.

PubMed Abstract | Publisher Full Text | Free Full Text

Shaw EK, Howard J, Clark EC, et al.: Decision-making processes of patients who use the emergency department for primary care needs. J Health Care Poor Underserved. 2013; 24(3): 1288-305.

PubMed Abstract | Publisher Full Text

Slota NEP, Martin D: Methodological considerations in Life Course Theory Research. Disabil Stud Q. 2003; 23(2): 19-29. **Publisher Full Text**

Ståhlnacke K, Söderfeldt B, Unell L, et al.: Changes over 5 years in utilization of dental care by a Swedish age cohort. Community Dent Oral Epidemiol. 2005;

PubMed Abstract | Publisher Full Text

Stoller EP, Grzywacz JG, Quandt SA, et al.: Calling the doctor: a qualitative study of patient-initiated physician consultation among rural older adults. J Aging Health. 2011; 23(5): 782–805. PubMed Abstract | Publisher Full Text | Free Full Text

Thurston WE, Coupal S, Jones CA, et al.: Discordant indigenous and provider frames explain challenges in improving access to arthritis care: a qualitative study using constructivist grounded theory. Int J Equity Health. 2014; 13(1): 46. PubMed Abstract | Publisher Full Text | Free Full Text

Toloo GS, Fitzgerald GJ, Aitken PJ, et al.: Ambulance use is associated with higher self-rated illness seriousness: user attitudes and perceptions. Acad Emerg Med. 2013; 20(6): 576-83.

PubMed Abstract | Publisher Full Text

Tricco AC, Lillie E, Zarin W, et al.: PRISMA Extension for Scoping Reviews (PRISMA-ScR): Checklist and Explanation. Ann Intern Med. 2018; 169(7): 467–473. PubMed Abstract | Publisher Full Text

Uscher-Pines L, Pines J, Kellermann A, et al.: Emergency department visits for nonurgent conditions: systematic literature review. Am J Manag Care. 2013; **19**(1): 47-59.

PubMed Abstract | Free Full Text

Wallace BB, MacEntee MI: Access to dental care for low-income adults: perceptions of affordability, availability and acceptability. J Community Health. 2012; 37(1): 32-39

PubMed Abstract | Publisher Full Text

Ward LS: Preliminary tests of an ecological model of Hispanic farmworker health. Public Health Nurs. 2007; 24(6): 554–564. PubMed Abstract | Publisher Full Text

Witucki JM, Wallace DC: Differences in functional status, health status, and community-based service use between black and white diabetic elders. *J Cult Divers*, 1998; 5(3): 94–100.

PubMed Abstract

Wong WC, Gray SA, Ling DC, et al.: Patterns of health care utilization and health behaviors among street sex workers in Hong Kong. Health Policy. 2006; 77(2): 140-8.

PubMed Abstract | Publisher Full Text

World Health Organization: A Health Telematics Policy in Support of WHO'S Health-For-All Strategy for Global Development. Report of the WHO Group Consultation on Health Telematics 11-16 December, Geneva, 1997. World Health

Reference Source

World Health Organisation: The implications for training of embracing a life course to health. 2000.

World Health Organisation: International Classification of Functioning, Disability and Health (ICF). Geneva: World Health Organisation, 2001.

World Health Organization: Active Aging: A policy Framework. World Health Organisation, 2002

World Health Organization: World report on ageing and health. World Health Organization; 2015.

Reference Source

Yalon-Chamovitz S: Invisible access needs of people with intellectual disabilities: a conceptual model of practice.. Intellect Dev Disabil. 2009; 47(5):

PubMed Abstract | Publisher Full Text

Xiao Q, Savage GT, Zhuang W: A longitudinal model of the dynamics between HMOs' consumer-friendliness and preventive health care utilization. J Health Hum Serv Adm. 2014; 37(1): 76-109.

PubMed Abstract

Open Peer Review

Current Peer Review Status:





Version 1

Reviewer Report 29 July 2019

https://doi.org/10.21956/hrbopenres.13975.r26640

© 2019 Numans M. This is an open access peer review report distributed under the terms of the Creative Commons Attribution Licence, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



Mattijs Numans (iii)



Department of Public Health and Primary Care, Leiden University Medical Centre, Leiden, The Netherlands

This is a very thoroughly carried out systematic review of existing literature on healthcare utilisation from a life course perspective. I have no relevant concerns on the methods and content nor on conclusions, it was very interesting to read. Although shortening would be attractive from a readers perspective, I don't believe much of the content can be missed to understand the conclusions. However, maybe further editing (by native speakers, which I am not) might help. I would be willing to accept this manuscript for indexing as is.

Are the rationale for, and objectives of, the Systematic Review clearly stated? Yes

Are sufficient details of the methods and analysis provided to allow replication by others? Yes

Is the statistical analysis and its interpretation appropriate?

Yes

Are the conclusions drawn adequately supported by the results presented in the review?

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Primary care, mixed methods, population health management.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Reviewer Report 24 April 2019

https://doi.org/10.21956/hrbopenres.13975.r26563

© 2019 Chambers D. This is an open access peer review report distributed under the terms of the Creative Commons Attribution Licence, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



Duncan Chambers

School of Health and Related Research (ScHARR), University of Sheffield, Sheffield, UK

The paper is a scoping/mapping review rather than a full systematic review. As such, some elements of systematic review methodology, for example assessment of risk of bias in included studies, are omitted. Relevant reporting guidelines were followed and the results are available in supplementary files. The authors don't say explicitly whether a protocol was developed in advance and I suggest that this could be clarified. The literature search was thorough, although the search strategy was basic, leading to a large number of records to be screened.

The inclusion criteria were clear, although it was a bit of a stretch to fit them to a PICO-type structure (Table 1). Many other review question formulations are available, for example the authors might like to look at BeHEMoTh (**Be**haviour; **H**ealth context; **E**xclusions; **M**odels **o**r **T**heories; Booth and Carroll; 2015¹).

Methods for identifying frameworks and mapping them against the Andersen model were clearly reported. A minor point is that Figure 2 moves from counting articles to counting frameworks at level 4. How many articles do the 164 frameworks represent?

It may be unavoidable but the process of meta-synthesis comes across as a bit of a 'black box' ('a process of sense-making and developing a cohesive structure'). Can the authors enlarge on this at all?

The presentation and discussion of the review findings follow logically from the structure of the life course domains and Andersen's model. The main conclusion about the need to develop a new model also follows logically from the review findings. The conclusion in the abstract that applying such a framework would be of benefit to researchers etc. may be overstated as this review is descriptive rather than evaluative.

References

1. Booth A, Carroll C: Systematic searching for theory to inform systematic reviews: is it feasible? Is it desirable?. *Health Info Libr J.* 2015; **32** (3): 220-35 PubMed Abstract | Publisher Full Text

Are the rationale for, and objectives of, the Systematic Review clearly stated? Yes

Are sufficient details of the methods and analysis provided to allow replication by others? Yes

Is the statistical analysis and its interpretation appropriate? Not applicable

Are the conclusions drawn adequately supported by the results presented in the review?

Yes

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Systematic reviews, including rapid and mapping reviews; SR-based briefings for decision-makers.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.