

Complete Genome Sequences of Six *Legionella pneumophila* Isolates from Two Collocated Outbreaks of Legionnaires' Disease in 2005 and 2008 in Sarpsborg/Fredrikstad, Norway

Marius Dybwad, Tone Aarskaug, Else-Marie Fykse, Elisabeth Henie Madslie, Janet Martha Blatny

Norwegian Defence Research Establishment FFI, Kjeller, Norway

Here, we report the complete genome sequences of *Legionella pneumophila* isolates from two collocated outbreaks of Legionnaires' disease in 2005 and 2008 in Sarpsborg/Fredrikstad, Norway. One clinical and two environmental isolates were sequenced from each outbreak. The genome of all six isolates consisted of a 3.36 Mb-chromosome, while the 2005 genomes featured an additional 68 kb-episome sharing high sequence similarity with the *L. pneumophila* Lens plasmid. All six genomes contained multiple mobile genetic elements including novel combinations of type-IVA secretion systems. A comparative genomics study will be launched to resolve the genetic relationship between the *L. pneumophila* isolates.

Received 13 October 2016 Accepted 24 October 2016 Published 15 December 2016

Citation Dybwad M, Aarskaug T, Fykse E-M, Henie Madslie E, Blatny JM. 2016. Complete genome sequences of six *Legionella pneumophila* isolates from two collocated outbreaks of Legionnaires' disease in 2005 and 2008 in Sarpsborg/Fredrikstad, Norway. *Genome Announc* 4(6):e01367-16. doi:10.1128/genomeA.01367-16.

Copyright © 2016 Dybwad et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Marius Dybwad, marius.dybwad@ffi.no.

Legionella pneumophila is an opportunistic bacterial pathogen capable of airborne transmission from contaminated freshwater systems to susceptible humans, resulting in a severe pneumonia known as Legionnaires' disease (1). Sequence-based typing is the standard subtyping method for *L. pneumophila* (2), however, recent cost reductions and increased availability have enabled whole-genome sequencing-based subtyping (3).

Here, we report complete genome sequences of *L. pneumophila* isolates from two collocated outbreaks of Legionnaires' disease in 2005 and 2008 in Sarpsborg/Fredrikstad, Norway (Table 1). One clinical and two environmental isolates were sequenced from each outbreak. Additional isolate information has been reported elsewhere (4–7).

Isolates were grown on buffered-charcoal-yeast-extract agar (72 h, 37°C). DNA was purified using Genomic-Tip 100/G (Qiagen, Hilden, Germany). Sequencing was done with PacBio RSII (Menlo Park, CA) and Illumina MiSeq (San Diego, CA). RSII library was prepared using the 20 kb-protocol and size selection

done with BluePippin (9 kb-cutoff). Sequencing was done using P6-C4 chemistry and one single-molecule real-time (SMRT) cell. MiSeq library (300 bp paired-end) was prepared with TruSeq PCR-free protocol. Approximately 90,000 RSII and 3,000,000 MiSeq reads were generated for each isolate. RSII reads were *de novo*-assembled with HGAP_v3.0. Minimus2 (AMOS_v3.1) was used for circularization and RS_Resequencing for mapping of RSII reads. MiSeq reads were mapped onto the final RSII assembly with Bionumerics_v7.6 (Applied Maths, Sint-Martens-Latem, Belgium). Annotation was done with NCBI PGAP_v3.3.

All genomes consisted of a 3.36 Mb-chromosome, while the 2005 genomes featured an additional 68 kb—episome showing high similarity (>97%) with a 39 kb-region of the 60 kb-episome of *L. pneumophila* Lens (Table 1). Average coverage was 284× (RSII) and 466× (MiSeq). All genomes showed conserved synteny with other *L. pneumophila* genomes and highest degree of similarity with Lens (>95%), in agreement with previous sequence-based typing (6). Average G+C content was 38.5% and number of protein-coding genes

TABLE 1 Isolate information and key genomic features of the *Legionella pneumophila* isolates subjected to whole-genome sequencing in this work

Strain/isolate	Source	Country	Yr	SG ^b	ST ^c	Genome size (bp)	Chromosome accession no.	Chromosome size (bp)	Episome accession no.	Episome size (bp)
FFI102	Clinical	Norway	2005	SG1	ST15	3,431,799	CP016868	3,363,654	CP016869	68,145
FFI103	Environmental	Norway	2005	SG1	ST15	3,431,761	CP016870	3,363,616	CP016871	68,145
FFI329	Environmental	Norway	2005	SG1	ST15	3,431,804	CP016874	3,363,658	CP016875	68,146
FFI104	Clinical	Norway	2008	SG1	ST462	3,362,494	CP016872	3,362,494	— ^d	—
FFI105	Environmental	Norway	2008	SG1	ST462	3,363,998	CP016873	3,363,998	—	—
FFI337	Environmental	Norway	2008	SG1	ST462	3,362,463	CP016876	3,362,463	—	—
Lens ^a	Clinical	France	2004	SG1	ST15	3,405,519	CR628337	3,345,687	CR628339	59,832

^a The genome sequence of *L. pneumophila* Lens was obtained from GenBank and included as a reference because sequence-based typing had previously showed that the 2005 isolates had the same ST as Lens (ST15) while the 2008 isolates had a different ST (ST462) but were still closely related to Lens.

^b SG, serotype/serogroup.

^c ST, sequence type.

^d —, not applicable.

2,900, both comparable to Lens (38.4% and 2,932 genes, respectively) (8). *L. pneumophila* often have a dynamic accessory genome consisting of mobile genetic elements, including integrative conjugative elements encoding type-IVA secretion systems (T4ASS), that may facilitate horizontal gene transfer, genome plasticity, and environmental adaptation potential (3, 9). All genomes contained Dot/Icm type-IVB secretion system (T4BSS) and genomic island-associated T4ASS (GI-T4ASS). None of the genomes contained Lvh T4ASS, which is present in Lens (3), while all genomes contained Trb (P-type) T4ASS, which is absent in Lens. The 2008 genomes contained a Trb similar to the one in Corby/Alcoy, while the 2005 genomes contained a Trb similar to the one in Lorraine (3). Only the 2005 genomes contained Tra (F-type) T4ASS, which also is present in Lens. Tra was same as in Lens located on the episome in the 2005 genomes. All genomes contained additional virulence-associated elements including RtxA (10) and clustered regularly interspaced short palindromic repeats (CRISPR)-associated systems (Cas) (11). The RtxA in the 2005 genomes shared high similarity (>92%) with the one in Lens, while the RtxA in the 2008 genomes was more similar to the one in Corby/Alcoy.

Comparative genomics will be used to resolve the genetic relationship between the sequenced isolates. Efforts to increase the availability of *L. pneumophila* genomes may serve as an important catalyst of advancements in this field.

Accession number(s). This whole-genome sequencing (WGS) project was deposited in GenBank under the accession numbers listed in Table 1.

ACKNOWLEDGMENTS

This work was funded by the Norwegian Defence Research Establishment FFI.

The sequencing service was provided by the Norwegian Sequencing Centre hosted by the University of Oslo and supported by the Research Council of Norway and the Southeastern Regional Health Authorities. We thank the Norwegian Institute of Public Health, the Health Authorities of the City of Fredrikstad, and Borregaard Industrier for providing us with the *L. pneumophila* isolates.

FUNDING INFORMATION

This work was funded by the Norwegian Defence Research Establishment FFI.

REFERENCES

- Zhan X-Y, Hu C-H, Zhu Q-Y. 2015. *Legionella* pathogenesis and virulence factors. *Ann Clin Lab Res* 3:1–16.
- Gaia V, Fry NK, Afshar B, Lück PC, Meugnier H, Etienne J, Peduzzi R, Harrison TG. 2005. Consensus sequence-based scheme for epidemiological typing of clinical and environmental isolates of *Legionella pneumophila*. *J Clin Microbiol* 43:2047–2052. <http://dx.doi.org/10.1128/JCM.43.5.2047-2052.2005>.
- Khodr A, Kay E, Gomez-Valero L, Ginevra C, Doublet P, Buchrieser C, Jarraud S. 2016. Molecular epidemiology, phylogeny and evolution of *Legionella*. *Infect Genet Evol* 43:108–122. <http://dx.doi.org/10.1016/j.meegid.2016.04.033>.
- Norwegian Institute of Public Health. 2008. Utbrudd av legionellose i Østfold juni-juli (in Norwegian). <https://www.fhi.no/globalassets/migrering/dokumenter/pdf/rapport-om-legionellautbrudd-2008.pdf>. Accessed September 2016.
- Nygård K, Werner-Johansen Ø, Rønsen S, Caugant DA, Simonsen Ø, Kanestrøm A, Ask E, Ringstad J, Ødegård R, Jensen T, Krogh T, Høiby EA, Ragnhildstveit E, Aaberge IS, Aavitsland P. 2008. An outbreak of legionnaires disease caused by long-distance spread from an industrial air scrubber in Sarpsborg, Norway. *Clin Infect Dis* 46:61–69. <http://dx.doi.org/10.1086/524016>.
- Olsen JS, Aarskaug T, Thrane I, Pourcel C, Ask E, Johansen G, Waagen V, Blatny JM. 2010. Alternative routes for dissemination of *Legionella pneumophila* causing three outbreaks in Norway. *Environ Sci Technol* 44:8712–8717. <http://dx.doi.org/10.1021/es1007774>.
- Wedeg E, Bolstad K, Borgen K, Fritzsønn E, Caugant DA. 2013. Molecular characterization of clinical and environmental isolates of *Legionella pneumophila* in Norway, 2001–2008. *Scand J Infect Dis* 45:59–64. <http://dx.doi.org/10.3109/00365548.2012.710855>.
- Cazalet C, Rusniok C, Brüggemann H, Zidane N, Magnier A, Ma L, Tichit M, Jarraud S, Bouchier C, Vandenesch F, Kunst F, Etienne J, Glaser P, Buchrieser C. 2004. Evidence in the *Legionella pneumophila* genome for exploitation of host cell functions and high genome plasticity. *Nat Genet* 36:1165–1173. <http://dx.doi.org/10.1038/ng1447>.
- Gomez-Valero L, Rusniok C, Jarraud S, Vacherie B, Rouy Z, Barbe V, Medigue C, Etienne J, Buchrieser C. 2011. Extensive recombination events and horizontal gene transfer shaped the *Legionella pneumophila* genomes. *BMC Genomics* 12:536. <http://dx.doi.org/10.1186/1471-2164-12-536>.
- Cirillo SLG, Bermudez LE, El-Etr SH, Duhamel GE, Cirillo JD. 2001. *Legionella pneumophila* entry gene rtxA is involved in virulence. *Infect Immun* 69:508–517. <http://dx.doi.org/10.1128/IAI.69.1.508-517.2001>.
- Gunderson FF, Cianciotto NP. 2013. The CRISPR-Associated gene cas2 of *Legionella pneumophila* is required for intracellular infection of amoebae. *mBio* 4:e00074-00013. <http://dx.doi.org/10.1128/mBio.00074-13>.