

Postnatal Expressional Patterns of Adipose-Associated Molecules in the Mouse Proximal Epididymal Fat

†Ki-Ho Lee

Dept. of Biochemistry and Molecular Biology, College of Medicine, Eulji University, Daejeon 34824, Korea

ABSTRACT : The epididymal fat pad is a male gonadal adipocyte tissue and is histochemically separated into distal and proximal parts. The development of epididymal fat during postnatal period has not been examined in detail. A previous research showed that expression of adipocyte-associated molecules in the distal epididymal fat of mouse is generally increased as postnatally aged. In the present study, expressional patterns of same adipocyte-associated molecules in the mouse proximal epididymal fat at 2, 5, 8, and 12 months of age were studied by quantitative real-time PCR analysis and were compared with those in the distal epididymal fat. The expressional levels of peroxisome proliferator-activated receptor gamma (*Pparg*), lipoprotein lipase (*Lpl*), and fatty acid synthase (*Fasn*) at 5 months of age were significantly lower than those at 2 months of age, while transcript level of leptin (*Lep*) at 5 months was higher than that at 2 months of age. The transcript levels of all molecules at 8 months of age were significantly increased, compared with those at 2 and 5 months of age. At 12 months of age, expression of delta like non-canonical Notch ligand 1 (*Dkl1*) was further significantly increased, while there was no change on the transcript level of *Pparg* and significant decreases of *Fabp4*, *Retn*, *Lpl*, *Lep*, *Fasn*, and adiponectin (*Adipoq*) transcript levels. The current findings show that expressional patterns of molecules associated with adipocyte in the proximal epididymal fat is somewhat different with those of the distal epididymal fat, suggesting the existence of regional variance in the epididymal fat.

Key words : Proximal epididymal fat, Differential gene expression, Adipose tissue, Postnatal development

INTRODUCTION

The epididymal fat is a kind of intraperitoneal white adipose tissue and is located at the head of epididymis (Fig. 1) (Niernalä, 2008; Berry et al., 2013). Like other adipose depot, the epididymal fat is composed of several cell types, including mature adipocytes in variable size, immune cells, endothelial cells, and preadipocytes (Cleary et al., 1977; Dani & Billon, 2012). The lipid-filled adipocyte in epididymal fat region of mouse appears as early as at 7 postnatal age and is originated from undifferentiated

mesenchymal cells (Han et al., 2011). *In vitro* experiment with cells isolated from primitive epididymal fat tissue reveals that the adipogenic differentiation of epididymal progenitor cells requires cell-to-cell and cell-to-matrix interaction and occurrence of normal angiogenesis controlled by macrophage and vascular endothelial growth factor systems (Han et al., 2011).

Unlike other tissues in the body which do not grow over the size at the adult, the fat tissue is capable to continuously expend its mass afterward (Hausman et al., 2001). The increase of epididymal fat weight becomes observed after

Manuscript received September 29, 2019, Received in revised form October 7, 2019, Accepted October 18, 2019

† Corresponding Author : Ki-Ho Lee, Ph.D., Dept. of Biochemistry and Molecular Biology, Eulji University, Daejeon 34824, Korea. Tel: +82-42-259-1643, Fax: +82-42-259-1649, E-mail: kiholee@eulji.ac.kr



This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.




Fig. 1. A representative photograph of the rodent male reproductive tract. Of the male reproductive tract, vas deference, seminal vesicle, and prostate are present, but not labelled.

the birth (Björntorp et al., 1979), and the postnatal growth of epididymal fat is associated with the increase of adipocyte in size and number, depending on the stage of development (Cleary et al., 1977). While the development of epididymal fat during early postnatal period is governed by the increase of adipocyte number, the increase in size of each adipocyte contributes to the expansion of epididymal fat at the later postnatal age (Knittle & Hirsch, 1968; Johnson & Hirsch, 1972; Cleary et al., 1977). In addition, the cellular composition of epididymal fat is substantially changed during postnatal period. The highest ratio of preadipocytes in the epididymal fat is found at the early prenatal period, followed by a drastic decrease of the ratio after the puberty (Björntorp et al., 1979). These findings suggest the presence of distinct regulatory mechanisms at different development stages, in accordance with a whole postnatal growth.

The differentiation and proliferation of preadipocyte into mature adipocyte require expression of a series of adipogenic molecules (Hausman et al., 2001; Dani & Billon, 2012). Differential expression of adipocyte-related genes in the epididymal fat during postnatal development

has been examined by our and other researches (Liu et al., 2011; Lee & Kim, 2018). Even though expressional patterns of some adipocyte-associated genes in the epididymal fat of rat and mouse are generally similar, expression of other genes related with adipocyte differentiation and proliferation is different. For example, expressional level of fatty acid binding protein 4 (*Fabp4*) in the mouse epididymal fat is gradually increased with age (Liu et al., 2011), while transcript level of *Fabp4* in the rat epididymal fat becomes decreased as aged (Lee & Kim, 2018). In contrast, expressional patterns of peroxisome proliferator-activated receptor gamma (*Pparg*) in the mouse and rat during postnatal period are opposite to those of *Fabp4* (Liu et al., 2011; Lee & Kim, 2018). In addition to differential expression of adipocyte-associated genes between species and at different postnatal ages, regional difference of the epididymal fat on gene expression has been observed. Tirard et al. (2007) have shown that histochemically distinct two epididymal regions, proximal and distal, are present. The proximal epididymal fat part is in direct contact with the epididymis, and the distal epididymal fat region is extended toward the abdominal cavity (Tirard et al., 2007). Moreover, different expressional patterns of aldo-keto reductase 1B7, cyclooxygenase 2, and GATA-3 genes between proximal and distal epididymal fat have been observed (Tirard et al., 2007). The comparison of findings from our previous and other researches also supports the existence of regional difference of epididymal fat on gene expression of adipocyte-associated molecules (Liu et al., 2011; Lee & Kim, 2019). However, direct comparison of expressional patterns of adipocyte-associated molecules between different epididymal fat region of mouse during postnatal period has not been carried out.

In the present research, expressional patterns of some adipocyte-associated genes in the proximal epididymal region of mouse at different postnatal ages had been evaluated with quantitative real-time polymerase chain reaction analysis. From the current study, we attempted to

reveal the regional variance in the mouse epididymal fat on gene expression of adipocyte-associated molecules.

MATERIALS AND METHODS

1. Experimental design and tissue collection

Male C57BL/6N mice at 1 month of postnatal age were obtained from Samtako (OSan, Korea). At the arrival, the mice were randomly assigned into four different groups, a young adult group (2 months of age), two fully mature adult groups (5 and 8 months of age), and an old group (12 months of age), and each experimental group had six mice. The animals were maintained in controlled conditions and permitted for food and water *ad libitum*. This work was carried out in accordance with the guide for the care and use of laboratory animals of National Research Council in Korea.

At the proper postnatal age, the animal was euthanized by CO₂ stunning in a chamber. An incision was made on lower abdominal part, and the male reproductive tract was pulled out through the incision (Fig. 1). In cold-PBS buffer, the whole epididymal fat pad was dissected out from the rest of male reproductive tract. The proximal epididymal fat surrounding the efferent ductules was carefully separated from the distal epididymal fat, the peripheral part of epididymal fat pad. Two parts of the epididymal fat pad were rapidly frozen in liquid nitrogen and separately stored in -80°C for further use.

2. Extraction of total RNA from the proximal epididymal fat and generation of complementary DNA (cDNA)

One milliliter of TRI REAGENT solution (Molecular Research Center, Inc, Cincinnati, OH) was applied to each proximal epididymal fat sample for homogenization. Then, chloroform and isopropanol were sequentially added to extract total RNA from tissue homogenate, and an acquired total RNA pellet was re-dissolved in DEPC-dH₂O. The

yield and concentration and quality of total RNA were evaluated by the NanoDrop Lite spectrophotometry (Thermo Scientific, Wilmington, DE) and 1.2% agarose gel electrophoresis, respectively.

One microgram of total RNA was used to construct the first-stranded complementary DNA (cDNA) by reverse transcription (RT) reaction in iScript™ Reverse transcription Supermix for RT-qPCR (Bio-Rad Laboratories, Inc., Hercules, CA). The total RNA was mixed with oligo-dT primer, RTase, 5X buffer, and nuclease-free dH₂O, and the final volume of a RT reaction was 20 µL. The RT reaction was performed in 25°C for 5 min, 46°C for 20 min, and 95°C for 1 min, and generated cDNA was directly utilized for quantitative real-time polymerase chain reaction (PCR).

3. Performance of quantitative real-time PCR and statistical analysis

Table 1 provides information of oligonucleotide primers for molecules analyzed by quantitative real-time PCR. Primer 3 software (<http://www.bioneer.co.kr/cgi-bin/primer/primer3.cgi>; Whitehead Institute/MIT Center for Genomes Research, USA) was used to design the PCR primers.

One microliter of cDNA was mixed with 10 pmol of each primer, 7 µL of iQ™ SYBR® Green Supermix (Bio-Rad Laboratories), and dH₂O to make 20 µL of PCR mixture. The condition of PCR was a pre-denaturation at 95°C for 5 min, cycles of a denaturation at 95°C for 30 sec, an annealing at T_m for 30 sec, and an extension at 72°C for 30 sec, and an additional extension at 72°C for 10 min. The expected size of PCR product was confirmed with 1.2% agarose gel electrophoresis.

As an internal control of PCR, 18S ribosomal RNA (*Rn18s*) was added for quality control purpose. In order to get a mean and standard error for expressional level of a particular gene at an age, independently quadrupled or quintupled RT reactions and PCRs were executed. The expressional levels of target genes were normalized with that of *Rn18s* and were expressed as a relative ratio against

Table 1. Oligonucleotide primer information

Gene	Primer sequence (5' → 3')	T _m (°C)	PCR product size (bps)
<i>Adipoq</i> (NM_009605.5)	(F) AGGCCGTTCTCTTCACCTACG (R) GACTTGGTCTCCCACCTCCAGA	62	95
<i>Dlk1</i> (NM_010052.5)	(F) GGCCATCGTCTTTCTCAACA (R) CTCCTCATCACCAGCCTCCT	60	181
<i>Fabp4</i> (NM_024406.3)	(F) CAAGCCCAACATGATCATCAGC (R) CACGCCAGTTTGAAGGAAATC	60	97
<i>Fasn</i> (NM_011480.3)	(F) GTGTGGACATGGTCACAGATG (R) GACCGCTTGGGTAATCCATA	60	67
<i>Lep</i> (NM_008493.3)	(F) TTCACACACGCAGTCGGTAT (R) TCATTGGCTATCTGCAGCAC	55	165
<i>Lpl</i> (NM_008509.2)	(F) TCAGAGCCAAGAGAAGCAGCA (R) TTGTGTTGCTTGCCATCCTCA	60	118
<i>Pparg</i> (U01841.1)	(F) GATTCATGACCAGGGAGTTC (R) GACTCTGGGTGATTCAGCTT	60	247
<i>Retn</i> (NM_022984.3)	(F) TTCCTTGCCCTGAACTGCT (R) TCAAGACTGCTGTGCCTTCT	55	187
<i>Rn18s</i> (NR_03278.3)	(F) ACTCAACACGGGAAACCTCA (R) AACCAGACAAATCGCTCCAC	55	123

Adipoq, adiponectin; *Dlk1*, delta like non-canonical Notch ligand 1; *Fabp4*, fatty acid binding protein 4; *Fasn*, fatty acid synthase; *Lep*, leptin; *Lpl*, lipoprotein lipase; *Pparg*, peroxisome proliferator-activated receptor, gamma; *Retn*, resistin; *Rn18s*, 18S ribosomal RNA.

Rn18s.

To determine statistical significance of expressional levels of the particular molecule among experimental groups at a postnatal age, one-way ANOVA analysis was first carried out. If there was a statistical significance present, Duncan's test, a post-hoc analysis, was followed. If *p* value was less than 0.05, it was considered that there was significant difference among experimental groups, presented in different letters.

RESULTS

1. Expression of *Pparg* and *Dlk1* in the mouse proximal epididymal fat pad at different postnatal ages

The expression of *Pparg* was significantly decreased at 5 months of age, compared with that at 2 months of age

(Fig. 2A). However, a surge, over 5-folds increase, of *Pparg* transcript level was observed at 8 months of age (Fig. 2A). There was no significant change of *Pparg* expressional level at 12 months of age, compared with at 8 months of age (Fig. 2A).

The change of transcript level of *Dlk1*, delta like non-canonical Notch ligand 1 (*Dlk1*) at 5 months of age was not significantly different with that at 2 months of age (Fig. 2B). But, a significant increase of *Dlk1* transcript level was detected at 8 months of age (Fig. 2B). A further increase of *Dlk1* expressional level was observed at 12 months of age (Fig. 2B).

2. Expression of *Fasn* and *Lpl* in the mouse proximal epididymal fat pad at different postnatal ages

Similar to *Pparg*, a significant decrease of fatty acid




Fig. 2. Expression of *Pparg* and *Dlk1* in the mouse proximal epididymal fat pad at different postnatal ages. The relative levels of *Pparg* (A) and *Dlk1* (B) transcripts are shown here. Different letters indicate statistical significances at $p < 0.05$. M, month; *Pparg*, peroxisome proliferator-activated receptor; *Dlk1*, delta like non-canonical Notch ligand 1.

synthase (*Fasn*) transcript level at 5 months of age was found (Fig. 3A). The expression *Fasn* was however greatly increased at 8 months of age, compared with that at 5 months of age (Fig. 3A). At 12 months of age, the level of *Fasn* transcript was significantly lower than that at 8 months of age, but still higher than that at 2 and 5 months of age (Fig. 3A).

The level of lipoprotein lipase (*Lpl*) transcript at 5 months of age was significantly lower than that at 2 months of age (Fig. 3B). But, the expressional level of *Lpl* at 8 months of age was much higher than those at 2 and 5 months of age (Fig. 3B). The transcript level of *Lpl* at 12 months of age was significantly decreased, but was higher than those at 2 and 5 month of age (Fig. 3B).




Fig. 3. Expression of *Fasn* and *Lpl* in the mouse proximal epididymal fat pad at different postnatal ages. The relative levels of *Fasn* (A) and *Lpl* (B) transcripts are shown here. Different letters indicate statistical significances at $p < 0.05$. M, month; *Fasn*, fatty acid synthase; *Lpl*, lipoprotein lipase.

3. Expression of *Lep* and *Retn* in the mouse proximal epididymal fat pad at different postnatal ages

The lowest level of leptin (*Lep*) in the proximal epididymal fat of mouse was detected at 2 months of age (Fig. 4A). The abundance of *Lep* transcript was significantly increased at 5 months of age, and an additional extension of *Lep* transcript level was observed at 8 months of age (Fig. 4A). However, expressional level of *Lep* at 12 months of age was significantly lower than that at 8 months of age and comparable with that at 5 months of age (Fig. 4A).

The level of *Retn* expression was not significantly changed between 2 months and 5 months of age (Fig. 4B). But, the level of *Retn* transcript was significantly increased at 8 months of age, followed by a considerable drop at 12 months of age, which was yet higher than those at 2 months and 5 months of age (Fig. 4B).

4. Expression of *Adipoq* and *Fabp4* in the mouse proximal epididymal fat pad at different postnatal ages

Expressional level of adiponectin (*Adipoq*) at 5 months of age was not significantly changed, compared with that at 2 months of age (Fig. 5A). But, a huge increase of *Adipoq* transcript level was detected at 8 months of age

(Fig. 5A). The level of *Adipoq* transcript was significantly lower than that at 8 months of age but still higher than those at 2 months and 5 months of age (Fig. 5A).

The expressional pattern of *Fabp4* was quietly similar with that of *Adipoq*, as shown in Fig. 5B. The level of *Fabp4* at 5 months of age was not significantly different with that at 2 months of age (Fig. 5B). However, over 100-fold increases of *Fabp4* transcript level was observed at 8 months of age (Fig. 5B). The level of *Fabp4* transcript at 12 months of age was significantly lower than that at 8 months of age but significantly higher than those at 2 months and 5 months of age (Fig. 5B).

DISCUSSION

Expressional levels of adipocyte-associated molecules in the proximal epididymal fat at the old age were mostly higher than that at the early age, even though expressional patterns of some genes were different with those of other genes. Except *Dkl1*, the highest transcript levels of other adipocyte-associated molecules in the proximal epididymal fat were detected at 8 months of postnatal age.

The leptin, resistin, and adiponectin are most prominent




Fig. 4. Expression of *Lep* and *Retn* in the mouse proximal epididymal fat pad at different postnatal ages. The relative levels of *Lep* (A) and *Retn* (B) transcripts are shown here. Different letters indicate statistical significances at $p < 0.05$. M, month; *Lep*, leptin; *Retn*, resistin.




Fig. 5. Expression of *Adipoq* and *Fabp4* in the mouse proximal epididymal fat pad at different postnatal ages. The relative levels of *Adipoq* (A) and *Fabp4* (B) transcripts are shown here. Different letters indicate statistical significances at $p < 0.05$. M, month; *Adipoq*, adiponectin; *Fabp4*, fatty acid binding protein 4.

adipocytokines secreted from adipose tissue, primarily involving in regulation of energy balance among tissues at various levels (Cao, 2014). It is well recognized that leptin regulates the release of secretory molecules from hypothalamus for control of food intake and energy spend (Cao, 2014). Leptin also exerts its effect on the male reproduction, including regulation of spermatogenesis, development of puberty, and sperm maturation (Zhang & Gong, 2018). Liu et al. (2011) have shown an increase of *Lep* in the epididymal fat of aging mouse, and our previous research has also revealed gradual increase of *Lep* in the distal epididymal fat of mouse with age (Lee & Kim, 2019). However, the present study detected that the level of *Lep* transcript in the proximal epididymal fat was significantly decreased at 12 months, compared to the level at 8 months of age. Together, these results suggest that an overall increase of *Lep* expression in the mouse epididymal fat at 12 months of age is likely influenced more by expressional change of *Lep* in the distal epididymal fat than by expressional change of *Lep* in the proximal epididymal fat.

Expressional levels of *Adipoq* and *Retn* in the rat whole epididymal fat are increased with age (Lee & Kim, 2018)

and in the mouse distal epididymal fat (Lee & Kim, 2019). Interestingly, the present research shows the highest transcript levels of *Adipoq* and *Retn* in the mouse proximal epididymal fat at 8 months of age, followed by significant decreases at 12 months of age. Ouchi et al. (2003) have shown a decrease of *Adipoq* expression in adipose tissue as body mass is increased. Indeed, Liu et al. (2011) have revealed a decreased expressional level of *Adipoq* in the mouse epididymal fat at 14 months of age, compared with that at 6 months of age. However, from our previous findings, the level of *Adipoq* transcript in the mouse distal epididymal fat at 12 months of age is significantly higher than that at 5 months of age and as close as that at 8 months of age (Lee & Kim, 2019). These observations suggest that an increase of *Adipoq* expression in the mouse epididymal fat until 8 months of age could be a synergic effect of proximal and distal epididymal fat, while a decrease of *Adipoq* transcript level at 12 months of age would be a results from the proximal epididymal fat overwhelming the effect observed in the distal epididymal fat. Expressional patterns of *Retn* in different mouse epididymal fat parts of mouse are very close to those of *Adipoq*. Thus, it is speculated that overall level of *Retn* transcript at 12 months of age

would be lower than that at 8 months of age. It is considered that adiponectin and resistin exert opposite effects on insulin-resistance and atherogenesis (Cao, 2014). However, based on the findings of similar expressional patterns of *Adipoq* and *Retn* in the epididymal fat, it is assumed that adiponectin and resistin would play a physiological role in the epididymal fat and/or male reproductive tract in unknown manner.

The expressional level of *Fabp4* in the mouse epididymal fat is decreased with age (Liu et al., 2011). However, our previous study has shown a gradual increase of *Fabp4* in the distal epididymal fat of mouse (Lee & Kim, 2019). It has been suggested that such disagreement on expressional patterns of *Fabp4* could be due to differential expression of *Fabp4* between the proximal and distal epididymal fat part (Lee & Kim, 2019). Indeed, the present study shows that the expressional level of *Fabp4* at 12 months of age is significantly lower.

Than that at 8 months of age. Thus, it is reasonable to consider that a decrease of *Fabp4* transcript level in the mouse epididymal fat at the old age would be a result of a reduction of *Fabp4* expression in the proximal part of mouse epididymal fat. As a fatty acid binding protein, FABP4 promotes the accumulation of lipid droplets in adipogenic cells (Kletzien et al., 1992). Because the size of fat cells becomes enlarged and the storage of lipid components are increased as aged (Cleary et al., 1977), it is thought that expression and/or activity of FABP4 in the epididymal fat at the old age would be higher than that at the young age. A future research is suggested to examine posttranscriptional regulation of *Fabp4* expression to retain FABP4 protein and/or activity level in the epididymal fat at the old age.

Dlk-1, a.k.a. preadipocyte factor-1(*Pref-1*), is an inhibitory factor of adipogenesis to suppress the differentiation of preadipocyte into mature adipocyte (Wang & Sul, 2009). Thus, expression of *Dlk-1* is higher in preadipocyte than in mature adipocyte (Wang & Sul, 2009). In the rat epididymal fat, expressional level of *Dlk-1* is extraordinarily high at

the early postnatal age, and basal level of *Dlk-1* transcript is detectable at 5 months of age (Lee & Kim, 2018). However, the expression of *Dlk-1* in the rat epididymal fat at old age is significantly increased (Lee & Kim, 2018). Interestingly, our previous (Lee & Kim, 2019) and current researches have demonstrated the basal expression of *Dlk-1* at 2 and 5 months of age and surges of *Dlk-1* expressional level at 8 and 12 months of age in the mouse distal and proximal epididymal fat regions. These observations imply that *Dlk-1* in the epididymal fat would act a factor with unknown function, other than inhibiting differentiation of preadipocyte to adipocyte. Further researches are suggested to examine possible regulatory effect of *Dlk-1* derived from the epididymal fat on the testicular functions.

Of several adipogenic factors, *Pparg* is a key regulator for differentiation of preadipocyte into mature adipocyte and preservation of differentiated status of mature adipocyte (Tamori et al., 2002). The expressional level of *Pparg* in the distal epididymal fat part of mouse is gradually increased with aging (Lee & Kim, 2019). Even though a similar expressional pattern of *Pparg* is also found in the proximal epididymal fat from the present research, a significant decrease of *Pparg* at 5 months of age in the proximal epididymal fat is a notable observation different with results acquired from the distal epididymal fat region. There is no clear explanation suggested for such change on *Pparg* expressional pattern. It is possible that expressional decrease of *Pparg* at 5 months of age in the proximal epididymal part would be negligible in the level of a whole epididymal fat. This suggestion is partly supported by Liu et al.'s findings (2011), showing a significantly higher level of *Pparg* transcript at 14 months of age than that at 6 months of age.

Lipolysis, a process of triglyceride hydrolysis, and lipogenesis, a synthetic pathway to store fatty acid, are found in adipocyte, and these two opposite mechanisms are regulated by various molecules, including LPL and FASN, respectively. Even though expressional level of *Lpl* in the rat epididymal fat is decreased as aged (Kirkland et

al., 1993), our previous (Lee & Kim, 2019) and current researches have shown the higher level of *Lpl* transcript level in both proximal and distal epididymal fat regions at the old age than that at early postnatal age. It is notable that expressional level of *Lpl* at 5 months of age in the proximal epididymal fat region is significantly lower than that at 2 months of age. This result is absolutely opposite to our expectation described from the previous report (Lee & Kim, 2019), assuming a significant increase of *Lpl* expression in the proximal epididymal fat at 5 months of age. Moreover, expressional level of *Lpl* in the proximal epididymal fat at 12 months of age is unexpectedly lower than at 8 months of age. Expressional pattern of *Fasn* in the proximal epididymal fat is quite similar with that of *Lpl*. However, unlike in the distal epididymal fat (Lee & Kim, 2019), expressional level of *Fasn* in the proximal epididymal fat at 12 months of age is substantially reduced, even though significant decreases of *Fasn* transcript level at 5 months of age are noticeable in both proximal and distal epididymal fat regions. Such unanticipated expressional patterns of *Lpl* and *Fasn* in the mouse epididymal fat at 5 and 12 months of age are not understandable at this point. A functional study of *Lpl* and *Fasn* in the mouse epididymal fat at a given postnatal age would provide possible explanation for such fluctuation of postnatal expression.

In summary, the current study exhibits the differential patterns of gene expression of adipocyte-associated molecules between the proximal and distal epididymal fat parts of mouse during postnatal period. The function of epididymal fat pad is not clearly elucidated yet. Chu et al. (2010) have demonstrated that the surgical removal of epididymal fat causes the cease of spermatogenesis in the testis, without a change of serum testosterone or luteinizing hormone concentration. These observations imply that locally-acting factor(s) derived from the epididymal fat would influence on the regulation of spermatogenesis in the testis. Our previous (Lee & Kim, 2019) and current researches have revealed expressional patterns of adipocyte-associated

molecules in different two epididymal fat regions of mouse during postnatal period. Such observations suggest the existence of different regulatory mechanisms for adipocyte-associated gene expression in these epididymal fat regions. Even though exact functions of these adipocyte-associated molecules in the epididymal fat have not been elucidated, it is carefully suggested that the secretory molecules derived from the epididymal fat, such as leptin, resistin, and adiponectin, would influence on the regulation of testicular functions.

ORCID

Ki-Ho Lee <https://orcid.org/0000-0002-3495-5126>

CONFLICT OF INTEREST

The authors declare no potential conflict of interest.

ACKNOWLEDGEMENTS

This research was partly supported by Basic Science Research Program through the National Research Foundation of Korea (NRF) funded by the Ministry of Education (2017R1D1A 1B03030209).

AUTHOR CONTRIBUTIONS

The article is prepared by a single author.

REFERENCES

- Berry DC, Stenesen D, Zeve D, Graff JM (2013) The developmental origins of adipose tissue. *Development* 140:3939-3949.
- Björntorp P, Karlsson M, Gustafsson L, Smith U, Sjöström L, Cigolini M, Storck G, Pettersson P (1979) Quantitation of different cells in the epididymal fat pad of the rat. *J*

- Lipid Res 20:97-106.
- Cao H (2014) Adipocytokines in obesity and metabolic disease. *J Endocrinol* 220:T47-T59.
- Chu Y, Huddleston GG, Clancy AN, Harris RB, Bartness TJ (2010) Epididymal fat is necessary for spermatogenesis, but not testosterone production or copulatory behavior. *Endocrinology* 151:5669-5679.
- Cleary MP, Greenwood MR, Brasel JA (1977) A multifactor analysis of growth in the rat epididymal fat pad. *J Nutr* 107:1969-1974.
- Dani C, Billon N (2012) Adipocyte precursors: Developmental origins, self-renewal, and plasticity. In: Symonds ME (ed), *Adipose Tissue Biology*. Springer, New York, pp 1-16.
- Han J, Lee JE, Jin J, Lim JS, Oh N, Kim K, Chang SI, Shibuya M, Kim H, Koh GY (2011) The spatiotemporal development of adipose tissue. *Development* 138:5027-5037.
- Hausman DB, DiGirolamo M, Bartness TJ, Hausman GJ, Martin RJ (2001) The biology of white adipocyte proliferation. *Obes Rev* 2:239-254.
- Johnson PR, Hirsch J (1972) Cellularity of adipose depots in six strains of genetically obese mice. *J Lip Res* 13:2-11.
- Kletzien RF, Foellmi LA, Harris PK, Wyse BM, Clarke SD (1992) Adipocyte fatty acid-binding protein: regulation of gene expression *in vivo* and *in vitro* by an insulin-sensitizing agent. *Mol Pharmacol* 42:558-562.
- Kirkland JL, Hollenberg CH, Gillon WS (1993) Ageing, differentiation, and gene expression in rat epididymal preadipocytes. *Biochem Cell Biol* 71:556-561.
- Knittle JL, Hirsch J (1968) Effect of early nutrition on the development of rat epididymal fat pads: Cellularity and metabolism. *J Clin Invest* 47:2091-2098.
- Lee KH, Kim NH (2018) Expressional patterns of adipocyte-associated molecules in the rat epididymal fat during postnatal development period. *Dev Reprod* 22:351-360.
- Lee KH, Kim NH (2019) Differential expression of adipocyte-related molecules in the distal epididymal fat of mouse during postnatal period. *Dev Reprod* 23:213-221.
- Liu LF, Shen WJ, Ueno M, Patel S, Kraemer FB (2011) Characterization of age-related gene expression profiling in bone marrow and epididymal adipocytes. *BMC Genomics* 12:212.
- Niemälä S, Miettinen S, Sarkanen JR, Ashammakhi N (2008) Adipose tissue and adipocyte differentiation: Molecular and cellular aspects and tissue engineering applications. In: Ashammakhi N, Reis R, Chiellini F (ed), *Topics in Tissue Engineering*, Vol. 4. Biomaterials and Tissue Engineering Group, Porto, Portugal. pp 1-26.
- Ouchi N, Kihara S, Funahashi T, Matsuzawa Y, Walsh K (2003) Obesity, adiponectin and vascular inflammatory disease. *Curr Opin Lipidol* 14:561-566.
- Tamori Y, Masugi J, Nishino N, Kasuga M (2002) Role of peroxisome proliferator-activated receptor-gamma in maintenance of the characteristics of mature 3T3-L1 adipocytes. *Diabetes* 51:2045-2055.
- Tirard J, Gout J, Lefrancois-Martinez AM, Martinez A, Begeot M, Naville D (2007) A novel inhibitory protein in adipose tissue, the aldo-keto reductase AKR1B7: Its role in adipogenesis. *Endocrinology* 148:1996-2005
- Wang Y, Sul HS (2009) Pref-1 regulates mesenchymal cell commitment and differentiation through Sox9. *Cell Metab* 9:287-302.
- Zhang J, Gong M (2018) Review of the role of leptin in the regulation of male reproductive function. *Andrologia* 50:e12965.