

# Effect of *Helix aspersa* extract on TNF $\alpha$ , NF- $\kappa$ B and some tumor suppressor genes in breast cancer cell line Hs578T

Ibtissem El Ouar, Cornelia Braicu<sup>1,2</sup>, Dalila Naimi, Alexendru Irimie<sup>2,3,4</sup>, Ioana Berindan-Neagoe<sup>1,2,5</sup>

Laboratory of Microbiological Engineering and Application, University Mentouri Constantine, Algeria, <sup>1</sup>Department of Functional Genomics and Experimental Pathology, Cancer Institute "Ion Chiricuta", Cluj-Napoca, <sup>2</sup>Research Center for Functional Genomics, Biomedicine and Translational Medicine, <sup>3</sup>Department of Surgery, <sup>4</sup>Department of Surgical Oncology, <sup>5</sup>Department of Immunology, Iuliu Hatieganu. University of Medicine and Pharmacy, Cluj-Napoca, Romania

Submitted: 30-03-2015

Revised: 26-06-2015

Published: 18-04-2017

## ABSTRACT

**Background:** The garden snail, *Helix aspersa*, is a big land snail widely found in the Mediterranean countries. It is one of the most consumed species and widely used in zootherapy. **Objective:** The present study was carried out to investigate for the first time the antitumor activity of an aqueous extract from *Helix aspersa*. **Materials and Methods:** The effect of *H. aspersa* extract was studied on a triple negative breast cancer cell line Hs578T. Firstly, the morphological changes and the mode of cell death induced by the extract have been evaluated by microscopy and acridine orange/ethidium bromide staining. The effect of the extract at dilution 0.1% and 1% was then tested on some genes, regulators of cell death and proliferation like tumor necrosis factor  $\alpha$  (TNF $\alpha$ ), NF- $\kappa$ B, and the tumor suppressor genes P53 and PTEN. **Results:** Data demonstrate that the extract induces necrosis in tumor cells. It enhances significantly the expression of TNF $\alpha$ ; mRNA levels were 20 and 10 times more important in treated cells compared to nontreated cells. NF- $\kappa$ B and PTEN were inhibited with the dilution 1% after 8 and 24 hours of treatment. P53 expression was further inhibited but only with the highest dose, after 4, 8, and 24 hours. **Conclusion:** Our results show that *H. aspersa* extract has an antitumor activity against Hs578T cells; it is a potent stimulator for TNF $\alpha$  and a good inhibitor for NF- $\kappa$ B.

**Key words:** Breast cancer, *Helix aspersa*, necrosis, NF-KB, TNF $\alpha$

**Abbreviations used:** AO: acridine orange; Bcl-2: B cell lymphoma 2. cDNA: complementary DNA; ELISA: enzyme linked immunosorbent assay; EB: ethidium bromide; IC50: the half maximal inhibitory concentration; mRNA: messenger RNA. MAPK: mitogen-activated protein kinase; NF- $\kappa$ B: nuclearfactor $\kappa$ B; PBS: phosphate buffered saline. PI3K: phospho-inositol 3 kinase; PTEN: phosphatase and tensin homolog; ROS: reactive oxygen species. RT-PCR: reverse transcription polymerase chain reaction; TNF $\alpha$ : tumor necrosis factor alpha. TNFR1: TNF receptor-1; TP53: tumor protein 53

## Correspondence:

Dr. EL OUAR Ibtissem.  
Faculty of Nature And Life Sciences,  
University Mentouri, road Ain el  
bey -Constantine, Algeria.  
E-mail: ibtissem\_elouar@yahoo.fr  
DOI: 10.4103/0973-1296.204618

Access this article online

Website: www.phcog.com

Quick Response Code:



## INTRODUCTION

After cardiovascular disease, cancer is the second leading cause of death in the world.<sup>[1]</sup> The treatment is basically on synthetics and chemotherapeutics; these kinds of treatments cause various harmful side effects to human beings.<sup>[2]</sup> For this reason, the search for new natural and safe drugs is the aim of the different laboratories in the world. Plant and marine species are the major source of natural drugs; few researches have been interested in purifying bioactive molecules from other terrestrial consumed species like snails.

Snails are a member of mollusk, the second largest phylum in the animal kingdom with about 100,000 living species. One of the most consumed species is the garden common snail, *Helix aspersa*; it is a small nutrient with high contents of proteins and minerals and low contents of fat and cholesterol.<sup>[3,4]</sup>

Snails have been used in medicine since antiquity and are prepared by several methods; they are recommended for stomach pain, vertigo, nephritis, and respiratory and cardiovascular diseases.<sup>[5]</sup> Nowadays, lectins from snails are used as a marker for metastatic tissues in breast and colon cancers.<sup>[6,7]</sup>

Initially, we have observed that the *Helix aspersa* extract had an antitumor effect on breast cancer cell line Hs578T.<sup>[8]</sup> This finding led us to investigate its mode of action and understand the mechanism involved. In particular, we focused on the expression of TNF $\alpha$ , a stimulator of the extrinsic pathway of apoptosis,<sup>[9]</sup> NF- $\kappa$ B, a cell proliferation regulator<sup>[10]</sup> and the tumor suppressor genes P53 and PTEN.<sup>[11]</sup>

TNF  $\alpha$  is a key cytokine that plays an important role in inflammation, immunity, and diverse cellular events. It modulates cell proliferation, necrosis, and apoptosis.<sup>[12]</sup> The major source of TNF $\alpha$  is activated macrophages; it can also be produced by other cells including fibroblasts, astrocytes, kupffer cells, smooth muscles, keratinocytes, and tumor cells. It contributes to the metastatic process, invading the host tissues, penetrating to blood vessel's endothelium and establishing tumor cell growth at secondary sites.<sup>[13]</sup>

NF- $\kappa$ B, a transcription factor that plays important roles in cancer development, appears to be involved in the regulation of cell proliferation, apoptosis control, angiogenesis promotion, and invasion/metastasis stimulation.<sup>[14]</sup>

P53 is encoded by the TP53 gene, located at 17p13; it plays an important role in mediating cell response to various stresses,<sup>[15]</sup> mainly inducing or repressing a number of genes involved in cell cycle arrest, senescence, apoptosis, DNA repair, and angiogenesis.<sup>[16]</sup> Indeed, P53 mutation is

This is an open access article distributed under the terms of the Creative Commons Attribution-Non Commercial-Share Alike 3.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as the author is credited and the new creations are licensed under the identical terms.

For reprints contact: reprints@medknow.com

**Cite this article as:** El Ouar I, Braicu C, Naimi D, Irimie A, Berindan-Neagoe I. Effect of *Helix aspersa* extract on TNF $\alpha$ , NF- $\kappa$ B and some tumor suppressor genes in breast cancer cell line Hs578T. Phcog Mag 2017;13:281-5.

associated with more aggressive disease and worse overall survival, like breast and prostate cancers.<sup>[17]</sup>

Next to P53, PTEN is the most common tumor suppressor to be lost or inactivated in human cancers like glioblastoma, prostate, and breast cancers.<sup>[18]</sup> The PTEN gene encodes a dual specificity lipid and protein phosphatase. PTEN modulates cell growth, migration, and survival by antagonizing the phospho-inositol 3 kinase (PI3K)/AKT signaling.<sup>[19]</sup>

## MATERIALS AND METHODS

**Cell culture:** Human breast cancer Hs578T cell line was obtained from the laboratory of functional genomics and experimental pathology of the oncologic institute "Ion Chiricuta" (Cluj Napoca- Roumania). The cells were cultured at 37°C and humidified atmosphere of 5% CO<sub>2</sub>, in Dulbecco's Modified Eagle's Medium high glucose supplemented with 1% nonessential amino acids, 1% L-glutamine (200 mM), 1% gentamicin (10 mg/ml), 1% insulin, and 10% fetal bovine serum. All reagents were purchased from Sigma Aldrich, Germany.

**Preparation of *H. aspersa* extract:** The *H. aspersa* snails were maintained fasting during at least 2 weeks in order to empty their gut. Before use the snails were washed in 3 or 4 bath of 10% NaCl solution (to completely eliminate their mucus). After removing the shell, the snails were homogenized. Three volumes of water per one volume of wet tissues were then added to the homogenate. The crude extract was filtered and centrifuged for 10 minutes at 5000 g. The supernatants were collected and sterilized using 22 µm Millipore filter and served as extract to treat cells. The final concentration of the extract was 0,33 g/l.

**Acridine orange/ Ethidium bromide staining:** Cells were seeded for 24 hours in the presence of *H. aspersa* extract; at 0.1% and 1% dilutions of crude extract. Cells were stained by Acridine orange (AO)/ethidium bromide (EB), the dye mix for staining was 100 µg/ml AO and 100 µg/ml EB in PBS (pH=7).<sup>[20]</sup>

## RT-PCR

To study gene expression, cells were treated with the extract at different times 4, 8 and 24 hours.

Total cellular RNA was isolated using TRIagen (Sigma, Germany) and converted to cDNA with The Random Hexamer Transcriptor First Strand cDNA Synthesis Kit (Roche Diagnosis, Germany) following manufacturer's instruction.

Quantitative RT-PCR was performed using TaqMan Master Kit and light cycler. All samples samples were run in triplicates and the β actin was amplified as an internal control. Primers used are presented in Table 1.

Quantification of relative gene expression was done using the competitive threshold cycle C<sub>T</sub> method. C<sub>T</sub> values were averaged for triplicate wells and subtracted from corresponding C<sub>T</sub> of internal reference RNA to obtain ΔC<sub>T</sub> values. The averaged control ΔCT was subtracted from the experimental ΔC<sub>T</sub> to yield ΔΔC<sub>T</sub>. The fold change was calculated as 2<sup>-ΔΔCT</sup> for experimental versus control.<sup>[21]</sup>

**ELISA Assay:** The concentration of TNFα in culture medium was determined by ELISA assay and using TNFα human Boster immunoleader kit (Cliniscience France).

**STATISTICS** Data were expressed as mean ± SD from at least three separate experiments performed on triplicate samples. The differences between experimental conditions and controls were analyzed using t test ( $p < 0.05$  is considered statistically significant). Statistical analyses were carried out using Graph Pad Prism software (free trial).


## RESULTS

### Morphological identification of cell death


Cells were treated with two doses of *H. aspersa* extract, the first dose corresponded to the 1% dilution (1:100 dilution of filtered extract) which is approximately equal to IC<sub>50</sub>.<sup>[8]</sup> The second dose was 10 times less than the first one.

To determine if *H. aspersa* extract induces necrosis or apoptosis in breast cancer cell line Hs578T, cells were stained by AO/EB.

AO/EB staining provides a reliable method to measure cells in different compartments of cell death; live, apoptotic, and necrotic cells were differentiated using fluorescence microscopy. AO (excitation: 502 nm; emission: 525 nm) permeates all cells and makes the nuclei appear green. It will also enter acidic compartments such as lysosomes and become protonated and sequestered. In low pH conditions such as this, AO will emit orange fluorescence (emission ~590 nm) when excited by blue light (475 nm). EB (excitation: 360 nm, emission: 590 nm) is only taken up



**Figure 1:** Microscope fluorescence image of Hs578T cells treated with *H. aspersa* extract and stained by AO/EB observed at 100X magnification: (A) Control. (B) Cells treated by the dilution 0.1%. (C) Cells treated by the dilution 1%. Live cell showed by a green arrow, apoptotic cell showed by a yellow arrow, necrotic cell showed by a red arrow.



**Figure 2:** Effect of *H. aspersa* extract on the expression of TNFα mRNA in breast cancer cell line Hs578T (\*significant  $p < 0.05$ , \*\*\* significant  $p < 0.001$ , compared to control)

**Table 1:** Primers used in RT-PCR

Genes	PrimerRight	PrimerLeft	Reference
TNFα	5' CAGCCTCTCTCCTTCTCTGAT	5' GCCAGAGGGCTGATTAGAGA	(TibMolBiolcode794483;794484)
NF-κB	5' GGCTGGCAGCTCTTCTCA	5' TCCAGGTCATAGAGAGGCTCA	(TibMolBiolcode934127;934128)
TP53	5' CCCTTTTGGACTTCAGGTG	5' AGGCCTTGGAACTCAAGGAT	(TibMolBiolcode934133;934134)
PTEN	5' TCCAGATGATTCTTTAACAGGTAGC	5' GGGGAAGTAAGGACCAGAGAC	(TibMolBiolcode:9341131,9341132)
B-actin	5'AGGAATGGAAGCTTGC GGTA	5'AATTTTCATGGTGGATGGTGC	(TibMolBiol,code881493;881492).

by cells when cytoplasm membrane integrity is lost and stains nuclei in red. Therefore, live cells (showed in Figure 1A by a green arrow) have normal green nuclei. Apoptotic cells (showed in Figure 1C by a yellow arrow) have a bright green nucleus with condensed or fragmented chromatin and cells that have died from necrosis have an orange/yellow nucleus (showed in Figures 1B and 1C by a red arrow).

The analysis of the fluorescence and comparison between the two doses [Figures 1B and 1C] indicates that the extract induces necrosis. At 1%, the presence of some apoptotic cells has been also noted. Thus, the extract induces necrosis rather than apoptosis in Hs578T cell line.

### Effect of *H. aspersa* extract on TNF $\alpha$ expression

TNF $\alpha$  mRNA (messenger RNA) levels were clearly elevated, after 4 and 8 hours, in Hs578T treated samples as indicated in Figure 2.

After 4 hours, TNF $\alpha$  expression was 20 and 10 times more important with dilutions 0.1% and 1%, respectively, after treatment. Statistically and compared to controls, the increase of TNF $\alpha$  mRNA was highly significant ( $p < 0.001$ ).


After 8 hours, mRNA levels were decreased, but they remained significantly higher than those registered in control cells, especially with the 1% dilution.

After 24 hours of treatment with the two doses, TNF $\alpha$  expression has been significantly inhibited ( $p < 0.05$ ).


Results from ELISA assay [Figure 3] showed that TNF $\alpha$  concentrations rise after 8 hours of exposition to *H. aspersa* extract, and still higher after 24 hours. This data confirms that the extract stimulates TNF $\alpha$  production. In addition, the dose 0.1% has a more stimulatory effect.

### Effect of *H. aspersa* extract on NF- $\kappa$ B expression


Treatment of Hs578T cells by *H. aspersa* extract at 0.1% decreases NK- $\kappa$ B expression after 4, 8, and 24 hours compared to control. At 1%, the effect




**Figure 3:** Effect of *H. aspersa* extract on TNF $\alpha$  concentration in culture supernatant of breast cancer cell line Hs578T (\* significant  $p < 0.05$ , \*\*\* significant  $p < 0.001$ , compared to control)



**Figure 4:** Effect of *H. aspersa* extract on the expression of NF- $\kappa$ B mRNA in breast cancer cell line Hs578T (\*\*significant  $p < 0.01$ , compared to control).



**Figure 5:** Effect of *H. aspersa* extract on the expression of PTEN mRNA in breast cancer cell line Hs578T (\*significant  $p < 0.05$ , compared to control)



**Figure 6:** Effect of *H. aspersa* extract on the expression of p53 mRNA in breast cancer cell line Hs578T (\*significant  $p < 0.05$ , \*\* significant  $p < 0.01$ , compared to control).

of the extract was observed after 8 and 24 hours, we noted a significant decrease of NF- $\kappa$ B expression compared to nontreated cells [Figure 4]. The highest dose (1%) had more inhibitory effect on NK- $\kappa$ B.

### Effect of *H. aspersa* extract on PTEN expression

Treatment of Hs578T cells with *H. aspersa* extract resulted in the diminution of PTEN expression levels with the two doses 0.1 and 1%. A modest increase in PTEN expression was observed after 4 hours of treatment with the lowest dose (0.1%). The highest dose (1%), however, had more inhibitory effect ( $p < 0.05$ ) [Figure 5].

### Effect of *H. aspersa* extract on P53 expression

From Figure 6, we note that, no significant change has been registered with the 0.1% dilution. At 1% the extract significantly decreases p53 mRNA levels after 8 and 24 hours of treatment.

## DISCUSSION

The main aim of this study was to determine the mode of action of the aqueous extract from *H. aspersa* on breast cancer cells. The fluorescence staining shows that *H. aspersa* induces necrosis in Hs578T cells; at the same time it enhances the TNF $\alpha$  expression significantly. This cytokine had been demonstrated to stimulate both apoptosis and necrotic cell death, depending on cell type and treatment conditions.<sup>[22]</sup> The necrosis induced by our extract seemed to be mediated by high levels of TNF $\alpha$ . The extract has stimulated the expression and secretion of TNF $\alpha$ , in culture medium. Therefore, the cytokine stimulates its specific receptor and act by autocrine mode.

The TNF $\alpha$  induced necrosis is a form of cell death called "necroptosis" that resulted from an incomplete execution of apoptosis program, in which caspases and NF- $\kappa$ B are inhibited. These characteristics are in accordance

with our results because *H. aspersa* extract blocked the apoptotic process and the complete apoptosis has been revealed only in some cases.

TNF $\alpha$  enhancement in our experiments may have resulted from the effect of *H. aspersa* lectins, and change of glycozylation at the surface of tumor cells. A lectin from *Helix pomatia*, a similar species of snails, has been shown to recognize diverse epitopes on tumor cells like O-linked  $\alpha$ -N-acetylgalactosamine on Tn epitopes [6–7], integrin  $\alpha 6$ , transcription factors heterogeneous nuclear ribo-nuclear protein (HnRNPs), heat shock protein 27 (Hsp27) and enolase 1 (ENO1) in breast and colon tumor cells. *H. pomatia* lectin binding was also observed in the golgi apparatus in the T47D and MCF7 cells.<sup>[23]</sup>

Indeed, a similar augmentation of TNF $\alpha$  expression had been registered with lectin extracted from mushroom.<sup>[24]</sup>

Another explanation is that *H. aspersa* extract has a pro-oxidant activity and may act by generation of reactive oxygen species (ROS). A high level of ROS can lead to necrotic cell death while, a low level leads to apoptotic cell death.<sup>[25]</sup> Moreover, Sakon *et al.*, 2003 have reported that TNF stimulation leads to accumulation of ROS, which is essential for prolonged mitogen-activated protein kinase (MAPK) activation and cell death.<sup>[26]</sup> TNF receptor-1 (TNFR1) has been shown to initiate necrotic cell death<sup>[27]</sup> and leads to the generation of ROS, which functions as second messengers in the necrotic cell death pathway.<sup>[28]</sup>

Furthermore, our results show that *H. aspersa* extract is a good inhibitor of NF-KB. Recent studies have reported that inhibition of NF-KB alone or in combination with cancer therapies leads to tumor cell death or growth inhibition.<sup>[14]</sup> Li *et al.*, 2003 have reported that the increased ROS generation might contribute to the induction of apoptosis and revealed that NF-KB activity was almost completely inhibited by preventing the degradation of I $\kappa$ B $\alpha$ . Additionally, the level of bcl-2 was decreased.<sup>[29]</sup>

The P53 and PTEN downregulation induced by *H. aspersa* extract is the result of TNF $\alpha$  overexpression. Different contradictory effects of TNF $\alpha$  have been reported on P53 expression. It was shown to increase P53 expression with HT29 cells and to decrease or minimally change its expression in HCT116 cells.<sup>[30]</sup>

Treatment of cancer cell lines with TNF $\alpha$  decreases PTEN expression and the overexpression of TNF $\alpha$  lowers PTEN expression via TNF $\alpha$ /NIK/NF-kB pathway.<sup>[31]</sup>

Currently, anticancer drugs are developed with the aim to maximize apoptosis. However, cancer cells eventually become resistant to these therapeutics and multiple mechanisms may contribute to resistance to apoptosis. So, necrosis is an alternative pathway with fewer mechanisms of resistance compared to apoptosis. Targeted necrosis has potential clinical utility because this cell death mechanism retains the cancer cell specificity of apoptosis and bypasses the apoptotic resistance by redirection into necrosis.<sup>[32]</sup>

In addition, Olofsson *et al.*, 2007 have shown that chemotherapy induces more necrotic than apoptotic cell death in breast cancer patients, and this necrotic response is associated with a better survival.<sup>[33]</sup> So the induction of necrosis by *H. aspersa* extract may be an interesting characteristic and can open a new perspective in cancer therapy.

## CONCLUSION

Our results demonstrate for the first time that *H. aspersa* extract has a high cytotoxicity against breast cancer cells. The extract can be used as an associated treatment, in chemotherapy that enhances tumor sensitivity by downregulating Bcl2<sup>[8]</sup> and NF-kB; because tumor resistance is usually correlated with overexpression of these factors. Although the propriety of *H. aspersa* extract as an inducer of necroptosis has to be further clarified, the present study opens new perspectives in the search for new natural anticancer drugs.

## Acknowledgements

The authors are thankful to Dr Bensouyad Abdelhalim from Constantine University for language revision.

## Financial support and sponsorship

Nil.

## Conflicts of interest

There are no conflicts of interest.

## REFERENCES

- Amin A, Moussa M. Merits of anti cancer plants from the Arabian Gulf region. *Cancer therapy* 2007;5:55-66.
- Boopathy NS, Kathiresan K. Anti cancer drugs from marine flora: An overview. *J. Oncol.* 2010;article ID 214186, 18 pages
- Galgitay F, Erkan N, Tosun D, Selcuk A. Amino acid, Fatty acid, vitamin and mineral contents of Edible garden snail (*Helix aspersa*). *Journal of Fisheries Sciences* 2011;5:354-63.
- y Ozogul, Ozygul F, Olgunoglu A. Fatty acid profile and mineral content of wild snail *Helix pomatia* from the region of south of the turkey. *Eur. Food. Res.Techol* 2004;221:547-49.
- Bonnemain B. Helix and drugs: Snails for western health care from antiquity to the present, *eCAM.* 2005;2:25-28.
- Walker RA. *Helix pomatia* and prognosis and prognosis of breast cancer. *Br J Cancer* 1993;68:453-54.
- Lescar J, Sanchez JF, Audfay A, Coll JL, Breton C, Mitchell EP, *et al.* Structural basis for recognition of breast and colon cancer epitopes Tn antigen and forssman disaccharide by helix pomatia lectin. *Glycobiology* 2007;17:1077-083.
- El Ouar I, Braicu C, Naimi D, Irimie A, Berindan-Neagoe I. Anti tumour effect of aqueous extract from *Helix aspersa*. *Int J Pharm Bio Sci.* 2013;4:1325-1332.
- Elmore S. Apoptosis: A Review of Programmed Cell Death. *Toxicol Pathol* 2007;35:495-516.
- Nishikori M. Their Roles in Lymphoid Malignancies. *J. Clin. Exp. Hematopathol* 2005;45:15-24.
- Liu J, Voisin V, Wang S, Wang DY, Jones RA, Datti A, *et al.* Combined deletion of PTEN and p53 in mammary epithelium accelerates triple-negative breast cancer with dependency on eEF2K. *EMBOMol Med* 2014;6:1542-1560.
- Liu ZG. Molecular mechanism of TNF signaling and beyond. *Cell Research* 2005;15:24-27.
- Szlosarek P, Charles KA, Balkwill FR. Tumor necrosis factor  $\alpha$  as tumor promoter. *Eur j cancer* 2006;42:745-50.
- Madonna G, Ullman D, Gentilcore G, Palmieri G, Ascierto PA. NF-kB as potencial target in the treatmnt of melanoma. *J Transl Med* 2012;10:53.
- Lacroix M, Toillon RA, Leclercq G. P53 and breast cancer, an update. *Endocrine Related Cancer* 2006;13:293-325.
- Turner N, Moretti E, Siclari O, Migliaccio I, Santarpia L, D'Incalci M, *et al.* Targeting triple negative breast cancer: Isp53 the answer? *Cancer Treat Rev* 2013;39:541-50.
- Gasco M, Shami S, Crook T. The P53 pathway in breast cancer. *Breast Cancer Research* 2002;4:70-76.
- Okumura N, Yoshida H, Kitagishi Y, Nishimura Y, Matsuda S. Alternative splincings on p53, BRCA1 and PTEN genes involved in breast cancer. *Biochemical and Biophysical Research Communications* 2011;413:395-99.
- Saal L H, Gruberger-Saal S K, Persson C, Lövgren K, Jumppanen M, Staaf J, *et al.* Recurrent gross mutations of the *PTEN* tumor suppressor gene in breast cancers with deficient DSB repair. *Nat Genet* 2008;40:102-07.
- Ribble D, Goldstein NB, Norris DA, Shellman YG. A simple technique for quantifying apoptosis in 96-well plates. *BMC Biotechnology* 2005;5:12.
- Livak KJ, Schmittgen TD. Analysis of relative gene expression data using real time PCR and the 2<sup>- $\Delta\Delta C_T$</sup>  Method. *Methods* 2001;25:402-08.
- Galluzi L, Kepp O, Kromer G. RIP kinases initiate programmed necrosis. *J. Mol. Cell. Biol* 2002;1:8-10.
- Rambaruth NDS, Greenwell P, Dwek MV. The lectin *Helix pomatia* agglutinin recognizes O-GlcNAc containing glycoproteins in human breast cancer. *Glycobiology* 2012;22:839-48.
- Mannel DN, Becker H, Gundlt A, Kist A, Franz H, *et al.* Induction of tumor necrosis factor expression by a lectin from *Viscum album*. *Cancer Immunol Immunother* 1991;33:177-82.

25. Kannan K, Jain SK. Oxidative stress and apoptosis. *Pathophysiology* 2000;7:153-63.
26. Sakon S, Xue X, Takekawa M, Sasazui T, Okazaki Kajima Y, Piao JH. NF-kB inhibits TNF induced accumulation of ROS that mediate prolonged MAPK activation and necrotic cell death. *EMBO journal* 2003;22:3838-909.
27. Organ MJ, Kim YS, Liu ZG. TNF alpha and reactive oxygen species in necrotic cell death. *Cell Res* 2008;18:343-49.
28. Shen HM, Pervaiz S. TNF receptor superfamily-induced cell death: redox-dependent execution. *FASEBJ* 2006;20:1589-598.
29. Li HL, Chen DD, Li XH, Zhang HW, Lü YQ, Ye CL, *et al.* Changes of NF-kB, p53, Bcl-2 and caspase in apoptosis induced by JTE-522 in human gastric adenocarcinoma cell line AGS cells: role of reactive oxygen species. *World j of Gastroentrol* 2003;8:431-5.
30. Pastor DM, Irby RB, Poritz LS. Tumor necrosis factor alpha induces p53 up regulated of apoptosis expression in colorectal cancer cell lines. *Dis Colon Rectum* 2010;53:257-63.
31. Kim S, Domon-Dekk C, Kang J, Chung DH, Freund JN, Mark Ever B, *et al.* Down-regulation of the tumor suppressor PTEN by the tumor necrosis factor-alpha/nuclear factor-kappaB (NF-kappaB)-inducing kinase/NF-kappaB pathway is linked to a default IkappaB-alpha autoregulatory loop. *J Biol Chem* 2004;279:4285-291.
32. Chaabane W, User SD, El-Gazzah M, Jaksik R, Sajjadi E, Rzeszowska-Wolny J, *et al.* Autophagy, Apoptosis, mitoptosis and Necrosis: Interdependence between Those Pathways and Effects on Cancer. *Arch Immunol Ther Exp* 2013;61:43-58.
33. Olofsson M.H, Ueno T, Pan Y. Cytokeratin-18 is a useful serum biomarker for early determination of response of breast carcinomas to chemotherapy. *Clin Cancer Res* 2007;13:3198-206.