

# circRNA-sponging: a pipeline for extensive analysis of circRNA expression and their role in miRNA sponging

Markus Hoffmann<sup>1,2,3,\*</sup>, Leon Schwartz<sup>1,\*</sup>, Octavia-Andreea Ciora<sup>1,\*</sup>, Nico Trummer<sup>1</sup>, Lina-Liv Willruth<sup>1</sup>, Jakub Jankowski<sup>3</sup>, Hye Kyung Lee<sup>3</sup>, Jan Baumbach<sup>4,5</sup>, Priscilla Furth<sup>6</sup>, Lothar Hennighausen<sup>2,3</sup>, and Markus List<sup>1,\*\*</sup>

<sup>1</sup>Big Data in BioMedicine Group, Chair of Experimental Bioinformatics, TUM School of Life Sciences, Technical University of Munich, Munich, Germany

<sup>2</sup>Institute for Advanced Study (Lichtenbergstrasse 2a, D-85748 Garching, Germany), Technical University of Munich, Germany

<sup>3</sup>National Institute of Diabetes, Digestive, and Kidney Diseases, Bethesda, MD 20892, United States of America

<sup>4</sup>Chair of Computational Systems Biology, University of Hamburg, Hamburg, Germany

<sup>5</sup>Computational BioMedicine Lab, University of Southern Denmark, Odense, Denmark

<sup>6</sup>Departments of Oncology & Medicine, Georgetown University, Washington, DC, USA

\*The authors wish to be known that in their opinion the first three authors should be considered as shared first authors;

\*\*corresponding authors

## ABSTRACT

**Motivation:** Circular RNAs (circRNAs) are long non-coding RNAs (lncRNAs) often associated with diseases and considered potential biomarkers for diagnosis and treatment. Among other functions, circRNAs have been shown to act as microRNA (miRNA) sponges, preventing the role of miRNAs that repress their targets. However, there is no pipeline to systematically assess the sponging potential of circRNAs.

**Results:** We developed circRNA-sponging, a nextflow pipeline that (1) identifies circRNAs via back-splicing junctions detected in RNA-seq data, (2) quantifies their expression values in relation to their linear counterparts spliced from the same gene, (3) performs differential expression analysis, (4) identifies and quantifies miRNA expression from miRNA-sequencing (miRNA-seq) data, (5) predicts miRNA binding sites on circRNAs, (6) systematically investigates potential circRNA-miRNA sponging events, (7) creates a network of competing endogenous RNAs, and (8) identifies potential circRNA biomarkers. We showed the functionality of the circRNA-sponging pipeline using RNA sequencing data from brain tissues where we identified two distinct types of circRNAs characterized by a distinct ratio of the binding site length. The circRNA-sponging pipeline is the first end-to-end pipeline to identify circRNAs and their sponging systematically with raw total RNA-seq and miRNA-seq files, allowing us to better indicate the functional impact of circRNAs as a routine aspect in transcriptomic research.

**Availability:** <https://github.com/biomedbigdata/circRNA-sponging>

**Contact:** [markus.daniel.hoffmann@tum.de](mailto:markus.daniel.hoffmann@tum.de); [markus.list@tum.de](mailto:markus.list@tum.de)

**Supplementary Material:** Supplementary data are available at Bioinformatics online.

## INTRODUCTION

Circular RNAs (circRNAs) are classified as long non-coding RNAs (lncRNAs) even though a few have been reported to encode proteins [1]. circRNAs are characterized by their loop structure, which makes them less prone to degradation [2,3]. The biogenesis of circRNAs is explained by the occurrence of a back-splicing event (see Suppl. Fig. 1) during the alternative splicing process of precursor messenger RNA (pre-mRNA), where the 5' terminus of an upstream exon and the 3' terminus of a downstream exon are covalently joined [2]. The difference between circRNAs and linear RNAs is the lack of a 5' cap and a 3' polyadenylation (poly(A)) tail along with its circular shape, which makes circRNAs more stable and, in most cases, resistant to exonuclease activity [4–6]. These circular molecules can be made up of exonic

and intronic regions of its spliced pre-mRNA and are thus found in a huge variety of sizes, ranging from 100 to >4,000 nucleotides [4,7]. circRNAs are conserved across species and their expression is tissue- and disease-specific [3,8,9]. Hence, they can play an important role as biomarkers and therapeutic targets [9–11]. Another type of non-coding RNA is microRNA (miRNA) which plays a role in post-transcriptional gene regulation [12–14] and is involved in many biological processes and diseases [15]. miRNAs bind their target genes via the RNA-induced silencing complex causing their degradation or preventing their translation [16].




Figure 1: Overview of the individual steps of the pipeline. total RNA-seq data processing is shown on top and miRNA-seq processing on the bottom. In the miRNA sponging step, these results are integrated for network analysis and biomarker detection.

The possible interplay between circRNAs, miRNAs, messenger RNAs (mRNAs) that code for proteins and other types of RNA that share miRNA binding sites gives rise to a large regulatory network. Salmena et al. proposed that any RNA that carries miRNA binding sites (e.g., mRNAs, circRNAs, pseudogenes, transcripts of 3' untranslated regions (UTRs), and lncRNAs) can act as a competing endogenous RNA (ceRNA) [17] that competes for the limited pool of available miRNAs in a cell. As a result of this competition, an overexpressed RNA can sponge away miRNAs required for the regulation of other RNAs, which can explain why non-coding RNAs such as circRNAs can be implicated in a phenotype.

The enhanced stability of circRNAs might allow them to work as buffers for miRNAs, by binding them until sufficient miRNAs are present to outnumber the circRNA binding sites [9]. The regulatory function of circRNAs and their alleged association with diseases are the main reasons why identifying sponging activity between circRNAs and miRNAs is of particular interest. The presence of an interaction between miRNAs and circRNAs has been repeatedly proven and several circRNAs (e.g., CDR1as/CiRS-7, SRY [18], and circNCX1 [19]) have been recognized as miRNA sponges. Even though individual studies confirmed the existence of circRNA sponges, further studies are needed to elucidate the role of circRNAs in miRNA-mediated gene regulation.

From a computational point of view, the detection of circRNAs is difficult due to their circular shape and the lack of poly(A) tail, which makes it unlikely to observe them in poly(A)-enriched RNA sequencing (RNA-seq) libraries [7]. Hence, circRNAs can only be robustly detected in libraries without poly(A) enrichment, such as ribosomal RNA (rRNA) depleted RNA-seq and total RNA-sequencing (total RNA-seq), which do not deplete circRNAs [7]. Identification of circRNAs relies on the detection of back-splicing junctions (BSJ) among the unmapped reads, which allows for the estimation of circRNA abundance. By focusing on the back-splicing junction alone, the expression of circRNAs in relation to their linear counterparts is typically underestimated [20].

Several approaches for circRNA analysis have been proposed. Chen et. al reviewed 100 existing circRNA-related tools for circRNA detection, annotation, downstream analysis, as well as network analysis [21]. They list a total of 44 circRNA

identification tools including, but not limited to, CIRCexplorer [22,23], find\_circ [24], CIRI [25], KNIFE [26], and circRNA\_finder [27]. They also present a total of 14 circRNA annotation databases collecting circRNA information from the literature, such as circBase [28] and CIRCpedia [22,29]. Other circRNA-related tools include databases for feature collection, and storing circRNA information related to disease and biomarkers. In addition, circRNA network identification tools model the interactions between circRNAs and miRNAs, lncRNAs, or RNA-binding proteins. Other tools for downstream analysis of circRNAs cover alternative splicing detection, circRNA assembly, and structure prediction and visualization [21]. However, to the best of our knowledge, none of the tools performs a comprehensive and automated circRNA sponging analysis integrating identification and quantification of methods both circRNAs and miRNAs, systematic investigation of potential circRNA-miRNA sponging events, and ceRNA network analysis. We developed “circRNA-sponging”, a nextflow pipeline integrating state-of-the-art methods to (1) detect circRNAs via identifying back-splicing junctions from total RNA-seq data, (2) quantify their expression values relative to linear transcripts, (3) perform differential expression analysis, (4) identify and quantify miRNA expression from miRNA-sequencing (miRNA-seq) data, (5) predict miRNA binding sites on circRNAs, (6) systematically investigate potential circRNA-miRNA sponging events, (7) create a ceRNA network, and (8) identify potential circRNA biomarkers using the ceRNA network (Fig. 1). We demonstrate the potential of the circRNA-sponging pipeline on matched rRNA-depleted RNA-seq and miRNA-seq data from mouse brain tissues.




Figure 2: Workflow of the circRNA-sponging pipeline. The pipeline consists of three modules: (1) the circRNA module, (2) the miRNA module, and (3) the sponging module. In (1), we detect circRNAs via identifying back-splicing junctions from total RNA-seq data, quantify their expression values, perform a differential expression analysis, and predict miRNA binding sites on circRNAs using a majority

vote between three state-of-the-art methods. In (2), we either detect and quantify miRNAs in raw miRNA-seq or directly process miRNA expression data. In (3), we systematically investigate circRNA-miRNA sponging events, create a ceRNA network and use it to identify potential circRNA biomarkers.

## MATERIALS AND METHODS

### Data

Using circRNA-sponging, we processed a total of 23 samples of matched single-end rRNA-depleted RNA-seq and miRNA-seq data for four brain regions (cerebellum, cortex, hippocampus, olfactory bulb). Samples include 2-3 replicates for wild-type (WT) and 3 CDR1 knock-out (KO) mouse replicates (GEO accessions: GSE100265, GSE93129) [30]. (see Suppl. Table 1). We use the mm10 genome version for mapping.

### Pipeline Architecture

The circRNA-sponging pipeline is implemented in R (v. 4.2.0) and python (v. 3.8.12) and wrapped with nextflow version 22.04.0.5697. It follows the nf-core guidelines [31] and encompasses several state-of-the-art techniques organized into three modules: (1) the circRNA module, (2) the miRNA module, and (3) the sponging module, the latter of which can only be performed if both other modules have been executed (Fig. 2). In the following, we provide a deeper insight into each module and highlight important components of the pipeline.

(1) *The circRNA module* addresses the identification, quantification, and miRNA binding site prediction of circRNAs. For read mapping, we employ the STAR [32] aligner, which provides support for the detection of splice-junction and fusion reads. The resulting unmapped split-reads are used by CIRCexplorer2, which uses a combination of methods (i.e., a de novo assembly approach to identify novel circRNA and a reference-based approach, which uses known exon-exon junctions to map back-splicing events to known genes) to increase the accuracy of its predictions [22] to identify back-splicing events. Next, raw read counts are normalized with DESeq2 [33] and circRNAs with low expression levels are excluded to reduce false positives. By default, only circRNAs with a normalized read count >5 in at least 20% of samples are retained. Database annotation is performed using circBase [28], which covers curated circRNAs with experimental evidence of several model organisms.

For quantifying expression values, we use psirc [20], which in turn employs kallisto [34] considering both linear and circular transcripts in the expectation-maximization step to yield comparable expression values. If the data has been sampled from different conditions (e.g., case and control), the quantified linear transcripts and circRNAs can be used to perform a differential expression analysis using DESeq2 [33]. The pipeline generates heatmaps, volcano plots, and principal component analysis (PCA) of the circRNAs and linear transcripts between conditions. To boost reliability, we predict circRNA-miRNA binding sites using a majority voting between miRanda [35], PITA [36], and TarPmiR [37] since each method has a distinct approach for predicting miRNA binding sites. We consider a circRNA-miRNA binding site as relevant if it is predicted by at least two out of the three methods. miRanda considers seed matching, conservation, and free energy and we consider predictions with a score above the 25% quantile. PITA additionally considers site accessibility and target-site abundance. TarPmiR further integrates machine learning to improve results for supported organisms [37]. We further incorporate experimentally validated



target sites from DIANA-LncBase v3 [38], miRTarBase [39,40], and miRWalk3.0 [41].

(2) *The miRNA module* covers the quantification and processing of miRNA expression. miRDeep2 [42] is used to obtain miRNA counts. Alternatively, already mapped miRNA expression data can be provided. Raw counts are normalized with DESeq2 [33] followed by a filtering step, where by default miRNAs with a normalized read count > 5 in at least 20% of samples are retained.

(3) *The sponging module* is used for the identification of crosstalk between circRNAs, miRNAs as well as ceRNA interactions of circRNAs with other transcripts. To identify potential sponging activity, we perform a correlation analysis of circRNA-miRNA pairs, where a negative correlation coefficient indicates a sponging relationship. For all circRNA-miRNA pairs with at least one binding site for a shared miRNA, we compute a Pearson correlation coefficient along with the normalized residual sum of squares and the adjusted p-value, after the Benjamini-Hochberg correction. Pairs are filtered (e.g., p-adjusted < 0.05, RMSE < 1.5, and optionally by the number of binding sites) and are considered potential sponging candidates. We further construct a ceRNA network using SPONGE [12] on matched gene and miRNA expression data. Finally, we apply spongEffects [14] to extract ceRNA modules consisting of circRNAs with a high node centrality score in the ceRNA network and their direct neighbors. For each module, spongEffects computes a sample-specific enrichment score allowing for differential analysis.

## RESULTS AND DISCUSSION

circRNAs are highly abundant and conserved in the mammalian brain [43,44]. For demonstrating the capabilities of the circRNA-sponging pipeline, we analyzed a public RNA-seq data set from the mouse brain. We focused on the sponging capacity of circRNAs and their potential role as ceRNAs.

### Comparing circRNA and host gene expression reveals changes in circRNA splicing

In total, we detected 46,380 and 27,390 circRNAs before and after filtering, respectively. This number is in line with the known high abundance of circRNAs in brain tissue [43,44]. We could annotate only 1,027 (~4%) of the circRNAs that passed the filter (Suppl. Fig. 2 a), likely as comparably few circRNAs have thus far been annotated in mice using circBase. psirc-estimated expression levels, which take reads mapping to parts other than the back-splicing junction of the ceRNA into account, are up to 6-fold higher compared to counts derived from back-splicing junctions only (Fig. 3 a, per tissue type: Suppl. Fig. 2 d-g). We observed a generally higher expression of circRNAs in the cerebellum compared to other brain regions, which could indicate a higher importance of the circRNAs in this brain region (Suppl. Fig. 2 b). The number of shared circRNAs across brain regions is comparably high with only very few circRNAs showing brain region-specific occurrence (Suppl. Fig. 2 c). However, expression levels of the circRNAs differ considerably such that samples cluster by brain region (Figs. 3 b-c). The difference between KO and WT samples is negligible, with the exception of the CDR1 region (mmu\_hsa\_circ\_0001878 in circBase annotation) that was targeted successfully (Suppl. Fig. 3. b). As expected, we observed a clear separation between the cerebellum and other brain regions, while the cortex and hippocampus are more similar [45]. Our analysis revealed a total of 33 circRNAs that show significantly different expression between brain regions (p-adjusted < 0.01, absolute log<sub>2</sub> fold change > 5, Suppl. Table 2).

Rybak-Wolf et al. reported circRNAs of twelve host genes (TULP4, RIMS2, ELF2, PHF21A, MYST4, CDR1, STAU2, SV2B, CPSF6, DYM, RMST, and RTN4), nine of which (all but TULP4, SV2B, and RMST) were detected in our analysis (Suppl. Table 3, Suppl. Fig. 3 c-d). They speculated on the importance of circRNAs originating from these genes for brain cell identity but we posit that a change in circRNA expression alone does not necessarily imply a functional role as circRNA expression is coupled to the expression of the host gene, as we expect the number of reads mapping to the black-splicing junction to correlate. However, by comparing the expression level of the circRNAs to the linear transcripts, as facilitated by psirc-quant, we can identify cases where the expression of circRNAs increases beyond the level suggested by the overall gene expression. Such cases offer in our opinion stronger evidence for the functional importance of a circRNA. For example, mmu\_circ\_0000595, a circRNA of host gene RIMS2 shows a higher than expected expression specifically in the cortex, suggesting a possible functional role for this circRNA in this brain region (Fig. 3 d, see also Suppl. Fig. 4 a-p).




Figure 3: circRNA results of the mouse brain regions data set. a) circRNAs shared between brain regions. b) expression of circRNAs across tissues and experimental conditions. c) psirc-quant estimated compared to CIRCexplorer2 counts. d) comparison between a circRNA originating from RIMS2 and expression of the linear RIMS2 gene.

## A ceRNA network reveals circRNAs acting as miRNA sponges

If matched total RNA and miRNA sequencing data are provided, circRNA-sponging infers a ceRNA network using the R package SPONGE [12] and visualizes the result (Fig. 4). Important players in this regulatory network are characterized by a large node degree, i.e., they indirectly regulate many of the connected RNAs via sequestering miRNA copies. Since the network inferred by SPONGE does not offer insights into individual samples or conditions, we subsequently computed spongeEffects [14] enrichment scores which capture the interaction of individual circRNAs and their target genes. As these scores are sample-specific, they can offer insights into condition-specific circRNA sponging activity. spongeEffects scores can also be used as features for machine learning tasks such as classification [14]. Since the number of available samples for training is rather small here, the random forest reported subset accuracy drops considerably on the holdout set in ten-fold cross-validation. While the cortex and hippocampus are difficult to differentiate, the cerebellum can be robustly distinguished from other brain regions (Suppl. Fig. 5). In particular two circRNAs, *chr10:9770449-9800068\_-* and *chr10:79860969-79862010\_+* stand out as distinctive features of the cerebellum (Suppl. Fig. 6-7). While the inferred ceRNA network shows that circRNAs in the mouse brain are regulatory active through miRNA sponging, a larger number of samples is likely needed to fully resolve brain region-specific sponging activity.





Figure 4: circRNA-ceRNA-subnetwork with the top 25 ceRNA modules ranked by the number of significant interactions (node degree in the network). For each ceRNA module consisting of the circRNA and its target genes, we computed spongeEffects enrichment scores and used them as input for a random forest model. The bottom-right corner shows the subset accuracy of this model in distinguishing different brain regions on the training and test set. The results of a model trained on random modules of the same size show random performance.

## Comparing the number of circRNA binding sites with respect to their length reveals two distinct clusters

miRNA sponging has long been considered a potential function of circRNAs [18]. To fulfill this function, it would be beneficial for circRNAs to carry a large number of miRNA binding sites and indeed, some known circRNAs such as CDR1as harboring



over 70 miRNA binding sites for miR-7 alone [46] fit the hypothesis well. To investigate if this is a general property of circRNAs, we systematically compared the length of a transcript to the number of binding sites, expecting to observe a larger ratio for circRNAs compared to the 3' untranslated regions of linear transcripts, where miRNA binding sites are predominantly located (Fig. 5). While linear transcripts show a very diverse picture, circRNA length correlates well with the number of binding sites. We observed two distinct clusters which can be explained by the different prediction methods we used. It appears that TarPmiR, which is based on machine learning, predicts binding sites in fewer

Figure 5: miRNA binding site versus transcript length for linear and circular RNA. For the 3'UTRs of mRNAs, the number of binding sites was inferred from miRWalk 3.0. circRNA-miRNA binding sites were counted if they were reported by two of the three prediction methods employed, i.e. miRanda, TarPmiR, and PITA. circRNAs form two clusters that can be explained by the different target site prediction methods used. Linear regression models were fit to each of the groups to show the trend of the association.

circRNAs compared to PITA or miRanda but for those, it predicts a considerably larger number of binding sites. Compared to the prediction in 3' UTRs (also based on TarPmiR), circRNAs show a comparably high ratio between the number of binding sites and the length. An interesting question is hence, if TarPmiR is able to differentiate between circRNAs that are active miRNA sponges and those that have other functions. Previous research has defined three types of circular RNAs based on structural features - exonic circular (ecirc) RNAs, circular intronic RNA (ciRNA), and exon-intron circRNA (ElciRNA) [47–49]. It has been suggested that ecircRNAs function predominantly through a miRNA sponging effect in the cytoplasm whereas other circular RNA forms (e.g. ciRNA and ElciRNA) function in the nucleus to regulate gene transcription [50–52]. Hence, circRNAs that are functional in the nucleus could have fewer miRNA binding sites. To test alternative explanations, we checked if clusters differed by (1) genesis, i.e. the splicing method of the circRNA (exonic or intronic), (2) biotype of the circRNA host gene (i.e., coding or non-coding gene, Suppl. Fig. 8. a-b) or (3) circRNA expression level (Suppl. Fig. 8 c). The observed clusters did not differ in any of these categories and further work is needed to elucidate if these results are related to structural features. It should also be noted that TarPmiR was not trained specifically on circRNAs and that a prediction method tailored towards circRNAs should be developed when suitable experimental data become available.

## CONCLUSION AND OUTLOOK

We developed a new circRNA processing and analysis pipeline consisting of three modules harboring multiple current state-of-the-art methods: 1) circRNA detection, 2)



miRNA detection, and 3) detection of sponging events between circRNAs, ceRNAs, and miRNAs. To the best of our knowledge, it is the first comprehensive circRNA pipeline to detect, quantify and annotate circRNAs as well as to determine their sponging activity. The latter allows users to bring circRNAs into a functional context with other RNAs such as mRNAs and lncRNAs through a joint ceRNA network which is mediated by miRNA sponging. Wen et al. recently highlighted the need for a further extensive investigation into circRNAs due to their enormous potential to explain human diseases like cardiovascular diseases, autoimmune diseases, and human cancers [53]. circRNAs are also known to be involved in brain development, brain cell differentiation, and neuronal signaling [30,43]. To demonstrate the capabilities of the circRNA-sponging pipeline, we hence re-analyzed a public data set where we investigated circRNAs across different mouse brain regions. Using our pipeline, we could offer novel insights into circRNA biology across tissues of the brain. We showed that differences in circRNA splicing can be revealed when considering the expression of circRNAs relative to the expression of a host gene, similar to how alternative splicing events are detected by considering exon or intron inclusion. We further placed our findings into the context of miRNA sponging, demonstrating that circRNA exerts regulatory control over a vast number of transcripts. Finally, we showed that the number of binding sites in circRNAs correlated well with their length and observed that TarPmiR's machine learning strategy identifies a subset of circRNAs that could indicate promising candidates for miRNA sponging. Further work is needed to investigate if these two classes represent structurally different circRNAs such as ecircRNAs, ciRNAs, or ElciRNAs or if this observation can be explained by differences in the miRNA prediction methods with no biological implication at all.

In the future, we plan to extend the circRNA-sponging pipeline with additional features. As various functions other than miRNA sponging have been attributed to circRNAs [54], we see room for expanding the features towards e.g. investigating the protein-coding potential of circRNAs [1]. We further seek to integrate circRNA-sponging into ongoing community efforts such as nf-core [31] to build up long-term support for maintaining and expanding this pipeline. In summary, the circRNA-sponging pipeline is a powerful tool to detect, investigate, and analyze circRNAs and their sponging effects and thus it helps researchers consider circRNAs as a routine aspect in RNA-seq and miRNA-seq data analysis.

## **AVAILABILITY**

Data are available in GEO under GSE100265 (rRNA-depleted RNA-seq) and GSE93129 (miRNA). circRNA-sponging is available under the GPL v.3.0 license at:

<https://github.com/biomedbigdata/circRNA-sponging>

The results presented in this manuscript are available as RData objects at:

<https://doi.org/10.6084/m9.figshare.21864948>

## **FUNDING**

With the support of the Technical University Munich – Institute for Advanced Study, funded by the German Excellence Initiative. JB was partially funded by his VILLUM Young Investigator Grant nr.13154.

## **CONFLICT OF INTEREST DISCLOSURE**

The authors declare no competing interests.

## ACKNOWLEDGMENTS

We thank Bahar Tercan, Mauro A. A. Castro, A. Gordon Robertson, Katherine A. Hoadley, and Victoria K. Cortessis for the fruitful discussions. We thank Christina Trummer for the logo design. Figures were created with <https://www.biorender.com>. Parts of the images were designed using resources from Flaticon.com.

## REFERENCES

1. Miao Q, Ni B, Tang J. Coding potential of circRNAs: new discoveries and challenges. *PeerJ*. 2021;9: e10718.
2. Yu C-Y, Kuo H-C. The emerging roles and functions of circular RNAs and their generation. *J Biomed Sci*. 2019;26: 29.
3. Jeck WR, Sharpless NE. Detecting and characterizing circular RNAs. *Nat Biotechnol*. 2014;32: 453–461.
4. Lasda E, Parker R. Circular RNAs: diversity of form and function. *RNA*. 2014;20: 1829–1842.
5. Suzuki H, Zuo Y, Wang J, Zhang MQ, Malhotra A, Mayeda A. Characterization of RNase R-digested cellular RNA source that consists of lariat and circular RNAs from pre-mRNA splicing. *Nucleic Acids Res*. 2006;34: e63.
6. Mester-Tonczar J, Hašimbegović E, Spannbauer A, Traxler D, Kastner N, Zlabinger K, et al. Circular RNAs in Cardiac Regeneration: Cardiac Cell Proliferation, Differentiation, Survival, and Reprogramming. *Front Physiol*. 2020;11: 580465.
7. Szabo L, Salzman J. Detecting circular RNAs: bioinformatic and experimental challenges. *Nat Rev Genet*. 2016;17: 679–692.
8. Jeck WR, Sorrentino JA, Wang K, Slevin MK, Burd CE, Liu J, et al. Circular RNAs are abundant, conserved, and associated with ALU repeats. *RNA*. 2013;19: 141–157.
9. Zhang Z, Yang T, Xiao J. Circular RNAs: Promising Biomarkers for Human Diseases. *EBioMedicine*. 2018;34: 267–274.
10. Kristensen LS, Hansen TB, Venø MT, Kjems J. Circular RNAs in cancer: opportunities and challenges in the field. *Oncogene*. 2018;37: 555–565.
11. Qu S, Yang X, Li X, Wang J, Gao Y, Shang R, et al. Circular RNA: A new star of noncoding RNAs. *Cancer Lett*. 2015;365: 141–148.
12. List M, Dehghani Amirabad A, Kostka D, Schulz MH. Large-scale inference of competing endogenous RNA networks with sparse partial correlation. *Bioinformatics*. 2019;35: i596–i604.
13. Hoffmann M, Pachi E, Hartung M, Stiegler V. SPONGEdb: a pan-cancer resource for competing endogenous RNA interactions. *Narodonaselenie*. 2021. Available: <https://academic.oup.com/narcancer/article-abstract/3/1/zcaa042/6066568>
14. Boniolo F, Hoffmann M, Roggendorf N, Tercan B, Baumbach J, Castro MAA, et al. spongeEffects: ceRNA modules offer patient-specific insights into the miRNA regulatory landscape. *bioRxiv*. 2022. p. 2022.03.29.486212. doi:10.1101/2022.03.29.486212
15. Kartha RV, Subramanian S. Competing endogenous RNAs (ceRNAs): new entrants to the intricacies of gene regulation. *Front Genet*. 2014;5: 8.
16. Weinstein JN, Collisson EA, Mills GB, Shaw KRM, Ozenberger BA, Ellrott K, et al. The Cancer Genome Atlas Pan-Cancer analysis project. *Nat Genet*. 2013;45: 1113–1120.
17. Salmena L, Poliseno L, Tay Y, Kats L, Pandolfi PP. A ceRNA hypothesis: the

- Rosetta Stone of a hidden RNA language? *Cell*. 2011;146: 353–358.
18. Hansen TB, Jensen TI, Clausen BH, Bramsen JB, Finsen B, Damgaard CK, et al. Natural RNA circles function as efficient microRNA sponges. *Nature*. 2013;495: 384–388.
  19. Li M, Ding W, Tariq MA, Chang W, Zhang X, Xu W, et al. A circular transcript of *ncx1* gene mediates ischemic myocardial injury by targeting miR-133a-3p. *Theranostics*. 2018;8: 5855–5869.
  20. Yu KH-O, Shi CH, Wang B, Chow SH-C, Chung GT-Y, Lung RW-M, et al. Quantifying full-length circular RNAs in cancer. *Genome Res*. 2021. doi:10.1101/gr.275348.121
  21. Chen L, Wang C, Sun H, Wang J, Liang Y, Wang Y, et al. The bioinformatics toolbox for circRNA discovery and analysis. *Brief Bioinform*. 2021;22: 1706–1728.
  22. Zhang X-O, Dong R, Zhang Y, Zhang J-L, Luo Z, Zhang J, et al. Diverse alternative back-splicing and alternative splicing landscape of circular RNAs. *Genome Res*. 2016;26: 1277–1287.
  23. Zhang X-O, Wang H-B, Zhang Y, Lu X, Chen L-L, Yang L. Complementary sequence-mediated exon circularization. *Cell*. 2014;159: 134–147.
  24. Memczak S, Jens M, Elefsinioti A, Torti F, Krueger J, Rybak A, et al. Circular RNAs are a large class of animal RNAs with regulatory potency. *Nature*. 2013;495: 333–338.
  25. Meng X, Chen Q, Zhang P, Chen M. CircPro: an integrated tool for the identification of circRNAs with protein-coding potential. *Bioinformatics*. 2017;33: 3314–3316.
  26. Szabo L, Morey R, Palpant NJ, Wang PL, Afari N, Jiang C, et al. Statistically based splicing detection reveals neural enrichment and tissue-specific induction of circular RNA during human fetal development. *Genome Biol*. 2015;16: 126.
  27. Westholm JO, Miura P, Olson S, Shenker S, Joseph B, Sanfilippo P, et al. Genome-wide analysis of drosophila circular RNAs reveals their structural and sequence properties and age-dependent neural accumulation. *Cell Rep*. 2014;9: 1966–1980.
  28. Glažar P, Papavasileiou P, Rajewsky N. circBase: a database for circular RNAs. *RNA*. 2014;20: 1666–1670.
  29. Dong R, Ma X-K, Li G-W, Yang L. CIRCpedia v2: An Updated Database for Comprehensive Circular RNA Annotation and Expression Comparison. *Genomics Proteomics Bioinformatics*. 2018;16: 226–233.
  30. Piwecka M, Glažar P, Hernandez-Miranda LR, Memczak S, Wolf SA, Rybak-Wolf A, et al. Loss of a mammalian circular RNA locus causes miRNA deregulation and affects brain function. *Science*. 2017;357. doi:10.1126/science.aam8526
  31. Ewels PA, Peltzer A, Fillinger S, Patel H, Aneberg J, Wilm A, et al. The nf-core framework for community-curated bioinformatics pipelines. *Nat Biotechnol*. 2020;38: 276–278.
  32. Dobin CA, Davis FS, J. Drenkow, C. Zaleski, S. Jha, P. Batut, M. Chaisson, and TR Gingeras. 2013. STAR: ultrafast universal RNA-seq aligner ...; 2013.
  33. Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol*. 2014;15: 550.
  34. Bray NL, Pimentel H, Melsted P, Pachter L. Erratum: Near-optimal probabilistic RNA-seq quantification. *Nat Biotechnol*. 2016;34: 888.
  35. Enright AJ, John B, Gaul U, Tuschl T, Sander C, Marks DS. MicroRNA targets in

- Drosophila*. *Genome Biol.* 2003;5: R1.
36. Kertesz M, Iovino N, Unnerstall U, Gaul U, Segal E. The role of site accessibility in microRNA target recognition. *Nat Genet.* 2007;39: 1278–1284.
  37. Ding J, Li X, Hu H. TarPmiR: a new approach for microRNA target site prediction. *Bioinformatics.* 2016;32: 2768–2775.
  38. Karagkouni D, Paraskevopoulou MD, Tastsoglou S, Skoufos G, Karavangeli A, Pierros V, et al. DIANA-LncBase v3: indexing experimentally supported miRNA targets on non-coding transcripts. *Nucleic Acids Res.* 2020;48: D101–D110.
  39. Hsu S-D, Lin F-M, Wu W-Y, Liang C, Huang W-C, Chan W-L, et al. miRTarBase: a database curates experimentally validated microRNA–target interactions. *Nucleic Acids Research.* 2011. pp. D163–D169. doi:10.1093/nar/gkq1107
  40. Huang H-Y, Lin Y-C-D, Li J, Huang K-Y, Shrestha S, Hong H-C, et al. miRTarBase 2020: updates to the experimentally validated microRNA-target interaction database. *Nucleic Acids Res.* 2020;48: D148–D154.
  41. Sticht C, De La Torre C, Parveen A, Gretz N. miRWalk: An online resource for prediction of microRNA binding sites. *PLoS One.* 2018;13: e0206239.
  42. Friedländer MR, Mackowiak SD, Li N, Chen W, Rajewsky N. miRDeep2 accurately identifies known and hundreds of novel microRNA genes in seven animal clades. *Nucleic Acids Res.* 2012;40: 37–52.
  43. Hanan M, Soreq H, Kadener S. CircRNAs in the brain. *RNA Biol.* 2017;14: 1028–1034.
  44. Rybak-Wolf A, Stottmeister C, Glažar P, Jens M, Pino N, Giusti S, et al. Circular RNAs in the Mammalian Brain Are Highly Abundant, Conserved, and Dynamically Expressed. *Mol Cell.* 2015;58: 870–885.
  45. Galea JM, Vazquez A, Pasricha N, de Xivry J-JO, Celnik P. Dissociating the roles of the cerebellum and motor cortex during adaptive learning: the motor cortex retains what the cerebellum learns. *Cereb Cortex.* 2011;21: 1761–1770.
  46. Jiang C, Zeng X, Shan R, Wen W, Li J, Tan J, et al. The Emerging Picture of the Roles of CircRNA-CDR1as in Cancer. *Front Cell Dev Biol.* 2020;8: 590478.
  47. Yang L, Fu J, Zhou Y. Circular RNAs and Their Emerging Roles in Immune Regulation. *Front Immunol.* 2018;9: 2977.
  48. Xiao J, Joseph S, Xia M, Teng F, Chen X, Huang R, et al. Circular RNAs Acting as miRNAs' Sponges and Their Roles in Stem Cells. *J Clin Med Res.* 2022;11. doi:10.3390/jcm11102909
  49. Zhang Y, Tian Z, Ye H, Sun X, Zhang H, Sun Y, et al. Emerging functions of circular RNA in the regulation of adipocyte metabolism and obesity. *Cell Death Discov.* 2022;8: 268.
  50. Zhang Y, Zhang X-O, Chen T, Xiang J-F, Yin Q-F, Xing Y-H, et al. Circular intronic long noncoding RNAs. *Mol Cell.* 2013;51: 792–806.
  51. Li Z, Huang C, Bao C, Chen L, Lin M, Wang X, et al. Exon-intron circular RNAs regulate transcription in the nucleus. *Nat Struct Mol Biol.* 2015;22: 256–264.
  52. Li Z, Huang C, Bao C, Chen L, Lin M, Wang X, et al. Corrigendum: Exon-intron circular RNAs regulate transcription in the nucleus. *Nat Struct Mol Biol.* 2017;24: 194.
  53. Wen G, Zhou T, Gu W. The potential of using blood circular RNA as liquid biopsy biomarker for human diseases. *Protein Cell.* 2021;12: 911–946.
  54. Nielsen AF, Bindereif A, Bozzoni I, Hanan M, Hansen TB, Irimia M, et al. Best practice standards for circular RNA research. *Nat Methods.* 2022;19: 1208–1220.