



# Association of serum uric acid level with coronary artery stenosis severity in Korean end-stage renal disease patients [Volume 36, Issue 3, September 2017, Pages 282–289]

Hye Yun Jeong<sup>1</sup>, Hye Jeong Cho<sup>1</sup>, Sang Hoon Kim<sup>2</sup>, Jun Chul Kim<sup>3</sup>, Mi Jung Lee<sup>1</sup>, Dong Ho Yang<sup>1</sup>, So-Young Lee<sup>1</sup>

*Divisions of <sup>1</sup>Nephrology and <sup>2</sup>Cardiology, Department of Internal Medicine, CHA Bundang Medical Center, CHA University, Seongnam, Korea*

*<sup>3</sup>Division of Nephrology, Department of Internal Medicine, CHA Gumi Medical Center, CHA University, Gumi, Korea*

The authors would like to publish this corrigendum to clarify the roles of some authors in the above article (<https://doi.org/10.23876/j.krcp.2017.36.3.282>).

Hye Yun Jeong and Hye Jeong Cho contributed equally to the above study as co-first authors.

---

**Correspondence:** So-Young Lee

Division of Nephrology, Department of Internal Medicine, CHA Bundang Medical Center, CHA University, 59 Yatap-ro, Bundang-gu, Seongnam 13496, Korea. E-mail: [ysyo119@cha.ac.kr](mailto:ysyo119@cha.ac.kr)

ORCID: <http://orcid.org/0000-0003-1877-2124>

Dong Ho Yang

Division of Nephrology, Department of Internal Medicine, CHA Bundang Medical Center, CHA University, 59 Yatap-ro, Bundang-gu, Seongnam 13496, Korea. E-mail: [dhyang@cha.ac.kr](mailto:dhyang@cha.ac.kr)

ORCID: <http://orcid.org/0000-0001-9404-0019>

Copyright © 2018 by The Korean Society of Nephrology

© This is an open-access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc-nd/4.0/>), which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.