

BMJ Open Short-term and long-term revision rates after lumbar spine discectomy versus laminectomy: a population-based cohort study

Feng-Chen Kao,^{1,2} Yao-Chun Hsu,^{3,4,5,6,7} Chang-Bi Wang,⁸ Yuan-Kun Tu,^{1,2} Pao-Hsin Liu⁹

To cite: Kao F-C, Hsu Y-C, Wang C-B, *et al.* Short-term and long-term revision rates after lumbar spine discectomy versus laminectomy: a population-based cohort study. *BMJ Open* 2018;**8**:e021028. doi:10.1136/bmjopen-2017-021028

► Prepublication history and additional material for this paper are available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2017-021028>).

Received 21 December 2017
Revised 3 May 2018
Accepted 6 June 2018

ABSTRACT

Background/objective Degenerative diseases of the lumbar spine were managed with discectomy or laminectomy. This study aimed to compare these two surgical treatments in the postoperative revision rates.

Design A population-based cohort study from analysis of a healthcare database.

Setting Data were gathered from the Taiwan National Health Insurance Research Database (NHIRD).

Participants We enrolled 16 048 patients (4450 women and 11 598 men) with a mean age of 40.34 years who underwent lumbar discectomy or laminectomy for the first time between 1 January 1997 and 31 December 2007. All patients were followed up for 5 years or until death.

Results Revision rate within 3 months of the index surgery was significantly higher in patients who underwent discectomy (2.75%) than in those who underwent laminectomy (1.18%; $p < 0.0001$). This difference persisted over the first year following the index surgery (3.38% vs 2.57%). One year afterwards, the revision rates were similar between the discectomy (9.75%) and laminectomy (9.69%) groups. The final spinal fusion surgery rates were also similar between the groups (11.25% vs 12.08%).

Conclusion The revision rate after lumbar discectomy was higher than that after laminectomy within 1 year of the index surgery. However, differences were not identified between patient groups for the two procedures with respect to long-term revision rates and the proportion of patients who required final spinal fusion surgery.

INTRODUCTION

The natural progression of a degenerative spine leads to primary disc herniation and lumbar spinal stenosis, and most patients with these conditions are treated through surgical interventions.^{1 2} Lumbar disc herniation is a common manifestation of degenerative lumbar disc disease^{3–5} that occurs early in the degenerative cascade and involves tensile failure of the annulus to contain the gel-like nuclear portion of the disc. Although treatment for lumbar herniated discs can be challenging, non-surgical treatment is effective in most cases.^{6 7} However, studies have

Strengths and limitations of this study

- This population-based cohort study encompassed all residents of Taiwan.
- The universal and compulsory national health insurance mitigated attrition bias as no patients were lost to follow-up.
- However, radiographic and pathological data were unavailable in the Taiwan National Health Insurance Research Database. Therefore, we could not ascertain the level and pathology of the treated spine.
- The physical conditions of the patients could not be evaluated, and unmeasured confounding was possible.

indicated that surgery provides superior results to non-surgical treatments, especially with respect to short-term pain relief.^{3 8}

Lumbar spinal stenosis is a progressive and dynamic disease that constitutes a continuum of pathological changes in the spinal column as a person ages. The likelihood of lumbar spinal stenosis increases during the fifth decade of life and ranges from 1.7% to 8% in the general population.⁹ Surgical treatment focuses on a patient's pathological anatomy and involves relieving neurological compression; surgical procedures are usually more complex than those performed to relieve simple compression.¹⁰

Revision surgery, which is required in many cases of spinal disease after initial surgical treatment,^{11 12} presents a challenge for spinal surgeons. Surgeons should be attuned to the clinical circumstances that are appropriate for additional surgery and should be technically qualified to address the anatomical and pathological obstacles involved in repeat surgery. Incidence of revision surgery after lumbar surgical discectomy varies from 0% to approximately 15%.¹ Frymoyer¹³ reported incidence of postdiscectomy instability



© Author(s) (or their employer(s)) 2018. Re-use permitted under CC BY. Published by BMJ.

For numbered affiliations see end of article.

Correspondence to

Dr Yao-Chun Hsu;
holdenhsu@ntu.edu.tw

requiring further spinal fusion surgery as high as 6.5%. Relatively few reports have specifically addressed revision surgery for lumbar spinal stenosis. Malter and colleagues¹² reported that the 5-year reoperation rate for patients with spinal stenosis was as high as 12%.

To investigate whether spinal reoperation rates differ after lumbar discectomy and laminectomy for lumbar spinal stenosis, we performed a population-based retrospective study of patients' 5-year follow-up data retrieved from the Taiwan National Health Insurance Research Database (NHIRD).

DATA SOURCE

We examined data from the Taiwan NHIRD which is released by the Taiwan National Health Research Institute (NHRI) for public use. The NHRI covers the medical claims of 22.9 million residents of Taiwan, accounting for >99% of the total population. The NHIRD contains claims data from 1997 to 2013. The Department of Health and the National Health Insurance (NHI) Bureau of Taiwan ensure the completeness and accuracy of the NHIRD. This study was exempt from an ethics review because the medical records released by the insurance authority are encrypted secondary data and have been approved for use in research.

This retrospective population-based cohort study examined data from the longitudinal NHIRD. Until the end of 2013, all sampled individuals were followed up for outcome identification by using International Classification of Diseases, Ninth Revision, Clinical Modification (ICD-9-CM) codes.

MATERIALS AND METHODS

Patient and public involvement

Our study cohort included patients from the NHIRD who underwent lumbar discectomy or laminectomy for the first time between 1 January 1997 and 31 December 2007. Those who received their first lumbar discectomy or laminectomy after 2007 were excluded because dynamic stabilisation systems such as the Wallis system¹⁴ were marketed in Taiwan after 2007. We also excluded individuals who were continually exposed to oral or injected forms of systemic corticosteroids for 6 months or longer, as well as those with diseases such as ankylosing spondylitis, systemic lupus erythematosus, rheumatoid arthritis, malignant cancers, spinal tumours, congenital spinal anomalies, spinal tuberculosis, spinal infections, spinal fractures, cervical spinal disease and thoracic spinal disease; the corresponding ICD-9-CM codes are listed in (online supplementary appendix 1).

We divided the study cohort into discectomy and laminectomy groups. Each patient's date of discharge from the hospital after their first lumbar discectomy or laminectomy was considered their index date. Revision lumbar spine surgery was defined as a second lumbar spine operation performed after the index date and

comprised the following types: lumbar spine discectomy, lumbar spine laminectomy (including laminotomy) and lumbar spinal fusion surgery (with or without instrumentation). The revision rates in the two surgical groups were evaluated and compared, and the groups were propensity-score matched at a ratio of 1:1 based on the baseline characteristics of the patients. We assessed unmatched and matched data in this study.

Comorbidities existing prior to the index date were classified based on Charlson Comorbidity Index scores,¹⁵ and incidences of mortality after the index dates were calculated for both groups. Mortality rates were considered when comparing revision rates to eliminate the influence of death on the calculated likelihood of revision surgery. We also calculated and compared the rates of final revision spinal fusion surgery in the two groups. All patients were followed up until death, withdrawal from the NHI programme or 31 December 2012.

STATISTICAL ANALYSIS

We use Pearson's χ^2 test and Yates's continuity correction to compare qualitative data, whereas the Student's t-test was employed for quantitative data. The annual revision rates were calculated with 95% CIs. The association between revision lumbar spine surgery between discectomy and laminectomy was explored by the Cox proportional hazard model that took into account age, gender and baseline comorbidity. Our study analysed the lumbar spine revision surgery rate by using the Fine and Gray regression model to calculate subdistribution hazards, and p values were determined using Gray's test. The subdistribution HR (sHR) was defined as significant when $p < 0.05$. All statistical tests and calculations were performed using Statistical Analysis Software V.9.4 (SAS Institute).

RESULTS

Baseline characteristics of the patients

Our study cohort consisted of 66 754 patients (31 964 women and 34 790 men). The discectomy group comprised 27 867 patients and the laminectomy group comprised 38 887 patients. The unmatched and matched baseline characteristics and comorbidities of all patients are listed in table 1. After propensity-score matching, a total of 8024 patients were enrolled in this study. Lumbar spine revision surgery was defined as any of the following types of lumbar surgery performed after initial lumbar surgery: lumbar spine discectomy, lumbar spine laminectomy (including laminotomy) and lumbar spinal fusion surgery (with or without instrumentation). Final spinal fusion surgery referred to lumbar spinal fusion surgery (with or without instrumentation) performed during the follow-up period.

REASONS OF LUMBAR SPINE REVISION SURGERY

Causes of lumbar spine revision surgeries are listed in online supplementary table S1.1 and S1.2. The prevalence

Table 1 Characteristics and primary outcomes of patients who received laminectomy or discectomy surgeries

1. Unmatched baseline	Discectomy n=27 867	Laminectomy n=38 887	P values
Age	47.83±15.58	59.91±14.02	<0.0001
Age group			<0.0001
<20	416 (1.49)	232 (0.60)	
20–39	8987 (32.25)	3667 (9.43)	
40–59	11 511 (41.31)	13 030 (33.51)	
60–79	6663 (23.91)	20 561 (52.87)	
≥80	290 (1.04)	1397 (3.59)	
Gender			<0.0001
Female	10 629 (38.14)	21 335 (54.86)	
Male	17 238 (61.86)	17 552 (45.14)	
Comorbidities			
Myocardial infarct	149 (0.53)	404 (1.04)	<0.0001
Congestive heart failure	436 (1.56)	1632 (4.20)	<0.0001
Peripheral vascular disease	196 (0.70)	630 (1.62)	<0.0001
Cerebrovascular disease	1320 (4.74)	4050 (10.41)	<0.0001
Dementia	199 (0.71)	632 (1.63)	<0.0001
Chronic lung disease	514 (1.84)	1620 (4.17)	<0.0001
Connective tissue disease	80 (0.29)	132 (0.34)	0.2357
Ulcer	5528 (19.84)	11 362 (29.22)	<0.0001
Chronic liver disease	2593 (9.30)	4768 (12.26)	<0.0001
Diabetes	2291 (8.22)	5741 (14.76)	<0.0001
Diabetes with end organ damage	761 (2.73)	2029 (5.22)	<0.0001
Hemiplegia	80 (0.29)	238 (0.61)	<0.0001
Moderate or severe kidney disease	545 (1.96)	1590 (4.09)	<0.0001
Tumour, leukaemia, lymphoma	20 (0.07)	49 (0.13)	0.0315
Moderate or severe liver disease	52 (0.19)	98 (0.25)	0.0784
Malignant tumour, metastasis			–
AIDS	4 (0.01)	3 (0.01)	0.4087
Spinal revision surgery (3 month)	765 (2.75)	459 (1.18)	<0.0001
Discectomy	449 (1.61)	128 (0.33)	<0.0001
Laminectomy	187 (0.67)	196 (0.5)	0.0048
Spinal instrumentation	129 (0.46)	135 (0.35)	0.0188
Spinal revision surgery (3 month~1 year)	941 (3.38)	999 (2.57)	<0.0001
Discectomy	389 (1.40)	186 (0.48)	<0.0001
Laminectomy	287 (1.03)	406 (1.04)	0.8587
Spinal instrumentation	265 (0.95)	407 (1.05)	0.2220
Spinal revision surgery (>1 year)	2718 (9.75)	3770 (9.69)	0.8006
Discectomy	844 (3.03)	485 (1.25)	<0.0001
Laminectomy	708 (2.54)	1282 (3.3)	<0.0001
Spinal instrumentation	1166 (4.18)	2003 (5.15)	<0.0001
Total spinal revision surgery	4424 (15.88)	5228 (13.44)	<0.0001
Discectomy	1682 (6.04)	799 (2.05)	<0.0001
Laminectomy	1182 (4.24)	1884 (4.84)	0.0002
Spinal instrumentation	1560 (5.60)	2545 (6.54)	<0.0001

Continued

Table 1 Continued

	Discectomy n=27 867	Laminectomy n=38 887	P values
1. Unmatched baseline			
Final spinal fusion	3136 (11.25)	4699 (12.08)	0.0010
Death	3900 (14.00)	8545 (21.97)	<0.0001
2. Matched baseline			
	Discectomy n=8024	Laminectomy n=8024	P values
Age	40.16±11.26	40.51±11.51	0.0536
Age group			0.3398
<20	195 (2.43)	217 (2.70)	
20–39	3621 (45.13)	3500 (43.62)	
40–59	3922 (48.88)	4023 (50.14)	
60–79	246 (3.07)	244 (3.04)	
≥80	40 (0.50)	40 (0.50)	
Gender			1.0000
Female	2225 (27.73)	2225 (27.73)	
Male	5799 (72.27)	5799 (72.27)	
Comorbidities			
Myocardial infarct	32 (0.40)	34 (0.42)	0.8051
Congestive heart failure	87 (1.08)	88 (1.10)	0.9394
Peripheral vascular disease	49 (0.61)	60 (0.75)	0.2904
Cerebrovascular disease	215 (2.68)	220 (2.74)	0.8080
Dementia	41 (0.51)	44 (0.55)	0.7442
Chronic lung disease	86 (1.07)	79 (0.98)	0.5838
Connective tissue disease	15 (0.19)	17 (0.21)	0.7234
Ulcer	1124 (14.01)	1129 (14.07)	0.9095
Chronic liver disease	705 (8.79)	693 (8.64)	0.7369
Diabetes	431 (5.37)	412 (5.13)	0.5014
Diabetes with end organ damage	150 (1.87)	144 (1.79)	0.7240
Hemiplegia	18 (0.22)	17 (0.21)	0.8656
Moderate or severe kidney disease	107 (1.33)	113 (1.41)	0.6838
Tumour, leukaemia, lymphoma	3 (0.04)	4 (0.05)	0.7054
Moderate or severe liver disease	7 (0.09)	10 (0.12)	0.4666
Malignant tumour, metastasis			–
AIDS			–
Spinal revision surgery (3 month)	208 (2.59)	123 (1.53)	<0.0001
Discectomy	128 (1.60)	48 (0.60)	<0.0001
Laminectomy	46 (0.57)	37 (0.46)	0.3220
Spinal instrumentation	34 (0.42)	38 (0.47)	0.6366
Spinal revision surgery (3 month~1 year)	241 (3.00)	189 (2.36)	0.0110
Discectomy	109 (1.36)	54 (0.67)	<0.0001
Laminectomy	58 (0.72)	63 (0.79)	0.6482
Spinal instrumentation	74 (0.92)	72 (0.90)	0.8679
Spinal revision surgery (>1 year)	675 (8.41)	665 (8.29)	0.7754
Discectomy	278 (3.46)	181 (2.26)	<0.0001
Laminectomy	132 (1.65)	164 (2.04)	0.0605
Spinal instrumentation	265 (3.30)	320 (3.99)	0.0205

Continued

Table 1 Continued

2. Matched baseline	Discectomy n=8024	Laminectomy n=8024	P values
Total spinal revision surgery	1124 (14.01)	977 (12.18)	0.0006
Discectomy	515 (6.42)	283 (3.53)	<0.0001
Laminectomy	236 (2.94)	264 (3.29)	0.2033
Spinal instrumentation	373 (4.65)	430 (5.36)	0.0390
Final spinal fusion	784 (9.77)	838 (10.44)	0.1573
Death	795 (9.91)	884 (11.02)	0.0217

of incidental durotomy was 0.04%. The proportions of postoperative haemorrhage and postoperative spine infection were 0.18% and 1.73%, respectively. Finally, the lumbar disc pathology rate was 40.74%.

TOTAL SPINAL SURGERY REVISION RATES

The annual revision rates in the discectomy and laminectomy groups were 5.63% (95% CI 5.15% to 6.16%) and 3.92% (95% CI 3.52% to 4.37%), respectively. Values representing cumulative incidence of revision spinal surgery are displayed in [figure 1](#). Significant differences in total revision spinal surgery rates between patients

who received lumbar discectomy and those who received lumbar laminectomy as initial surgery were identified. In the unmatched data, the revision spinal surgery rates in the discectomy and laminectomy groups were 15.88% and 13.44%, respectively ($p < 0.0001$). In the matched data, the corresponding rates were 14.01% and 12.18%, respectively ($p < 0.001$).

RATES FOR REVISION SURGERY PERFORMED WITHIN 3 MONTHS OF INITIAL SPINAL SURGERY

The rates for revision spinal surgery performed within 3 months of initial spinal surgery significantly differed




Figure 1 The cumulated incidence of total revision spinal surgery after the first time spinal surgeries.

between the two groups ($p < 0.0001$). Based on the unmatched data, the revision spinal surgery rates in the discectomy and laminectomy groups were 2.75% and 1.18%, respectively. In the matched data, the corresponding rates were 2.59% and 1.53%, respectively.

RATES FOR REVISION SURGERY PERFORMED BETWEEN 3 MONTHS AND 1 YEAR AFTER INITIAL SPINAL SURGERY

The rates for revision spinal surgery performed between 3 months and 1 year after initial spinal surgery also significantly differed between patients who initially received lumbar discectomy and those who initially received lumbar laminectomy. In the unmatched data, the revision spinal surgery rates in the discectomy and laminectomy groups were 3.38% and 2.57%, respectively ($p < 0.0001$). In the matched data, the corresponding rates were 3.00% and 2.36%, respectively ($p < 0.05$).

RATES FOR REVISION SURGERY PERFORMED MORE THAN 1 YEAR AFTER INITIAL SPINAL SURGERY

The rates for revision spinal surgery performed more than 1 year after initial spinal surgery did not significantly differ between patients who initially received

lumbar discectomy and those who initially received lumbar laminectomy. In the unmatched data, the revision spinal surgery rates in the discectomy and laminectomy groups were 9.75% and 9.69%, respectively. In the matched data, the corresponding rates were 8.41% and 8.29%, respectively.

DIFFERENCES IN MULTIVARIATE-ADJUSTED TOTAL REVISION SPINAL SURGERY RATES BETWEEN DISCECTOMY AND LAMINECTOMY GROUPS

A multivariate-adjusted Cox proportional hazards model revealed independent differences in the unmatched and matched data (adjusted sHRs 0.81 and 0.86, respectively; 95% CIs 0.78 to 0.85 and 0.79 to 0.94, respectively; [table 2](#)) between the discectomy and laminectomy groups. Analysis of the unmatched data ([table 2](#)) revealed that age (sHR 1.01; 95% CI 1.00 to 1.01), sex (sHR 1.09; 95% CI 1.05 to 1.14), peripheral vascular disease (sHR 0.73; 95% CI 0.59 to 0.91) and diabetes mellitus (DM; sHR 1.09; 95% CI 1.01 to 1.17) were the risk factors responsible for differences in spinal revision rates between the discectomy and laminectomy groups. Analysis of the matched data indicated that age (sHR

Table 2 Multivariate Cox proportional hazard models for revision lumbar spine surgical rates between discectomy and laminectomy with or without matched data

	Unmatched		Matched	
	sHR (95% CI)	P values	sHR (95% CI)	P values
Laminectomy vs discectomy	0.81 (0.78 to 0.85)	<0.0001	0.86 (0.79 to 0.94)	0.0007
Age	1.01 (1.00 to 1.01)	<0.0001	1.01 (1.00 to 1.01)	0.0007
Male vs female	1.09 (1.05 to 1.14)	<0.0001	1.09 (0.99 to 1.20)	0.0937
Comorbidities				
Myocardial infarct	1.15 (0.94 to 1.42)	0.1825	1.21 (0.69 to 2.14)	0.5097
Congestive heart failure	1.04 (0.92 to 1.18)	0.4979	1.30 (0.89 to 1.90)	0.1751
Peripheral vascular disease	0.73 (0.59 to 0.91)	0.0046	0.82 (0.48 to 1.41)	0.4788
Cerebrovascular disease	0.99 (0.91 to 1.07)	0.7413	0.97 (0.73 to 1.28)	0.8136
Dementia	1.11 (0.92 to 1.33)	0.2813	1.19 (0.69 to 2.05)	0.5300
Chronic lung disease	1.05 (0.94 to 1.18)	0.4067	1.00 (0.67 to 1.50)	0.9833
Connective tissue disease	1.16 (0.83 to 1.61)	0.3925	1.73 (0.86 to 3.49)	0.1262
Ulcer	0.96 (0.92 to 1.01)	0.1285	1.12 (0.99 to 1.27)	0.0854
Chronic liver disease	0.99 (0.92 to 1.06)	0.7486	1.14 (0.98 to 1.33)	0.0917
Diabetes	1.09 (1.01 to 1.17)	0.0263	1.14 (0.92 to 1.42)	0.2392
Diabetes with end organ damage	1.12 (1.00 to 1.25)	0.0590	0.99 (0.70 to 1.40)	0.9436
Hemiplegia	1.18 (0.88 to 1.57)	0.2672	1.18 (0.52 to 2.72)	0.6897
Moderate or severe kidney disease	1.09 (0.97 to 1.23)	0.1319	0.83 (0.57 to 1.22)	0.3431
Tumour, leukaemia, lymphoma	1.40 (0.80 to 2.47)	0.2434	NA	
Moderate or severe liver disease	1.36 (0.90 to 2.06)	0.1399	1.34 (0.43 to 4.22)	0.6124
Malignant tumour, metastasis	NA		NA	–
AIDS	1.11 (0.16 to 7.90)	0.9149	NA	–

sHR, subdistribution HR.




Figure 2 The cumulated incidence of final spinal fusion surgery after the first time spinal surgeries.

1.01; 95% CI 1.00 to 1.01) was the risk factor responsible for differences in spinal revision rates between the two groups.

RATES FOR FINAL SPINAL FUSION SURGERY PERFORMED AFTER INITIAL SPINAL SURGERY

The annual revision rates in the discectomy and laminectomy groups were 2.38% (95% CI 2.07% to 2.75%) and 2.16% (95% CI 1.86% to 2.51%), respectively. The value representing cumulative incidence of final spinal fusion surgery performed after initial spinal surgery is displayed in [figure 2](#). No significant differences in the rates for final spinal fusion surgery performed after initial surgery were identified between patients who initially received lumbar discectomy and those who initially received lumbar laminectomy. In the unmatched data, the final spinal fusion surgery rates in the discectomy and laminectomy groups were 11.25% and 12.08%, respectively. In the matched data, the corresponding rates were 9.77% and 10.44%, respectively.

DIFFERENCES IN MULTIVARIATE-ADJUSTED RATES OF FINAL SPINAL FUSION SURGERY PERFORMED AFTER INITIAL SPINAL SURGERY BETWEEN DISCECTOMY AND LAMINECTOMY GROUPS

The multivariate-adjusted Cox proportional hazards model revealed no differences in the unmatched data between the discectomy and laminectomy groups (adjusted sHR 1.05; 95% CI 1.00 to 1.10; [table 3](#)). However, the model revealed independent differences in the matched data between the groups (adjusted sHR 1.11; 95% CI 1.01 to 1.22). In the unmatched data analysis ([table 3](#)), age (sHR 1.00; 95% CI 1.00 to 1.01), chronic lung disease (sHR 1.15; 95% CI 1.01 to 1.30), ulcer (sHR 1.18; 95% CI 1.12 to 1.24), chronic liver disease (sHR 1.21; 95% CI 1.13 to 1.30), DM (sHR 1.29; 95% CI 1.19 to 1.39) and moderate or severe kidney disease (sHR 1.20; 95% CI 1.06 to 1.36) were the risk factors for different final spinal fusion rates between the discectomy and laminectomy groups. In the matched data analysis, age (sHR 1.02; 95% CI 1.01 to 1.02), ulcer (sHR 1.34; 95% CI 1.16 to 1.55) and chronic liver disease (sHR 1.37; 95% CI 1.16 to 1.62) were the corresponding risk factors.

Table 3 Multivariate Cox proportional hazard models for final revision lumbar spine fusion rates between discectomy and laminectomy with or without matched data

	Unmatched		Matched	
	sHR (95% CI)	P values	sHR (95% CI)	P values
Laminectomy vs discectomy	1.05 (1.00 to 1.10)	0.0524	1.11 (1.01 to 1.22)	0.0377
Age	1.00 (1.00 to 1.01)	<0.0001	1.02 (1.01 to 1.02)	<0.0001
Comorbidities				
Myocardial infarct	1.16 (0.92 to 1.45)	0.2131	0.95 (0.47 to 1.91)	0.8832
Congestive heart failure	1.06 (0.93 to 1.21)	0.4071	1.16 (0.75 to 1.78)	0.5045
Peripheral vascular disease	0.96 (0.78 to 1.18)	0.6927	0.90 (0.52 to 1.57)	0.7183
Cerebrovascular disease	1.04 (0.95 to 1.13)	0.3858	1.07 (0.80 to 1.45)	0.6419
Dementia	1.13 (0.93 to 1.38)	0.2320	0.87 (0.45 to 1.69)	0.6863
Chronic lung disease	1.15 (1.01 to 1.30)	0.0351	0.95 (0.61 to 1.50)	0.8351
Connective tissue disease	0.89 (0.59 to 1.34)	0.5653	1.09 (0.46 to 2.60)	0.8492
Ulcer	1.18 (1.12 to 1.24)	<0.0001	1.34 (1.16 to 1.55)	<0.0001
Chronic liver disease	1.21 (1.13 to 1.30)	<0.0001	1.37 (1.16 to 1.62)	0.0002
Diabetes	1.29 (1.19 to 1.39)	<0.0001	1.19 (0.93 to 1.54)	0.1730
Diabetes with end organ damage	1.11 (0.98 to 1.25)	0.0887	0.95 (0.65 to 1.40)	0.7954
Hemiplegia	1.12 (0.80 to 1.56)	0.5194	0.43 (0.10 to 1.80)	0.2471
Moderate or severe kidney disease	1.20 (1.06 to 1.36)	0.0042	1.04 (0.71 to 1.53)	0.8466
Tumour, leukaemia, lymphoma	1.31 (0.71 to 2.41)	0.3819	NA	
Moderate or severe liver disease	1.36 (0.87 to 2.13)	0.1778	1.02 (0.26 to 4.01)	0.9728
Malignant tumour, metastasis	NA		NA	–
AIDS	1.91 (0.32 to 11.39)	0.4762	NA	–

sHR, subdistribution HR.

DISCUSSION

Lumbar disc herniation is one of the most common lumbar spine disorders.¹⁶ In 1934, Mixter and Barr¹⁷ identified a link between sciatica and lumbar disc herniation; since this discovery, discectomy through limited laminotomy has been the most common form of surgical management for lumbar disc prolapse in cases of conservative management failure.¹⁸ The efficacy of lumbar discectomy for treating lumbar disc herniation has been demonstrated^{19 20}; however, unsatisfactory outcomes after lumbar discectomy have been reported in 5%–20% of cases.^{21–24} The Spine Patient Outcomes Research Trial reported that in patients with lumbar disc herniation, the proportions of reoperation within 4 and 8 years of index procedures were as high as 9% for discectomy patients and 13% for laminectomy patients.¹⁹ The most common cause of ongoing disability after lumbar discectomy is recurrent lumbar disc herniation which occurs in 5%–15% of patients (this incidence proportion increases over time).^{21 23 25–28} In our study cohort, the rates for revision spinal surgery performed within 3 months and 1 year of lumbar discectomy were 2.75% and 3.38%, respectively; those for revision surgery performed after 1 year and of total revision surgery were 9.75% and 15.88%, respectively.

Lumbar stenosis is caused by spondylotic changes in the facet joints, spinal instability or a congenitally small spinal canal.²⁹ Laminectomy remains the standard treatment for spinal stenosis when the spine does not exhibit instability.²⁹ Despite adequate lumbar decompression, substantial postoperative back and leg pain occur in 10%–15% of patients.³⁰ Historically, a high proportion of lumbar laminectomies fail, and the proportion of patients who experience recurrent back pain may reach 47%.^{31 32} No reoperation rates after lumbar laminectomy without spinal fusion surgery have been reported. In our study, the rates for revision spinal surgery performed within 3 months and 1 year of lumbar laminectomy were 1.18% and 2.57%, respectively; those for revision surgery performed after 1 year and for total revision surgery were 9.69% and 13.44%, respectively.

Spinal structures that contribute to spinal stability in certain proportions of patients are as follows: facet capsule 39%, disc and annulus 29%, supraspinous and intraspinal ligaments 19% and ligamentum flavum 13%.³³ Interventions at the hemilamina and ligamentum flavum can change both the load-bearing and kinematic characteristics of the spine and lead to spinal segment hypermobility and accelerated bone degeneration.^{34 35} Even microdiscectomy can increase the risk of single-level

instability.³⁶ Extensive laminectomy can also potentiate spinal instability.^{37 38} Lai *et al*³⁹ reported that sacrificing supraspinous ligaments or tendon insertion points in spinous processes can accelerate development of adjacent instability. Incidences of adjacent instability increase with the number of destructed laminae, and far more posterior spinal complexes are destructed in lumbar laminectomy than in lumbar discectomy. Hence, theoretically, lumbar laminectomy causes greater spinal instability than does lumbar discectomy, leading to a higher reoperation rate after lumbar laminectomy.

In contrast to the theoretically expected outcomes, our study revealed independent differences in reoperation rates based on the unmatched and matched data (adjusted sHR 0.81 and 0.86; 95% CI 0.78 to 0.85 and 0.79 to 0.94, respectively) between the discectomy and laminectomy groups. Based on the unmatched data, revision spinal surgery rates in the discectomy and laminectomy groups were 15.88% and 13.44%, respectively ($p < 0.0001$). According to the matched data, the corresponding rates in the discectomy and laminectomy groups were 14.01% and 12.18%, respectively ($p < 0.001$). Compared with the laminectomy group, the discectomy group had higher rates of reoperation within 3 months and between 3 months and 1 year after initial surgery ($p < 0.05$). However, beyond 1 year, the reoperation rates did not significantly differ between the laminectomy and discectomy groups.

Numerous reasons for reoperation after discectomy have been suggested. Early recurrence may be due to reherniation, infection or arachnoiditis, whereas late recurrence may be attributed to foraminal stenosis, a painful disc, epidural fibrosis, iatrogenic segmental instability, progressive facet degeneration or sacroiliac joint pain.^{40–42} Outcomes based on natural degeneration of the lumbar spine more than 1 year after initial lumbar spine surgery were similar in the discectomy and laminectomy groups.

North *et al*⁴³ reported that incidence of instability increased from 12.5% after initial revision surgery to 50% after the fourth surgery. Fusion of the symptomatic spinal segment during revision spinal surgery is related to successful outcomes.^{44–47} In our study, no significant differences were observed in the final spinal fusion surgery rates after initial spinal surgery between patients who received lumbar discectomy (11.25%) and those who received lumbar laminectomy (12.08%).

Our study had some limitations. First, the laboratory, radiographic and pathological data of the patients were unavailable in the NHIRD. Thus, we were unable to differentiate between true lumbar disc prolapse and spinal canal stenosis. Second, the physical conditions of the study cohort patients could not be evaluated; this may have led to healthy patient bias. Nevertheless, this stringent definition would have biased the results towards a null association rather than creating a spurious one. In addition, the potential influence of body weight, habitual cigarette smoking, alcohol consumption and dietary habits could not be assessed because related information

was unavailable in the NHIRD. We were also unable to acquire direct information on these factors because linking the NHIRD with external databases is strictly prohibited for privacy protection. However, an advantage of the NHIRD is its inclusion of information on 99% of the residents of Taiwan, and no patients in our NHIRD study cohort were lost to follow-up. The complete follow-up in this study was particularly attributable to hospital accessibility.

In conclusion, rates for reoperation within 1 year were higher after lumbar discectomy than after lumbar laminectomy. Beyond 1 year after initial lumbar surgery, reoperation rates and final lumbar spinal fusion surgery rates were similar in the discectomy and laminectomy groups.

Author affiliations

¹Department of Orthopedics, E-Da Hospital, Kaohsiung, Taiwan

²School of Medicine for International Students, I-Shou University, Kaohsiung, Taiwan

³School of Medicine, Fu-Jen Catholic University, New Taipei, Taiwan

⁴Big Data Research Center, Fu-Jen Catholic University, New Taipei, Taiwan

⁵Division of Gastroenterology, Fu-Jen Catholic University Hospital, New Taipei, Taiwan

⁶Graduate Institute of Clinical Medicine, China Medical University, Taichung, Taiwan

⁷Division of Gastroenterology and Hepatology, E-Da Hospital, Kaohsiung, Taiwan

⁸Graduate Institute of Public Health, China Medical University, Taichung, Taiwan

⁹Department of Biomedical Engineering, I-Shou University, Kaohsiung, Taiwan

Contributors Substantial contributions to the conception or design of this work or the acquisition, analysis, or interpretation of data for this work: Y-CH and Y-KT. Drafting the work or revising it critically for important intellectual content: F-CK, C-BW and P-HL. Final approval of the version to be published: C-BW and P-HL. Agreement to be accountable for all aspects of this work in ensuring that questions related to the accuracy or integrity of any part of this work are appropriately investigated and resolved: Y-CH and Y-KT.

Funding This work was supported by the Center for Database Research, E-DA HEALTHCARE GROUP, and E-Da Hospital (EDAHP103048).

Competing interests None declared.

Patient consent Not required.

Ethics approval Institutional Review Board of E-Da hospital (EMRP-104-04) and the Taiwan National Health Research Institute (NHIRD-104-167).

Provenance and peer review Not commissioned; externally peer reviewed.

Data sharing statement No additional data available.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution 4.0 Unported (CC BY 4.0) license, which permits others to copy, redistribute, remix, transform and build upon this work for any purpose, provided the original work is properly cited, a link to the licence is given, and indication of whether changes were made. See: <https://creativecommons.org/licenses/by/4.0/>.

REFERENCES

- Hoffman RM, Wheeler KJ, Deyo RA. Surgery for herniated lumbar discs: a literature synthesis. *J Gen Intern Med* 1993;8:487–96.
- McCulloch JA. Focus issue on lumbar disc herniation: macro- and microsurgery. *Spine* 1996;21(24 Suppl):45s–56.
- Atlas SJ, Keller RB, Chang Y, *et al*. Surgical and nonsurgical management of sciatica secondary to a lumbar disc herniation: five-year outcomes from the Maine Lumbar Spine Study. *Spine* 2001;26:1179–87.
- DePalma AF, Rothman RH. Surgery of the lumbar spine. *Clin Orthop Relat Res* 1969;63:162–70.
- Fisher RG, Saunders RL. Lumbar disc protrusion in children. *J Neurosurg* 1981;54:480–3.

6. Saal JA, Saal JS. Nonoperative treatment of herniated lumbar intervertebral disc with radiculopathy. An outcome study. *Spine* 1989;14:431–7.
7. Saal JA, Saal JS, Herzog RJ. The natural history of lumbar intervertebral disc extrusions treated nonoperatively. *Spine* 1990;15:683–6.
8. Weber H, herniation Ldisc, controlled A. prospective study with ten years of observation. *Spine* 1983;8:131–40.
9. Rosen CD, Kahanovitz N, Bernstein R, et al. A retrospective analysis of the efficacy of epidural steroid injections. *Clin Orthop Relat Res* 1988;228:270–2.
10. Porter RW. Spinal stenosis and neurogenic claudication. *Spine* 1996;21:2046–52.
11. Keskimäki I, Seitsalo S, Osterman H, et al. Reoperations after lumbar disc surgery: a population-based study of regional and interspecialty variations. *Spine* 2000;25:1500–8.
12. Malter AD, McNeney B, Loeser JD, et al. 5-year reoperation rates after different types of lumbar spine surgery. *Spine* 1998;23:814–20.
13. Frymoyer JW. Back pain and sciatica. *N Engl J Med* 1988;318:291–300.
14. Sénégas J. Mechanical supplementation by non-rigid fixation in degenerative intervertebral lumbar segments: the Wallis system. *Eur Spine J* 2002;11(Suppl 2):S164–9.
15. Charlson ME, Pompei P, Ales KL, et al. A new method of classifying prognostic comorbidity in longitudinal studies: development and validation. *J Chronic Dis* 1987;40:373–83.
16. Li Z, Tang J, Hou S, et al. Four-year follow-up results of transforaminal lumbar interbody fusion as revision surgery for recurrent lumbar disc herniation after conventional discectomy. *J Clin Neurosci* 2015;22:331–7.
17. Mixer WJ, Barr JS. Rupture of the Intervertebral Disc with Involvement of the Spinal Canal. *N Engl J Med Overseas Ed* 1934;211:210–5.
18. Morgan-Hough CV, Jones PW, Eisenstein SM. Primary and revision lumbar discectomy. A 16-year review from one centre. *J Bone Joint Surg Br* 2003;85:871–4.
19. Lurie JD, Tosteson TD, Tosteson AN, et al. Surgical versus nonoperative treatment for lumbar disc herniation: eight-year results for the spine patient outcomes research trial. *Spine* 2014;39:3–16.
20. Yorimitsu E, Chiba K, Toyama Y, et al. Long-term outcomes of standard discectomy for lumbar disc herniation: a follow-up study of more than 10 years. *Spine* 2001;26:652–7.
21. Cinotti G, Roysam GS, Eisenstein SM, et al. Ipsilateral recurrent lumbar disc herniation. A prospective, controlled study. *J Bone Joint Surg Br* 1998;80:825–32.
22. Cheng J, Wang H, Zheng W, et al. Reoperation after lumbar disc surgery in two hundred and seven patients. *Int Orthop* 2013;37:1511–7.
23. Aizawa T, Ozawa H, Kusakabe T, et al. Reoperation for recurrent lumbar disc herniation: a study over a 20-year period in a Japanese population. *J Orthop Sci* 2012;17:107–13.
24. Carragee EJ, Han MY, Suen PW, et al. Clinical outcomes after lumbar discectomy for sciatica: the effects of fragment type and anular competence. *J Bone Joint Surg Am* 2003;85-A:102–8.
25. Swartz KR, Trost GR. Recurrent lumbar disc herniation. *Neurosurg Focus* 2003;15:1–4.
26. Ruetten S, Komp M, Merk H, et al. Recurrent lumbar disc herniation after conventional discectomy: a prospective, randomized study comparing full-endoscopic interlaminar and transforaminal versus microsurgical revision. *J Spinal Disord Tech* 2009;22:122–9.
27. Ambrossi GL, McGirt MJ, Sciubba DM, et al. Recurrent lumbar disc herniation after single-level lumbar discectomy: incidence and health care cost analysis. *Neurosurgery* 2009;65:574–8.
28. Miwa S, Yokogawa A, Kobayashi T, et al. Risk factors of recurrent lumbar disk herniation: a single center study and review of the literature. *J Spinal Disord Tech* 2015;28:E265–9.
29. Herkowitz H, Garfin E. *Bell & Balderston Rothman-Simeone the spine*. 6th edn, 2011.
30. Booth KC, Bridwell KH, Eisenberg BA, et al. Minimum 5-year results of degenerative spondylolisthesis treated with decompression and instrumented posterior fusion. *Spine* 1999;24:1721–7.
31. Jackson RK. The long-term effects of wide laminectomy for lumbar disc excision. A review of 130 patients. *J Bone Joint Surg Br* 1971;53:609–16.
32. Fritsch EW, Heisel J, Rupp S. The failed back surgery syndrome: reasons, intraoperative findings, and long-term results: a report of 182 operative treatments. *Spine* 1996;21:626–33.
33. Adams MA, Hutton WC. The mechanical function of the lumbar apophyseal joints. *Spine* 1983;8:327–30.
34. Kaigle AM, Holm SH, Hansson TH. Experimental instability in the lumbar spine. *Spine* 1995;20:421–30.
35. Nachemson A. Lumbar spine instability. A critical update and symposium summary. *Spine* 1985;10:290–1.
36. Schaller B. Failed back surgery syndrome: the role of symptomatic segmental single-level instability after lumbar microdiscectomy. *Eur Spine J* 2004;13:193–8.
37. Ebara S, Harada T, Hosono N, et al. Intraoperative measurement of lumbar spinal instability. *Spine* 1992;17(3 Suppl):44–50.
38. Iida Y, Kataoka O, Sho T, et al. Postoperative lumbar spinal instability occurring or progressing secondary to laminectomy. *Spine* 1990;15:1186–9.
39. Lai PL, Chen LH, Niu CC, et al. Relation between laminectomy and development of adjacent segment instability after lumbar fusion with pedicle fixation. *Spine* 2004;29:2527–32. Discussion 32.
40. Slipman CW, Shin CH, Patel RK, et al. Etiologies of failed back surgery syndrome. *Pain Med* 2002;3:200–14. Discussion 14–7.
41. Burton CV, Kirkaldy-Willis WH, Yong-Hing K, et al. Causes of failure of surgery on the lumbar spine. *Clin Orthop Relat Res* 1981;157:191–9.
42. Waguespack A, Schofferman J, Slosar P, et al. Etiology of long-term failures of lumbar spine surgery. *Pain Med* 2002;3:18–22.
43. North RB, Campbell JN, James CS, et al. Failed back surgery syndrome: 5-year follow-up in 102 patients undergoing repeated operation. *Neurosurgery* 1991;28:685–91. Discussion 90–1.
44. Lakkol S, Bhatia C, Taranu R, et al. Efficacy of less invasive posterior lumbar interbody fusion as revision surgery for patients with recurrent symptoms after discectomy. *J Bone Joint Surg Br* 2011;93:1518–23.
45. Wong CB, Chen WJ, Chen LH, et al. Clinical outcomes of revision lumbar spinal surgery: 124 patients with a minimum of two years of follow-up. *Chang Gung Med J* 2002;25:175–82.
46. Laus M, Alfonso C, Tigani D, et al. Failed back syndrome: a study on 95 patients submitted to reintervention after lumbar nerve root decompression for the treatment of spondylotic lesions. *Chir Organi Mov* 1994;79:119–26.
47. Duggal N, Mendiondo I, Pares HR, et al. Anterior lumbar interbody fusion for treatment of failed back surgery syndrome: an outcome analysis. *Neurosurgery* 2004;54:636–44. Discussion 43–4.