


Retention of Knowledge and Skills After a Basic Life Support Course for Schoolchildren: A Prospective Study

INQUIRY: The Journal of Health Care Organization, Provision, and Financing
Volume 59: 1–6
© The Author(s) 2022
Article reuse guidelines:
sagepub.com/journals-permissions
DOI: 10.1177/00469580221098755
journals.sagepub.com/home/inq


Vesna Borovnik Lesjak, MD¹ , Andrej Šorgo, PhD^{2,3}, and Matej Strnad, MD, PhD^{1,4,5}

Abstract

Courses on basic life support (BLS) and automated external defibrillator (AED) in schools lead to increase in knowledge but its retention is less well explored. We aimed to explore the long-term retention of knowledge and practical skills among schoolchildren after a BLS and AED course to be able to tailor future courses accordingly. Study was conducted in 3 parts and included 823 seventh and ninth graders from different elementary schools in Maribor, Slovenia. In Study 1 (n=611) we assessed students' baseline knowledge and immediate knowledge gain after our BLS and AED course with a validated questionnaire; in Study 2 (n=116) we assessed retention of gained knowledge and skills after 5 months with a modified Cardiff test and Little Anne QCPR manikin; in Study 3 (n=96) we assessed retention of knowledge 2 years after the course. Mean differences in knowledge before and after the course in Study 1 and between studies were analyzed using paired t-tests and independent t-tests. Differences between individual question scores at different time points were compared using Mann – Whitney U test. A two-sided P<0,05 was considered significant. Practical skills retention was presented with descriptive statistics. Knowledge gain was significant immediately after the course with 83% correct answers compared to 60% at baseline. Scores dropped significantly after 5 months (73%) and after 2 years (75%), but remained significantly better than at baseline (P<0.001). Practical skills performance score as per Cardiff test after 5 months was 63%. Overall BLS performance score as per QCPR app was 59%, with an overall cardio score of 77% (average compression rate: 124/min and depth: 52 mm) and ventilation score of 44%. This study showed that long term retention of theoretical knowledge was satisfying whereas poor practical skills performance after 5 months calls for a more intense practical training on repeat courses.

Keywords

automated external defibrillator, cardiopulmonary resuscitation, knowledge, out-of-hospital cardiac arrest, schoolchildren, teaching

¹Emergency Medical Services Unit, Zdravstveni Dom Dr Adolfa Drolca Maribor, Slovenia

²Faculty of Natural Sciences and Mathematics, University of Maribor, Slovenia

³Faculty of Electrical Engineering and Computer Science, University of Maribor, Slovenia

⁴Faculty of Medicine, University of Maribor, Slovenia

⁵Emergency Department, Univerzitetni Klinični Center Maribor, Slovenia

Corresponding Author:

Vesna Borovnik Lesjak, Emergency Medical Services Unit, Zdravstveni Dom dr Adolfa Drolca Maribor, Cesta proletarskih brigad 21, 2000, Maribor 2001, Slovenia.

Email: vesnick@gmail.com



Creative Commons Non Commercial CC BY-NC: This article is distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 License (<https://creativecommons.org/licenses/by-nc/4.0/>) which permits non-commercial use, reproduction and distribution of the work without further permission provided the original work is attributed as specified on the SAGE and

Open Access pages (<https://us.sagepub.com/en-us/nam/open-access-at-sage>).

What do we already know about this topic?

A well-established fact is that knowledge and skills decline over time and there are suggested time intervals for repeat courses but there is no definitive data on how to conduct repeat courses on basic life support (BLS) and use of automated external defibrillators (AED).

How does your research contribute to the field?

Our findings add valuable data to scarce existing literature on retention of knowledge and skills after BLS and AED courses for schoolchildren.

What are your research's implications towards theory, practice, or policy?

Our findings suggest that repeat BLS and AED courses should be more focused on practical skills training as retention of theoretical knowledge is satisfactory.

Introduction

The importance of educating a wide population and of imparting a positive attitude towards cardiopulmonary resuscitation (CPR) has been recognized for some time now.^{1,2} CPR training in schools is an effective and vital part of disseminating knowledge, skills, and positive attitude towards CPR among laypersons.^{3,4} Schoolchildren are particularly susceptible and motivated for learning CPR and can be easily accessed and quickly taught.⁵⁻⁷ They serve not only as potential rescuers but also as multipliers of CPR knowledge and positive attitude towards action in cases of OHCA in their environment.^{8,9}

It has long been established that BLS and AED courses improve knowledge immediately after the course whereas retention of knowledge and skills is less well explored.¹⁰⁻¹² Existing studies advocate annual training.^{3,7,10,13} The »Kids save lives« statement endorsed by The World Health Organization recommends 2 hours of BLS and AED training annually from the age of 12 years.¹⁴ However, implementation of such initiatives and other solutions depend on the motivation of individual countries. Unfortunately, only 5 European Union countries (Belgium, Denmark, France, Italy, and Portugal) have CPR education in schools mandated by law, whereas in 16 countries (including Slovenia), CPR education is merely a suggestion.¹⁵

In our study we aimed to explore the longitude of knowledge retention and the magnitude of practical skills retention among schoolchildren after our BLS and AED course to be able to tailor future and repeat courses to make them more efficient and accessible.

Methods

Our study was divided into 3 parts. In Study 1 we assessed students' baseline knowledge and immediate knowledge gain after our BLS and AED course; In Study 2 we assessed retention of knowledge and practical skills after 5 months in a smaller sample of the population from Study 1; In Study 3 we assessed long-term retention of knowledge in a sample of students that completed our BLS and AED course 2 years ago.

Description of the Course

The course was organized by the Emergency Medical Services Unit of the Maribor Community Healthcare Center in cooperation with the City of Maribor Municipality, which financed the courses for all interested public elementary schools in the Municipality of Maribor. Maribor is the second biggest city in Slovenia with approximately 110.000 inhabitants.

The course was designed for individual classes consisting of up to 30 seventh- and ninth-grade students of Slovenian compulsory 9 year basic school. It was divided into 2 parts, each lasting 1 academic hour (45 minutes). In the first part, an emergency physician gave students an interactive lecture on principles of BLS and AED. The course was continued with a practical workshop, led by emergency technicians (paramedics) and physicians, where each student practiced on their own training torso manikin (Prestan Professional Adult Manikin, Prestan, Mayfield Village, OH, USA) with an AED prop – a cardboard sample with adhesive paper electrodes. A single real training AED (Defibtech Trainer AED, Defibtech, Guilford, CT, USA) was used for guidance and cardiac arrest simulation.

Sample

Sampling in all 3 studies was based on the incidental application of any of the twenty eligible schools in the pre-determined time frame of the studies: from October to December 2018 for Study 1 and in March and April 2019 for Study 2. Study 3 was conducted in October and November 2019 with students that completed the same BLS and AED course 2 years prior and were not included in Studies 1 or 2.

In Study 1 we investigated schoolchildren's knowledge of BLS and AED before and immediately after the course (t_0 and t_1 , respectively). The sample consisted of 623 students from 9 elementary schools but only 611 of them (283 seventh- and 328 ninth graders) fulfilled both the pre- and post-course questionnaire and were therefore eligible to be included in the study. Of those, 310 were boys and 299 were girls, two students refrained from stating their gender. Only 148 students have received any prior BLS training.

In Study 2, we conducted a knowledge and skills test 5 months after the course (t_2) in 2 elementary schools that had previously been included in Study 1. Only students that had attended the course 5 months prior were included. One hundred and sixteen students from 2 seventh and 2 ninth grades correctly fulfilled questionnaires and completed practical testing. Of those, 67 were seventh graders and 49 ninth graders and 63 girls and 53 boys, respectively. The average age of children in seventh and ninth grade of the 9 years compulsory elementary schools in Slovenia is 12 and 14 years, respectively.

In the third study, we investigated long-term retention of knowledge after our BLS and AED course at 2 years (t_3). Five ninth grades from 3 different schools that completed our BLS and AED course 2 years prior were included. Only students that attended the course 2 years ago (in seventh grade), completed the questionnaire. Forty-nine girls and 47 boys were included. Study design is presented in Figure 1.

Measuring Instrument

The first part of the instrument contained demographic data (gender, school grade, attendance to any previous BLS courses). The second part was a validated knowledge test consisting of ten multiple-choice questions about BLS and AED. Each question offered 5 answers of which only 1 was correct, therefore allowing students to achieve a maximal score of ten points.¹¹

Questionnaires were handed out immediately before the lecture and immediately after the practical workshop in Study 1. In Study 2, they were handed out and fulfilled before the practical evaluation. In Study 3, they were handed out before the repeat BLS and AED course. The content of the paper and pencil test was identical throughout all studies. It did not

require provision of a name to assure anonymity. In Study 1, students were asked to come up with individual codes that were used on the pre- and post-course test for the purpose of pairing tests of the same individuals for further analyses. A written informed consent was obtained from each student prior to fulfilling the questionnaires in each study. As per Slovenian rules for such type of educational research general written consent of parents is collected in advance at the beginning of each school year. Request for approval of the National Medical Ethics Committee had been filed and granted (dated October 23, 2018; filed under 0120-549/2017/9).

Practical Skills Testing

Practical skills were tested only in Study 2. Students were presented with a theoretical cardiac arrest scenario - they were witnesses to someone collapsing. Practical skills evaluation was conducted on analytical manikins (Little Anne QCPR, Laerdal Medical, Stavanger, Norway). They were connected via Bluetooth connection to a registered application on smartphones (Little Anne QCPR Instructor App, version 3.4.11, Laerdal Medical, Stavanger, Norway). Two instructors (emergency medicine specialists) observed and evaluated their skills with the use of a modified Cardiff test.¹⁶

Statistical Analyses

Statistical significance of the mean differences in knowledge before and after the course in Study 1 were calculated using paired t-test. Differences between unpaired test scores from Study 2 and 3 were compared using independent t-tests. Differences between individual question scores before and after intervention and at different time points were compared using Mann – Whitney U test. A two-sided $P < 0,05$ was considered significant.

SPSS software (SPSS Inc, Chicago, IL, USA) was used for statistical analysis.

Results

As expected, knowledge gain was significant immediately after a BLS and AED course (Table 1). Although the overall level of knowledge significantly decreased after 5 months and also after 2 years (test score declined from 83% to 73% and




Figure 1. Flowsheet summarizing study design and timeline.

Table 1. Comparison of scores on knowledge tests at all time points (no missing data).

		N	Mean Score (SD)
Study 1	t_0	611	6.02 (2.06)
	t_1	611	8.27 (1.61)*
Study 2	t_2	116	7.33 (2.19)**
Study 3	t_3	96	7.51 (1.54)**

Notes: *, $P < 0.001$ compared to t_0 ; **, $P < 0.001$ compared to t_1 .

75%, respectively) it was still significantly better than at baseline (60% before the course). Level of knowledge did not change significantly from 5 months to 2 years after the course (Table 1). Recognition of cardiac arrest remained at a comparable level 2 years after the course compared to baseline (66% vs 68%, respectively). Even in case of uncertainty whether a person was in a cardiac arrest or not most of the students that provided an answer would theoretically act correctly (95% compared to 32% at baseline). However, the difference was not statistically significant. After 2 years, almost 80% of students theoretically remembered how to check breathing and deliver artificial breaths which was significantly better than at baseline when about 65% knew the correct answer. Theoretical recognition of agonal breathing was also significantly better than at baseline (89% after 2 years compared to 64% at baseline). They remained knowledgeable about the correct ratio of chest compression to artificial breaths

(69% after 2 years compared to 52% at baseline) and the location of chest compressions (around 67% after 2 years vs 48% at baseline). Most of the students were still familiar with the use of an AED after 5 months and 2 years (75% and 85% compared to 45% at baseline). Scores on individual questions on the knowledge tests are shown in [Supplementary material](#).

Retention of practical skills, however, did not reflect the satisfying retention of theoretical knowledge about certain skills (Table 2). The average total score on the modified Cardiff test was 28 out of 44 total (63%). A lot of the steps of the BLS and AED algorithm were neglected or omitted. The subjective and objective assessment of chest compressions (with the modified Cardiff test and QCPR app, respectively) was outstanding compared to other steps, especially ventilations. Hand location was well adapted. Compression rate was mostly out of desired range (too slow on average). However, the quality of chest compressions was satisfying

Table 2. Correct performance of practical skills 5 months after the course as per modified Cardiff test (t_2 ; $n = 116$).

BLS Step	Activity	Performance, N (%)	
		Correct	Incorrect/None
Safety	Ensure safety for self, victim, and bystanders	5 (4.3)	111 (95.7)
Responsiveness	Gently shake victim	59 (50.9)	57 (49.1)
	Look for help	8 (6.9)	108 (93.1)
Airway and breathing	Open airway: Tilt head and lift chin	20 (17.2)	63 (54.3)
	Check airway	24 (20.7)	92 (79.3)
	Check breathing (look, listen, feel) ≤ 10 s	23 (19.8)	63 (54.3)
Shout for help, call 112	Phone EMS	6 (5.2)	110 (94.8)
	Shout for help, send for AED	5 (4.3)	111 (95.7)
CPR (2 minutes; 5 \times 30:2)	Hand position	89 (76.7)	7 (6)
	Open airway: Tilt head and lift chin	39 (33.6)	39 (33.6)
	Close nostrils	63 (54.3)	53 (45.7)
	Blow in victim's mouth and check for chest rise	25 (21.6)	91 (78.4)
AED	Open/turn on AED	92 (79.3)	24 (20.7)
	Attach adhesive pads on skin	92 (79.3)	0 (0)
	Visual and verbal hands-off check during AED analysis	45 (38.8)	71 (61.2)
	Visual and verbal hands-off check before pushing button and pushing button	34 (29.3)	63 (54.3)

Table 3. Quality of BLS as per Little Anne QCPR app (sample size = 116).

		N	Mean (SD)	Min	Max
Overall score	Overall score* (%)	116	59 (19.67)	0	99
	Flow fraction (%)	116	74 (9.89)	54	100
	Compressions per cycle	108	33 (7.44)	11	68
	Ventilations per cycle	34	1 (.82)	0	2
Cardio score	Overall (%)	116	77 (23.46)	0	99
	Compressions fully released (%)	116	96 (10.75)	34	100
	Average depth (mm)	116	50 (3.35)	30	51
	Adequate depth (%)	116	80 (31.31)	0	100
	Average rate (/min)	116	124 (14.95)	82	171
	Adequate rate (%)	116	29 (32.67)	0	98
Ventilation score	Overall (%)	34	44 (32.33)	7	100
	Ventilations with adequate chest rise (%)	34	99 (5.21)	75	100
	Too much chest rise (%)	34	1 (5.21)	0	25

Notes: N, number of recorded and analyzed actions by the application; *, based on compressions, ventilations and flow.

with adequate depth and release in more than 80% (Table 3). They were also confident about AED use, but mostly forgot about safety measures.

Ventilations were omitted by most students. Consequently, the ventilation score (based on quality and quantity of ventilations) was below 50%. However, the few ventilations that were delivered and detected by the app were adequate (Table 3).

Discussion

Immediate knowledge gain after the course was significant and expected as this has been previously reported.^{4,10,12,17} Retention of gained theoretical knowledge was long lasting - up to 2 years after our BLS and AED course. Understandably, there was a decrease in gained knowledge after approximately 5 months compared to immediately after the course but the level of knowledge remained at a comparable level up to 2 years after the course and remained better than at baseline which is similar to reports from other studies.^{6,12,17-19} Students had not attended any other official BLS and AED course as part of their school curriculum during the 2 years between the 2 tests. It was actually noted that students after 2 years scored even better percentage wise than after 5 months on some questions. This could be attributed to a better understanding of the matter as the students grow older.¹¹ We can speculate that the content of the course was well presented and memorable.

Practical skills were more prone to decline. Plotnikoff and Moore had long before reported that good BLS knowledge acquisition does not correlate well with the quality of BLS skills performance.¹⁹ Skills deterioration has repeatedly been reported by numerous authors^{1,12,19} and results from our study confirm their findings. Five months after the course safety measures were omitted in most cases. This could partially be due to mental mismatch in the idea of a cardiac arrest scenario and a safe school environment in which testing was conducted. A manikin in place of a real person could be the reason for the low rate of checking responsiveness and breathing in a victim. Namely, students after 5 months and 2 years in theory still knew what to do and how to perform responsiveness and breathing checks according to the results of the knowledge test. Most students were focused on performing chest compressions and using an AED and their performance of this segment remained adequate in theory and practice throughout the entire study and comparable to reports from other studies.^{9,20} A rather poor retention of overall skills 5 months after the course (mean score on Cardiff test was 63% and overall score of BLS performance on the QCPR app was 59%) indicate an important decline in practical skills with time. This finding conforms to the European Resuscitation Council's (ERC) consensus on education for resuscitation.²¹ That is to be expected as practical skills in any activity need regular training for adequate performance. The ERC guidelines summarize evidence that frequent training improves CPR skills but the actual retraining periods are yet to be determined.^{21,22}

Limitations

Courses were led by 3 different instructors and their assistants (registered nurses working as paramedics) which was a potential limitation of the study. The content of the course was predetermined and based strictly on the latest resuscitation guidelines with which the instructors, all of which are emergency physicians - and their paramedic assistants are perfectly familiar with as per their profession. Therefore, there could not be any inter-instructor variability in this respect. Practical evaluation was performed by 2 emergency physicians and was done only in Study 2. Lack of objective testing of practical skills immediately after the course and re-testing for retention of skills after 2 years are great limitations of this study.

Even though the authors have acquired all reasonable measures to alleviate possible bias, the sole process of educating and training includes a variety of uncontrollable factors, such as personal attractiveness, assertiveness, communication skills of the instructor on 1 hand, and students' momentary and general attentiveness on the other. However, the sum of all factors' variability reflects the real-life situation where a number of instructors will be involved in the process.

Unfortunately, it has been rather challenging to include even a free BLS and AED course on schools' busy schedules, let alone include such a course in official school curricula on a national level. Based on our findings and considering the current organizational situation, a valid suggestion for educators designing BLS and AED courses for schoolchildren would be to place more emphasis on practical training on repeat courses after 2 years (where annual courses are not feasible). Given the satisfactory knowledge retention even after 2 years, repeat courses could potentially be shorter in duration - less than 2 hours as is suggested by the "Kids save lives" initiative.²³ The theoretical lecture could be shortened to an overview of key aspects of BLS and AED and more emphasis given to practical training. Thereby, the least amount of valuable pedagogical time would be taken from schools, but the children would nevertheless get the optimal amount of training needed for adequate performance of BLS and AED skills.

Conclusion

Baseline knowledge of participating schoolchildren improved significantly after the BLS and AED course. Level of knowledge expectedly decreased after 5 months but remained at a comparable level even after 2 years. Drop in retention of practical skills after 5 months, however, calls for a more intense training of practical skills on the repeat courses. Therefore, we suggest that on repeat courses on BLS and AED the instructors put greater emphasis on practical training, with only a quick theoretical overview of key BLS and AED steps.

Acknowledgments

We would like to thank all participating elementary schools in Maribor and the instructors for making this study possible.

Declaration of conflicting interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Funding

The author(s) received no financial support for the research, authorship, and/or publication of this article.

Ethics approval and consent to participate

Approval from the National Medical Ethics Committee (Ministry of Health of republic of Slovenia) has been granted (dated October 23, 2018; filed under 0120-549/2017/9).

ORCID iD

Vesna Borovnik Lesjak  <https://orcid.org/0000-0002-5416-7266>

Supplemental Material

Supplemental material for this article is available online.

References

- Lester CA, Weston CFM, Donnelly PD, Assar D, Morgan MJ. The need for wider dissemination of CPR skills: are schools the answer? *Resuscitation*. 1994;28:233-237.
- Parnell MM, Pearson J, Galletly DC, Larsen PD, Larsen P. Knowledge of and attitudes towards resuscitation in New Zealand high-school students. *Emerg Med J*. 2006;23:899-902. doi:10.1136/emj.2006.041160.
- Plant N, Taylor K. How best to teach CPR to schoolchildren: A systematic review. *Resuscitation*. 2013;84(4):415-421. doi:10.1016/j.resuscitation.2012.12.008.
- Kitamura T, Nishiyama C, Murakami Y, et al. Compression-only CPR training in elementary schools and student attitude toward CPR. *Pediatr Int*. 2016;58(8):698-704. doi:10.1111/ped.12881.
- Goldberg RJ, Gore JM, Love DG, Ockene JK, Dalen JE. Layperson CPR - are we training the right people? *Ann Emerg Med*. 1984;13:701-704.
- Moore PJ, Plotnikoff RC, Preston GD. A study of school students' long term retention of expired air resuscitation knowledge and skills. *Resuscitation*. 1992;24:17-25.
- Boettiger B, Bossaert L, Castren M, et al. Kids Save Lives - ERC position statement on school children education in CPR.: "Hands that help - Training children is training for life". *Resuscitation*. 2016;105:A1-A3.
- Stroobants J, Monsieurs KG, Devriendt B, Dreezen C, Vets P, Mols P. Schoolchildren as BLS instructors for relatives and friends: Impact on attitude towards bystander CPR. *Resuscitation*. 2014;85(12):1769-1774. doi:10.1016/j.resuscitation.2014.10.013.
- Watanabe K, Lopez-colon D, Shuster JJ, Philip J. Efficacy and retention of Basic Life Support education including Automated External Defibrillator usage during a physical education period. *Preventive Medicine Reports*. 2017;5:263-267. doi:10.1016/j.pmedr.2017.01.004.
- Bohn A, Van Aken HK, Möllhoff T, et al. Teaching resuscitation in schools: Annual tuition by trained teachers is effective starting at age 10. A four-year prospective cohort study. *Resuscitation*. 2012;83(5):619-625. doi:10.1016/j.resuscitation.2012.01.020.
- Borovnik Lesjak V, Šorgo A, Strnad M. Development, validation and assessment of the test on knowledge about basic life support and use of automated external defibrillator among schoolchildren. *Scand J Trauma Resuscitation Emerg Med*. 2019;27(1):114. doi:10.1186/s13049-019-0683-6.
- Connolly M, Toner P, Connolly D, McCluskey DR. The 'ABC for life' programme-Teaching basic life support in schools. *Resuscitation*. 2007;72(2):270-279. doi:10.1016/j.resuscitation.2006.06.031.
- Böttiger B, Van Aken H. Training children in cardiopulmonary resuscitation worldwide. *Lancet*. 2015;385:2353.
- Böttiger B, Van Aken H. Kids save lives - Training school children in cardiopulmonary resuscitation worldwide is now endorsed by the World Health Organization (WHO). *Resuscitation*. 2015;94:A5-A7.
- Schroeder DC, Ecker H, Wingen S, Semeraro F, Böttiger BW. "Kids Save Lives" - Wiederbelebungstrainings für Schulkinder. *Anaesthesist*. 2017;66:589-597. doi:10.1007/s00101-017-0319-z.
- Whitfield RH, Newcombe RG, Woollard M. Reliability of the Cardiff Test of basic life support and automated external defibrillation version 3.1. *Resuscitation*. 2003;59:291-314. doi:10.1016/S0300-9572(03)00246-6.
- Monteiro MLRBP, Ferraz AIB, Rodrigues FMP. ASSESSMENT OF KNOWLEDGE AND SELF EFFICACY BEFORE AND AFTER TEACHING BASIC LIFE SUPPORT TO SCHOOLCHILDREN. *Revista paulista de pediatria : orgao oficial da Sociedade de Pediatria de Sao Paulo*. 2021;39:e2019143.
- Bollig G, Wahl HA, Svendsen MV. Primary school children are able to perform basic life-saving first aid measures. *Resuscitation*. 2009;80(6):689-692. doi:10.1016/j.resuscitation.2009.03.012.
- Plotnikoff R, Moore PJ. Retention of cardiopulmonary resuscitation knowledge and skills by 11- and 12-year-old children. *Med J Aust*. 1989;150:296-302.
- Meissner TM, Kloppe C, Hanefeld C. Basic life support skills of high school students before and after cardiopulmonary resuscitation training: a longitudinal investigation. *Scand J Trauma Resuscitation Emerg Med*. 2012;20(1):31. doi:10.1186/1757-7241-20-31.
- Greif R, Bhanji F, Bigham BL, et al. Education, Implementation, and Teams: 2020 International Consensus on Cardiopulmonary Resuscitation and Emergency Cardiovascular Care Science With Treatment Recommendations. *Resuscitation*. 2020;156:A188A239. doi:10.1016/j.resuscitation.2020.09.014.
- Greif R, Lockey A, Breckwoldt J, et al. European Resuscitation Council Guidelines 2021: Education for resuscitation. *Resuscitation*. 2021;161:388-407. doi:10.1016/j.resuscitation.2021.02.016.
- Böttiger BW, Semeraro F, Wingen S. "Kids Save Lives": Educating Schoolchildren in Cardiopulmonary Resuscitation Is a Civic Duty That Needs Support for Implementation. *J Am Heart Assoc*. 2017;6(3):e005738. doi:10.1161/JAHA.117.005738.