

Methodology article

Open Access

Method for determination of (-102C>T) single nucleotide polymorphism in the human manganese superoxide dismutase promoter

Robert CG Martin*^{1,3}, Kalista Hughes¹, Mark A Doll², Qing Lan⁴, Benjamin D Martini², Jolanta Lissowska⁵, Nathaniel Rothman⁴ and David W Hein^{2,3}

Address: ¹Departments of Surgery, University of Louisville School of Medicine, Louisville, KY, USA, ²Pharmacology & Toxicology, University of Louisville School of Medicine, Louisville, KY, USA, ³James Graham Brown Cancer Center, University of Louisville School of Medicine, Louisville, KY, USA, ⁴Division of Cancer Epidemiology and Genetics, National Cancer Institute, NIH, DHHS, Bethesda, MD, USA and ⁵Division of Cancer Epidemiology and Prevention, Cancer Center and M. Slodowska-Curie Institute of Oncology, Warsaw, Poland

Email: Robert CG Martin* - robert.martin@louisville.edu; Kalista Hughes - Kalista.hughes@louisville.edu; Mark A Doll - Mark.doll@louisville.edu; Qing Lan - qingl@mail.nih.gov; Benjamin D Martini - b.martini@louisville.edu; Jolanta Lissowska - lissowsj@coi.waw.pl; Nathaniel Rothman - rothmann@exchange.nih.gov; David W Hein - david.hein@louisville.edu

* Corresponding author

Published: 14 December 2004

Received: 14 April 2004

BMC Genetics 2004, 5:33 doi:10.1186/1471-2156-5-33

Accepted: 14 December 2004

This article is available from: <http://www.biomedcentral.com/1471-2156/5/33>

© 2004 Martin et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Background: Manganese superoxide dismutase (MnSOD) plays a critical role in the detoxification of mitochondrial reactive oxygen species constituting a major cellular defense mechanism against agents that induce oxidative stress. The *MnSOD* promoter contains an activator protein-2 (AP-2) binding site that modifies transcription of *MnSOD*. Mutations have been identified in the proximal region of the promoter in human tumor cell lines. One of these mutations (-102C>T) has been shown to change the binding pattern of AP-2 leading to a reduction in transcriptional activity. The aim of our study was to develop a method to identify and determine the frequency of this (-102C>T) polymorphism in human tissues.

Results: A new TaqMan allelic discrimination genotype method was successfully applied to genomic DNA samples derived from blood, buccal swabs, snap frozen tissue and paraffin blocks. The polymorphism was shown to be in Hardy-Weinberg Equilibrium in an evaluation of 130 Caucasians from Warsaw, Poland: 44 (33.8%) were heterozygous and 6 (4.6%) were homozygous for -102T.

Conclusion: This report represents the first description of the *MnSOD* -102C>T polymorphism in human subjects by a novel Taqman allelic discrimination assay. This method should enable molecular epidemiological studies to evaluate possible associations of this polymorphism with malignancies and other diseases related to reactive oxygen species.

Background

Antioxidant enzymes such as superoxide dismutase (SOD) protect cells from oxidative stress. Generation of reactive oxygen species (ROS) has been implicated in the etiology of a diversity of human diseases, including cancer[1], aging[2], atherosclerosis[3] and neurodegenerative diseases[4,5]. Superoxide dismutase catalyzes the dismutation of superoxide radical (O_2^-) to H_2O_2 and O_2 . Three distinct types of SODs have been identified in human cells: 1) a homodimeric cytosolic CuZnSOD [6], 2) an extracellular homotetrameric glycosylated SOD [7], and 3) a mitochondrial matrix homotetrameric manganese superoxide dismutase (MnSOD) [8].

Numerous reports indicate a relative deficiency of superoxide dismutase catalytic activity, including mitochondrial MnSOD, in many types of solid tumors [9,10]. Interest in this relative deficiency of SOD activity has been greatly increased by observations that over-expression of SOD in tumor cells will suppress cell division in culture and tumor growth in vivo[11]. In addition recent reports have suggested a possible association between decreased SOD activity and malignant phenotype[12]. While the precise reasons for this relationship between tumor cell growth rate and intracellular SOD activity are not known, these findings support the general idea that decreased expression of SOD may promote tumor growth. In fact, as a result of these and other observations, MnSOD is considered a tumor suppressor gene[1].

Further evaluation of MnSOD suggests that it is critically important in maintenance of mitochondrial function. Mice with deficiency of this enzyme exhibit progressive cardiomyopathy, neurodegeneration and perinatal death[13]. These studies went on to confirm that transgenic mice that express human MnSOD in the mitochondria are protected from environmental oxygen-induced lung injury [14] and adriamycin-induced cardiac toxicity[15]. In contrast, disruption of the other two SODs yielded viable mice which were normal in non-stressful conditions [16]. Thus the mitochondrial MnSOD represents a major cellular defense against oxidative stress.

Genetic polymorphism in the *MnSOD* mitochondrial targeting sequence has been associated with risks to various diseases including breast cancer[17,18], lung cancer[19], cardiomyopathy[20] and Parkinson's disease[21].

A reduction of MnSOD activity has been shown to exist in many types of human cancer cells when compared to normal cells [22]. A recent report has also demonstrated the possible association between decreased SOD activity and malignant phenotype[12]. A recent report demonstrated a new mutation L60F, in exon 3 of the mature protein in the Jurkat human T-cell leukemia-derived cell line that

reduced MnSOD [12]. Thus, it appears that reduced levels of MnSOD activity in human cancer cells can be associated with coding region mutations that alter protein sequence as well as promoter region mutations that alter gene expression [23].

The human *MnSOD* gene is localized to chromosome 6 (6q25). The *MnSOD* promoter region is characterized by a lack of TATA or CAAT boxes but the presence of a GC rich region containing multiple SP-1 and AP-2 binding sites [24]. Further work identified one cause for the reduced expression of MnSOD in some human tumor lines; the occurrence of three heterozygous mutations in the upstream promoter region of this gene [25]. One of these mutations in the *MnSOD* promoter sequence (MnSOD -102C>T) has been shown to change the binding pattern of AP-2 leading to a reduction in transcriptional activity. However the presence of this polymorphism has not been reported in human tissue.

In this study we developed a TaqMan allelic discrimination assay to reliably genotype DNA from many tissues (i.e. blood, buccal swabs, paraffin blocks, and snap frozen tissue) for the -102C>T polymorphism in the *MnSOD* promoter.

Results

We confirmed the presence of the -102C>T single nucleotide polymorphism in human subjects and submitted the sequence variant to Genbank[26]. The genotyping success rate with this technique in the Polish Caucasian population was 85%. An evaluation of 130 DNA samples successfully genotyped from Polish Caucasians not known to have cancer, demonstrated 80 (61.5%) were homozygous (-102C), 44 (33.8%) were heterozygous (-102CT) and 6 (4.6%) were homozygous (-102TT). This distribution is consistent with the Hardy-Weinberg Law. The success rate with this technique in an additional American control population was blood (95%), buccal swabs (90%), snap frozen tissue (80%) and paraffin-embedded samples (75%). The success rates were influenced by DNA quality, DNA extraction technique, and the ability to acquire enough DNA from the buccal swab.

Discussion

Reactive oxygen species in the form of superoxide radicals, hydrogen peroxide, and hydroxy-radicals are formed during incomplete reduction of molecular oxygen during normal respirations. The production of reactive oxygen species remains relatively stable during normal physiologic respirations. A significant increase in the production of reactive oxygen species such as superoxide radicals can be greatly increased as a result of metabolic disorders or more commonly from exposure to toxins such as cigarette

smoke, well-cooked meat, urban residency, and excessive alcohol consumption.

Under normal physiologic conditions, superoxide radicals are detoxified by superoxide dismutase. Among the three SODs, MnSOD has been demonstrated to be the only form that has been essential for survival of aerobic life [27]. Inactivation of the *MnSOD* gene in *E. coli* significantly increased mutation frequency and cell death when bacteria were grown under aerobic conditions [28]. This has been further demonstrated in the evaluation in mammals in which the inactivation of *MnSOD* gene has led to detrimental effects. Polymorphisms of the human *MnSOD* gene have been found in the promoter region, the sequence coding for mature protein, and the mitochondrial targeting sequence. Initial evaluation of the five prime flanking regions from human tumor cell lines indicated that there were no major additions or deletions in the five prime flanking regions of the human *MnSOD* gene [29]. However, there were three mutations that were identified in these tumor cell lines: a C to a T at the - 102 position; a C to a G at the - 38 and an insertion of an A in 11 straight Gs at the - 93 position in relation to the transcription initiation site. The significance of these mutations was felt to be important because this region includes multiple binding sites for SP-1 proteins as well as AP-2 binding sites. Further evaluation of these mutations identified that the C to T change at the - 102 position effected the overall transcription of the *MnSOD* gene [30]. This change in transcription may result from an effect on the AP-2 binding site. Although the -102 C to T mutation was reported in human tumor cell lines [25], no evaluation has been assessed in human subjects.

Evaluation of the -102C>T polymorphism is complicated by difficulty in PCR because of the excessive GC rich region in which this polymorphism exists. This location, upstream from the transcription start site was extremely difficult to identify through multiple PCR-restriction fragment length polymorphism (RFLP) assays, which failed to adequately digest at this polymorphism site, and led to multiple false negative results. We found only the highest quality DNA (i.e. blood) was able to be evaluated using a PCR RFLP assay with only 50% genotyping success. This failure to accurately reproduce the PCR-RFLP assay [31], led us to the development of this TaqMan allelic discrimination assay.

The TaqMan allelic discrimination assay provided results that were confirmed by automated DNA sequencing and blind repeat genotyping. Although we did not test it use on DNA from multiple tissues from the same individual, it was successful for DNA samples derived from buccal swabs and paraffin blocks. It has significant advantages over RFLP analysis, allele-specific amplification, allele-

specific hybridization, and oligo-nucleotide ligation assay techniques. The reasons for this advantage come from the reduction in labor intensive work up, the lack of need for special handling of radioactive probes, and the ability to modify this technique to evaluate multiple polymorphisms in this gene. In addition as more significant polymorphisms within the *MnSOD* gene are discovered, this technique will facilitate detection within the *MnSOD* gene.

The limitations of this technique ultimately come from the quality of DNA that is available and the significant initial expense that is required for a TaqMan assay instrumentation.

Conclusions

This report represents the first description of the *MnSOD* -102C>T polymorphism in human subjects by a novel Taqman allelic discrimination assay. This method should enable molecular epidemiological studies to evaluate possible associations of this polymorphism with malignancies and other diseases related to reactive oxygen species.

Methods

DNA sources

Most DNA samples (130) were isolated from buffy coats of Caucasian controls derived from a population-based case-control study of stomach cancer carried out in Warsaw, Poland as previously described [32]. To test the utility of the method to genotype DNA from various tissue sources, peripheral blood (20 samples), buccal swabs (40 samples), paraffin blocks (40 samples), and snap frozen tissues (15 samples) were collected from research subjects in the USA (Louisville, Kentucky).


DNA extraction from paraffin sections was performed after tissue sections (10 sections, 10 μ m thick) were cut from paraffin blocks. Samples were removed from paraffin through a sequential extraction with histaclear, 100% ethanol and acetone, and dried under vacuum. The pellet was incubated overnight with proteinase K (200 μ g/ml in 50 mM Tris-HCL pH 8.5, 1 mM EDTA and 0.5% Tween-20) at 55 °C. After heating at 100 °C for 10 min, digestion was sequentially extracted with phenol, phenol/chloroform and chloroform. DNA was precipitated with the addition of 3X volume 95% ethanol.

Primer design

SNP-specific polymerase chain reaction (PCR) primers and fluorogenic probes (Table 1) were designed using Primer Express (Version 1.5; Applied Biosystems, Foster City, CA). This technique has been utilized extensively in genotyping other candidate genes with multiple single nucleotide polymorphisms[33,34]. The fluorogenic probes were labeled with a reporter dye (either FAM or

Table 1: Primers and Fluorogenic Probes for -102C>T *MnSOD* Allelic Discrimination

Primers	
-102-Forward Primer (-252 to -234)	5'-gcagacaggcagcgagggtt-3'
-102-Reverse Primer (35 to 19) [287 bp]	5'-ctgaagccgctgccgaa-3'
Probes	
-102C-Taqman Probe (-97 to -107)	fam-ccgcgggccccc
-102T-Taqman Probe (-97 to -107)	vic-ccgcgagccccc

**Figure 1**

Fluorescence ratios of FAM-labeled/VIC-labeled fluorogenic probes specific for -102C>T polymorphism in *MnSOD*. Each bar represents mean \pm standard error for determinations in DNA from 3 human subjects. Open bar represents DNA samples homozygous for -102C. Solid bars represent DNA samples homozygous for -102T. Crossed bars represent DNA samples heterozygous for the SNP. The fluorescence ratios differed significantly ($p < 0.05$) among homozygous and heterozygous genotypes.

VIC) and are specific for one of the two possible bases (-102 C or T) in the *MnSOD* promoter region. A MGB quencher probe was utilized on the 3' end by a linker arm. TaqMan Universal PCR Master Mix (Applied Biosystems) was used to prepare the PCR. The 2X mix was optimized for TaqMan reactions and contained AmpliTaq-Gold DNA polymerase, AmpErase, UNG, dNTPs with UTP and a Passive Reference. Primers, probes and genomic DNA were added to final concentrations of 300 nM, 100 nM, and 0.5–2.5 ng/ μ l respectively. Controls (no DNA template) were run to ensure there was no amplification of contaminating DNA. Reference control DNA was also utilized to verify the polymorphisms identified. The amplification reactions were carried out in an ABI Prism 7700 Sequence Detection System (Applied Biosystems) with two initial hold steps (50°C for 2 min, followed by 95°C

for 10 min) and 50 cycles of a two step PCR (95°C for 15 sec, 60°C for 1 min). The fluorescence intensity of each sample was measured at each temperature change to monitor amplification of the 278 base pair *MnSOD* promoter region. The -102 nucleotide was determined by the fluorescence ratio of the two SNP-specific fluorogenic probes. The fluorescence signal increases when the probe with the exact sequence match binds to the single stranded template DNA and is digested by the 5'-3' exonuclease activity of AmpliTaq-Gold DNA polymerase (Applied Biosystems). Digestion of the probe releases the fluorescent reporter dye (either FAM or VIC) from the quencher dye. As shown in figure 1, the method readily distinguishes between at C or T at -102 in the *MnSOD* promoter region.

Twenty samples with genotypes C/T (4 samples), T/T (3 samples), and C/C (13 samples), some of which were derived from paraffin-embedded tissues, were all confirmed by automated DNA sequencing. These sequence-confirmed samples served as reference standards for the remaining samples. In addition, 10% of the samples were genotyped blind a second time with identical results obtained.

Author contributions

RM: Participated in design of study and manuscript preparation

KH: Participated in genotyping samples

MD: Participated in design of methods of assay

QL: Participated in statistical analysis

BM: Participated in design of methods of assay

JL: Participated in sample collection

NR: Contributed to the study design and the analysis and interpretation of the data

DH: Participated in design of study and manuscript preparation

Acknowledgements

This study was partially supported by USPHS grants CA34627 and CA97942 from the National Cancer Institute.

References

1. St Clair D, Wan XS, Kuroda M, Vichitbandha S, Tsuchida E, Urano M: **Suppression of tumor metastasis by manganese superoxide dismutase is associated with reduced tumorigenicity and elevated fibronectin.** *Oncol Rep* 1997, **4**:753-757.
2. Ku HH, Brunk UT, Sohal RS: **Relationship between mitochondrial superoxide and hydrogen peroxide production and longevity of mammalian species.** *Free Radic Biol Med* 1993, **15**:621-627.

3. Halliwell B: **The role of oxygen radicals in human disease, with particular reference to the vascular system.** *Haemostasis* 1993, **23**(Suppl 1):118-126.
4. Ferrante RJ, Browne SE, Shinobu LA, Bowling AC, Baik MJ, MacGarvey U, Kowall NW, Brown RH Jr, Beal MF: **Evidence of increased oxidative damage in both sporadic and familial amyotrophic lateral sclerosis.** *J Neurochem* 1997, **69**:2064-2074.
5. Fahn S, Cohen G: **The oxidant stress hypothesis in Parkinson's disease: evidence supporting it.** *Ann Neurol* 1992, **32**:804-812.
6. McCord JM, Fridovich I: **Superoxide dismutase. An enzymic function for erythrocyte (hemocuprein).** *J Biol Chem* 1969, **244**:6049-6055.
7. Marklund SL: **Human copper-containing superoxide dismutase of high molecular weight.** *Proc Natl Acad Sci U S A* 1982, **79**:7634-7638.
8. Weisiger RA, Fridovich I: **Mitochondrial superoxide dismutase. Site of synthesis and intramitochondrial localization.** *J Biol Chem* 1973, **248**:4793-4796.
9. Zhao Y, Xue Y, Oberley TD, Kiningham KK, Lin SM, Yen HC, Majima H, Hines J, St Clair D: **Overexpression of manganese superoxide dismutase suppresses tumor formation by modulation of activator protein-1 signaling in a multistage skin carcinogenesis model.** *Cancer Res* 2001, **61**:6082-6088.
10. Zhao Y, Oberley TD, Chaiswing L, Lin SM, Epstein CJ, Huang TT, StClair D: **Manganese superoxide dismutase deficiency enhances cell turnover via tumor promoter-induced alterations in AP-1 and p53-mediated pathways in a skin cancer model.** *Oncogene* 2002, **21**:3836-3846.
11. St Clair DK, Wan XS, Kuroda M, Vichitbandha S, Tsuchida E, Urano M: **Suppression of tumor metastasis by manganese superoxide dismutase is associated with reduced tumorigenicity and elevated fibronectin.** *Oncol Rep* 1997, **4**:753-757.
12. Hernandez-Saavedra D, McCord JM: **Paradoxical effects of thiol reagents on Jurkat cells and a new thiol-sensitive mutant form of human mitochondrial superoxide dismutase.** *Cancer Res* 2003, **63**:159-163.
13. Li Y, Huang TT, Carlson EJ, Melov S, Ursell PC, Olson JL, Noble LJ, Yoshimura MP, Berger C, Chan PH: **Dilated cardiomyopathy and neonatal lethality in mutant mice lacking manganese superoxide dismutase.** *Nat Genet* 1995, **11**:376-381.
14. Wispe JR, Warner BB, Clark JC, Dey CR, Neuman J, Glasser SW, Crapo JD, Chang LY, Whitsett JA: **Human Mn-superoxide dismutase in pulmonary epithelial cells of transgenic mice confers protection from oxygen injury.** *J Biol Chem* 1992, **267**:23937-23941.
15. Yen HC, Oberley TD, Vichitbandha S, Ho YS, St Clair DK: **The protective role of manganese superoxide dismutase against adriamycin-induced acute cardiac toxicity in transgenic mice.** *J Clin Invest* 1996, **98**:1253-1260.
16. Yen HC, Oberley TD, Gairola CG, Szweda LI, St Clair DK: **Manganese superoxide dismutase protects mitochondrial complex I against adriamycin-induced cardiomyopathy in transgenic mice.** *Arch Biochem Biophys* 1999, **362**:59-66.
17. Ambrosone CB, Freudenheim JL, Thompson PA, Bowman E, Vena JE, Marshall JR, Graham S, Laughlin R, Nemoto T, Shields PG: **Manganese superoxide dismutase (MnSOD) genetic polymorphisms, dietary antioxidants, and risk of breast cancer.** *Cancer Res* 1999, **59**:602-606.
18. Mitrunen K, Sillanpaa P, Kataja V, Eskelinen M, Kosma VM, Benhamou S, Uusitupa M, Hirvonen A: **Association between manganese superoxide dismutase (MnSOD) gene polymorphism and breast cancer risk.** *Carcinogenesis* 2001, **22**:827-829.
19. Wang LI, Miller DP, Sai Y, Liu G, Su L, Wain JC, Lynch TJ, Christiani DC: **Manganese superoxide dismutase alanine-to-valine polymorphism at codon 16 and lung cancer risk.** *J Natl Cancer Inst* 2001, **93**:1818-1821.
20. Hiroi S, Harada H, Nishi H, Satoh M, Nagai R, Kimura A: **Polymorphisms in the SOD2 and HLA-DRB1 genes are associated with nonfamilial idiopathic dilated cardiomyopathy in Japanese.** *Biochem Biophys Res Commun* 1999, **261**:332-339.
21. Shimoda-Matsubayashi S, Matsumine H, Kobayashi T, Nakagawa-Hattori Y, Shimizu Y, Mizuno Y: **Structural dimorphism in the mitochondrial targeting sequence in the human manganese superoxide dismutase gene. A predictive evidence for conformational change to influence mitochondrial transport and a study of allelic association in Parkinson's disease.** *Biochem Biophys Res Commun* 1996, **226**:561-565.
22. Oberley LW, Buettner GR: **Role of superoxide dismutase in cancer: a review.** *Cancer Res* 1979, **39**:1141-1149.
23. St Clair DK, Holland JC: **Complementary DNA encoding human colon cancer manganese superoxide dismutase and the expression of its gene in human cells.** *Cancer Res* 1991, **51**:939-943.
24. Wan XS, Devalaraja MN, St Clair DK: **Molecular structure and organization of the human manganese superoxide dismutase gene.** *DNA Cell Biol* 1994, **13**:1127-1136.
25. Xu Y, Krishnan A, Wan XS, Majima H, Yeh CC, Ludewig G, Kasarskis EJ, St Clair DK: **Mutations in the promoter reveal a cause for the reduced expression of the human manganese superoxide dismutase gene in cancer cells.** *Oncogene* 1999, **18**:93-102.
26. Martin RCG, Hughes KH, Doll MA, Rothman N, Hein DW: **Homo sapiens manganese superoxide dismutase gene, 5' flanking sequence.** *Genbank AY397775* 2003. Ref Type: Abstract
27. Carlizo A, Touati D: **Isolation of superoxide dismutase mutants in *Escherichia coli*: is superoxide dismutase necessary for aerobic life?** *EMBO J* 1986, **5**:623-630.
28. Farr SB, D'Ari R, Touati D: **Oxygen-dependent mutagenesis in *Escherichia coli* lacking superoxide dismutase.** *Proc Natl Acad Sci U S A* 1986, **83**:8268-8272.
29. Xu Y, Krishnan A, Wan XS, Majima H, Yeh CC, Ludewig G, Kasarskis EJ, St Clair DK: **Mutations in the promoter reveal a cause for the reduced expression of the human manganese superoxide dismutase gene in cancer cells.** *Oncogene* 1999, **18**:93-102.
30. Xu Y, Krishnan A, Wan XS, Majima H, Yeh CC, Ludewig G, Kasarskis EJ, St Clair DK: **Mutations in the promoter reveal a cause for the reduced expression of the human manganese superoxide dismutase gene in cancer cells.** *Oncogene* 1999, **18**:93-102.
31. Xu Y, Krishnan A, Wan XS, Majima H, Yeh CC, Ludewig G, Kasarskis EJ, St Clair DK: **Mutations in the promoter reveal a cause for the reduced expression of the human manganese superoxide dismutase gene in cancer cells.** *Oncogene* 1999, **18**:93-102.
32. Chow WH, Swanson CA, Lissowska J, Groves FD, Sobin LH, Nasierowska-Guttmejer A, Radziszewski J, Regula J, Hsing AW, Jagannatha S, Zatonski W, Blot WJ: **Risk of stomach cancer in relation to consumption of cigarettes, alcohol, tea and coffee in Warsaw, Poland.** *Int J Cancer* 1999, **81**:871-876.
33. Doll MA, Hein DW: **Comprehensive human NAT2 genotype method using single nucleotide polymorphism-specific polymerase chain reaction primers and fluorogenic probes.** *Anal Biochem* 2001, **288**:106-108.
34. Doll MA, Hein DW: **Rapid genotype method to distinguish frequent and/or functional polymorphisms in human N-acetyltransferase-1.** *Anal Biochem* 2002, **301**:328-332.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

