



Covid-19 related oncologist's concerns about breast cancer treatment delays and physician well-being (the CROWN study)

Katharine A. Yao¹ · Deanna Attai² · Richard Bleicher³ · Kristine Kuchta⁴ · Meena Moran⁵ · Judy Boughey⁶ · Lee G. Wilke⁷ · Jill R. Dietz⁸ · Randy Stevens⁹ · Catherine Pesce¹ · Katherine Kopkash¹ · Scott Kurtzman¹⁰ · Terry Sarantou¹¹ · David Victorson¹²

Received: 2 December 2020 / Accepted: 9 January 2021 / Published online: 31 January 2021
© The Author(s), under exclusive licence to Springer Science+Business Media, LLC part of Springer Nature 2021

Abstract

Purpose To examine how treatment delays brought on by the COVID-19 pandemic impacted the physical and emotional well-being of physicians treating these patients.

Methods A cross-sectional survey of physician breast specialists was posted from April 23rd to June 11th, 2020 on membership list serves and social media platforms of the National Accreditation Program for Breast Centers and the American Society of Breast Surgeons. Physician well-being was measured using 6 COVID-19 burnout emotions and the 4-item PROMIS short form for anxiety and sleep disturbance. We examined associations between treatment delays and physician well-being, adjusting for demographic factors, COVID-19 testing and ten COVID-19 pandemic concerns.

Results 870 physicians completed the survey, 61% were surgeons. The mean age of physicians was 52 and 548 (63.9%) were female. 669 (79.4%) reported some delay in patient care as a result of the COVID-19 pandemic. 384 (44.1%) and 529 (60.8%) of physicians scored outside normal limits for anxiety and sleep disturbance, respectively. After adjusting for demographic factors and COVID-19 testing, mean anxiety and COVID-19 burnout scores were significantly higher among physicians whose patients experienced either delays in surgery, adjuvant chemotherapy, radiation, breast imaging or specialty consultation. A multivariable model adjusting for ten physician COVID-19 concerns and delays showed that “delays will impact my emotional well-being” was the strongest concern associated with anxiety, sleep disturbance and COVID-19 burnout factors.

Conclusions Breast cancer treatment delays during the initial surge of the COVID-19 pandemic in the United States were associated with a negative impact on physician emotional wellness.

Keywords Breast cancer delays · Physician well being · COVID-19 treatment delays

✉ Katharine A. Yao
kyao@northshore.org

¹ Division of Surgical Oncology, Department of Surgery, Clinical Professor of Surgery, Pritzker School of Medicine, NorthShore University HealthSystem, University of Chicago, 2650 Ridge Ave, Evanston, IL 60201, USA

² Department of Surgery, David Geffen School of Medicine, University of California, Los Angeles, Westwood, CA, USA

³ Division of Surgical Oncology, Fox Chase Cancer Center, Philadelphia, PA, USA

⁴ Biostatistical Core, NorthShore University HealthSystem Research Institute, Evanston, IL, USA

⁵ Yale Medicine, New Haven, CT, USA

⁶ Department of Surgery, Mayo Clinic, Rochester, MN, USA

⁷ Department of Surgery, University of Wisconsin, Madison, WI, USA

⁸ Case Western Reserve University, Cleveland, OH, USA

⁹ White Plains Hospital Center for Cancer Care, Westchester, NY, USA

¹⁰ Waterbury Hospital, Waterbury, CT, USA

¹¹ Department of Surgery, Carolinas Medical Center, Charlotte, NC, USA

¹² Department of Medical Social Sciences, Northwestern University Feinberg School of Medicine, Chicago, IL, USA

Background

On January 20, 2020 the first infection with SARS CoV-2 in the United States was reported in Washington State [1]. Since then, over 9 million cases have been reported with over 220,000 deaths [2]. The COVID-19 pandemic required health systems and physicians to prioritize which patients required the most urgent care and resulted in many delays in standard treatment algorithms particularly for patients with cancer. Additionally, on March 13, 2020 the American College of Surgeons (ACS) issued a recommendation to cancel all elective surgeries [3] to conserve personal protective equipment and other resources for COVID-19 positive patients. Although it was questionable whether cancer surgery was considered “elective”, these recommendations were quickly followed by consensus statements from many different societies on how to manage delays in cancer surgery including patients with breast cancer [4–6]. A recent survey [7] concluded that 50% of breast centers in Europe reported altered systemic treatments resulting from the COVID-19 pandemic and 20% of patients experienced delays in radiation therapy. However, data on delays for patients with breast cancer in the United States have not yet been reported.

At the time that elective cases were being canceled or delayed, several studies were published that showed that patients with cancer who were infected with SARS CoV-2 were experiencing high rates of death from COVID-19 [8, 9]. Breast physician specialists had to weigh delaying treatment for these patients to conserve resources or proceed with treatment and potentially increase the risk of infection with SARS CoV-2 for their patients and its negative sequelae on outcomes. However, little empirical evidence exists to describe how these delays impacted the physical and emotional well-being of physicians treating these patients. Studies conducted outside the United States have shown that physicians directly caring for patients infected with SARS CoV-2 impacts physician well-being. A study of over 1200 frontline healthcare workers in China in January 2020 reported depression was experienced in approximately 50%, anxiety in 44% and insomnia in 30% [10]. Similarly, another study of 376 healthcare workers from Italy found high rates of emotional exhaustion and somatic symptoms from frontline care of COVID-19 positive patients [11]. A recent meta-analysis of 13 studies showed that just over 20% of frontline workers experienced depression and anxiety [12]. However, these studies examined physicians and other healthcare workers taking direct care of COVID-19 infected patients. There is little empirical evidence examining physicians who were not necessarily caring directly for patients infected by SARS CoV-2 but who were nonetheless experiencing downstream effects of the COVID-19 pandemic such as

an inability to provide timely and standard care for their patients with a diagnosed cancer. The impact on physician well-being could potentially affect the future medical care of patients (in the post-pandemic period) since physician well-being and burnout emotions are directly associated with medical errors, absenteeism, relationship discord and substance abuse [13].

To further explore pandemic-related breast cancer treatment delays and potential effects on physicians treating breast cancer, we used a cross sectional survey administered to breast specialists across the country during the initial surge of COVID-19 cases. First, we assessed the scope of treatment delays for breast cancer patients. Second, we assessed the association of these delays with physician anxiety, sleep disturbance and emotions associated with physician burnout.

Methods

Study design and setting

A cross sectional survey was created using a convenience sample of multispecialty physicians specializing in breast cancer. The survey was distributed using RedCap to members of NAPBC-accredited breast centers and ASBrS via social media posts, email and list serves of the ACS and the ASBrS from April 23rd to June 11th, 2020 (See supplementary materials for full version of the survey). Survey questions were created by a group of ten breast physician specialists from across the country based on personal conversations with colleagues, newspaper or social media articles commenting on physician wellness at the time of the study or stories in the news or radio. The 27-item survey was composed of four domains: COVID-19 testing questions, delays or altered treatment plans, COVID-19 concerns and demographic factors (see Appendix for full survey). COVID testing questions focused on coronavirus testing, direct care of infected patients and personal knowledge of persons (non-patients) infected by SARS CoV-2 during that period. Physicians were asked about their perception of how delayed treatments impacted their patient’s overall survival and delivery of care. COVID-19 concerns included how the pandemic was impacting physicians personal and professionally. Demographic factors including number of breast cancer patients seen weekly, age, gender, time in practice, location of practice (by State), type of practice and specialty were also collected.

This study was approved by the Institutional Review Board at NorthShore University HealthSystem. Completion of the survey was considered consent to participate in the study and all responses were de-identified such that responses could not be linked to individual respondents or centers.

Participants

Only physicians who treated breast cancer patients in some capacity and were members of the aforementioned medical societies were eligible to participate. All specialties in breast cancer care were represented, including: surgery, medical and radiation oncology, breast imaging, geneticists, pathology and survivorship.

Outcome measures

Physician anxiety and sleep disturbance were measured with the Patient Reported Outcomes Measurement Information System (PROMIS) 4-item Anxiety and Sleep Disturbance short forms [14, 15]. These validated measures are based on a t-score metric with a mean of 50 and standard deviation of 10. A brief measure of COVID-19 related burnout symptoms was created for this study and was based on known emotions related to physician burnout, including emotional exhaustion, depression, helplessness, and anxiety. This was scored by computing a summary score of these 6 items which were based on a Likert scale of 0 = not at all to 4 = very much with a possible score range from 0 to 24. In this study, this measure had a normal distribution, with a mean (\pm SD) of 8.9 ± 6.3 . It was strongly correlated with the PROMIS Anxiety scale ($r = 0.72$) and weakly correlated with the PROMIS Sleep Disturbance scale ($r = 0.20$). The 6 item set demonstrated strong internal consistency reliability (Cronbach's $\alpha = 0.91$), which was not improved by removing any single item.

Statistical analysis

Descriptive statistics and measures of central tendency (e.g., frequencies, percentages, means, standard deviations (SD)) were used to summarize the survey results. Pearson correlation coefficients and Cronbach's α were used to assess preliminary psychometric characteristics of the COVID-19 related burnout item set that was administered. Bivariate differences in outcomes between groups were assessed with the chi-square test or univariate general linear models. Cohen's d effect sizes were calculated to determine the magnitude of effect of delays on physician emotional well being ($d = 0$ –0.19 trivial effect, 0.20–0.49 small effect, 0.50–0.79 medium effect and > 0.80 large effect). Multivariate general linear models controlling for physician age, gender, years in practice, testing positive for SARS CoV-2 and caring for COVID-19 positive patients using physician wellness and opinions about delays as the outcomes were used to assess differences between physicians whose patients did and did not experience treatment delays. Multiple comparisons between groups were made using the Tukey–Kramer adjustment. Multivariable general linear models with standardized

regression coefficients and backward selection method using $p < 0.05$ to remain in the model were used to identify the treatment delays and COVID-19 concerns most strongly associated with outcomes. Analysis of items with missing data were performed with non-missing observations only. All statistical analyses were performed using SAS 9.4 (SAS Institute, Cary, NC). p values < 0.05 were considered significant and all tests were two sided.

Results

Demographics of the physician cohort

870 physicians completed the survey. Missing data did not exceed 5% for any individual survey item. Responding physician demographics are listed in Table 1. Mean age was 52 ± 11 and 63.9% were female. Surgeons comprised the largest group ($n = 547$, 61.7%). Participants came from all nine U.S. Census divisions with the greatest proportion from the East North Central division ($n = 197$, 22.6%) (Fig. 1). 113 (13%) of physicians underwent testing for SARS CoV-2 and only eleven (9.8%) tested positive.

Treatment delays in breast cancer care

Treatment delays and mean length of delays were examined by facility location, practice setting and specialty type (Supplementary Table 1). Overall, 669 (79.4%) physicians reported at least some delay in either breast surgery, systemic or radiation therapy, breast imaging or specialty consultation. Delays in breast surgery ($n = 479$, 59.4%) and breast imaging ($n = 427$, 51.6%) were most common. Most respondents ($n = 317$, 43.3%) reported average delays of one month, 230 (31.4%) reported 2 months, 156 (21.3%) reported 3 months and 29 (4.0%) reported 4 months or longer.

Physician concerns about the COVID-19 pandemic and treatment delays

Surgeons were asked to choose which factors concerned them the most about treatment delays. “Patients becoming anxious” was the factor chosen with the greatest frequency ($n = 579$, 66.6%), “delays will impact overall survival of patients” was the second most frequent factor ($n = 345$, 39.7%) and “cancer will get larger” ($n = 286$, 32.9%) was the third most frequent factor (Supplementary Fig. 1). When physicians were asked about how certain factors related to the pandemic were impacting their practice or personal well-being, 437 (51.3%) of physicians stated they were very or somewhat concerned “that my emotional well-being will be impacted by delays” (Fig. 2).

Table 1 Demographics and COVID-19 testing factors of the physicians

Total participants [N]	870
Age [Mean ± SD]	52 ± 11
Gender [N (%)]	
Female	548 (63.9)
Male	307 (35.8)
Other	2 (0.2)
Years in practice [N (%)]	
Less than 1 year	24 (2.8)
1–5 years	75 (8.7)
6–10 years	113 (13.1)
11–15 years	143 (16.6)
16 years of longer	509 (58.9)
Number of breast cancer patients normally seen per week [Median (Q1–Q3)]	15 (6–30)
Primary practice setting [N (%)]	
Academic employed/university setting	138 (15.9)
Academic employed/non-university setting	111 (12.8)
Hospital or health plan employed/community setting	365 (42.0)
Private practice	262 (30.1)
Veterans administration hospital or government employed	6 (0.7)
Other	14 (1.6)
Specialty [N (%)]	
Medical oncology	94 (10.8)
Radiation oncology	70 (8.0)
Surgery	537 (61.7)
Plastic surgery	22 (2.5)
Pathology	46 (5.3)
Genetics	9 (1.0)
Radiology	78 (9.0)
Survivorship	8 (0.9)
Other	21 (2.4)
Tested for SARS CoV-2 [N (%)]	
No	756 (87.0)
Yes	113 (13.0)
Did you test positive? (N = 113) [N (%)]	
No	101 (90.2)
Yes	11 (9.8)
Taken care of COVID-19 positive patients [N (%)]	
No	408 (47.0)
Yes	296 (34.1)
Not Sure	165 (19.0)
Know anyone personally who is COVID-19 Positive? [N (%)]	
No	319 (38.2)
Yes	517 (61.8)



Physician outcomes of anxiety, sleep disturbance and COVID-19 related burnout

Based on recommended cut-off points for PROMIS Anxiety and Sleep Disturbance scores (0–54 = normal, 55–60 = mild, 61–70 = moderate, > 70 = severe) [16], 384 (44.1%) and 529 (60.8%) of the sample were outside normal limits for anxiety and sleep disturbance respectively. A quarter of the sample ($n = 220$, 25.3%) reported mild symptoms of anxiety with roughly twice as many reporting symptoms of mild sleep disturbance 466 (53.6%). Amongst the COVID-19 related burnout emotions, “emotional exhaustion” was reported with the highest frequency ($n = 291$, 33.6%), “anxious” was the second most common ($n = 225$, 26.1%) and “burnout” was the third most common ($n = 190$, 22.1%).

Physician anxiety, sleep disturbance and COVID-19 related burnout were significantly associated with demographic factors including COVID-19 testing factors (Table 2). Anxiety and COVID-19 related burnout were significantly associated with physician age, gender and years in practice and sleep disturbance was significantly associated with physician age. There were no significant differences in physician anxiety, sleep disturbance and COVID-19 related burnout between different facility locations, practice settings or physician specialty type. Physicians who took care of COVID-19 positive patients reported significantly higher anxiety, sleep disturbance and COVID-19 related burnout than physicians who did not take care of COVID-19 positive patients ($54.5 + 8.6$ vs $51.4 + 8.2$, $p \leq 0.001$; $55.2 + 3.7$ vs $54.3 + 3.8$, $p = 0.03$; $10.2 + 6.7$ vs $7.6 + 5.9$, $p \leq 0.001$ respectively). Furthermore, physicians who tested positive for SARS CoV-2 reported significantly lower anxiety ($48.1 + 8.8$ vs $54.5 + 9.1$, $p = 0.03$) and sleep disturbance scores ($51.3 + 7.3$ vs $54.5 + 3.7$, $p = 0.02$) than those who tested negative for SARS CoV-2.

Physicians whose patients experienced delays in either surgery, adjuvant therapy, breast imaging, radiation therapy, or specialty consultation reported significantly higher anxiety ($53.4 + 8.3$ vs $50.5 + 8.3$, $p \leq 0.001$) and COVID-19-related burnout ($9.5 + 6.3$ vs $7.0 + 6.2$, $p \leq 0.001$) than those who did not experience delays after adjusting for demographic factors, testing positive for SARS CoV-2 and taking care of COVID-19 patients (Table 2). Examination of Cohen’s d effect sizes suggests that the majority of significant group differences were small in magnitude, however, burnout score differences between groups who experienced delays in specialty consultations were medium in size ($d = 0.55$).

Nearly half ($n = 391$, 47.4%) of physicians somewhat or strongly agreed that treatment delays would impact their patient’s overall survival but the majority ($n = 623$, 73.0%) somewhat or strongly agreed that they were still able to deliver the same high quality care during the COVID-19

Fig. 1 Map of facility location of physician practices**Fig. 2** Physician concerns about impact of the COVID-19 pandemic on their practice and well being

pandemic as before the pandemic (Table 3). Physicians who perceived the delays would impact their patient's survival reported significantly higher anxiety, COVID-19 related burnout and sleep disturbance than physicians who did not feel the delays would impact survival, after adjusting for the aforementioned confounders ($54.3 + 8.2$ vs $51.2 + 8.0$, $p \leq 0.001$; $54.9 + 3.8$ vs $54.0 + 4.0$, $p = 0.005$;

$10.2 + 6.5$ vs $7.6 + 5.9$, $p \leq 0.001$, respectively) (Table 3). Physicians who felt they could deliver the same high-quality care reported significantly lower anxiety and COVID-19 related burnout compared with physicians who did not feel they could deliver high-quality care ($52.2 + 8.2$ vs $53.6 + 9.7$, $p \leq 0.001$; $8.5 + 6.2$ vs $9.4 + 7.1$, $p = 0.002$, respectively) (Table 3).

Table 2 Physician anxiety, sleep disturbance and COVID-19 related burnout stratified by physician demographic factors and Covid-19 testing factors

	All N (%)	Physician Anxiety		Physician sleep disturbance		COVID-19 Related Burnout	
		Mean ± SD	p value	Mean ± SD	p value	Mean ± SD	p value
All patients	870	52.8 ± 8.4	–	54.6 ± 3.9	–	8.9 ± 6.3	–
Demographics							
Age, years			< 0.001		0.001		0.001
< 45	235 (28.0)	54.2 ± 9.0	*	55.1 ± 4.0	*	9.8 ± 6.6	*
45–59	364 (43.4)	52.8 ± 8.5	~	54.7 ± 4.0	~	9.1 ± 6.3	~
≥ 60	239 (28.5)	51.2 ± 7.2	*~	53.8 ± 3.7	*~	7.7 ± 5.8	*~
Gender			< 0.001		0.592		< 0.001
Female	548 (64.1)	53.7 ± 8.5		54.6 ± 4.1		9.9 ± 6.3	
Male	307 (35.9)	51.1 ± 7.8		54.4 ± 3.6		7.2 ± 6.0	
Years in practice			< 0.001		0.197		0.018
Less than 6 years	99 (11.5)	55.7 ± 9.0	*~	55.3 ± 4.0		10.1 ± 6.1	
6–10 years	113 (13.1)	54.3 ± 8.9		54.7 ± 3.8		10.0 ± 7.2	
11–15 years	143 (16.6)	52.1 ± 8.8	*	54.5 ± 4.1		9.0 ± 6.4	
16 years of longer	509 (58.9)	52.1 ± 7.9	~	54.4 ± 3.9		8.4 ± 6.1	
Number of breast cancer patients normally seen per week			0.933		0.655		0.911
< 10	254 (33.6)	52.8 ± 8.6		54.4 ± 4.0		8.9 ± 6.7	
10–29	277 (36.6)	52.6 ± 8.4		54.5 ± 3.9		9.0 ± 6.3	
≥ 30	226 (29.9)	52.7 ± 8.0		54.7 ± 4.0		9.2 ± 6.0	
Facility location			0.064		0.970		0.529
New England	57 (7.0)	51.4 ± 7.6		54.6 ± 3.0		9.2 ± 5.4	
Mid-Atlantic	144 (17.8)	54.2 ± 7.3		54.8 ± 3.6		9.2 ± 6.3	
South Atlantic	150 (18.5)	52.4 ± 8.8		54.6 ± 3.9		8.0 ± 6.3	
East North Central	197 (24.3)	53.1 ± 8.4		54.5 ± 3.8		9.3 ± 6.3	
East South Central	36 (4.4)	49.3 ± 8.4		54.4 ± 4.6		7.3 ± 6.5	
West North Central	50 (6.2)	51.2 ± 8.6		54.0 ± 4.5		8.3 ± 6.7	
West South Central	54 (6.7)	52.6 ± 8.4		55.0 ± 3.8		8.7 ± 5.8	
Mountain	35 (4.3)	52.3 ± 8.5		54.7 ± 3.8		8.8 ± 7.1	
Pacific	88 (10.9)	52.5 ± 8.0		54.6 ± 4.1		8.6 ± 6.1	
Primary practice setting			0.112		0.640		0.614
Academic employed/university setting	138 (16.2)	54.1 ± 8.0		54.7 ± 4.3		9.4 ± 6.3	
Academic employed/non-university setting	108 (12.7)	53.1 ± 8.0		54.8 ± 3.4		9.1 ± 5.8	
Hospital/health plan employed/community setting	355 (41.8)	52.8 ± 8.5		54.7 ± 4.0		9.0 ± 6.4	
Private practice	249 (29.3)	52 ± 8.4		54.3 ± 3.6		8.5 ± 6.6	
Specialty			0.445		0.490		0.094
Medical oncology	94 (11.4)	53.5 ± 7.1		55.1 ± 4.0		9.7 ± 6.2	
Radiation oncology	70 (8.5)	52.8 ± 7.4		54.0 ± 3.1		7.8 ± 6.2	
Surgery	537 (65.1)	52.6 ± 8.8		54.5 ± 4.0		9.2 ± 6.4	
Pathology	46 (5.6)	51.4 ± 8.5		54.8 ± 3.4		7.4 ± 5.5	
Radiology	78 (9.5)	53.9 ± 8.2		54.4 ± 3.9		8.4 ± 6.6	
COVID-19 testing factors							
Tested for SARS CoV-2			0.117		0.313		< 0.001
No	756 (87.0)	52.6 ± 8.2		54.6 ± 3.9		8.6 ± 6.2	
Yes	113 (13.0)	54.0 ± 9.2		54.2 ± 4.3		11.1 ± 6.9	
Did you test positive? (N = 113)			0.029		0.017		0.428
No	101 (90.2)	54.5 ± 9.1		54.5 ± 3.7		11.2 ± 6.7	
Yes	11 (9.8)	48.1 ± 8.8		51.3 ± 7.3		9.5 ± 8.6	
Taken care of COVID-19 positive patients			< 0.001		0.026		< 0.001

Table 2 (continued)

	All N (%)	Physician Anxiety		Physician sleep disturbance		COVID-19 Related Burnout	
		Mean ± SD	<i>p</i> value	Mean ± SD	<i>p</i> value	Mean ± SD	<i>p</i> value
No	408 (47.0)	51.4 ± 8.2	* ~	54.3 ± 3.8	~	7.6 ± 5.9	* ~
Yes	296 (34.1)	53.8 ± 8.1	*	54.6 ± 4.1		10.0 ± 6.4	*
Not Sure	165 (19.0)	54.5 ± 8.6	~	55.2 ± 3.7	~	10.2 ± 6.7	~
Know anyone personally who is COVID-19 Positive?			0.099		0.734		0.084
No	319 (38.2)	52.1 ± 8.6		54.5 ± 4.2		8.4 ± 6.4	
Yes	517 (61.8)	53.1 ± 8.3		54.6 ± 3.7		9.2 ± 6.3	

Statistically significant values are highlighted in bold

* ~ Indicate significant statistical differences between groups using the Tukey–Kramer adjustment for multiple comparisons ($p < 0.05$)

Of the 10 physician concerns about the COVID-19 pandemic (Fig. 2) and the five different treatment delays, “that my emotional well-being will be impacted by delays” was most strongly associated physician concern with anxiety, sleep disturbance and COVID-19 related burnout on multivariable analysis adjusting for physician demographic factors (Supplementary Table 2).

Discussion

For the first time in many physicians’ careers, physicians were forced to delay or significantly alter treatments for their patients because of the COVID-19 pandemic. Our study of over 800 breast specialists across the United States demonstrates that delays for breast cancer patients were quite common with nearly 80% of physicians reporting some type of delay in either screening, systemic treatment or surgery for their patients. During the study period, these same physicians also reported an impact in their physical well-being more than emotional well-being; approximately 44% of all responding physicians had anxiety scores outside the normal ranges, compared to 60% of physicians reporting sleep disturbance scores. We also showed that physicians whose patients experienced delays had higher anxiety and more frequent COVID-19 burnout emotions such as emotional and physical exhaustion, depression and helplessness than physicians who did not have patients experiencing delays. However, the magnitude of these differences was not large. The reasons for these low effect sizes may be multifactorial and not necessarily all captured in our survey. The majority of physicians in this study did not have direct contact with COVID-19 positive patients, there was regional variation in hospital response to the pandemic which would directly impact physician wellness and regional variation in prevalence of the pandemic at the time of the study. Of note, burnout rates were relatively low in this physician population

likely because many physicians were not working on the “frontline”. Nonetheless, our data show that the COVID-19 pandemic had at least some impact on physician wellness and in most cases a negative impact.

Physicians expressed many concerns about pandemic-related treatment delays and the direct impact these delays on their patients’ breast cancer outcomes. Nearly half of physicians thought that treatment delays would impact their patients’ survival outcomes. Although the survey did not specifically question which type of treatment delay was perceived to impact survival, our data suggests that the most common delays that occurred were for definitive surgery, corresponding to the majority of respondents being surgeons. However, surgery is often not the first method of treatment for a newly diagnosed breast cancer patient and there are short term alternative treatments for some types of breast cancer besides immediate surgery. For patients with hormone receptor negative or Her2/neu over-expressed disease, neoadjuvant chemotherapy or HER2/neu targeted therapy is preferred and recommended by guidelines [17], with no negative impact on survival. In fact, a recent consortium recommendation statement outlined which breast cancer surgical cases could be safely delayed to later in the course of treatment [4] and a recent multidisciplinary framework states that surgery for older women with breast cancer can be safely delayed because patients can be placed on hormonal therapy [18]. For those patients with hormone receptor positive disease, randomized studies have shown no survival difference between hormone therapy alone versus hormone therapy plus surgery over a limited timeframe [19, 20]. Many recommendation statements issued during the early stages of the pandemic recommended upfront hormonal therapy for early-stage hormone receptor positive patients and consideration of neoadjuvant chemotherapy and/or Her2/neu targeted therapy for hormone receptor negative or HER2neu positive disease [3, 4, 6]. However, it is possible that some physicians may have felt uncomfortable administering systemic therapy given the uncertain mortality

Table 3 Association of physician anxiety, sleep disturbance and COVID-19 related burnout with treatment delays after adjusting for known confounders

	All patients		Physician anxiety		Physician sleep disturbance		COVID-19 related burnout		
	N (%)	Mean ±SD	Unadjusted <i>p</i> value	Adjusted <i>p</i> value†	Mean ±SD	Unadjusted <i>p</i> value	Mean ±SD	Unadjusted <i>p</i> value	Adjusted <i>p</i> value‡
Delays regarding care									
Delay in breast surgery									
No	328 (40.6)	52.0 ± 8.5	0.014	0.042	54.4 ± 4.2	0.232	0.370	0.001	0.002
Yes	479 (59.4)	53.5 ± 8.2			54.7 ± 3.8				
Effect Size (Cohen's <i>d</i>)		<i>d</i> = 0.18^			<i>d</i> = 0.09^				<i>d</i> = 0.25^
Delay in starting adjuvant therapy									
No	666 (89.5)	52.4 ± 8.4			54.5 ± 4.1	0.540	0.702	0.002	< 0.001
Yes	78 (10.5)	56.0 ± 8.0			54.8 ± 3.2				
Effect Size (Cohen's <i>d</i>)		<i>d</i> = 0.43^			<i>d</i> = 0.07^				<i>d</i> = 0.37^
Delay in breast imaging									
No	400 (48.4)	51.6 ± 8.3			54.4 ± 4.2	0.180	0.161	< 0.001	< 0.001
Yes	427 (51.6)	53.9 ± 8.4			54.7 ± 3.7				
Effect Size (Cohen's <i>d</i>)		<i>d</i> = 0.27^			<i>d</i> = 0.09^				<i>d</i> = 0.34^
Delay in specialty consultation									
No	704 (90.1)	52.4 ± 8.2			54.4 ± 4.0	0.052	0.100	< 0.001	< 0.001
Yes	77 (9.9)	56.3 ± 9.3			55.4 ± 3.5				
Effect Size (Cohen's <i>d</i>)		<i>d</i> = 0.47^			<i>d</i> = 0.23^				<i>d</i> = 0.55^
Delay in starting radiation therapy									
No	575 (76.6)	52.3 ± 8.3	0.002	0.002	54.5 ± 4.2	0.233	0.320	0.002	0.001
Yes	176 (23.4)	54.5 ± 8.6			54.9 ± 3.2				
Effect Size (Cohen's <i>d</i>)		<i>d</i> = 0.27^			<i>d</i> = 0.10^				<i>d</i> = 0.27^
Cumulative delay									
No	174 (20.6)	50.5 ± 8.3			54.1 ± 4.3	0.078	0.114	< 0.001	< 0.001
Yes	669 (79.4)	53.4 ± 8.3			54.7 ± 3.8				
Effect size (Cohen's <i>d</i>)		<i>d</i> = 0.35^			<i>d</i> = 0.15^				<i>d</i> = 0.40^
Physician opinions									
I worry that treatment delays will impact my patient's overall survival									
Somewhat or strongly agree	391 (47.4)	54.3 ± 8.2	*	*~	54.9 ± 3.8	*	*	*	*
Neither agree or disagree	122 (14.8)	52.9 ± 8.9	~	~	54.6 ± 4.0			~	~

Table 3 (continued)

	All patients		Physician anxiety		Physician sleep disturbance		COVID-19 related burnout	
	N (%)	Mean ±SD	Unadjusted p value	Adjusted p value†	Mean ±SD	Unadjusted p value	Adjusted p value	Adjusted p value‡
Somewhat or strongly disagree	312 (37.8)	51.2 ± 8.0	*	*	54.0 ± 4.0	*	7.6 ± 5.9	* ~ *
I feel that I have been able to deliver the same high quality care			< 0.001	< 0.001	0.395	0.336	0.002	0.002
Somewhat or strongly agree	623 (73.0)	52.2 ± 8.2	*	*	54.4 ± 4.1	*	8.5 ± 6.2	* *
Neither agree or disagree	33 (3.9)	53.6 ± 9.7			55.2 ± 3.4		9.4 ± 7.1	
Somewhat or strongly disagree	198 (23.2)	55.0 ± 8.5	*	*	54.7 ± 3.6	*	10.4 ± 6.5	* *

Statistically significant values are highlighted in bold

†Adjusted for physician age, gender, years in practice, tested positive for SARS CoV-2, taken care of COVID-19 positive patients

* ~ Indicate significant statistical differences between groups using the Tukey–Kramer adjustment for multiple comparisons ($p < 0.05$)

^Cohen's d effect sizes (for interpretive guide see Methods section)

risks with infection of SARS CoV-2 and neutropenia due to systemic treatment [8, 9, 21, 22].

Prior to the pandemic, delays in treatment have been reported to impact patient outcomes. It will be many years before we know if delays brought on by the COVID-19 pandemic have impacted survival outcomes of breast cancer patients. Physicians in our study reported a mean delay of two months. In one study, time from diagnosis to first treatment was up to 60 days in 18% of Medicare patients and 24% of patients in the National Cancer Data Base [23]. Time to first treatment also varies by region in the United States and by type of operation [24, 25]. Retrospective studies have shown that delays beyond just 30 days can be associated with incremental decreases in overall survival [26, 27] However, although some aspect of care (such as surgery) was delayed, it is important to remember that many patients in this study were undergoing some type of other treatment such as endocrine therapy.

Interestingly, COVID testing factors impacted physician well-being in an unexpected way. Physicians who tested positive for SARS CoV-2 reported significantly lower anxiety and sleep disturbance than those who tested negative for SARS CoV-2 although the number of physicians who tested positive was low. Perhaps the anticipation of testing positive produces more anxiety and sleep disturbance than actually knowing you are positive. Physicians who tested negative for SARS CoV-2 may not have trusted the test results and would still have to worry about possibly testing positive in the future. There was a lot of uncertainty about testing for COVID-19 at the time of the study. Furthermore, those who were tested for SARS CoV-2 had higher COVID-19 burnout emotions than those who were not tested. This finding may stem from the reason the testing was ordered in the first place such as presence of symptoms, exposure to potential COVID-19 positive patients or having to work in areas of the hospital unfamiliar to breast specialists- all factors that could increase COVID-19 related burnout emotions.

There were several physician-related demographic factors that were independently associated with higher levels of anxiety and COVID-19-related burnout emotions. Compared with older, more experienced physicians and male physicians, female physicians and physicians practicing for shorter periods experienced greater anxiety and burnout emotions. These findings are similar to previous reports of studies suggesting female surgical residents experiencing more burnout than male physicians during their training [28, 29] or during the pandemic [30]. Studies have reported that physicians in their early career are more susceptible to burnout [31] and mid-career physicians often have the highest rates of burnout and emotional exhaustion [32]. Investigators have hypothesized that female physicians suffer more burnout due to more challenging work-life balance issues related to home and childcare responsibilities, as well as harassment

and discrimination. While some or all of these factors may have also played a role in the responses from our female physician respondents, deciphering confounding factors in this survey tool is beyond the scope of this study. Nevertheless, our findings are consistent with other studies suggesting that females and less experienced attending physicians reported more burnout rates.

Our study showed a trend toward regional variance in some of the physician well-being outcomes across the country during the study period. Higher physician anxiety and COVID-19 burnout emotions did correlate with regions with higher prevalence rates during the study period. For example, the mid-atlantic region and east north central regions had the highest and second highest rates, respectively, of infection with SARS CoV-2, respectively, during the study period and physician anxiety and COVID-19 burnout scores were highest in these regions. However, sleep disturbance scores fell into higher categories across all regions suggesting that the initial surge of the COVID-19 pandemic had a higher impact on sleep than anxiety for all physicians regardless of location.

Our study is not without limitations. Our data capture one point in time; we have no data on these physician's anxiety and sleep disturbance prior to the pandemic. Due to the design of our study and time sensitive nature, we are not able to calculate a known response rate. We used social media, email and other networks to quickly recruit breast specialists to report their delays. The majority of respondents were surgeons perhaps because surgeons had to delay many of their patient's surgeries because of the pandemic. Many respondents had also been in practice for 16 years or longer. Since our findings are subject to selection bias, they are not necessarily generalizable to the entire oncology community who cares for breast cancer patients. While the survey attempted to control for expected increases in anxiety, sleep disturbances, and feelings of burnout from predictable confounding factors such as taking care of COVID+ patients, or knowing someone who tested positive for SARS CoV-2 and even important COVID-19 concerns, it is beyond the scope of this survey to be able to account for all such environmental, personal and geographic confounding factors. As such, while our study clearly demonstrates an association between physician self-reported well-being symptoms, the onset of the pandemic and physicians' perceptions of compromised survival outcomes for their breast cancer patients due to treatment delays, these associations cannot be taken as causal. The COVID-19 pandemic is ongoing, and our survey captures just the beginning of the initial surge of COVID-19 cases in the United States. Since the time of our study, additional knowledge has been gained about how to manage patients infected with SARS CoV-2 and trends in COVID-19 infections are increasing. Physician wellness may be different now compared to the time period of our study.

In conclusion, these findings suggest that physicians taking care of breast cancer patients also experienced compromised well-being despite not having front-line COVID patient care responsibilities. These data can serve as an important guide to those physicians who are currently experiencing the COVID-19 pandemic on a scale similar to the initial surge experienced in the Northeast and other parts of the country. COVID-19 cases continue to increase across the country and more treatment delays may again become a reality for many physicians. Perhaps this study will motivate other hospitals and cancer centers to provide psychosocial support and interventions for physicians that may not have been available during the initial surge of the COVID-19 pandemic. Health systems that deal with these issues directly are the most likely to benefit both in terms of their physician well-being, as well as the safety of their patients.

Supplementary Information The online version of this article (<https://doi.org/10.1007/s10549-021-06101-1>) contains supplementary material, which is available to authorized users.

Funding There was no funding source for this study.

Data availability The datasets generated during and/or analyzed during the current study are available from the corresponding author on reasonable request.

Compliance with ethical standards

Conflicts of interest The authors have no disclosures or conflicts to report.

References

1. Holshue ML, DeBolt C, Lindquist S et al (2020) First Case of 2019 Novel Coronavirus in the United States. *The New England journal of medicine* 382(10):929–936
2. COVID CDC gov statistics. https://covid.cdc.gov/covid-data-tracker/?utm_source=morning_brew#cases. 2020.
3. ACS Elective surgery guidelines. <https://www.facs.org/covid-19/clinical-guidance/elective-surgery>. 2020.
4. Dietz JR, Moran MS, Isakoff SJ et al (2020) Recommendations for prioritization, treatment, and triage of breast cancer patients during the COVID-19 pandemic. *The COVID-19 pandemic breast cancer consortium. Breast Cancer Res Treat* 181(3):487–497
5. ASCO Coronavirus resources. <https://www.asco.org/asco-coronavirus-resources/care-individuals-cancer-during-covid-19/disease-specific-information>. 2020.
6. SSO Breast resource during COVID-19 <https://www.surgonc.org/wp-content/uploads/2020/03/Breast-Resource-during-COVID-19-3.30.20.pdf>. 2020.
7. Gasparri ML, Gentilini OD, Lueftner D, Kuehn T, Kaidar-Person O, Poortmans P (2020) Changes in breast cancer management during the Corona Virus Disease 19 pandemic: an international survey of the European Breast Cancer Research Association of Surgical Trialists (EUBREAST). *Breast* 52:110–115

8. Kuderer NM, Choueiri TK, Shah DP et al (2020) Clinical impact of COVID-19 on patients with cancer (CCC19): a cohort study. *Lancet* 395(10241):1907–1918
9. Robilotti EV, Babady NE, Mead PA et al (2020) Determinants of COVID-19 disease severity in patients with cancer. *Nat Med* 26(8):1218–1223
10. Lai J, Ma S, Wang Y et al (2020) Factors associated with mental health outcomes among health care workers exposed to coronavirus disease 2019. *JAMA Netw Open* 3(3):e203976
11. Barello S, Palamenghi L, Graffigna G (2020) Burnout and somatic symptoms among frontline healthcare professionals at the peak of the Italian COVID-19 pandemic. *Psychiatry Res* 290:113129
12. Pappa S, Ntella V, Giannakas T, Giannakoulis VG, Papoutsi E, Katsaounou P (2020) Prevalence of depression, anxiety, and insomnia among healthcare workers during the COVID-19 pandemic: a systematic review and meta-analysis. *Brain Behav Immun* 88:901–907
13. Shanafelt TD, Balch CM, Bechamps G et al (2010) Burnout and medical errors among American surgeons. *Ann Surg* 251(6):995–1000
14. Pilkonis PA, Choi SW, Reise SP, Stover AM, Riley WT, Cella D (2008) The Development of Scales for Emotional Distress from the Patient-Reported Outcomes Measurement Information System (PROMIS): Depression, Anxiety, and Anger
15. Buysse DJ, Yu L, Moul DE et al (2010) Development and validation of patient-reported outcome measures for sleep disturbance and sleep-related impairments. *Sleep* 33(6):781–792
16. PROMIS site health measures. www.healthmeasures.net/score-and-interpret/interpret-scores/promis/promis-score-cut-points. 2020. Accessed Nov 2020.
17. NCCN guidelines. <https://www.nccn.org>. 2019.
18. Freedman RA, Sedrak MS, Bellon JR, et al. (2020) Weathering the storm: managing older adults with breast cancer amid COVID-19 and beyond. *J Natl Cancer Instit*
19. Masuda N, Sagara Y, Kinoshita T et al (2012) Neoadjuvant anastrozole versus tamoxifen in patients receiving goserelin for premenopausal breast cancer (STAGE): a double-blind, randomised phase 3 trial. *Lancet Oncol* 13(4):345–352
20. Spring LM, Gupta A, Reynolds KL et al (2016) Neoadjuvant endocrine therapy for estrogen receptor-positive breast cancer: a systematic review and meta-analysis. *JAMA Oncol* 2(11):1477–1486
21. Desai A, Sachdeva S, Parekh T, Desai R (2020) COVID-19 and cancer: lessons from a pooled meta-analysis. *JCO Glob Oncol* 6:557–559
22. Dai M, Liu D, Liu M et al (2020) Patients with cancer appear more vulnerable to SARS-CoV-2: a multicenter study during the COVID-19 outbreak. *Cancer Discov* 10(6):783–791
23. Bleicher RJ, Ruth K, Sigurdson ER et al (2012) Preoperative delays in the US Medicare population with breast cancer. *J Clin Oncol: Off J Am Soc Clin Oncol* 30(36):4485–4492
24. Bleicher RJ, Chang C, Wang CE et al (2019) Treatment delays from transfers of care and their impact on breast cancer quality measures. *Breast Cancer Res Treat* 173(3):603–617
25. Kupstas AR, Hoskin TL, Day CN, Habermann EB, Boughey JC (2019) Effect of Surgery Type on Time to Adjuvant Chemotherapy and Impact of delay on breast cancer survival: a national cancer database analysis. *Ann Surg Oncol* 26(10):3240–3249
26. Bleicher RJ, Ruth K, Sigurdson ER et al (2016) Time to surgery and breast cancer survival in the United States. *JAMA Oncol* 2(3):330–339
27. Yung R, Ray RM, Roth J et al (2020) The association of delay in curative intent treatment with survival among breast cancer patients: findings from the Women’s Health Initiative. *Breast Cancer Res Treat* 180(3):747–757
28. Dahlke AR, Johnson JK, Greenberg CC et al (2018) Gender differences in utilization of duty-hour regulations, aspects of burnout, and psychological well-being among general surgery residents in the United States. *Ann Surg* 268(2):204–211
29. Hu YY, Ellis RJ, Hewitt DB et al (2019) Discrimination, abuse, harassment, and burnout in surgical residency training. *Engl J Med* 381(18):1741–1752
30. Coleman JR, Abdelsattar JM, Glocker RJ, Force RACT (2020) COVID-19 Pandemic and the Lived Experience of Surgical Residents, Fellows, and Early-Career Surgeons in the American College of Surgeons. *J Am College Surg*.
31. Del Carmen MG, Herman J, Rao S et al (2019) Trends and factors associated with physician burnout at a multispecialty academic faculty practice organization. *JAMA Netw Open* 2(3):e190554
32. Dyrbye LN, Varkey P, Boone SL, Satele DV, Sloan JA, Shanafelt TD (2013) Physician satisfaction and burnout at different career stages. *Mayo Clin Proc* 88(12):1358–1367

Publisher’s Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.