

Available online at www.sciencedirect.com

ScienceDirect

journal homepage: www.elsevier.com/locate/ajur

Editorial

Gene expression profiles and treatments for metastatic renal cell carcinoma: What does still need to be defined?



Approximately one third of renal cell carcinoma (RCC) patients have metastasis at the time of diagnosis, while one third of localized patients inevitably develop disease recurrence [1,2]. To date, there are four drug combinations with immunotherapies which have been approved by the US Food and Drug Administration as first-line therapy in metastatic RCC (mRCC): pembrolizumab (anti-programmed cell death protein 1 [anti-PD-1] antibody) plus axitinib (vascular endothelial growth factor receptor [VEGF-R] tyrosine kinase inhibitor [TKI]), avelumab (anti-PD-L1 antibody) plus axitinib, nivolumab (anti-PD-1 antibody) plus cabozantinib (multi-target TKI), and nivolumab plus ipilimumab (anti-cytotoxic T lymphocyte antigen 4 antibody), replacing monotherapies with TKIs targeting the VEGF-R pathway [1]. Therefore, one of the main unmet needs is how to select the best first-line regimen for treatment-naïve mRCC patients. Heng score has been validated as a useful prognostic score [3] and it has been used to stratify treatment-naïve mRCC patients enrolled in clinical trials, being low-score patients usually treated with anti-angiogenic TKIs and intermediate/high-score ones with combination therapy; however, it could not be useful on its own for choosing between immuno-immuno and immuno-TKI combination therapies. Genetic alterations, *i.e.*, *PBRM1* mutations, have also been interrogated as predictive factors for immunotherapy, with conflicting results [4].

In this scenario, it is crucial to develop biomarkers able to predict response or lack of response to available therapies.

Motzer et al. [5] evaluated gene expression profiles (GEPs) on tumor specimens from patients enrolled in the JAVELIN Renal 101 clinical trial (avelumab+axitinib vs. sunitinib) and correlated them to progression-free survival (PFS) in both arms of treatment. The 26-gene Renal 101 Immuno-signature—comprising regulators of adaptive and innate immune responses, cell trafficking, and inflammation—was able to stratify patients only in the combination

arm, with an improvement in PFS in patients who had greater than or equal to the median expression of this signature. In a similar but opposite way, a 26-gene angiogenesis discriminated patients in the sunitinib arm—but not in the combination one—with a prolonged PFS.

GEPs from the IMmotion150 phase 2 trial—atezolizumab (anti-PD-L1 antibody) alone or in combination with bevacizumab (anti-VEGF) versus sunitinib as first-line therapy in mRCC—identified three main signatures: angiogenesis, T-effector, and myeloid signatures [6]. Interestingly, sunitinib efficacy was greater in highly angiogenic tumors (high angiogenesis signature), whilst atezolizumab showed lower responses in case of myeloid inflammation (high myeloid signature) even if the addition of bevacizumab in such cases may overcome resistance to immune checkpoint inhibitors.

The GEPs of 469 clear cell RCC patients from The Cancer Genome Atlas have been studied by D'Costa and colleagues [7], who have identified a 66-gene signature that clusters patients into three groups: angiogenesis, T-effector, and mixed signature. Differently from the use of IMmotion150 32-gene signature applied to the same population, this 66-gene signature was able to stratify patients with a better overall survival and disease-free survival for angiogenesis signature compared to the others.

The 66-gene signature was also applied to 316 RCC patients as described in the abstract by Barata and colleagues [8], showing significantly different mutations and immune profiles between angiogenesis and T-effector signatures. In details, biomarkers of potential response to immunotherapy such as PD-L1, tumor mutational burden, and deficient mismatch repair were more frequent in the T-effector signature, even if only the difference of PD-L1 expression was statistically significant.

Sequencings of bulk and individual RNA transcripts are important tools to study not only tumor characteristics but also to inquire tumor microenvironment and to dissect biology of specific cancer types, identifying subgroups

<https://doi.org/10.1016/j.ajur.2022.03.005>

2214-3882/© 2022 Editorial Office of Asian Journal of Urology. Production and hosting by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

which may behave differently also in relation to therapies. In mRCC patients, however, the studies so far presented, were not able to validate a specific signature (*i.e.*, immune signature vs. angiogenic ones) as a predictive biomarker of response to treatments. In fact, even if the T-effector signature predicted high responses in patients who received combination with immunotherapy, these same patients were also treated with TKIs; on the other hand, the angiogenic signature would not clarify the role of VEGF-R targeting agents since patients were treated with TKIs in both of the arms of the aforementioned trials.

Therefore, to clinically validate and explore the prognostication of GEPs in order to identify specific subset of patients who benefit from a specific therapy, clinical trials should be designed using GEPs as stratification factors. In this way, they would be able to assess the specific impact of those signatures in relation to the treatment received by the patients.

In conclusion, GEPs are a potentially useful tool to predict response to specific therapies and could guide clinicians in choosing among different therapeutic options, but the lack of clinical validations of “real world” patients’ stratification based on GEPs data and the lack of GEPs-led clinical trials makes their use difficult at the moment.

Author contributions

Study concept and design: Pasquale Rescigno.

Drafting of manuscript: Emilio F. Giunta.

Critical revision of the manuscript: Roberto Pili.

Conflicts of interest

The authors declare no conflict of interest.

References

- [1] Tran J, Ornstein MC. Clinical review on the management of metastatic renal cell carcinoma. *JCO Oncol Pract* 2022;18:187–96.
- [2] Vartolomei MD, Matei DV, Renne G, Tringali VM, Crişan N, Musi G, et al. Long-term oncologic and functional outcomes

after robot-assisted partial nephrectomy in elderly patients. *Minerva Urol Nefrol* 2019;71:31–7.

- [3] Heng DY, Xie W, Regan MM, Warren MA, Golshayan AR, Sahi C, et al. Prognostic factors for overall survival in patients with metastatic renal cell carcinoma treated with vascular endothelial growth factor-targeted agents: results from a large, multicenter study. *J Clin Oncol* 2009;27:5794–9.
- [4] Dias Carneiro APC, Marques Monteiro FS, Soares A. *PBRM1* mutations as a predictive biomarker for immunotherapy in metastatic renal cell carcinoma: a systematic review. *Kidney Cancer* 2021;5:79–92.
- [5] Motzer RJ, Robbins PB, Powles T, Albiges L, Haanen JB, Larkin J, et al. Avelumab plus axitinib versus sunitinib in advanced renal cell carcinoma: biomarker analysis of the phase 3 JAVELIN Renal 101 trial. *Nat Med* 2020;26:1733–41.
- [6] McDermott DF, Huseni MA, Atkins MB, Motzer RJ, Rini BI, Escudier B, et al. Clinical activity and molecular correlates of response to atezolizumab alone or in combination with bevacizumab versus sunitinib in renal cell carcinoma. *Nat Med* 2018;24:749–57.
- [7] D’Costa NM, Cina D, Shrestha R, Bell RH, Lin YY, Asghari H, et al. Identification of gene signature for treatment response to guide precision oncology in clear-cell renal cell carcinoma. *Sci Rep* 2020;10:2026. <https://doi.org/10.1038/s41598-020-58804-y>.
- [8] Barata PC, Gulati S, Elliott A, Rao A, Hammers HJ, Quinn DI, et al. Angiogenic and T-effector subgroups identified by gene expression profiling (GEP) and propensity for *PBRM1* and *BAP1* alterations in clear cell renal cell carcinoma (ccRCC). *J Clin Oncol* 2021;39(Suppl. 6):343. https://ascopubs.org/doi/abs/10.1200/JCO.2021.39.6_suppl.343.

Emilio F. Giunta

Pasquale Rescigno*

Interdisciplinary Group for Translational Research and Clinical Trials, Urological Cancers (GIRT-Uro), Candiolo Cancer Institute, FPO-IRCCS, Candiolo, Turin, Italy

Roberto Pili

Division of Hematology and Oncology, Jacobs School of Medicine and Biomedical Sciences, University at Buffalo, Buffalo, NY, USA

*Corresponding author.

E-mail address: pasquale.rescigno@ircc.it (P. Rescigno)

27 January 2022