

Impact of genomic stability on protein expression in endometrioid endometrial cancer

MI Lomnytska^{*,1,2}, S Becker², T Gemoll^{2,3}, C Lundgren⁴, J Habermann³, A Olsson⁵, I Bodin⁵, U Engström⁶, U Hellman⁶, K Hellman⁴, A-C Hellström⁴, S Andersson¹, M Mints¹ and G Auer²

¹Department of Obstetrics and Gynaecology, Karolinska University Hospital, Solna, Karolinska Institute, SE-17176 Stockholm, Sweden; ²Unit of Cancer Proteomics, Department of Oncology and Pathology, Karolinska Institute, Karolinska University Hospital, SE-17176 Stockholm, Sweden; ³Department of Surgery, University of Lübeck, D-23538 Lübeck, Germany; ⁴Department of Gynaecological Oncology, Radiumhemmet, Karolinska University Hospital, Solna, SE-17176 Stockholm, Sweden; ⁵Department of Oncology and Pathology, Karolinska Institute, Karolinska University Hospital, SE-17176 Stockholm, Sweden; ⁶Ludwig Institute for Cancer Research Ltd, Uppsala University, Box 595, Uppsala SE-75124, Sweden

BACKGROUND: Genomic stability is one of the crucial prognostic factors for patients with endometrioid endometrial cancer (EEC). The impact of genomic stability on the tumour tissue proteome of EEC is not yet well established.

METHODS: Tissue lysates of EEC, squamous cervical cancer (SCC), normal endometrium and squamous cervical epithelium were subjected to two-dimensional (2D) gel electrophoresis and identification of proteins by MALDI TOF MS. Expression of selected proteins was analysed in independent samples by immunohistochemistry.

RESULTS: Diploid and aneuploid genomically unstable EEC displayed similar patterns of protein expression. This was in contrast to diploid stable EEC, which displayed a protein expression profile similar to normal endometrium. Approximately 10% of the differentially expressed proteins in EEC were specific for this type of cancer with differential expression of other proteins observed in other types of malignancy (e.g., SCC). Selected proteins differentially expressed in 2D gels of EEC were further analysed in an EEC precursor lesion, that is, atypical hyperplasia of endometrium, and showed increased expression of CLIC1, EIF4A1 and PRDX6 and decreased expression of ENO1, ANXA4, EMD and Ku70.

CONCLUSION: Protein expression in diploid and aneuploid genomically unstable EEC is different from the expression profile of proteins in diploid genomically stable EEC. We showed that changes in expression of proteins typical for EEC could already be detected in precursor lesions, that is, atypical hyperplasia of endometrium, highlighting their clinical potential for improving early diagnostics of EEC.

British Journal of Cancer (2012) 106, 1297–1305. doi:10.1038/bjc.2012.67 www.bjcancer.com

Published online 13 March 2012

© 2012 Cancer Research UK

Keywords: endometrioid endometrial cancer; genomic stability; marker protein patterns; atypical endometrioid hyperplasia

Endometrial cancer (EC) is the fourth most common gynaecologic malignancy in Europe and Northern America. Even if detected at stage I, EC relapses in the majority of these cases (Creasman *et al*, 2006). Thus, diagnostics for detecting asymptomatic EC and precancer lesions is of paramount importance (Buchanan *et al*, 2009; Seebacher *et al*, 2009).

EC is divided into oestrogen-dependent endometrioid EC (EEC) that develops from atypical hyperplasia of endometrium (AH) and oestrogen-independent nonendometrioid EC that is usually characterised by a poorer prognosis (Bokhman, 1983; Horn *et al*, 2007). An important factor that defines the aggressiveness of malignancies, including EC, is chromosomal stability. More than half of the cases of EC are genomically stable and diploid (Lundgren *et al*, 2002, 2004). In comparison, all squamous cervical cancers (SCCs) and the vast majority of epithelial ovarian cancers are genomically unstable

and aneuploid. Expression of proteins in diploid and aneuploid EC differs significantly (Lundgren *et al*, 2004). Characterisation of these proteins may provide new biomarkers for improved early diagnostics of EC and precursor lesions.

Proteomics is a potential method in the search for new cancer markers (Pitteri and Hanash, 2010; Sharon *et al*, 2010). Several proteomics-based studies of EC revealed important information about the endometrium, that is, the impact of genomic instability in endometrial cancer on protein expression (Lundgren *et al*, 2006), the proteome involved in myometrial invasion of endometrial cancer (Monge *et al*, 2009), and new insights into the secretome of endometrium (Casado-Vela *et al*, 2009). Unfortunately, only a few of the proteins identified in these studies were further analysed for their clinical value. Also, in many cases a comparison is only made between cancer and the respective normal tissue, without comparison with other closely related malignancies. Thus, the cancer specificity of the identified proteins could not be determined (Pettrak *et al*, 2008). Furthermore, the similarities observed between protein expression in EEC and precursor lesions may be used for early detection of EEC.

*Correspondence: Dr MI Lomnytska; E-mail: marta.lomnytska@ki.se
Received 22 December 2011; revised 10 February 2012; accepted 13 February 2012; published online 13 March 2012

Table 1 Description of clinical material used for (a) 2D gel electrophoresis and (b) immunohistochemical analysis

(a)					
No.	Sample ID	TNM	Stage, FIGO, 1988	Ploidy	Age
<i>I. Endometrioid endometrial cancer</i>					
<i>II. Genomically stable</i>					
1	Gs1	T1aN0G1	IA	DS	54
2	Gs2	T1aN0G2	IA	DS	82
3	Gs3	T1aN0G2	IA	AS	51
4	Gs4	T1bN0G1	IB	DS	69
5	Gs5	T1bN0G1	IB	DS	86
6	Gs6	T1bN0G1	IB	AS	84
7	Gs7	T1cN0G3	IC	DS	69
				70.7 ± 14.2	
<i>III. Genomically unstable</i>					
8	Gu1	T1bN0G1	IB	DU	85
9	Gu2	T1bN0G1	IB	DU	52
10	Gu3	T1bN0G1	IB	DU	80
11	Gu4	T1bN0G2	IB	DU	52
12	Gu5	T1cN0G1	IC	DU	41
13	Gu6	T1cN0G1	IC	DU	79
14	Gu7	T1cN0G2	IC	AU	71
15	Gu8	T3N1G3	III	DU	54
				64.3 ± 16.4	
Stage, FIGO, 1994					
All:					67.3 ± 15.2
<i>II. Squamous cervical cancer</i>					
16	CC1	T1b1N0G2	IB1		65
17	CC2	T1bN0G3	IB1		52
18	CC3	T1b2N0G2	IB2		45
19	CC4	T1b2N0G2	IB2		39
20	CC5	T1b2N0G2	IB2		59
21	CC6	T1b2N0G3	IB2		53
22	CC7	T2aN0G2	IIA		69
23	CC8	T2aN0G2	IIA		44
24	CC9	T2aN0G3	IIA		89
25	CC10	T2aN0G3	IIA		63
26	CC11	T2bN0G3	IIIB		45
27	CC12	T3N0G3	III		60
28	CC13 ^a	T1b2N0	IB2		41
				55.7 ± 14.0	
<i>III. Endometrium</i>					
E9, E10, E11, E13, E13, E16, E29, E35					50.6 ± 2.7
<i>IV. Cervical mucosa</i>					
M1, M2, M4, M5					49.5 ± 7.1
(b)					
No.	Age	TNM	Stage, FIGO, 1988	Relapse, months	Overall survival, months
<i>Endometrioid endometrial cancer</i>					
<i>Diploid stable, n = 2</i>					
2	78	T1bNxM0G1	IB	—	72
3	62	T1cNxM0G1	IC	—	72
<i>Diploid unstable, n = 13</i>					
1	47	T1bNxM0G2	IB	—	64
4	73	T1bNxM0G1	IB	—	58
5	75	T3NxM1G3	IV	38 (distant)	70
6	72	T1bNxM0G1	IB	—	72
7	78	T1bNxM0G1	IB	10 (local)	72
8	71	T1aNxM0G1	IA	—	72
9	71	T1cNxM0G1	IC	18 (distant)	33
10	65	T1bNxM0G2	IB	—	72
11	63	T1cNxM0G3	IC	—	72
12	58	T1bNxM0G2	IB	—	72
13	57	T3aNxM0G2	IIIA	—	72
14	58	T1aNxM0G1	IA	—	72
15	63	T1bNxM0G1	IB	—	72

Table 1 (Continued)

(b)					
No.	Age	TNM	Stage, FIGO, 1988	Relapse, months	Overall survival, months
<i>Aneuploid unstable, n = 4</i>					
16	82	T1cNxM0G3	IC	22 (local)	<u>24</u>
17	75	T1bNxM0G3	IB	10 (distant)	<u>38</u>
18	58	T2bNxM0G3	IIIB	—	72
19	54	T1bNxM0G1	IB	—	72
				66.5 ± 9.3	
<i>Atypical hyperplasia of endometrium</i>					
				Diploid unstable, n = 8	
				Diploid stable, n = 7	
<i>Normal endometrium</i>					
				50.1 ± 3.7	n = 15

Abbreviations: 2D = two dimensional; AS = aneuploid stable; AU = aneuploid unstable; DS = diploid stable; DU = diploid unstable; FIGO = International Federation of Gynaecology and Obstetrics (Fédération Internationale de Gynécologie et d'Obstétrique); Gs = genomically stable endometrioid endometrial cancer; Gu = genomically unstable endometrioid endometrial cancer; TNM = Tumour, Node, and Metastasis. ^aAdenocarcinoma of cervix uteri. Underlined entries for overall survival correspond to deceased patients.

Finally, identification of proteins correlated with genomic instability has the potential to improve malignancy grading.

In the present study, we expand the current knowledge about the expression of proteins in EEC with respect to DNA ploidy as a measure of genomic stability and the relevance of these proteins to EEC carcinogenesis.

MATERIALS AND METHODS

Clinical material

Clinical material (Table 1a and b) was collected at the Department of Obstetrics and Gynaecology, Karolinska University Hospital, Huddinge; the Department of Gynaecologic Oncology, Radiumhemmet, Karolinska University Hospital, Solna, Sweden; and the Department of Oncology and Medical Radiology, Lviv National Medical University, Lviv, Ukraine, with informed consent and approval from the local ethics committees (Stockholm County Council – Dnr. 97-244 (1998-03-02), 00-068 (200-06-05), 2006/649-31/4, Ethics Committee of Lviv National Medical University – protocol no. 2).

Tissue biopsies of EEC (15 cases), SCC (13 cases) and control tissue from patients with nonmalignant gynaecological diseases (e.g., myoma and menorrhagia) consisting of normal endometrium (E; 8 cases) and squamous epithelium of cervical mucosa (SE; 4 cases) were collected before treatment for two-dimensional gel electrophoresis (2D; Table 1a). The tissue biopsies were snap frozen in liquid nitrogen and stored at -70°C . Histopathological diagnosis was performed in all cases. Formalin-fixed paraffin-embedded (FFPE) tissue samples for immunohistochemical (IHC) analysis consisted of independent cases of EEC (19 cases), AH (15 cases) and normal endometrium (15 cases; Table 1b).

DNA cytometry

Tissue biopsies of EEC (Table 1a) and an independent group of FFPE samples of EEC and AH (Table 1b) were analysed for DNA ploidy. The former were analysed in imprint cytological samples and the latter in $6\mu\text{m}$ thick tissue cuts. The prepared slides were stained according to the Feulgen method and the DNA content in




Figure 1 Description of the clinical material used in this study. **(A)** DNA histograms of diploid stable EEC showing narrow stem line in the 2c region, diploid unstable EEC with a broad stem line that expands from the 2c to the 4c region and aneuploid unstable EEC with a broad peak outside the 2c region and additional peaks exceeding the 4c region. **(B)** Examples of analysed 2D gels of EEC and endometrium. **(C)** Principal component analysis of the analysed 2D gels indicating similarity between the expression of protein spots in genomically unstable EEC and SCC, genomically stable EEC and normal endometrium as well as difference between the expression of protein spots in genomically stable and unstable EEC. **(D)** Clustering of identified proteins according to their function with numbers corresponding to the amount of detected proteins. **(E)** Distribution of selected proteins according to gains (to the right) and losses (to the left) on the chromosomes where the orange colour corresponds to early chromosomal changes during EEC carcinogenesis. A shaded pattern depicts chromosomal changes related to a bad prognosis for patients.

single cells was measured by means of image cytometry (Steinbeck *et al*, 1999). Histograms with a narrow stem line in the 2c region represented a diploid genomically stable subtype and those with a broad stem line in the 2c region that expanded towards the 4c region were classified as diploid genomically unstable (Figure 1A). Histograms with a narrow peak outside the 2c region were considered to be aneuploid genomically stable, whereas those with a broad peak outside the 2c region and additional peaks exceeding the 4c region were classified as aneuploid genomically unstable (Figure 1A).

Two-dimensional gel electrophoresis and MALDI TOF mass spectrometry

Tissue proteins were extracted and solubilised in lysis buffer: 9 M urea (Bio-Rad, Sundbyberg, Sweden), 2 M thiourea (USB, Cleveland, OH, USA), 5% Resolyte (BDH, Poole, Dorset, UK), 65 mM DTT (Bio-Rad), 1 M EDTA (Merck, Darmstadt, Germany), 0.5% v/v Nonidet P-40 (USB), 25 mM CHAPS, 0.1% PMSF, 0.01% benzamidine, 0.01% BHT, and 35 mM NaOH (Sigma, St Louis, MO, USA) (Hellman *et al*, 2009). Protein concentration was determined using the Bradford protein assay (Bradford, 1976). The IEF, SDS-PAGE, staining with silver nitrate and excision of spots were performed as previously described (Lomnytska *et al*, 2010). Expression of protein spots was analysed by Progenesis SameSpot software (Nonlinear Dynamics, Newcastle upon Tyne, UK). Protein spots with a relative expression difference of 1.5-fold (ANOVA with $P < 0.05$ and power > 0.8) were selected for MALDI TOF MS. All steps were performed as previously described (Lomnytska *et al*, 2010).

Western blot

In order to verify the identity of the proteins after MALDI TOF MS analysis, the same tissue protein lysates that were used for 2D gel analysis (Figure 2A) were subjected to western blot (Figure 2B). Equal concentration of protein lysates was applied to 10.5–14.0% SDS-PAGE (Criterion gels, Bio-Rad). The following commercial antibodies were used for western blot: EIF4A1 (1:2000; ab31217-100, rabbit polyclonal; Abcam, Cambridge, UK), CLIC1 (1:500; ab77214-100, mouse monoclonal; Abcam), PRDX6 (1:4000; ab59543, rabbit polyclonal; Abcam), CLIC4 (1:50; ab67593, rabbit polyclonal; Abcam), ENO1 (1:1000; ab85086, rabbit polyclonal; Abcam), ANXA4 (1:1000; ab109900, mouse monoclonal; Abcam), EMD (1:1000; ab54996, mouse monoclonal; Abcam) and Ku70 (1:1000; S5C11, mouse monoclonal; Abcam). All antibodies were diluted in Pierce (Rockford, IL, USA) Protein-Free T20 (PBS) Blocking Buffer (Thermo Scientific, Middletown, VA, USA) and incubated for 12 h at 4 °C. As positive controls, lysates of cell lines that contain corresponding antigens were used, that is, HeLa cell lysate for EIF4A1, CLIC1, PRDX6 and EMD, placenta lysate for ANXA4 and MCF7 cell lysate for CLIC4, ENO1 and Ku70 (Figure 2B). The membranes were incubated in secondary antibody of the corresponding species, diluted 1:15 000 in Pierce Protein-Free T20 (PBS) Blocking Buffer for 1.5 h at room temperature, followed by 4 washes of 15 min in PBS-T. Finally, the proteins were visualised by ECL. The secondary antibodies used were HRP-linked anti-mouse (NXA931) and HRP-linked anti-rabbit antibodies (NA934VS, GE Healthcare, Chalfont St Giles, UK). All steps were performed as described before (Lomnytska *et al*, 2010).




Figure 2 Expression of EIF4A1, CLIC1, PRDX6, CLIC4, ENO1, ANXA4, EMD, and Ku70 in 2D gels of endometrium, genomically stable EEC and genomically unstable EEC. **(A)** Selected areas of the 2D gels. Arrows indicate spots from which the selected proteins were identified. The numbers below indicate the normalised spot volume. Abbreviations: E = endometrium; Gu = genomically unstable endometrioid endometrial cancer. **(B)** Western blot (WB) analysis verifying protein expression patterns in the same samples as shown in **(A)**. The numbers below the bands represent the densitometrical analysis.

Immunohistochemistry

An immunohistochemical analysis was carried out on FFPE samples of EEC, AH and E of an independent group of patients in order to study the expression of the identified proteins during EEC carcinogenesis (Table 1b). Immunohistochemistry was performed using the two-step streptavidin–biotin method. Tissue slides were incubated overnight with the primary antibodies in 1% BSA at 4 °C. Antibodies used previously for western blot were applied in following dilutions for IHC: EIF4A1 (1:200), CLIC1 (1:10), PRDX6 (1:1000), ENO1 (1:200) and Ku70 (1:400). In addition, staining against ANXA4 (1:200; sc-1930, goat polyclonal; Santa Cruz Biotechnology, Santa Cruz, CA, USA), CLIC4 (1:30; HPA008019, rabbit polyclonal; Sigma-Aldrich, St Louis, MO, USA) and EMD (1:3000; HPA000609, rabbit polyclonal; Sigma-Aldrich) was performed (Figure 3A). Antibodies used for western blot against ANXA4, CLIC4 and EMD were also used for IHC for confirmation of specificity (data not shown). Several visualising systems were used: VectaStain (Vector, Peterborough, UK) ABC-Po-kit and DAB (positive stain was brown), LSAB+ (DAKO, Glostrup, Denmark) (positive stain was red). Control tissues that contained corresponding antigens were also utilised: placenta tissue for ANXA4, placenta and tonsillar tissue for EIF4A1, tonsillar and ovarian tissue for CLIC1, tonsillar and placenta tissue for PRDX6, tonsillar, placenta and breast cancer tissue for CLIC4, colon cancer and tonsillar tissue for EMD, breast cancer and tonsillar tissue for Ku70 and breast cancer and kidney tissue for ENO1. Images were captured with a Leica DM4500B (camera

DFC320, ocular 10 ×, objectives 20 ×/0.50 HC PL and 40 ×, 506145) and the Leica Application Suite software, version 2.4.0 (Wetzlar, Germany) as 16-bit depth .tif format images with 48-bit image resolution, and expression of the analysed proteins was scored as previously described (Cheng *et al*, 2008; Lomnytska *et al*, 2011).

Statistical analysis

We used the inbuilt statistical chapter of SameSpot Nonlinear software (PCA, ANOVA, power, *t*-test), MedCalc, version 11.1.1.0 (Mariakerke, Belgium) (receiver-operator-characteristic (ROC) curves) and Statistica 6.0 (Tulsa, OK, USA), (correlation, *t*-test, χ^2 test). A difference of $P < 0.05$ was considered statistically significant.

RESULTS

Expression of protein spots in analysed 2D gels

A total of 42 2D gels were generated from tissue biopsies of 40 patients with EEC, SCC or nonmalignant gynaecological diseases (Table 1a), with each gel containing ~2000 protein spots (Figure 1B). DNA cytometry was performed on all EEC samples in order to identify their genomic stability (Table 1a and Figure 1A).




Figure 3 Analysis of the expression of EIF4A1, CLIC1, PRDX6, CLIC4, ENO1, ANXA4, EMD and Ku70. **(A)** Examples of the immune staining in endometrium (a), atypical hyperplasia of endometrium (b) and endometrioid endometrial cancer (c). Inserts indicate an $\times 400$ magnification of the indicated areas. **(B)** Comparison between expression of proteins (panels a–h) in endometrium (15 cases), atypical hyperplasia of endometrium (15 cases), genomically stable endometrioid endometrial cancer (2 cases) and genomically unstable endometrioid endometrial cancer (17 cases) as evaluated by immunohistochemistry. Horizontal lines indicate statistically significant differences between the protein expression in compared groups (ANOVA, Kruskal–Wallis, $P < 0.05$). Abbreviations: AH = atypical hyperplasia of endometrium; E = endometrium; Gs = genomically stable endometrioid endometrial cancer; Gu = genomically unstable endometrioid endometrial cancer. **(C)** Sensitivity and specificity for discrimination between (a) endometrium and atypical hyperplasia of endometrium, (b) endometrium and genomically unstable endometrioid endometrial cancer, and (c) atypical hyperplasia of endometrium and genomically unstable endometrioid endometrial cancer as evaluated by receiver-operator curves.

Based on the DNA pattern, EEC cases were divided into two major groups – genomically stable EEC that included five diploid stable cases and two aneuploid stable cases and genomically unstable EEC that consisted of seven diploid unstable cases and one aneuploid unstable case. We performed a principal component analysis (PCA) that considered expression of all protein spots in a 2D gel (Figure 1C). Squamous cervical cancer was included in the comparison as a discriminative cancer with a different pathogenesis and that is characterised by genomic instability. According to the analysis, genomically stable EEC (7 cases), genomically unstable EEC (8 cases), SCC (13 cases), normal endometrium (8 cases) and squamous cervical mucosa (4 cases) clustered separately. Some proximity was observed between the genomically unstable EEC and SCC and between the genomically stable EEC and normal endometrium (Figure 1C).

We identified 121 differentially expressed proteins (Tables 2 and 3 and Supplementary Table S1). The majority of the proteins were overexpressed in the studied cancers. By comparison of EEC and SCC, we extracted 12 proteins explicitly overexpressed in genomically unstable EEC (Tables 2 and 3). Proteins overexpressed in EEC included those that were more expressed in genomically unstable EEC than in genomically stable EEC (44 proteins) and proteins that were more expressed in genomically unstable EEC than in SCC (29 proteins). We did not identify any proteins that


were overexpressed in genomically stable EEC in comparison with normal endometrium. Only a relative overexpression of 27 proteins in genomically stable EEC was observed in comparison with genomically unstable EEC and SCC (Tables 2 and 3 and Supplementary Table S1).

Function relevance of the identified proteins

Functional activity of the identified proteins was analysed using the NCBI/Protein and OMIM databases. We divided the proteins in the major functional groups, that is, regulators of cell cycle and apoptosis, migration and adhesion, metabolism, transcription and translation, maintainers of DNA, members of extracellular matrix and scaffold proteins. We observed that the representation of the proteins that were over- or under-expressed in EEC in the studied functional groups was unequal ($P = 0.0006$; Figure 1D).

Verification of protein identification

In order to confirm the accuracy of the protein identification with MALDI TOF MS, the tissue protein lysates used for 2D gels were immunoblotted using commercially available antibodies against eight selected proteins: EIF4A1, CLIC1, PRDX6, CLIC4, ENO1,

Table 2 Overview of the expression of identified proteins

Abbreviations: EEC = endometrioid endometrial cancer; SCC = squamous cervical cancer.

ANXA4, EMD and Ku70 (Figure 2). Expression of the protein spots in 2D gels of two selected cases of EEC and two controls is shown (Figure 2A). Concomitantly, their expression was verified using western blot for these respective cases (Figure 2B), confirming their expression profile in EEC. The expression profile of EIF4A1 and ANXA4 could not be confirmed by western blot but, conversely, on 2D gels their observed molecular weight was lower than expected. This could be because of cancer-specific overexpression of truncated forms of these proteins.

Expression of EIF4A1, CLIC1, PRDX6, CLIC4, ENO1, ANXA4, EMD and Ku70 in genomically stable and unstable EEC, AH and endometrium as evaluated by IHC

As these identified proteins have not been previously analysed in connection with EEC, their expression was investigated in greater detail. Therefore, a set of independent cases was subjected to IHC, encompassing normal endometrium, AH, a precursor lesion of EEC and genomically stable and unstable EEC (Table 1b). According to the IHC analysis, expression of EIF4A1 and CLIC1 increased in the nuclei of atypical cells and cytoplasmic expression of PRDX6 was enhanced in AH and genomically unstable EEC. A tendency towards decreased cytoplasmic expression of CLIC4 was observed in genomically stable and unstable EEC. Although ENO1 was not significantly overexpressed in 2D gels of EEC (Table 3 and Figure 2A), its cytoplasmic expression was low in AH and genomically unstable EEC (Figure 3Be). Also, low cytoplasmic expression of ANXA4 was observed in genomically stable and unstable EEC (Figure 3Bf). Interestingly, only the N-terminal part of ANXA4 was significantly overexpressed in 2D gels of EEC. This fragment migrated at 17 kDa whereas the molecular mass of the full-length protein is 34 kDa (Figure 2). Nuclear expression of EMD was low in AH, genomically stable and unstable EEC. Expression of Ku70 was highly abundant in endometrium and low in genomically stable and unstable EEC (Figure 3A and B).

Using ROC curves (Figure 3C), we determined that the expression of CLIC1, EIF4A1 and PRDX6 displayed the highest sensitivity and specificity for discrimination between E and AH (Figure 3Ca). Expression of EMD, Ku70 and ANXA4 depicted the highest sensitivity and specificity for discrimination between E, AH and genomically unstable EEC (Figure 3Cb and c). Thus, we demonstrated that changes in protein expression observed in EEC

can already be detected on the level of AH. No statistically significant difference was found between the expression of the proteins in genomically stable and unstable AH.

DISCUSSION

Malignancies are classically divided into diploid and aneuploid based on DNA ploidy. However, it has been shown in breast cancer that further subclassification into stable and unstable diploid and aneuploid tumours provides more accurate prognosis (Kronenwett *et al*, 2006). Our analysis of the tissue proteome of EEC offered a possibility for re-classification of this malignancy into stable and unstable subtypes. In particular, our analysis of 2D gels did not show any difference between the expression of proteins in diploid and aneuploid genomically unstable EEC, but showed a clear difference with diploid genomically stable EEC. In addition, similarities were observed between protein expression in genomically unstable SCC and genomically unstable EEC, suggesting an impact of genomic instability on protein expression. By comparing EEC and SCC, we identified changes in protein expression specific for EEC while excluding proteins commonly overexpressed in most malignancies (Petрак *et al*, 2008).

We also confirmed the identity of several proteins previously found to be overexpressed in endometrial cancer. One interesting example was CAPS (Li *et al*, 2008a), a protein related to low differentiation and worse survival of patients with endometrial cancer (Li *et al*, 2008b). Among the proteins linked to proliferation and invasion of endometrial cancer (Yi *et al*, 2009), we identified HSPA1, TPM2, PDIA, ENO and HNRNPK. Among the proteins downregulated in EEC in connection to invasion into myometrium (Monge *et al*, 2009), we identified MSN (family of EZR), TUBA1B, ANXA1, HNRNPH3 and TALDO1. We also observed a high expression of HSP90AA1, PTGES3 and ATP5B in relation to the stage of EEC (Supplementary Table S2).

Our study focussed on the analysis of protein expression in EEC whereas other groups have analysed chromosomal changes in EEC and in AH (Sonoda *et al*, 1997; Suzuki *et al*, 1997; Kiechle *et al*, 2000; Baloglu *et al*, 2001; Schulten *et al*, 2004; Levan *et al*, 2006; O'Toole *et al*, 2006) and CIN3 and SCC (Heselmeyer *et al*, 1996, 1997) (Supplementary Table S3 and Figure 1E). Once synthesised, proteins generally undergo numerous post-translational modifications in order to become functionally active. We observed

Table 3 Expression of identified proteins in genomically stable and unstable EEC in comparison with SCC

I) Proteins overexpressed in genomically unstable EEC

Protein spot №	GO	Chromosome	Sensitivity, fold			Specificity, fold		
			Genomically unstable EEC compared with E	Genomically stable EEC compared with E	SCC compared with SE	Genomically unstable EEC compared with genomically stable EEC	Genomically unstable EEC compared with SCC	Genomically stable EEC compared to SCC
1	2	3	4	5	6	7	8	9
1757	ANXA4	2p14	7.0699		<i>3.694</i>	10.27676337	2.46501	<i>0.23986</i>
1527	ACP1	2p25.3	2.29794		<i>3.155</i>	1.944458217	1.56593	
849	ADK	10q22.2	1.59812				2.48709	<i>0.20392</i>
935	ALB	4q13.3	2.25781			<i>1.613713545</i>	1.75274	
1378	APOA1	11q23.3	1.98253			2.171906395		<i>0.5843</i>
1603	ASRG1	11q12.3	1.91081		<i>2.9</i>	2.445327832	1.87986	
1300	Atp5b	12q13.3	1.75328			1.692193469		<i>0.40983</i>
1440	CAPS	19p13.3	1.65552			1.891565922	4.13314	<i>2.18504</i>
1461	CAPS	19p13.3	1.62705		<i>1.935</i>		1.93163	<i>1.5037</i>
1217	CLIC1	6p21.33	1.68594			2.268964361		<i>0.52454</i>
1473	CLIC1	6p21.33	4.61806	<i>2.096</i>	<i>1.644</i>	2.20363008	3.51021	<i>1.3929</i>
1814	CLIC1	6p21.33	1.50435			1.756903283		<i>0.61689</i>
1551	EEF1A1	6q13	4.52482		<i>1.576</i>	4.071797985	2.23519	<i>0.54894</i>
1411	EIF4A1	17p13.1	4.19021			3.103980246	2.02587	<i>0.6527</i>
1416	GGCT	7p15.1	2.53028	<i>1.721</i>			1.73179	
1326	GNAI2	3p21.31	3.32624			3.322799134		<i>0.36641</i>
1360	GPD2	2q24.1	1.53774			1.623945126		
1664	NDUFS8	11q13.2	1.53315					<i>0.46004</i>
1012	PDIAS	3q21.1	2.21771			2.551676336		
1174	PDLIT	16q21.3	2.29944			2.11508645	1.6633	
1327	PGAM1	10q24.1	3.25838			2.63807803	1.58352	<i>0.6003</i>
1710	PP1A	7p13	1.55226					
1701	PRDX6	1q25.1	4.58295		<i>1.831</i>	4.713111393	2.51816	<i>0.5343</i>
1456	PTGES3	12q13.3	1.84661		<i>0.652</i>		2.09376	<i>1.68624</i>
1843	RAB1A	2p14	1.59638			2.259058786		<i>0.58553</i>
1845	RAB2A	8q12.1	1.65241			1.557825756	1.95076	
1494	TPT2	13q12.11	1.80436			2.099729929	1.69109	
1538	TUBA4A	19p13.3	3.03048		<i>3.5</i>	2.929321266	1.68441	<i>0.57502</i>
1772	TUBA4A	19p13.3	1.57541		<i>2.248</i>	2.127389193	1.3376	
1820	TUBB	6p21.33	2.61983		<i>0.579</i>	3.243123607	1.6588	<i>0.3113</i>
1568	VDAC2	10q22.2	3.04905			3.132841478	1.81551	<i>0.5795</i>
1470	YWHAE	7p13.3	2.27989		<i>1.614</i>	2.060215246	2.02364	
1435	ZNF510	9q22.33	5.07176			3.65061239	1.5289	<i>0.4188</i>
1427	ZNF510	9q22.33	4.14702		<i>1.654</i>	3.174886752	1.5297	<i>0.4813</i>

II) Proteins overexpressed in EEC and SCC

1	2	3	4	5	6	7	8	9
1205	ACTB	7p22.1	2.09231		2.162	2.295153111	1.51492	<i>0.6601</i>
1529	ACTG1	17q25.3	1.892		2.603	1.762737285		
1250	ACTG1	17q25.3	2.702		<i>1.582</i>	1.876597096		<i>0.6297</i>
1268	ACTG	17q25.3	1.9861			2.468789409		<i>0.53461</i>
1410	ALB	4q13.3	2.11729		<i>1.571</i>	2.35046163		<i>0.40046</i>
1549	ANXA5	4q27	1.97592		2.243	2.542085261		<i>0.56335</i>
1560	ANXA5	4q27	2.06354		3.205	2.195061127	1.63245	
1496	CSNK2A2	16q21	1.64901		1.568	1.894204961		<i>0.60618</i>
1540	EEFIG	11q12.3	2.28445		2.026	2.210725438	1.72025	
1313	HSP90AB1	6p21.1	2.5436		2.214	3.132856488		<i>0.40069</i>
1439	HSP90AA	14q32.31	3.30753		<i>1.699</i>	3.177939769	1.8147	<i>0.571</i>
1837	HSP90AA1	14q32.31	1.53929			1.988560182		<i>0.64644</i>
1192	LAP3	4p15.32	1.63974		2.057	2.233210198		<i>0.49598</i>
1487	NME1	17q21.33	1.55736		1.84			
1499	PACAP	5q31.2	2.45683	<i>0.597</i>	1.852	4.112702544		<i>0.30417</i>
1571	PP1A	7p13	2.29466		2.196	1.786645631		
1331	VIM	10p12.33	2.21312		2.479	1.806580215	1.59759	

III) Proteins overexpressed in SCC

1	2	3	4	5	6	7	8	9
1196	ACTB	7p22.1			2.514			
1189	ACTB	7p22.1			1.924			
1171	ACTB	7p22.1			1.608			
661	ALB	4q13.3	0.62393		1.725	0.660901852	0.5266	
1717	ALDH9A1	1q24.1	0.52235		<i>1.618</i>	0.538209304		
1703	ATPSB	12q13.3	0.48291		<i>1.849</i>	0.57040057	0.50087	
1265	COMT	22q11.21			1.524			
1614	ENO1	1p36.23			<i>1.526</i>			<i>0.53244</i>
1614	FGB	4q32.1			<i>1.526</i>			<i>0.53244</i>
1764	HNRNP	10q11.21	0.6115		1.707		0.61694	
708	HNRNP	9q21.32	0.5867		<i>1.522</i>		0.63126	

Note:

- 1 - Number of a protein spot on 2D gel,
- 2 - Gene ontology name,
- 3 - Gene location on the chromosome,
- 4 - 9: Ratio between the expression of a protein spot where green corresponds to under-expression and red to over-expression (cutoff 1.5, $P < 0.05$):
- 4 - In genomically unstable EEC and endometrium (E),
- 5 - In genomically stable EEC and endometrium (E),
- 6 - In SCC and squamous cervical epithelium (SE),

III) Continue

Protein spot №	GO	Chromosome	Sensitivity, fold			Specificity, fold		
			Genomically unstable EEC compared with E	Genomically stable EEC compared with E	SCC compared with SE	Genomically unstable EEC compared with genomically stable EEC	Genomically unstable EEC compared with SCC	Genomically stable EEC compared with SCC
1	2	3	4	5	6	7	8	9
524	HPX	11p15.4					1.8102	
836	HSPA1A	6p21.33	0.47795				2.0645	
1706	KRT19	17q21.2					0.65241504	0.5357
738	KRT7	12q13.13	0.58145				1.802	
583	LMNB2	19p13.3	0.45832				2.2899	<i>0.666687</i>
584	LMNB2	19p13.3	0.56573				2.081	<i>0.6581</i>
1684	PPP2R1A	19q13.33	0.42397	<i>0.65</i>			1.539	0.6502095
1394	PRDX2	19p13.3					1.553	0.4579
659	SERPINA1	14q32.13	0.39525				1.516	0.49704436
927	SERPINB4	18q21.33						0.6491
1614	SERPINB5	18q21.33					1.526	
1256	SEF	1p36.11						0.5903
1032	TUBA1B	12q13.12					2.1731	0.466914812
708	TUBA1B	12q13.12	0.567				1.522	0.599480212
1084	TUBB2B	6p25.2		<i>0.67</i>			2.3793	0.6313

IV) Proteins partially overexpressed in genomically stable EEC

1	2	3	4	5	6	7	8	9
1766	AIDA	1q41				0.59854996		1.68972448
767	ARHGAP1	11p11.2	0.35985			0.47633145	<i>0.5116</i>	
1724	ANXA5	4q27	0.55344		0.6207	0.5909041		
832	EIF4A3	17q25.3	0.56255		0.6338	0.5107165		1.650582271
855	ERP44	9q31.1	0.4527			0.57666867	<i>0.5459</i>	
676	FBLN5	14q32.12	0.35468		0.4002	0.35120243	<i>0.4968</i>	
515	MSN	Xq11.1			0.666	0.65505196		2.19067482
1128	OGN	9q22.31			0.3633	0.48279406		2.449319102
1618	OGN	9q22.31	0.51672		0.259	0.42482732		2.365572109
1720	PDIA6	2p25.1	0.41745			0.51331209	<i>0.4016</i>	
1821	PP1R7	2q37.3	0.6172			0.62020647		
816	PSMC3	11p11.2	0.44319			0.60637016	<i>0.5544</i>	
923	SUCLG2	3p14.1	0.51342			0.57168192		

V) Proteins underexpressed in SCC

1	2	3	4	5	6	7	8	9
770	CSTF1	20q13.31			0.5856		1.7065	1.659624412
1604	Coor108	6p21.1					1.5212	1.537295552
1223	DDAH2	6p21.33						1.551842189
1019	DCPS	11q24.2	0.63688		0.4306			1.511295
1150	EMD	Xq28			0.3334		1.9417	1.687296397
1020	HDFG	1q23.1			0.4608			1.748306816
972	HNRNP	14q11.2			0.489		1.5506	1.97324742
1612	PPA1	10q22.1						1.649371878
1047	PPP1CB	2p23.2					1.5559	2.161791808
1163	RNF8	6p21.2			0.4866		1.5945	1.66559039
1043	TALDO1	11p15.5			0.6568			1.62284667
1197	TBC1	19q13.12					1.8155	2.118814513
1098	TXNL1	18q21.31			0.5519			1.980964187
1607	SEPT2	2q37.3					1.5318	1.747028989

VI) Proteins underexpressed in EEC and SCC

1	2	3	4	5	6	7	8	9
528	ANXA1	9q21.13	0.57115		0.5102			
1110	GNB2	7q22.1	0.63599	0.664	0.3878			
528	HNRNP3	10q21.3	0.57115		0.5102			
824	SERPINF1	17p13.3	0.65806		0.613			
1103	SFRS1	17q22	0.63987		0.3586			
1103	SFRS7	2p22.1	0.63987		0.3586			

VII) Proteins underexpressed in SCC

1	2	3	4	5	6	7	8	9
1095	EEF1G	8q24.3			0.5052			
841	EEF1G	11q12.3			0.6411			
1817	TPM2	9p13.3			0.6161			
967	VIM	10p12.33			0.4607			
1674	VIM	10p12.33	0.43008		0.5235	0.57977779		1.526384246
592	WDR1	4p16.1			0.6298			
527	XRCC6	22q13.2			0.6217			

VIII) Proteins underexpressed in EEC

1	2	3	4	5	6	7	8	9
664	CCT8	21q21.3	0.56951					
1255	CLIC4	1p36.11	0.62428					
708	VIM	10p12.33	0.567			1.522		0.6313
1712	VIM	10p12.33	0.55433					

underexpression of ENO1 and CLIC4 in both EEC and AH. Interestingly, loss of a specific part of the 1p chromosome is a key event during EEC carcinogenesis and this deleted region is responsible for the synthesis of ENO1 and CLIC4 (Kiechle *et al*, 2000; Baloglu *et al*, 2001). Other early events during EEC carcinogenesis are gains in the entire long arm of the 1q chromosome that contains the gene coding for PRDX6 and losses at 22q chromosome that disrupt the synthesis of Ku70 (XRCC6) (Kiechle *et al*, 2000; Baloglu *et al*, 2001), which also corresponds to our findings on the protein level in EEC and AH. In addition, EEC is characterised by gains at the 2p, 6p, 17p and Xq chromosomes (Suzuki *et al*, 1997) and those are responsible for the synthesis of ANXA4, CLIC1, EIF4A1 and EMD, respectively. In contrast to this, we observed decreased expression of ANXA4 in AH and EEC according to our IHC data, whereas we confirmed increased expression on our 2D gels. This discrepancy can be explained by the fact that the molecular weight of ANXA4 detected on the 2D gels was lower than expected and the protein was represented only by the NH2 domain. This can be due to cancer-specific truncation of the NH2 domain, leading to malfunction of the full-length protein (Gerke and Moss, 2002). EMD was also underexpressed in EEC and AH, which corresponds to its functional role in maintaining chromosomal stability.

For the first time, our paper describes EIF4A1, CLIC1, PRDX6, CLIC4, ENO1, ANXA4, EMD and Ku70 in relation to EEC, although their role is well established in other cancers. *EIF4A1* is overexpressed in hepatocellular carcinoma (Yoon *et al*, 2006) and is an early marker of distant metastases of non-small cell lung cancer (Ji *et al*, 2003). Similarly, we find it overexpressed in AH, suggesting that *EIF4A1* expression could also be used as an early marker of EEC. *CLIC1* is involved in invasion, cancer cell motility (Wang *et al*, 2009) and development of chemoresistance (Kang and Kang, 2008). It is overexpressed in nasopharyngeal carcinoma (Chang *et al*, 2009), colorectal cancer (Petrova *et al*, 2008) and hepatocellular cancer (Huang *et al*, 2004). *PRDX6* protects against oxidative injury, it is overexpressed in endometriosis (Stephens *et al*, 2010) and it increases the invasiveness of breast cancer (Chang *et al*, 2007). *CLIC4* is a chloride intracellular channel that translocates to the nucleus in response to DNA damage and is associated with growth arrest and apoptosis. Moreover, loss of the expression of *CLIC4* in cells and upregulation in stroma is associated with malignant progression (Suh *et al*, 2007a, b). *ENO1*

is a glycolytic enzyme that binds to the promoter of the oncogene *c-myc* and acts as a transcriptional repressor (Feo *et al*, 2000). Therefore, we hypothesise that loss of *ENO1* leads to increased *c-myc* expression, which is known to promote carcinogenesis. The transcription and translation of *ANXA4* in endometrium is regulated by progesterone, an important regulator of cyclic changes in endometrium (Ponnampalam and Rogers, 2006). *EMD* belongs to the inner nuclear membrane proteins that bind chromatin modifiers (Shaklai *et al*, 2007). Its loss in ovarian cancer is considered to be the basis for aneuploidy (Capochichi *et al*, 2009). *Ku70*, or *XRCC6*, is a nuclear complex involved in the repair of double-strand non-homologous DNA breaks. Malfunction of the *XRCC6* gene is observed in ovarian cancer (Kim *et al*, 2010) and breast cancer (Willems *et al*, 2009).

In summary, we analysed the tissue proteome of EEC with respect to genomic stability, one of the most important prognostic markers (Lundgren *et al*, 2002, 2004), and identified differentially expressed proteins. We showed that changes in protein expression could already be detected in precursor lesions, that is, atypical hyperplasia of endometrium, which could provide significant improvement in early detection of EEC.

ACKNOWLEDGEMENTS

This study was supported by the Swedish Cancer Foundation (070623, CAN 2007/1044), KI Cancer Strategic Grants (5888/05-722), Swedish Research Council (521-2008-2899), Medical Research Council, Cancer Society in Stockholm, Stockholm County Council, Swedish Labour Market Insurance, EU-grant FLUODIAMON, Neoproteomics AB and Eurosund AB. We thank Carmen Flores-Staino for technical assistance, Sofia Lundgren Hinnerdal for technical assistance, Pädraig Darcy for proof-reading and Michael Vanlandewijck for supplying lysates of HeLa and MCF7 cells.

Conflict of interest

The authors declare no conflict of interest.

Supplementary Information accompanies the paper on British Journal of Cancer website (<http://www.nature.com/bjc>)

REFERENCES

- Baloglu H, Cannizzaro L, Jones J, Koss L (2001) Atypical endometrial hyperplasia shares genomic abnormalities with endometrioid carcinoma by comparative genomic hybridization. *Hum Pathol* **32**: 616–622
- Bokhman JV (1983) Two pathogenetic types of endometrial carcinoma. *Gynecol Oncol* **15**: 10–17
- Bradford MM (1976) A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal Biochem* **72**: 248–254
- Buchanan EM, Weinstein LC, Hillson C (2009) Endometrial cancer. *Am Fam Physician* **80**: 1075–1080
- Capochichi CD, Cai KQ, Testa JR, Godwin AK, Xu XX (2009) Loss of GATA6 leads to nuclear deformation and aneuploidy in ovarian cancer. *Mol Cell Biol* **29**: 4766–4777
- Casado-Vela J, Rodriguez-Suarez E, Iloro I, Ametzazurra A, Alkorta N, García-Velasco JA, Matorras R, Prieto B, González S, Nagore D, Simón L, Elortza F (2009) Comprehensive proteomic analysis of human endometrial fluid aspirate. *J Proteome Res* **8**: 4622–4632
- Chang XZ, Li DQ, Hou YF, Wu J, Lu JS, Di GH, Jin W, Ou ZL, Shen ZZ, Shao ZM (2007) Identification of the functional role of peroxiredoxin 6 in the progression of breast cancer. *Breast Cancer Res* **9**: R76
- Chang YH, Wu CC, Chang KP, Yu JS, Chang YC, Liao PC (2009) Cell secretome analysis using hollow fiber culture system leads to the discovery of *CLIC1* protein as a novel plasma marker for nasopharyngeal carcinoma. *J Proteome Res* **8**: 5465–5474
- Cheng AL, Huang WG, Chen ZC, Zhang PF, Li MY, Li F, Li JL, Li C, Yi H, Peng F, Duan CJ, Xiao ZQ (2008) Identifying cathepsin D as a biomarker for differentiation and prognosis of nasopharyngeal carcinoma by laser capture microdissection and proteomic analysis. *J Proteome Res* **7**: 2415–2426
- Creasman WT, Odicino F, Maisonneuve P, Quinn MA, Beller U, Benedet JL, Heintz AP, Ngan HY, Pecorelli S (2006) Carcinoma of the corpus uteri. FIGO 26th Annual Report on the Results of Treatment in Gynecological Cancer. *Int J Gynaecol Obstet* **95**(Suppl 1): S105–S143
- Feo S, Arcuri D, Piddini E, Passantino R, Giallongo A (2000) *ENO1* gene product binds to the *c-myc* promoter and acts as a transcriptional repressor: relationship with Myc promoter-binding protein 1 (MBP-1). *J FEBS Lett* **473**: 47–52
- Gerke V, Moss SE (2002) Annexins: from structure to function. *Physiol Rev* **82**: 331–371
- Hellman K, Alaiya A, Becker S, Lomnytska M, Schedvins K, Steinberg W, Hellström A-C, Andersson S, Hellman U, Auer G (2009) Differential tissue-specific protein markers of vaginal carcinoma. *Br J Cancer* **100**: 1303–1314
- Heselmeyer K, Macville M, Schröck E, Blegen H, Hellström AC, Shah K, Auer G, Ried T (1997) Advanced-stage cervical carcinomas are defined

- by a recurrent pattern of chromosomal aberrations revealing high genetic instability and a consistent gain of chromosome arm 3q. *Genes Chromosomes Cancer* 19: 233–240
- Heselmeyer K, Schröck E, du Manoir S, Blegen H, Shah K, Steinbeck R, Auer G, Ried T (1996) Gain of chromosome 3q defines the transition from severe dysplasia to invasive carcinoma of the uterine cervix. *Proc Natl Acad Sci USA* 93: 479–484
- Horn LC, Meinel A, Handzel R, Einkenkel J (2007) Histopathology of endometrial hyperplasia and endometrial carcinoma: an update. *Ann Diagn Pathol* 11: 297–311
- Huang JS, Chao CC, Su TL, Yeh SH, Chen DS, Chen CT, Chen PJ, Jou YS (2004) Diverse cellular transformation capability of overexpressed genes in human hepatocellular carcinoma. *Biochem Biophys Res Commun* 315: 950–958
- Ji P, Diederichs S, Wang W, Böing S, Metzger R, Schneider PM, Tidow N, Brandt B, Buerger H, Bulk E, Thomas M, Berdel WE, Serve H, Müller-Tidow C (2003) MALAT-1, a novel noncoding RNA, and thymosin beta 4 predict metastasis and survival in early-stage non-small cell lung cancer. *Oncogene* 22: 8031–8041
- Kang MK, Kang SK (2008) Pharmacologic blockade of chloride channel synergistically enhances apoptosis of chemotherapeutic drug-resistant cancer stem cells. *Biochem Biophys Res Commun* 373: 539–544
- Kiechle M, Hinrichs M, Jacobsen A, Lüttges J, Pfisterer J, Kommos F, Arnold N (2000) Genetic imbalances in precursor lesions of endometrial cancer detected by comparative genomic hybridization. *Am J Pathol* 156: 1827–1833
- Kim YS, Hwan JD, Bae S, Bae DH, Shick WA (2010) Identification of differentially expressed genes using an annealing control primer system in stage III serous ovarian carcinoma. *BMC Cancer* 10: 576
- Kronenwett U, Ploner A, Zetterberg A, Bergh J, Hall P, Auer G, Pawitan Y (2006) Genomic instability and prognosis in breast carcinomas. *Cancer Epidemiol Biomarkers Prev* 15: 1630–1635
- Levan K, Partheen K, Osterberg L, Helou K, Horvath G (2006) Chromosomal alterations in 98 endometrioid adenocarcinomas analysed with comparative genomic hybridization. *Cytogenet Genome Res* 115: 16–22
- Li Z, Huang C, Bai S, Pan X, Zhou R, Wei Y, Zhao X (2008a) Prognostic evaluation of epidermal fatty acid-binding protein and calcyphosine, two proteins implicated in endometrial cancer using a proteomic approach. *Int J Cancer* 123: 2377–2383
- Li Z, Zhao X, Bai S, Wang Z, Chen L, Wei Y, Huang C (2008b) Proteomics identification of cyclophilin a as a potential prognostic factor and therapeutic target in endometrial carcinoma. *Mol Cell Proteomics* 7: 1810–1823
- Lomnyska M, Becker S, Hellman K, Hellström A-C, Souchelnytskyi S, Mints M, Hellman U, Andersson S, Auer G (2010) Diagnostic marker protein patterns in squamous cervical cancer. *Proteomics Clin Appl* 4: 17–31
- Lomnyska MI, Becker S, Bodin I, Olsson A, Hellman K, Hellström AC, Mints M, Hellman U, Auer G, Andersson S (2011) Differential expression of ANXA6, HSP27, PRDX2, NCF2, and TPM4 during uterine cervix carcinogenesis: diagnostic and prognostic value. *Br J Cancer* 104: 110–119
- Lundgren C, Auer G, Frankendal B, Moberger B, Nilsson B, Nordstrom B (2002) Nuclear DNA content, proliferative activity, and p53 expression related to clinical and histopathologic features in endometrial carcinoma. *Int J Gynecol Cancer* 12: 110–118
- Lundgren C, Auer G, Frankendal B, Nilsson B, Nordstrom B (2004) Prognostic factors in surgical stage I endometrial carcinoma. *Acta Oncol* 43: 49–56
- Lundgren C, Lahmann J, Becker S, Roblick U, Schedvins K, Boman K, Frankendal B, Nordström B, Auer G (2006) 2-DE protein expression in endometrial carcinoma. *Acta Oncol* 45: 685–694
- Monge M, Colas E, Doll A, Gil-Moreno A, Castellvi J, Diaz B, Gonzalez M, Lopez-Lopez R, Xercavins J, Carreras R, Alameda F, Canals F, Gabrielli F, Reventos J, Abal M (2009) Proteomic approach to ETV5 during endometrial carcinoma invasion reveals a link to oxidative stress. *Carcinogenesis* 30: 1288–1297
- O'Toole SA, Dunn E, Shepard BL, Klocker H, Bektic J, Smyth P, Martin C, Sheils O, O'Leary JJ (2006) Genome-wide analysis of deoxyribonucleic acid in endometrial cancer using comparative genomic hybridization microarrays. *Int J Gynecol Cancer* 16: 834–842
- Petrak J, Ivanek R, Toman O, Cmejla R, Cmejlova J, Vyoral D, Zivny J, Vulpe CD (2008) Déjà vu in proteomics. A hit parade of repeatedly identified differentially expressed proteins. *Proteomics* 8: 1744–1749
- Petrova DT, Asif AR, Armstrong VW, Dimova I, Toshev S, Yaramov N, Oellerich M, Toncheva D (2008) Expression of chloride intracellular channel protein 1 (CLIC1) and tumor protein D52 (TPD52) as potential biomarkers for colorectal cancer. *Clin Biochem* 41: 1224–1236
- Pitteri S, Hanash S (2010) A systems approach to the proteomic identification of novel cancer biomarkers. *Dis Markers* 28: 233–239
- Ponnampalam AP, Rogers PA (2006) Cyclic changes and hormonal regulation of annexin IV mRNA and protein in human endometrium. *Mol Hum Reprod* 12: 661–669
- Schulten H-J, Gunawan B, Enders Ch, Donhuijsen K, Emons G, Fuzes L (2004) Overrepresentation of 8q in carcinosarcomas and endometrial adenocarcinomas. *Am J Clin Pathol* 122: 546–551
- Seebacher V, Schmid M, Polterauer S, Hefler-Frischmuth K, Leopold H, Concin N, Reinthaller A, Hefler L (2009) The presence of postmenopausal bleeding as prognostic parameter in patients with endometrial cancer: a retrospective multi-centre study. *BMC Cancer* 9: 460
- Shaklai S, Amariglio N, Rechavi G, Simon AJ (2007) Gene silencing at the nuclear periphery. *FEBS J* 274: 1383–1392
- Sharon D, Chen R, Snyder M (2010) Systems biology approaches to disease marker discovery. *Dis Markers* 28: 209–224
- Sonoda G, du Manoir S, Godwin AK, Bell DW, Liu Z, Hogan M, Yakushiji M, Testa JR (1997) Detection of DNA gains and losses in primary endometrial carcinomas by comparative genomic hybridization. *Genes Chromosomes Cancer* 18: 115–125
- Steinbeck RG, Auer GU, Zetterberg AD (1999) Reliability and significance of DNA measurements in interphase nuclei and division figures in histological sections. *Eur J Cancer* 35: 787–795
- Stephens AN, Hannan NJ, Rainczuk A, Meehan KL, Chen J, Nicholls PK, Rombauts LJ, Stanton PG, Robertson DM, Salomonsen LA (2010) Post-translational modifications and protein-specific isoforms in endometriosis revealed by 2D DIGE. *J Proteome Res* 9: 2438–2449
- Suh KS, Crutchley JM, Koochek A, Ryscavage A, Bhat K, Tanaka T, Oshima A, Fitzgerald P, Yuspa SH (2007a) Reciprocal modifications of CLIC4 in tumour epithelium and stroma mark malignant progression of multiple human cancers. *Clin Cancer Res* 13: 121–131
- Suh KS, Malik M, Shukla A, Yuspa SH (2007b) CLIC4, skin homeostasis and cutaneous cancer: surprising connections. *Mol Carcinog* 8: 599–604
- Suzuki A, Fukushige S, Nagase S, Ohuchi N, Satomi S, Horii A (1997) Frequent gains on chromosome arms 1q and/or 8q in human endometrial cancer. *Hum Genet* 100: 629–636
- Wang JW, Peng SY, Li JT, Wang Y, Zhang ZP, Cheng Y, Cheng DQ, Weng WH, Wu XS, Fei XZ, Quan ZW, Li JY, Li SG, Liu YB (2009) Identification of metastasis-associated proteins involved in gallbladder carcinoma metastasis by proteomic analysis and functional exploration of chloride intracellular channel 1. *Cancer Lett* 281: 71–81
- Willems P, De Ruyck K, Van den Broecke R, Makar A, Perletti G, Thierens H, Vral A (2009) A polymorphism in the promoter region of Ku70/XRCC6, associated with breast cancer risk and oestrogen exposure. *J Cancer Res Clin Oncol* 135: 1159–1168
- Yi Z, Jingting C, Yu Z (2009) Proteomics reveals protein profile changes in cyclooxygenase-2 inhibitor-treated endometrial cancer cells. *Int J Gynecol Cancer* 19: 326–333
- Yoon SY, Kim JM, Oh JH, Jeon YJ, Lee DS, Kim JH, Choi JY, Ahn BM, Kim S, Yoo HS, Kim YS, Kim NS (2006) Gene expression profiling of human HBV- and/or HCV-associated hepatocellular carcinoma cells using expressed sequence tags. *Int J Oncol* 29: 315–327

This work is published under the standard license to publish agreement. After 12 months the work will become freely available and the license terms will switch to a Creative Commons Attribution-NonCommercial-Share Alike 3.0 Unported License.