

OPEN

# STROBE-Long-Term Exposure to Ambient Fine Particulate Air Pollution and Hospitalization Due to Peptic Ulcers

Chit-Ming Wong, PhD, Hilda Tsang, MPhil, Hak-Kan Lai, PhD, Thuan-Quoc Thach, PhD, G. Neil Thomas, PhD, King-Pan Chan, MPhil, Siu-Yin Lee, MD, Jon G. Ayres, MD, Tai-Hing Lam, MD, and Wai K. Leung, MD

**Abstract:** Little is known about the effect of air pollution on the gastrointestinal (GI) system. We investigated the association between long-term exposures to outdoor fine particles (PM<sub>2.5</sub>) and hospitalization for peptic ulcer diseases (PUDs) in a large cohort of Hong Kong Chinese elderly.

A total of 66,820 subjects aged ≥65 years who were enrolled in all 18 Government Elderly Health Service centers of Hong Kong participated in the study voluntarily between 1998 and 2001. They were prospectively followed up for more than 10 years. Annual mean exposures to PM<sub>2.5</sub> at residence of individuals were estimated by satellite data through linkage with address details including floor level. All hospital admission records of the subjects up to December 31, 2010 were retrieved from the central database of Hospital Authority. We used Cox regression to estimate the hazard ratio (HR) for PUD hospitalization associated with PM<sub>2.5</sub> exposure after adjustment for individual and ecological covariates.

A total of 60,273 subjects had completed baseline information including medical, socio-demographic, lifestyle, and anthropometric data at recruitment. During the follow-up period, 1991 (3.3%) subjects had been hospitalized for PUD. The adjusted HR for PUD hospitalization per 10 μg/m<sup>3</sup> of PM<sub>2.5</sub> was 1.18 (95% confidence interval: 1.02–1.36, *P* = 0.02). Further analysis showed that the associations with PM<sub>2.5</sub> were significant for gastric ulcers (HR 1.29; 1.09–1.53, *P* = 0.003) but not for duodenal ulcers (HR 0.98; 0.78 to 1.22, *P* = 0.81).

Long-term exposures to PM<sub>2.5</sub> were associated with PUD hospitalization in elder population. The mechanism underlying the PM<sub>2.5</sub> in the development of gastric ulcers warrants further research.

(*Medicine* 95(18):e3543)

**Abbreviations:** COPD = chronic obstructive pulmonary disease, GI = gastrointestinal, HR = hazard ratio, NSAID = nonsteroidal inflammatory drug, PM = particulate matter, PM<sub>2.5</sub> = particles less

Editor: Bing-Fang Hwang.

Received: July 27, 2015; revised: March 9, 2016; accepted: April 6, 2016. From the School of Public Health, Li Ka Shing Faculty of Medicine, The University of Hong Kong, Faculty of Medicine Building, Pokfulam (CMW, HT, HKL, TQT, KPC, THL); Institute of Applied Health Research, The University of Birmingham, Edgbaston, Birmingham, UK (GNT, JGA); Department of Health, Wu Chung House, Wan Chai (SYL); and Department of Medicine, Li Ka Shing Faculty of Medicine, The University of Hong Kong, Queen Mary Hospital, Pokfulam Road, Hong Kong (WKL). Correspondence: Thuan-Quoc Thach, School of Public Health, Li Ka Shing Faculty of Medicine, The University of Hong Kong, 5/F William MW Mong Block, Faculty of Medicine Building, 21 Sassoon Road, Hong Kong (e-mail: thach@hku.hk).

Supplemental Digital Content is available for this article.

This work was supported by the Wellcome Trust [#094330/Z10/Z].

The authors have no conflicts of interest to disclose.

Copyright © 2016 Wolters Kluwer Health, Inc. All rights reserved.

This is an open access article distributed under the Creative Commons Attribution License 4.0, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. ISSN: 0025-7974

DOI: 10.1097/MD.00000000000003543

than 2.5 μm in aerodynamic diameter, PUD = peptic ulcer disease, SD = standard derivation, SEC = surface extinction coefficients.

## INTRODUCTION

Air pollution is a major public health hazard, particularly in developing countries.<sup>1</sup> The World Health Organization estimated substantial premature death attributable to ambient air pollution in 2012.<sup>2</sup> In particular, there is a causal association between air pollution and morbidity or mortality from cardiorespiratory diseases.<sup>3</sup> Among various World Health Organization criterion pollutants, particulate matter (PM) which is emitted from burning of fossil fuels in vehicles, shipping and power generators, is often considered the most relevant to public health intervention.<sup>4</sup> PM is classified by its aerodynamic diameter of less than 2.5 μm into PM<sub>2.5</sub> (particles less than 2.5 μm in aerodynamic diameter), which can enter the blood stream via the alveolar capillaries of the lungs with a higher risk to cause serious health problems.<sup>5</sup> By using chemistry model to estimate exposure based on emission source categories, air pollution, mostly PM<sub>2.5</sub>, in urban and rural environments, would lead to 3.3 million premature deaths per year worldwide, predominantly in Asia.<sup>6</sup>

Recent long-term studies have estimated the horizontal spatial variation of PM using geospatial or dispersion modeling and satellite information as proxy indicators of exposure in individual residential areas.<sup>7–10</sup> However, measurement errors in estimating the exposure exist if the spatial variation in the vertical dimension has not been taken into account. Considering the vertical dimension is particularly important for studies in populations mainly living in high-rise buildings like Hong Kong. People who stay near the ground level could be exposed to as much as 5-time higher pollutant concentrations than those staying on floors above ground level, depending on the street canyon characteristics.<sup>11</sup>

Peptic ulcer disease (PUD) is a very common gastrointestinal (GI) disease in Hong Kong. It is estimated that more than 18.6% of all hospitalizations for GI diseases were due to PUD in Hong Kong in 2011.<sup>12</sup> Although the majority of PUD is caused by chronic *Helicobacter pylori* (*H pylori*) infection and the use of nonsteroidal inflammatory drugs (NSAIDs) or aspirin, the “*H pylori*-negative ulcers” in non-NSAID users appear to be on the rise, particularly in Asia.<sup>13</sup> Factors other than *H pylori* and NSAIDs may play an important but yet undefined role on the pathogenesis of this group of PUD.<sup>14</sup>

In the present study, we studied the effects of long-term exposure to PM<sub>2.5</sub> on hospital admissions for PUD in a large group of elderly persons from all 18 districts of Hong Kong. These subjects had participated in the Government Elderly Health Service during the recruitment period and were followed up for 10 to 13 years. The 1st hazard admissions to hospital due

to PUD during the follow-up period were regressed on the PM<sub>2.5</sub> concentration estimated at their residential addresses, taking into account both the horizontal and vertical spatial variations.

## METHODS

### Subjects

From July 1998 to December 2001, 66,820 adults aged 65 years or above were enrolled by the Elderly Health Service of the Department of Health of the Hong Kong Government for participation in a client-oriented primary health care service that aimed to promote the health of elderly population and to enhance self-care ability so as to minimize illness and disability ([www.info.gov.hk/elderly](http://www.info.gov.hk/elderly)). These Elderly Health Centers were located one in each of all the 18 districts in Hong Kong, which provide clinical services of health assessment, physical check-up, counseling, curative treatment, and health education to the elderly clients. All participants had their socio-demographic data, lifestyles, health, and anthropometric measurements recorded at baseline.<sup>10</sup> During the recruitment, subjects agreed that the data they provided could be used in research to improve the health of elderly population. The database was stored and managed by the Department of Health, and the anonymous data were available for public health research. Pseudo-identifiers were used to match up records from the Elderly Health Centre to the Hospital Authority data. The investigators and the research workers of the project were blinded to the link between the identity and personal data of the individual subjects. The study protocol was approved by the Institutional Review Board of The University of Hong Kong/Hospital Authority Hong Kong West Cluster and the Ethics Committee of Department of Health of Hong Kong Government.

### Exposure Model to Air Pollution and Estimation of Residential Exposure

The annual mean concentrations (in  $\mu\text{g}/\text{m}^3$ ) of PM<sub>2.5</sub> from 1998 to 2011 were measured hourly in 5 urban locations by tapered element oscillating microbalance of the Environmental Protection Department of the Hong Kong Government (<http://epic.epd.gov.hk/EPICDI/air/station/>). All the selected air pollutant monitoring stations ( $n=5$ ) were located throughout the metropolitan areas with height ranging from 11 to 27 m above the ground. In all the monitoring locations, we determined vertical height (m) above the mean sea level and the satellite-derived surface extinction coefficients (SEC) (within the  $1 \times 1$  km grid area of the station).<sup>15</sup> We then fitted regression models to estimate annual PM concentrations using SEC and inverse vertical height as independent variables and validated the approach by comparing with data estimated from leaving out 1 station and by comparing with subset of the data estimated from an independent study (eAppendix, <http://links.lww.com/MD/A925>).

We matched all residential addresses of the subjects with the SEC data by geocoding and estimated the vertical height of the residence based on the floor level on which they live. Using the exposure model, we determined the annual mean exposures to PM<sub>2.5</sub> for each residential location.

### Hospital Admissions for PUD

Hospitalization records of all participants from 1998 to 2010 were retrieved from the electronic health record system of the Hospital Authority of Hong Kong, which manages all public hospitals in Hong Kong, that accounts for 88% of all hospital

beds and 78% of total hospital admissions locally.<sup>16</sup> We captured the primary cause of hospitalization using the International Statistical Classification of Diseases (ICD9-CM) codes for specific site of PUD (531–532) and its subcategories: gastric ulcer (531) and duodenal ulcer (532) during the study period of all enrolled subjects.<sup>17</sup> We also included in the analysis another GI disease: reflux esophagitis (530.1×, 530.2, 530.81, 530.82, and 530.85). To assess the specificity of effects, we assessed the causes of hospital admissions which were considered to be unrelated to PM exposure including poisoning and injuries (800–999), diseases of skin and subcutaneous (680–709), and oral structures (520–529). To assess the sensitivity of our spatial exposure model of PM in this cohort, hospital admissions due to cardiovascular and respiratory diseases (390–459, 460–519), which are known to be associated with PM exposure, were included.

### Individual, Ecological, and Environmental Covariates

We determined the impact of individual, ecological, and environmental covariables as potential confounding factors for hospital admissions. Individual covariates were age, gender, body mass index, and the presence of medical illnesses that require regular medical care (including hypertension, heart diseases, diabetes, chronic obstructive pulmonary disease [COPD], stroke, and other diseases), smoking status (never, quit, and current smoker), exercise frequency (days per week), education level (secondary or above, primary, and below primary), and personal monthly expenditure (US\$ < 128, 128–384, and  $\geq 385$ ) that were available in the questionnaire (eAppendix Table 1, <http://links.lww.com/MD/A925>).<sup>18</sup> Based on the Census Statistics of Hong Kong in 2001, we determined ecological covariates including percentage (%) of elderly subjects (aged 65 or elderly), % of tertiary education, and average monthly domestic household income in the 197 small areas, namely Tertiary Planning Units of Hong Kong. Environmental covariates included % of smokers in 18 districts of Hong Kong as proxy for exposure to environmental tobacco smoke in each year.<sup>19,20</sup>

### Statistical Analysis

We used Cox proportional hazards model to estimate the hazard ratio (HR) for every  $10 \mu\text{g}/\text{m}^3$  increase of long-term exposure to PM<sub>2.5</sub> concentration on hospital admissions related to 1st diagnosis of PUD after the baseline year with adjustments for individual, ecological, environmental covariates, and the year-to-year changes in exposure to take account of potential changes from the baseline year. We put into the model, the baseline year PM<sub>2.5</sub> as time-independent variable and the year-to-year change which was the difference between the annual average at the year of hospitalization and at baseline year, as the time-dependent variable. We used time-on-study as a timescale and the estimated exposure in the baseline year to represent long-term exposure.

### Sensitivity Analysis

To reduce the possibility that the association between PUD and PM<sub>2.5</sub> was confounded by comorbidities of the subjects, we performed 4 different sensitivity analyses. In the 1st model (covariate adjustment I), we adjusted with an indicator variable for previous hospitalizations due to ischemic heart disease (ICD-9 410–414), stroke (430–438), COPD (490–496), or type 2 diabetes mellitus (250.0, 250.2). These diseases have

been previously shown to be associated with air pollution or the use of antiplatelet therapy that increases the risk of PUD.<sup>21–24</sup> For the 2nd model (covariate adjustment II), we adjusted in the main model another indicator variable for the presence of cardiopulmonary diseases and diabetes as reported by subjects at baseline, while not including other self-reported diseases. For the 3rd model (stratification I), we removed all subjects from the analyses with previous hospitalizations due to the 4 preexisting diseases described in the 1st model. For the 4th model (stratification II), we removed all subjects from the analyses with history of hospitalizations due to PUD during the baseline year. Cox models were performed using the command PHREG in SAS 9.2 (SAS Institute, Inc., Cary, NC). All authors had access to the study data and had reviewed and approved the final manuscript.

## RESULTS

### Concentrations of PM<sub>2.5</sub>

The overall mean concentrations of PM<sub>2.5</sub> recorded by the Environmental Protection Department monitors from 1998 to 2010 were 37.7  $\mu\text{g}/\text{m}^3$  (standard derivation [SD] 7.1  $\mu\text{g}/\text{m}^3$ ), and the annual mean concentrations (95% central range) estimated at individual residential locations in the baseline year was 33.7  $\mu\text{g}/\text{m}^3$  (29.5–39.8  $\mu\text{g}/\text{m}^3$ ) (Figure 1). The patterns of PM<sub>2.5</sub> concentrations were similar among the 5 monitoring stations during the study period (eAppendix Figure 1, <http://links.lww.com/MD/A925>).

### Hospitalizations for PUD

A total of 66,820 subjects were included in the initial study cohort. Among them, completed geocoding and satellite information was available in 60,273 subjects (90.1%; missing data due to missing individual-level covariates: 0.2%, problems in geo-coding: 8.1%, or problems in satellite data: 1.5%) who were included in the Cox model. The average age of these subjects was 72.1 years. There were 66% female and 71% never smokers (Table 1). The mean follow-up in the study was 10.1 (SD 2.9) years and the total follow-up was 664,927 person-years.

During the follow-up period, 1991 (3.3%) subjects had been hospitalized for the diagnosis of PUD, with a mean length of stay of 7.9 (SD 12.3) days. The factors which were significantly associated with PUD hospitalizations included PM<sub>2.5</sub>, elderly age, male gender, body mass index in 4th quartile ( $>26.3 \text{ kg}/\text{m}^2$ ), ever smoked (current or former smokers), low education level (primary or below), coexisting medical diseases that required regular medical care, and living in a community with low education level and with more smokers (Table 2).

After adjusting for potential confounding variables, the HR (95% confidence interval) estimate for hospitalization per 10  $\mu\text{g}/\text{m}^3$  increase of PM<sub>2.5</sub> was 1.18 (1.02–1.36) for PUD ( $n = 1991$ ), 1.29 (1.09–1.53) for gastric ulcer ( $n = 1175$ ), and 0.98 (0.78–1.22) for duodenal ulcer ( $n = 816$ ) (Table 3). There was however no significant association between peptic ulcer bleeding and PM<sub>2.5</sub> levels (HR 1.01; 0.81–1.27). For all other control health outcomes considered, the estimates were not statistically different from unity, while there was a significant association between PM<sub>2.5</sub> exposure and cardiovascular and respiratory diseases (Table 3).

### Sensitivity Analyses

The association between PM<sub>2.5</sub> and PUD remained significant after controlling for preexisting medical diseases by 2 different covariate adjustments. The magnitude of HR for PUD

was similar to the original values after adjusting for coexisting medical diseases or after removing subjects with comorbidities or hospitalized during the baseline year (Table 4). In particular, the HR remained around 1.28 to 1.29 for gastric ulcer.


## DISCUSSION

Ambient PM<sub>2.5</sub> exposure has been found to be one of the leading causes of death and disability in the world.<sup>25</sup> We have demonstrated for the 1st time an association between long-term exposure to PM<sub>2.5</sub> and PUD among elderly people in Hong Kong. Specifically, a significant association for gastric ulcer has been shown both before and after adjustment for other potential confounding factors including baseline medical conditions and hospitalization due to other diseases. Plots of the monthly counts of hospitalization for PUD along with PM<sub>2.5</sub> and particles less than 10 micrometers in aerodynamic diameter (PM<sub>10</sub>) monthly concentration from 2001 to 2011 (eAppendix Figure 2, <http://links.lww.com/MD/A925>) suggested that there were seasonal variations showing winter (December–February) peaks. The patterns were congruent to the global patterns of PUD seasonality and lent support to the hypothesis of this study that PUD hospitalizations are related to environmental factors.<sup>26</sup> However, the covariations between PUD and PM<sub>2.5</sub> concentration should be studied with adjustment for confounding.

In this study, we used a reliable and comprehensive prospective health database from the Department of Health, which include a large cohort of 66,820 elderly people who participated in the study and were being followed up for more than 10 years. The baseline medical and socio-economic information of the participants were collected and their subsequent hospital admission records were retrieved from the central database of the Hospital Authority, which captures all hospital admission records of public hospitals in Hong Kong.<sup>27</sup> We believe that the data are very comprehensive and would be representative of local situations in Hong Kong as the majority of elderly patients are admitted to public hospitals through the accident and emergency department when needed. With the aging population and the air pollution in Hong Kong, the potential impact on health care utilization should not be underestimated.

PM<sub>2.5</sub> typically originates from all types of organic material combustion in motor vehicles, power plants, and industrial processes in Hong Kong. PM<sub>10</sub> (comprising about 70% PM<sub>2.5</sub>) mainly contains chemical compounds (nitrate [27.9%], sulfate [19.5%], and ammonium [8.6%]), as well as carbonaceous (OC [15.9%] and EC [7.6%]) and metal species (Na<sup>+</sup>, Cl, Ca, K<sup>+</sup>, Fe, Mg, Al, Pb, Mn, V, As, Ni, and Cd) (eAppendix Table 2, <http://links.lww.com/MD/A925>). The chronic stimulation from long-term PM<sub>2.5</sub> exposure can induce systemic inflammatory responses and oxidative stress, which is triggered by acidity, transition metals or the ultrafine fraction in particles.<sup>5</sup> Due to the relatively small size, PM<sub>2.5</sub> can escape phagocytosis.<sup>28</sup> PM<sub>2.5</sub> has been linked to inflammation in blood vessels, which increases the risk of plaque deposition and atherosclerosis.<sup>29</sup> However, the mechanistic pathway linking PM<sub>2.5</sub> and peptic ulcer development remains elusive. Yet particles, which block sunlight and prevent absorption of vitamin D useful in enhancing intestinal calcium absorption and mobilizing osteoclastic activities, would lead to development of peptic diseases.<sup>30</sup> Although air pollutants affect the respiratory or circulation system directly, the potential effects on the GI tract is biological plausible. First, human experimental studies





**FIGURE 1.** Subject's residential location and the estimated particles less than 2.5 micrometers in aerodynamic diameter (PM<sub>2.5</sub>) annual mean concentrations at lower (14 m) and higher (60 m) heights above ground level at baseline. Subject's residential location and the estimated annual mean PM<sub>2.5</sub> concentrations at different heights above ground level at baseline. The lower (14 m) and higher (60 m) heights represent the 25th and 75th percentiles of height above ground level for the cohort's residential locations. Height from mean sea level to the ground level was taken into account in the exposure model.

have shown that air pollutants can enter the aerodigestive tract through the swallowing of inhaled PMs cleared from the lungs into the intestine,<sup>31</sup> as well as through ingestion of contaminated food and water.<sup>32,33</sup> Second, by means of a combination of factors including increased gut permeability,<sup>34</sup> decreased colonic motility and clearance,<sup>31,35</sup> altered gut microbial composition and metabolic function,<sup>29,36</sup> air pollutants including PM can induce adverse effects in the GI tracts.<sup>37</sup> In epidemiology studies, long-term exposure to higher concentrations of air pollutants including PM has been associated with an increased risk of early-onset Crohn disease,<sup>32</sup> increased risk of hospitalizations for inflammatory bowel diseases,<sup>38,39</sup> and increased standardized mortality ratio for peptic ulcer.<sup>40</sup> Recently a case cross-over analysis showed that PM<sub>2.5</sub> was inversely associated with upper GI bleeding secondary to PUD.<sup>41</sup> However, the significant estimates were not replicated in another data of the study and the assessment was for short-term instead of for long-term exposure.

Apart from *H pylori*, aspirin and NSAIDs, stress is another important cause of PUD, particularly in critically ill patients.<sup>42</sup> Recent studies from Japan demonstrated that both physical and

psychological stresses related to earthquakes play a pivotal role on the pathogenesis of PUD.<sup>43</sup> We hypothesize that the association between PM<sub>2.5</sub> exposure and PUD, particularly for gastric ulcer, is also mediated through stress and inflammation. It is interesting to note that only gastric ulcer appears to be associated with the long-term PM<sub>2.5</sub> exposure in the present study. Although duodenal ulcer is principally considered to be an acid related disease, the pathogenesis of gastric ulcer is likely related to breakdown of mucosal defense such as disruption of prostaglandin synthesis and even local irritation.<sup>44,45</sup>

Although air pollution is a major health hazard in many developing countries including China, our findings may suggest a novel risk factor for gastric ulcer. Further studies are needed to examine the role of PM<sub>2.5</sub> on peptic ulcer development. In keeping with our current observation, we have previously shown that about 4% of patients with bleeding PUD were negative for *H pylori* and NSAIDs in Hong Kong.<sup>13</sup> There has also been a marked increase in these "idiopathic ulcers" in the past years, with a rise from 4.2% in 1997 to 1998 to 18.8% in 2000.<sup>46</sup> Consistent findings were reported across countries in

**TABLE 1.** Characteristics of Subjects in Baseline (1998–2001)

Characteristics, %	All Subjects (N = 60,273)	%
PM <sub>2.5</sub> concentration, μg/m <sup>3</sup> (mean ± SD)	33.7 ± 3.2	
Age, year (mean ± SD)	72.1 ± 5.6	
Gender		
Male	20,493	34.0
Female	39,780	66.0
BMI quartiles		
Q2–Q3 (21.6–26.3)	30,558	50.7
Q1 (<21.6)	13,863	23.0
Q4 (>26.3)	15,852	26.3
Smoking		
Never	42,854	71.1
Quit	11,633	19.3
Current	5786	9.6
Exercise, days/week (mean ± SD)	5.8 ± 2.4	
Education		
Secondary or above	10,307	17.1
Primary	22,180	36.8
Below primary	27,786	46.1
Expenses/month		
Low*	9945	16.5
Medium	41,347	68.6
High	8981	14.9

BMI = body mass index, PM<sub>2.5</sub> = particles less than 2.5 μm in aerodynamic diameter, SD = standard deviation.

\*Expense per month, “low” represented <HKD\$1000 (US\$128), “medium” represented HKD\$1000 to 2999 (US\$128–384), “high” represented ≥HKD\$3000 (US\$385).

Asia with the proportion of patients with idiopathic ulcers ranging from 10% to 30%.<sup>14</sup> These studies point to a potential unidentified cause of idiopathic ulcers, which may be partly attributable to PM<sub>2.5</sub> exposure.

We merged several large databases to derive the estimation of PM<sub>2.5</sub> pollutant levels in this study. We estimated exposure in individual residential locations using both satellite information and information from the addresses including vertical height (estimated from floor level) of the subjects. In Hong Kong, it has been demonstrated that the concentration of PM is inversely related to the height above ground in an exponential decay pattern (eAppendix, <http://links.lww.com/MD/A925>). By taking into account the spatial variation of exposure, the measurement can minimize the underestimation of adverse health effects.<sup>8,47</sup> A thorough examination of the health effects of PM exposure by this spatial model has confirmed the presence of commonly detectable health effects of PM including hospitalization for cardiovascular and respiratory diseases, and the lack of associations between PM exposure and diseases previously shown to be unrelated to air pollution. All of these collectively provide support for the validity of our new findings of the associations between PUD and long-term PM<sub>2.5</sub> exposure.

This study has some limitations. This study only included elderly subjects who were previously enrolled in the Elderly Health Service. However, most of the hospitalizations related to PUD in Hong Kong are attributed to elderly patients. The presence of cardiovascular disease or

**TABLE 2.** Hazard Ratios (95% CI) of Individual, Ecological, and Environmental Covariates on Hospital Admission for PUD

Characteristics	Hazard Ratio	P Value
PM <sub>2.5</sub> concentration (per 10 μg/m <sup>3</sup> )	1.18 (1.02–1.36)	0.02
Change of PM <sub>2.5</sub> from baseline*	0.98 (0.95–1.00)	0.10
Individual covariates		
Age, year	1.06 (1.06–1.07)	<0.001
Gender		
Male	1	
Female	0.68 (0.60–0.76)	<0.001
BMI quartiles		
Q2–Q3 (21.6–26.3)	1	
Q1 (<21.6)	0.91 (0.81–1.02)	0.12
Q4 (>26.3)	1.17 (1.06–1.30)	0.002
Smoking		
Never	1	
Quit	1.20 (1.06–1.36)	0.004
Current	1.59 (1.37–1.84)	<0.001
Exercise, days/week	0.99 (0.95–1.03)	0.66
Education		
Secondary or above	1	
Primary	1.20 (1.04–1.37)	0.011
Below primary	1.42 (1.23–1.64)	<0.001
Expenses/month†		
Low	0.91 (0.77–1.08)	0.28
Medium	1.02 (0.90–1.16)	0.70
High	1	
Coexisting medical illnesses‡	1.39 (1.24–1.56)	<0.001
Ecological covariates		
Percentage (%) of older subjects (age ≥ 65)	0.99 (0.97–1.00)	0.075
% of education as tertiary level	0.99 (0.98–1.00)	0.058
% of income/month ≥US\$1923	1.00 (0.99–1.01)	0.99
Environmental covariates		
Environmental tobacco smoker (in term of % of smokers)	1.24 (1.01–1.53)	0.042

BMI = body mass index, CI = confidence interval, PM<sub>2.5</sub> = particles less than 2.5 μm in aerodynamic diameter, PUD = peptic ulcer disease.

\*Changes of PM<sub>2.5</sub> from baseline: it is the year-to-year changes in exposure from the baseline year, for which the PM<sub>2.5</sub> level in baseline is subtracted from the PM<sub>2.5</sub> level of the current year, while the temporal linear trend of PM<sub>2.5</sub> concentration having been adjusted for.

†Expense per month, “low” represented <HKD\$1000 (US\$128), “medium” represented HKD\$1000 to 2999 (US\$128–384), and “high” represented ≥HKD\$3000 (US\$385).

‡Coexisting medical illnesses: self-reported active diseases including hypertension, heart diseases, diabetes, COPD, stroke, and other diseases that require regular medical care.

stroke would increase the consumption of aspirin which increase the risk of ulcer related hospitalization. To compensate for this, we have adjusted for baseline medical illnesses including cardiorespiratory diseases and diabetes, or hospitalization for ischemic heart diseases, stroke, COPD, and type 2 diabetes that would increase the use of aspirin in different sensitivity analyses. Moreover, we did not have any information on the *H pylori* status of the subjects including those individuals who were hospitalized for PUD since this information was not available in the central hospital database. Also, we did not have access to individual endoscopy findings. Although we have demonstrated a significant

**TABLE 3.** Hazard Ratios (95% CI) of Hospitalization Due to PUD or Other Causes Per 10  $\mu\text{g}/\text{m}^3$  of  $\text{PM}_{2.5}$  (N = 60,273)

Hospitalization Causes*	n	Hazard Ratios	P Value
Hospitalization due to PUD			
Peptic ulcer	1991	1.18 (1.02–1.36)	0.022
Gastric ulcer	1175	1.29 (1.09–1.53)	0.003
Duodenal ulcer	816	0.98 (0.78–1.22)	0.81
Peptic ulcer bleeding	948	1.01 (0.81–1.27)	0.70
Hospitalization due to other causes			
Cardiovascular	15,518	1.03 (1.00–1.06)	0.0498
Respiratory	11,563	1.04 (1.00–1.07)	0.045
Oral cavity, salivary glands, jaws	186	1.04 (0.60–1.81)	0.90
Reflux esophagitis	453	1.01 (0.70–1.45)	0.95
Skin and subcutaneous tissue	2449	1.02 (0.86–1.20)	0.83
Poisoning and injuries	10,481	1.02 (0.95–1.09)	0.64

CI = confidence interval, ICD = International Statistical Classification of Diseases,  $\text{PM}_{2.5}$  = particles less than 2.5  $\mu\text{m}$  in aerodynamic diameter, PUD = peptic ulcer disease.

\*ICD-9 codes: peptic ulcer (531–532), gastric ulcer (531), duodenal ulcer (532), peptic ulcer bleeding (531.0, 531.2, 531.4, 531.6, 532.0, 532.2, 532.4, 532.6, 533.0, 533.2, 533.4, 533.6, 534.0, 534.2, 534.4, and 534.6), cardiovascular (390–459), respiratory (460–519), oral cavity, salivary glands, and jaws (520–529), reflux esophagitis (530.1 $\times$ , 530.2, 530.81, 530.82, and 530.85), skin and subcutaneous tissue (680–709), and poisoning and injuries (800–999).

association between PUD hospitalizations and  $\text{PM}_{2.5}$  exposure, there was no correlation between PUD bleeding and  $\text{PM}_{2.5}$  exposure. The exact reason for this finding remains uncertain. Although bleeding is only one of the complications related to PUD, not all hospitalization related to PUD are due to bleeding. Intuitively, it may be necessary to have a larger sample size to demonstrate the potential effect of  $\text{PM}_{2.5}$  on ulcer bleeding. Also, we did not consider the effects of other pollutants such as sulfur dioxides, nitrogen oxides, or carbon monoxide since we utilized a unique spatial method in measuring PM exposure, which could not be extended to other pollutants. Personal exposure measurement based on personal monitor is needed instead of one based on

monitoring stations. We were also not able to adjust for other environmental factors including noise and meteorological conditions which vary among geographical areas.

We have demonstrated for the 1st time an association between long-term  $\text{PM}_{2.5}$  exposure and hospitalization for PUD in elderly people. Further studies in different patient cohorts and regions are needed to clarify the potential role of  $\text{PM}_{2.5}$  on the development of PUD.

#### ACKNOWLEDGMENTS

The authors thank Mr Darren SZ Sun for the help on the literature search and the following Hong Kong SAR

**TABLE 4.** Hazard Ratios (95% CI) of PUD Hospitalization per 10  $\mu\text{g}/\text{m}^3$  of  $\text{PM}_{2.5}$  After Controlling for Preexisting Comorbidities

<b>(a) Covariate Adjustment</b>						
Hospitalization Causes	n	Covariate Adjustment I	P Value	Covariate Adjustment II	P Value	
Peptic ulcer	1991	1.18 (1.02–1.35)	.025	1.17 (1.01–1.35)	0.031	
Gastric ulcer	1175	1.29 (1.09–1.53)	.004	1.28 (1.08–1.52)	0.006	
Duodenal ulcer	816	0.97 (0.77–1.21)	.80	0.97 (0.77–1.20)	0.80	
<b>(b) Stratification</b>						
Hospitalization Causes	n	Stratification I	P Value	n	Stratification II	P Value
Peptic ulcer	1744	1.16 (1.00–1.36)	0.054	1808	1.15 (0.99–1.34)	0.062
Gastric ulcer	1015	1.29 (1.08–1.55)	0.007	1087	1.28 (1.07–1.54)	0.006
Duodenal ulcer	729	0.95 (0.74–1.21)	0.65	721	0.92 (0.73–1.15)	0.45

“Covariate adjustment I” adjusted for presence of hospitalization due to IHD, stroke, COPD or Type 2 diabetes at baseline. “Covariate adjustment II” adjusted for self-reported heart diseases, stroke, COPD, and diabetes as active diseases at baseline. “Stratification I” removed subjects with hospitalization of one of these four preexisting diseases during the baseline years. “Stratification II” removed subjects with hospitalization of one of these peptic ulcer diseases during the baseline years. CI = confidence interval, COPD = chronic obstructive pulmonary disease, ICD = International Statistical Classification of Diseases, IHD = ischemic heart disease,  $\text{PM}_{2.5}$  = particles less than 2.5  $\mu\text{m}$  in aerodynamic diameter, PUD = peptic ulcer disease.

Government departments including the Department of Health (Elderly Health Services) for cohort data and mortality data, and the Environmental Protection Department for the PM data. The authors also thank to the Hong Kong University of Science and Technology for satellite information and the Department of Geography of University of Hong Kong for geocoding information.

## REFERENCES

- HEI International Scientific Oversight Committee. Outdoor Air Pollution and Health in the Developing Countries of Asia: A Comprehensive Review Boston, MA: Health Effects Institute; 2010: Special Report 18.
- WHO Public Health Social Environmental Determinants of Health Department. Burden of Disease From Ambient Air Pollution for 2012. Geneva, Switzerland: World Health Organization; 2014.
- Schwartz J, Austin E, Bind MA, et al. Estimating causal associations of fine particles with daily deaths in Boston. *Am J Epidemiol*. 2015;182:644–650.
- WHO. Health effects of transport-related air pollution. 2005. Krzyzanowski M, Kuna-Dibbert B, Schneider J, Eds. Copenhagen, World Health Organization Regional Office for Europe. Available: [http://www.euro.who.int/\\_\\_data/assets/pdf\\_file/0006/74715/E86650.pdf](http://www.euro.who.int/__data/assets/pdf_file/0006/74715/E86650.pdf). [Accessed: 17 December, 2014].
- Anderson JO, Thundiyil JG, Stolbach A. Clearing the air: a review of the effects of particulate matter air pollution on human health. *J Med Toxicol*. 2012;8:166–175.
- Lelieveld J, Evans JS, Fnais M, et al. The contribution of outdoor air pollution sources to premature mortality on a global scale. *Nature*. 2015;525:367–371.
- Atkinson RW, Carey IM, Kent AJ, et al. Long-term exposure to outdoor air pollution and incidence of cardiovascular diseases. *Epidemiology*. 2013;24:44–53.
- Jerrett M, Burnett RT, Ma R, et al. Spatial analysis of air pollution and mortality in Los Angeles. *Epidemiology*. 2005;16:727–736.
- Kloog I, Coull BA, Zanutti A, et al. Acute and chronic effects of particles on hospital admissions in New-England. *PLoS One*. 2012;7:e34664.
- Wong CM, Lai HK, Tsang H, et al. Satellite-based estimates of long-term exposure to fine particles and association with mortality in elderly Hong Kong residents. *Environ Health Perspect*. 2015;123:1167–1172.
- Vardoulakis S, Fisher BEA, Pericleous K, et al. Modelling air quality in street canyons: a review. *Atmos Environ*. 2003;37:155–182.
- Hospital Authority (HA). Hospital Authority Statistical Report 2011–2012. Table 1.2. 2013. [http://www.ha.org.hk/upload/publication\\_15/471.pdf](http://www.ha.org.hk/upload/publication_15/471.pdf). [Accessed: 20 May, 2014].
- Chan HL, Wu JC, Chan FK, et al. Is non-Helicobacter pylori, non-NSAID peptic ulcer a common cause of upper GI bleeding? A prospective study of 977 patients. *Gastrointest Endosc*. 2001;53:438–442.
- Iijima K, Kanno T, Koike T, et al. Helicobacter pylori-negative, non-steroidal anti-inflammatory drug: negative idiopathic ulcers in Asia. *World J Gastroenterol*. 2014;20:706–713.
- Li CC, Lau AKH, Mao JT, et al. Retrieval, validation, and application of the 1-km aerosol optical depth from MODIS measurements over Hong Kong. *IEEE Trans Geosci Rem Sens*. 2005;43:2650–2658.
- Hospital Authority (HA). Hospital Authority Statistical Report 2010–2011. 2012. [http://www.ha.org.hk/upload/publication\\_15/411.pdf](http://www.ha.org.hk/upload/publication_15/411.pdf). [Accessed 27 June, 2014].
- Pisa F, Castellsague J, Drigo D, et al. Accuracy of International Classification of Diseases, 9th Revision, Clinical Modification codes for upper gastrointestinal complications varied by position and age: a validation study in a cohort of nonsteroidal anti-inflammatory drugs users in Friuli Venezia Giulia, Italy. *Pharmacoepidemiol Drug Saf*. 2013;22:1195–1204.
- Wong CM, Tsang H, Lai HK, et al. Cancer mortality risks from long-term exposure to ambient fine particle. *Cancer Epidemiol Biomarkers Prev*. in press. Accepted on 24 February 2016.
- Census and Statistics Department. Hong Kong 2004 population and household statistics analysed by district council district. 2005. Available: <http://www.statistics.gov.hk/pub/B11303012004AN04B0100.pdf>. [Accessed 26 September, 2014].
- Census and Statistics Department. Thematic household survey Report No. 48. 2011. Available: [http://smokefree.hk/UserFiles/resources/Statistics/Thematic\\_Household\\_Survey\\_No48.pdf](http://smokefree.hk/UserFiles/resources/Statistics/Thematic_Household_Survey_No48.pdf). [Accessed 26 September, 2014.]
- Balluz L, Wen XJ, Town M, et al. Ischemic heart disease and ambient air pollution of particulate matter 2.5 in 51 counties in the U.S. *Public Health Rep*. 2007;122:626–633.
- Brook RD, Cakmak S, Turner MC, et al. Long-term fine particulate matter exposure and mortality from diabetes in Canada. *Diabetes Care*. 2013;36:3313–3320.
- Leiva GMA, Santibañez DA, Ibarra ES, et al. A five-year study of particulate matter (PM<sub>2.5</sub>) and cerebrovascular diseases. *Environ Pollut*. 2013;181:1–6.
- Tsai SS, Chang CC, Yang CY. Fine particulate air pollution and hospital admissions for chronic obstructive pulmonary disease: a case-crossover study in Taipei. *Int J Environ Res Public Health*. 2013;10:6015–6026.
- Lim SS, Vos T, Flaxman AD, et al. A comparative risk assessment of burden of disease and injury attributable to 67 risk factors and risk factor clusters in 21 regions, 1990–2010: a systematic analysis for the Global Burden of Disease Study 2010. *Lancet*. 2012;380:2224–2260.
- Fares A. Global patterns of seasonal variation in gastrointestinal diseases. *J Postgrad Med*. 2013;59:203–207.
- Lam TH, Li ZB, Ho SY, et al. Smoking, quitting and mortality in an elderly cohort of 56,000 Hong Kong Chinese. *Tob Control*. 2007;16:182–189.
- Ritz B, Wilhelm M, Hoggatt KJ, et al. Ambient air pollution and preterm birth in the environment and pregnancy outcomes study at the University of California, Los Angeles. *Am J Epidemiol*. 2007;166:1045–1052.
- Kish L, Hotte N, Kaplan GG, et al. Environmental particulate matter induces murine intestinal inflammatory responses and alters the gut microbiome. *PLoS One*. 2013;8:e62220.
- Holick MF, Chen TC. Vitamin D deficiency: a worldwide problem with health consequences. *Am J Clin Nutr*. 2008;87:1080S–1086S.
- Möller W, Hussinger K, Winkler-Heil R, et al. Mucociliary and long-term particle clearance in the airways of healthy nonsmoker subjects. *J Appl Physiol*. 2004;97:2200–2206.
- Beamish LA, Osornio-Vargas AR, Wine E. Air pollution: an environmental factor contributing to intestinal disease. *J Crohns Colitis*. 2011;5:279–286.
- De Brouwere K, Buekers J, Cornelis C, et al. Assessment of indirect human exposure to environmental sources of nickel: oral exposure and risk characterization for systemic effects. *Sci Total Environ*. 2012;419:25–36.
- Arrieta MC, Bistriz L, Meddings JB. Alteration in intestinal permeability. *Gut*. 2006;55:1512–1520.

35. Kaplan GG, Szyszkowicz M, Fichna J, et al. Non-specific abdominal pain and air pollution: a novel association. *PLoS One*. 2012;7:e47669.
36. Van de Wiele T, Vanhaecke L, Boeckart C, et al. Human colon microbiota transform polycyclic aromatic hydrocarbons to estrogenic metabolites. *Environ Health Perspect*. 2005;113:6–10.
37. Salim SY, Kaplan GG, Madsen KL. Air pollution effects on the gut microbiota: a link between exposure and inflammatory disease. *Gut Microbes*. 2014;5:215–219.
38. Kaplan GG, Hubbard J, Korzenik J, et al. The inflammatory bowel diseases and ambient air pollution: a novel association. *Am J Gastroenterol*. 2010;105:2412–2419.
39. Ananthakrishnan AN, McGinley EL, Binion DG, et al. Ambient air pollution correlates with hospitalizations for inflammatory bowel disease: an ecologic analysis. *Inflamm Bowel Dis*. 2011;17:1138–1145.
40. Knox EG. Atmospheric pollutants and mortalities in English local authority areas. *J Epidemiol Community Health*. 2008;62:442–447.
41. Quan S, Yang H, Tanyingoh D, et al. Upper gastrointestinal bleeding due to peptic ulcer disease is not associated with air pollution: a case-crossover study. *BMC Gastroenterol*. 2015;15:131.
42. Levenstein S, Ackerman S, Kiecolt-Glaser JK, et al. Stress and peptic ulcer disease. *JAMA*. 1999;281:10–11.
43. Kanno T, Iijima K, Abe Y, et al. Peptic ulcers after the Great East Japan earthquake and tsunami: possible existence of psychosocial stress ulcers in humans. *J Gastroenterol*. 2013;48:483–490.
44. Chan FK, Leung WK. Peptic-ulcer disease. *Lancet*. 2002;360:933–941.
45. Laine L, Takeuchi K, Tarnawski A. Gastric mucosal defense and cytoprotection: bench to bedside. *Gastroenterology*. 2008;135:41–60.
46. Hung LC, Ching JY, Sung JJ, et al. Long-term outcome of *Helicobacter pylori*-negative idiopathic bleeding ulcers: a prospective cohort study. *Gastroenterology*. 2005;128:1845–1850.
47. Pope CA 3rd, Dockery DW. Health effects of fine particulate air pollution: lines that connect. *J Air Waste Manag Assoc*. 2006;56:709–742.