

Expression of eukaryotic translation initiation factor 3 subunit B in liver cancer and its prognostic significance

QING YUE^{1*}, LINGYU MENG^{2*}, BAOXING JIA² and WEI HAN²

Departments of ¹Oncology and ²Hepatobiliary and Pancreatic Surgery,
The First Hospital of Jilin University, Changchun, Jilin 130021, P.R. China

Received July 16, 2019; Accepted December 19, 2019

DOI: 10.3892/etm.2020.8726

Abstract. Liver cancer is one of the major malignancies with the worst prognosis among all solid tumor types. It is therefore ponderable to explore prognostic biomarkers and therapeutic targets for liver cancer. Eukaryotic translation initiation factor 3 subunit B (EIF3B) is closely linked to the transcription initiation of cancer-associated genes. In the present study, EIF3B was indicated to be a potential prognostic biomarker of liver cancer. The mRNA expression level of EIF3B in liver cancer was assessed by analyzing the Cancer Genome Atlas dataset. χ^2 and Fisher's exact tests were used to assess the association of EIF3B expression with clinical parameters. Receiver-operating characteristic curve analysis was used for evaluating the diagnostic value of EIF3B. Overall and relapse-free survival were assessed using Kaplan-Meier curves to determine the association between EIF3B expression and survival. Univariate and multivariate Cox regression analysis were performed to identify the factors affecting overall/relapse-free survival. Gene set enrichment analysis (GSEA) was used to identify signaling pathways associated with EIF3B in liver cancer. It was revealed that EIF3B was highly expressed in liver cancer tissues and it had a promising diagnostic ability. Furthermore, the survival analysis indicated that patients with high EIF3B expression generally had shorter overall as well as relapse-free survival. Univariate and multivariate Cox analysis suggested that high EIF3B mRNA expression may serve as an independent biomarker for the prognostication of patients with liver cancer. GSEA suggested that MYC-V1 (HALLMARK_MYC_TARGETS_V1 geneset; P=0.009), MYC-V2 (HALLMARK_MYC_TARGETS_V2 geneset; P=0.004) and DNA repair pathways (HALLMARK_DNA_REPAIR geneset; P<0.001) were differentially enriched

in high EIF3B expression and low EIF3B expression groups. In conclusion, high EIF3B expression was indicated to be an independent prognostic biomarker for patients with liver cancer.

Introduction

Liver cancer is a common malignant tumor type with high morbidity and mortality (1). Although various treatments have been improved, the mortality rate of liver cancer is still high and the prognosis remains poor (2,3). Therefore, prognostic biomarkers of liver cancer have become one of the hotspots of current research (4). The discovery of accurate prognostic biomarkers may contribute to clinical guidance in order to improve the evaluation system of liver cancer.

The family of eukaryotic translation initiation factors (EIFs) participates in eukaryotic translation by regulating the interaction between ribosomes and RNA. It is the rate-limiting step of protein synthesis and participates in numerous processes that are deregulated in cancer cells, including DNA repair and proliferation, cell cycle and apoptosis (5). EIF 3 subunit B (EIF3B) is an important member of the family of EIFs and has been observed to be overexpressed in numerous cancer types, including clear cell renal cell carcinoma (6), esophageal squamous cell carcinoma (7), glioblastoma (8), ovarian cancer (9), osteosarcoma (10) and lung cancer (11), and has an important role in the progression and prognosis of several cancer types (12-14). In addition, Golob-Schwarzl *et al* (15) reported that EIF3B was upregulated in hepatitis C virus (HCV)-associated hepatocellular carcinoma (HCC). However, the precise role of EIF3B in liver cancer has remained elusive.

To further evaluate the roles of EIF3B in patients with liver cancer, the expression of EIF3B was examined in a dataset from The Cancer Genome Atlas (TCGA) database. The χ^2 and Fisher's exact tests were used to assess the association of EIF3B with clinicopathological parameters and demographic features. Receiver operating characteristics (ROC) curve analysis was used for evaluating the diagnostic value of EIF3B. Kaplan-Meier overall survival and relapse-free survival analysis were performed to determine the association between EIF3B expression and survival. Univariate and multivariate Cox regression analysis were performed to identify the factors affecting overall survival and relapse-free survival. Furthermore, gene set enrichment analysis (GSEA) was used to explore EIF3B-associated signaling pathways.

Correspondence to: Dr Wei Han, Department of Hepatobiliary and Pancreatic Surgery, The First Hospital of Jilin University, 71 Xinmin Street, Changchun, Jilin 130021, P.R. China
E-mail: weihan_1989@126.com

*Contributed equally

Key words: eukaryotic translation initiation factor 3 subunit B, liver cancer, prognosis, The Cancer Genome Atlas, biomarker

Table I. Demographic and clinical characteristics of the cohort from The Cancer Genome Atlas-liver hepatocellular carcinoma dataset.

Characteristics	N (373)
Age (years)	
<55	117 (31.45)
≥55	255 (68.55)
NA	1 (0.00)
Sex	
Female	121 (32.44)
Male	252 (67.56)
Histological type	
Fibrolamellar carcinoma	3 (0.80)
Hepatocellular carcinoma	363 (97.32)
Hepatocholangiocarcinoma (mixed)	7 (1.88)
Histologic grade	
NA	5 (1.34)
G1	55 (14.75)
G2	178 (47.72)
G3	123 (32.98)
G4	12 (3.22)
Stage	
NA	24 (6.43)
I	172 (46.11)
II	87 (23.32)
III	85 (22.79)
IV	5 (1.34)
T classification	
NA	2 (0.54)
T1	182 (48.79)
T2	95 (25.47)
T3	80 (21.45)
T4	13 (3.49)
TX	1 (0.27)
N classification	
NA	1 (0.27)
N0	253 (67.83)
N1	4 (1.07)
NX	115 (30.83)
M classification	
M0	267 (71.58)
M1	4 (1.07)
MX	102 (27.35)
Radiation therapy	
NA	25 (6.70)
No	340 (91.15)
Yes	8 (2.14)
Residual tumor	
NA	7 (1.88)
R0	326 (87.40)
R1	17 (4.56)
R2	1 (0.27)
RX	22 (5.90)

Table I. Continued.

Characteristics	N (373)
Vital status	
Deceased	130 (34.85)
Alive	243 (65.15)
Sample type	
Primary tumor	371 (99.46)
Recurrent tumor	2 (0.54)
Overall survival (ten years)	
No	237 (64.58)
Yes	130 (35.42)
Relapse-free survival (ten years)	
No	179 (55.94)
Yes	141 (44.06)
EIF3B	
High	109 (29.22)
Low	264 (70.78)

NA, not available; EIF3B, eukaryotic translation initiation factor 3 subunit B; G1-4, grade relating to degree of differentiation; T1-4, size and or extension of the primary tumor; TX, tumor could not be assessed; N0, no regional lymph node metastasis; N1, regional lymph node metastasis present; NX, lymph nodes could not be assessed; M0, no distant metastasis; M1, metastasis to distant organs; MX, metastasis could not be assessed; R0, no residual tumor visible under the microscope; R1, residual tumor visible under the microscope; R2, residual tumor visible to the naked eye; RX, residual tumor could not be assessed.

Materials and methods

Data source. The clinical information of patients and their RNAseq data were obtained from TCGA (<https://cancergenome.nih.gov/>). All patients from the liver hepatocellular carcinoma (LIHC) cohort were screened based on TCGA inclusion and exclusion pre-selection criteria.

Statistical analysis. R (version 3.6.1; The R Foundation) (16) was used for statistical analysis (t-test, Kruskal-Wallis with Dunn's post-hoc test, Wilcoxon sum-rank test) and generation of images. The ggplot2 package (17) was used to draw boxplots of the EIF3B expression in subgroups by clinical characteristics. χ^2 and Fisher's exact tests were applied to estimate the significance of the association between EIF3B expression and clinicopathological or demographic characteristics. The pROC package (18) was used to plot the ROC curves and assess the diagnostic ability of EIF3B, and patients were divided into a high expression group and low expression group according to the best operating system cut-off value determined by the Youden index. The survival package (19) was used to draw survival curves. A univariate Cox linear regression model was utilized to select correlative variables affecting survival time. Multivariate Cox regression analysis was employed to evaluate the independent influencing factors of survival time.




Figure 1. Differences in EIF3B expression among subgroups of patients. Boxplots showing differences in EIF3B expression according to (A) tissue type ($P<0.001$ vs. normal tissue controls), (B) age, (C) sex, (D) histologic grade ($P<0.001$), (E) histological type, (F) T classification, (G) N classification, (H) M classification, (I) radiation therapy, (J) residual tumor classification, (K) sample type, (L) clinical stage and (M) vital status ($P=0.005$). EIF3B, eukaryotic translation initiation factor 3 subunit B. G1-4, grade relating to degree of differentiation; T1-4, size and or extension of the primary tumor; TX, tumor could not be assessed; N0, no regional lymph node metastasis; N1, regional lymph node metastasis present; NX, lymph nodes could not be assessed; M0, no distant metastasis; M1, metastasis to distant organs; MX, metastasis could not be assessed; R0, no residual tumor visible under the microscope; R1, residual tumor visible under the microscope; R2, residual tumor visible to the naked eye; RX, residual tumor could not be assessed.

GSEA. GSEA may be used to determine whether a predefined set of genes is able to indicate significant, consistent differences between two biological states (20). In the present study, GSEA was performed in the 'h.all.v6.2.symbols.gmt' and 'c2_cp.biocarta.v6.2.symbols.gmt' gene sets using GSEA3.0 software. The standardized enrichment fraction was obtained by 1,000 permutation analyses.

Results

Patient characteristics. In Table I, the clinical features of the 373 patient cohort, including sex (female 121, male 252), age (16-90 years old, median 61, mean 59.47), use of radiation

therapy, residual tumor, relapse-free survival, histological type, stage, vital status, survival data, T/N/M classification and EIF3B expression are provided.

Expression of EIF3B in liver tissues. Boxplots revealed that EIF3B was significantly upregulated in liver cancer compared with that in normal liver tissues (Fig. 1A; $P<0.001$). Furthermore, EIF3B was also differentially expressed between subgroups by vital status ($P=0.005$) and histologic grade ($P<0.001$; Fig. 1).

Diagnostic capability of EIF3B. To assess the diagnostic performance of EIF3B in liver cancer, ROC curve analysis was used. The area under the curve (AUC) was 0.821, indicating

Table II. Association between the expression of EIF3B and the clinicopathological characteristics of patients with liver cancer.

Clinical characteristics	No. of patients	EIF3B expression		χ^2	P-value
		High	Low		
Age (years)				0.089	0.765
<55	117	36 (33.03)	81 (30.80)		
≥55	255	73 (66.97)	182 (69.20)		
Sex				0.484	0.486
Female	121	32 (29.36)	89 (33.71)		
Male	252	77 (70.64)	175 (66.29)		
Histological type				1.251	0.534
Fibrolamellar carcinoma	3	0 (0.00)	3 (1.14)		
Hepatocellular carcinoma	363	107 (98.17)	256 (96.97)		
Hepatocholangiocarcinoma (mixed)	7	2 (1.83)	5 (1.89)		
Histologic grade				17.796	<0.001
G1	55	9 (8.33)	46 (17.69)		
G2	178	45 (41.67)	133 (51.15)		
G3	123	46 (42.59)	77 (29.62)		
G4	12	8 (7.41)	4 (1.54)		
Stage				4.532	0.209
I	172	43 (40.95)	129 (52.87)		
II	87	32 (30.48)	55 (22.54)		
III	85	28 (26.67)	57 (23.36)		
IV	5	2 (1.9)	3 (1.23)		
T classification				7.720	0.102
T1	182	44 (40.37)	138 (52.67)		
T2	95	34 (31.19)	61 (23.28)		
T3	80	29 (26.61)	51 (19.47)		
T4	13	2 (1.83)	11 (4.2)		
TX	1	0 (0.00)	1 (0.38)		
N classification				1.936	0.379
N0	253	77 (70.64)	176 (66.92)		
N1	4	0 (0.00)	4 (1.52)		
NX	115	32 (29.36)	83 (31.56)		
M classification				2.882	0.236
M0	267	83 (76.15)	184 (69.70)		
M1	4	2 (1.83)	2 (0.76)		
MX	102	24 (22.02)	78 (29.55)		
Radiation therapy				<0.001	>0.999
No	340	92 (97.87)	248 (97.64)		
Yes	8	2 (2.13)	6 (2.36)		
Residual tumor				5.116	0.163
R0	326	98 (91.59)	228 (88.03)		
R1	17	5 (4.67)	12 (4.63)		
R2	1	1 (0.93)	0 (0.00)		
RX	22	3 (2.80)	19 (7.34)		
Vital status				17.505	<0.001
Deceased	130	56 (51.38)	74 (28.03)		
Alive	243	53 (48.62)	190 (71.97)		
Sample type				0.017	0.895
Primary tumor	371	109 (100)	262 (99.24)		
Recurrent tumor	2	0 (0.00)	2 (0.76)		

Table II. Continued.

Clinical characteristics	No. of patients	EIF3B expression		χ^2	P-value
		High	Low		
Overall survival (ten years)				18.690	<0.001
No	237	50 (47.17)	187 (71.65)		
Yes	130	56 (52.83)	74 (28.35)		
Relapse-free survival (ten years)				1.018	0.312
No	179	42 (50.6)	137 (57.81)		
Yes	141	41 (49.4)	100 (42.19)		

EIF3B, eukaryotic translation initiation factor 3 subunit B; G1-4, grade relating to degree of differentiation; T1-4, size and or extension of the primary tumor; TX, tumor could not be assessed; N0, no regional lymph node metastasis; N1, regional lymph node metastasis present; NX, lymph nodes could not be assessed; M0, no distant metastasis; M1, metastasis to distant organs; MX, metastasis could not be assessed; R0, no residual tumor visible under the microscope; R1, residual tumor visible under the microscope; R2, residual tumor visible to the naked eye; RX, residual tumor could not be assessed.




Figure 2. Receiver-operating characteristic curves for EIF3B in The Cancer Genome Atlas-liver hepatocellular carcinoma dataset. The ability of EIF3B to distinguish between the following was assessed: (A) Non-tumor vs. tumor sample; non-tumor sample vs. tumor sample of (B) stage I, (C) stage II, (D) stage III and (E) stage IV. AUC, area under the curve; EIF3B, eukaryotic translation initiation factor 3 subunit B.

that EIF3B has moderate diagnostic ability. In addition, similar AUCs were obtained for distinguishing normal liver tissues from liver cancer at specific stages (stage I, 0.785; stage II, 0.850; stage III, 0.853; stage IV, 0.916; Fig. 2).

Association between EIF3B expression and clinical features of patients with liver cancer. As indicated in Table II, the

vital status of the patients with liver cancer ($P<0.001$), overall survival ($P<0.001$; duration ten years) and the histologic grade ($P<0.001$) were associated with the expression of EIF3B.

High expression of EIF3B is associated with poor overall survival of patients with liver cancer. Kaplan-Meier analysis indicated that high expression of EIF3B was significantly




Figure 3. Kaplan-Meier analysis of the influence of EIF3B expression on overall survival. (A) All patients. Subgroup analysis for (B) females, (C) males, (D) younger patients (<55), (E) older patients (≥55), (F) no lymph node dissection (N0), (G) lymph node dissection (R1/R2/RX), (H-M) histological grade, (H) G1/G2, (I) G3/G4, (J) G1, (K) G2, (L) G3, (M) G4, (N-Q) clinical stage (N) I, (O) II, (P) III, (Q) IV and (R) HCC. EIF3B, eukaryotic translation initiation factor 3 subunit B; HCC, hepatocellular carcinoma; G1-4, grade relating to degree of differentiation; T1-4, size and/or extension of the primary tumor; TX, tumor could not be assessed; N0, no regional lymph node metastasis; N1, regional lymph node metastasis present; NX, lymph nodes could not be assessed; M0, no distant metastasis; M1, metastasis to distant organs; MX, metastasis could not be assessed; R0, no residual tumor visible under the microscope; R1, residual tumor visible under the microscope; R2, residual tumor visible to the naked eye; RX, residual tumor could not be assessed. The number of results censored (removed from surviving) is indicated below the survival curve.

associated with poor overall survival ($P < 0.001$; Fig. 3). Subgroup analysis provided similar results, particularly in female ($P < 0.001$), younger (<55; $P = 0.008$) and older

subjects (≥55; $P < 0.001$), and in patients with R0 ($P < 0.001$), G2 ($P = 0.003$), G3 ($P < 0.0001$), stage I ($P < 0.001$) and stage III ($P < 0.001$).




Figure 4. Kaplan-Meier analysis of the influence of EIF3B expression on relapse-free survival. (A) All patients. (B-R) Subgroup analysis for (B) females, (C) males, (D) younger patients (<55), (E) older patients (≥ 55), (F) no lymph node dissection (R0), (G) lymph node dissection (R1/R2/RX), (H-K) histological grade (H) G1, (I) G2, (J) G3, (K) G4, (L-N) clinical stage (L) I, (M) II, (N) III and (O) HCC. EIF3B, eukaryotic translation initiation factor 3 subunit B; HCC, hepatocellular carcinoma; G1-4, grade relating to degree of differentiation; T1-4, size and/or extension of the primary tumor; TX, tumor could not be assessed; N0, no regional lymph node metastasis; N1, regional lymph node metastasis present; NX, lymph nodes could not be assessed; M0, no distant metastasis; M1, metastasis to distant organs; MX, metastasis could not be assessed; R0, no residual tumor visible under the microscope; R1, residual tumor visible under the microscope; R2, residual tumor visible to the naked eye; RX, residual tumor could not be assessed. The number of results censored (removed from surviving) is indicated below the survival curve.

As presented in Table III, T classification, stage, residual tumor and EIF3B expression were variables associated with overall survival according to univariate Cox regression analysis. In addition, multivariate Cox regression indicated that high EIF3B expression, T classification and residual tumor were independent risk factors for overall survival of patients with liver cancer [hazard ratio (HR)=2.44, 95% CI=1.71-3.47, $P < 0.001$].

High expression of EIF3B is associated with poor relapse-free survival of patients with liver cancer. Kaplan-Meier analysis indicated that patients with high expression of EIF3B had significantly poorer relapse-free survival ($P = 0.013$; Fig. 4). Subgroup analysis provided similar results, particularly in female ($P = 0.001$) and older (≥ 55 ; $P = 0.032$) subjects and in patients with R0 ($P = 0.020$), G2 ($P < 0.001$) and stage III ($P = 0.004$).

Table III. Summary of univariate and multivariate Cox regression analyses for overall survival duration (ten years).

Parameters	Univariate analysis			Multivariate analysis		
	Hazard ratio	95% CI	P-value	Hazard ratio	95% CI	P-value
Age (≥ 55 / < 55 years)	1.00	0.69-1.45	0.997			
Sex (male/female)	0.80	0.56-1.14	0.220			
Histological type (hepatocarcinoma/ hepatocellular, hepatocellular /fibrolamellar)	0.99	0.27-3.66	0.986			
Histologic grade (G4/G3/G2/G1)	1.04	0.84-1.30	0.698			
Stage (IV/III/II/I)	1.38	1.15-1.66	0.001	0.81	0.65-1.01	0.060
T classification (T4/T3/T2/T1/NX)	1.66	1.39-1.99	<0.001	1.91	1.51-2.42	<0.001
N classification (N1/N0/NX)	0.73	0.51-1.05	0.086			
M classification (M1/M0/MX)	0.72	0.49-1.04	0.077			
Radiation therapy (yes/no)	0.51	0.26-1.03	0.060			
Residual tumor classification (RX/R2/R1/R0)	1.42	1.13-1.80	0.003	1.45	1.13-1.87	0.004
EIF3B (high/low)	2.41	1.70-3.42	<0.001	2.44	1.71-3.47	<0.001

EIF3B, eukaryotic translation initiation factor 3 subunit B; G1-4, grade relating to degree of differentiation; T1-4, size and or extension of the primary tumor; TX, tumor could not be assessed; N0, no regional lymph node metastasis; N1, regional lymph node metastasis present; NX, lymph nodes could not be assessed; M0, no distant metastasis; M1, metastasis to distant organs; MX, metastasis could not be assessed; R0, no residual tumor visible under the microscope; R1, residual tumor visible under the microscope; R2, residual tumor visible to the naked eye; RX, residual tumor could not be assessed.

Table IV. Summary of univariate and multivariate Cox regression analyses or relapse-free survival duration.

Parameters	Univariate analysis			Multivariate analysis		
	Hazard ratio	95% CI	P-value	Hazard ratio	95% CI	P-value
Age (≥ 55 / < 55 years)	0.90	0.63-1.28	0.550			
Sex (male/female)	0.99	0.70-1.41	0.966			
Histological type (hepatocarcinoma/ hepatocellular, hepatocellular /fibrolamellar)	2.02	0.66-6.24	0.220			
Histologic grade (G4/G3/G2/G1)	0.98	0.80-1.21	0.883			
Stage (IV/III/II/I)	1.66	1.38-1.99	<0.001	1.10	0.85-1.42	0.473
T classification (T4/T3/T2/T1/TX)	1.78	1.49-2.12	<0.001	1.67	1.28-2.18	<0.001
N classification (N1/N0/NX)	0.97	0.67-1.40	0.874			
M classification (M1/M0/MX)	1.17	0.79-1.74	0.432			
Radiation therapy (yes/no)	0.74	0.26-2.16	0.584			
Residual tumor classification (RX/R2/R1/R0)	1.28	1.01-1.61	0.042	1.36	1.07-1.73	0.012
EIF3B (high/low)	1.58	1.10-2.28	0.014	1.54	1.06-2.23	0.022

EIF3B, eukaryotic translation initiation factor 3 subunit B; G1-4, grade relating to degree of differentiation; T1-4, size and/or extension of the primary tumor; TX, tumor could not be assessed; N0, no regional lymph node metastasis; N1, regional lymph node metastasis present; NX, lymph nodes could not be assessed; M0, no distant metastasis; M1, metastasis to distant organs; MX, metastasis could not be assessed; R0, no residual tumor visible under the microscope; R1, residual tumor visible under the microscope; R2, residual tumor visible to the naked eye; RX, residual tumor could not be assessed.

As presented in Table IV, T classification, stage, residual tumor and EIF3B expression were variables associated with relapse-free survival according to the univariate Cox regression analysis. In addition, high EIF3B expression, T classification and residual tumor were independent risk factors for relapse-free survival of patients with liver cancer

in the multivariate Cox regression analysis (HR=1.54, 95% CI=1.06-2.23, P=0.022).

Signaling pathways associated with EIF3B. To identify the signaling pathways associated with EIF3B in liver cancer, GSEA was performed between the low EIF3B expression

Table V. Gene sets enriched in phenotype high.

Molecular signatures database collection	Gene set name	NES	NOM P-value	FDR q-value
h.all.v6.2.symbols.gmt	HALLMARK_MYC_TARGETS_V2	2.154	0.004	0.006
h.all.v6.2.symbols.gmt	HALLMARK_MYC_TARGETS_V1	2.028	0.009	0.010
h.all.v6.2.symbols.gmt	HALLMARK_DNA_REPAIR	2.006	<0.001	0.009

Gene sets with NOM P-value <0.050 and FDR q-value <0.250 were considered as significant. FDR, false discovery rate; NES, normalized enrichment score; NOM, nominal.




Figure 5. Enrichment plots from GSEA. The GSEA results indicated that (A) the MYC-V1, (B) the MYC-V2 and (C) the DNA repair pathway are differentially enriched in high EIF3B expression and low EIF3B expression groups. GSEA, gene set enrichment analysis.

dataset and the high EIF3B expression dataset. The enrichment of the molecular signatures database (MSigDB) determined by GSEA was significantly different (nominal P-value <0.050, false discovery rate <0.250; Table V). As presented in Fig. 5 and Table V, GSEA indicated that MYC-V1 (HALLMARK_MYC_TARGETS_V1 geneset; P=0.009), MYC-V2 (HALLMARK_MYC_TARGETS_V2 geneset; P=0.004) and DNA repair pathways (HALLMARK_DNA_REPAIR geneset; P<0.001) were differentially enriched in high EIF3B expression and low EIF3B expression groups.

Discussion

Liver cancer is associated with a high mortality rate worldwide; the development of this cancer type may be influenced by viral infection, diet and environmental factors (21-24). In recent years, with the continuous progression of molecular biology and treatments, including chemotherapeutic drugs and surgical technology, the understanding of cancer biology and the treatment of liver cancer have made great progress. However, the prognosis of liver cancer remains poor. The World Health Organization/International Classification of Diseases-10 classifies diseases according to their etiology, pathology, clinical manifestations and anatomical location. Cancer is a gene-associated disease and molecular typing is required to deepen our understanding of the underlying mechanisms of disease development (25). Therefore, novel biomarkers are urgently required. Our research group has been exploring novel biomarkers for a

number of years (26-38). The present study focused on EIF3B and indicated that EIF3B is a potential and independent prognostic biomarker for liver cancer.

EIF3B is closely linked to cancer progression. Consistent with previous studies, it was indicated that EIF3B was highly expressed in patients with liver cancer. Although Golob-Schwarzl *et al* (15) reported that EIF3B was upregulated in HCV-associated HCC, all of their patients were Asians. The patients assessed in the present study were from all over the world and covered other types of liver cancer that may be related to HCV. All of these results indicate that EIF3B has an important role in cancer-associated processes. A previous study suggested that EIF3B is involved in the proliferation and metastasis of gastric cancer (39). In addition, the present study further determined that EIF3B is associated with the histologic grade and survival status of patients with liver cancer. Therefore, in-depth studies using experimental and bioinformatics methods are required.

EIF3B has been indicated to have a marked influence on the prognosis of patients with cancer. In the present study, it was observed that upregulation of EIF3B was associated with poor overall/relapse-free survival of patients with liver cancer. In addition, patients with high EIF3B expression in clear cell renal cell carcinoma, esophageal squamous cell carcinoma and non-small cell lung cancer had a shorter survival time (6,7,11). In order to further explore the association between EIF3B and clinical characteristics of patients with liver cancer, a subgroup analysis was performed. As the TCGA-LIHC dataset does not have a Barcelona Clinic Liver Cancer staging system, TNM

staging was used. Kaplan-Meier subgroup analysis indicated that high expression of EIF3B was associated with poor overall survival in the subgroups of females, younger (<55) or older (≥55) patients, R0, G2, G3, stage I and stage III. Furthermore, high expression of EIF3B was associated with poor relapse-free survival in the subgroups of females, older patients, R0, G2 and stage III. However, Tian *et al* (11) reported that upregulation of EIF3B was associated with tumor depth, TNM stage and lymph node metastasis in patients with esophageal squamous cell carcinoma. However, the results of the present study indicated that EIF3B was related to histological grade and survival status. This may be linked to the heterogeneity of tumor types and individual differences, which may help to select personalized treatments. Owing to the TCGA-LIHC data not including the body mass index and the presence of diabetes mellitus as variables, it is not possible to calculate their association with the prognosis of patients.

In the GSEA analysis, high EIF3B expression was indicated to be associated with MYC-V1, MYC-V2 and DNA repair in liver cancer. The MYC oncogene is an important regulator of liver cancer progression. Previous studies have indicated that MYC is able to promote the proliferation, metastasis and metabolism of liver cancer by regulating signaling pathways including AKT/mTOR and RAS/mitogen-activated protein kinase (40-42). In addition, each replication of DNA in cancer cells may cause a large amount of damage, including DNA substitutions or deletions (43). Therefore, DNA repair mechanisms (damage induction, signal transduction, signal response) are particularly important. This may explain why EIF3B may promote the progression of liver cancer through MYC-V1/V2 and DNA repair pathways.

The present study mainly uncovered the prognostic value of the EIF3B mRNA expression in liver cancer. Along with other studies on EIF3B, the present study contributed to a better understanding of the role of EIF3B, as well as the great possibility for precise prognostication. However, the underlying mechanisms remain to be fully elucidated and require further exploration by scientific research. In the future, the mechanisms of EIF3B will be studied at a deeper level.

In conclusion, the present study investigated the prognostic value of EIF3B in patients with liver cancer. High EIF3B expression was proved to be a potential and independent prognostic biomarker for liver cancer. Future work will include *in vivo* and *in vitro* experiments to explore the biological functions of EIF3B and the underlying mechanisms.

Acknowledgements

Not applicable.

Funding

No funding was received.

Availability of data and materials

Patient information was obtained from the open TCGA database (<https://portal.gdc.cancer.gov/>). The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Authors' contributions

WH designed the study. QY and LM were responsible for extracting data, conducted data analysis and wrote the first draft of the manuscript. BJ participated in the analysis of data and critical modification of important knowledge content. WH and BJ critically revised the manuscript and gave final approval for submission. All authors read and approved the final manuscript.

Ethics approval and consent to participate

Not applicable.

Patient consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

References

1. Siegel RL, Miller KD and Jemal A: Cancer statistics, 2019. *CA Cancer J Clin* 69: 7-34, 2019.
2. Ryerson AB, Ehemann CR, Altekruse SF, Ward JW, Jemal A, Sherman RL, Henley SJ, Holtzman D, Lake A, Noone AM, *et al*: Annual Report to the Nation on the Status of Cancer, 1975-2012, Featuring the Increasing Incidence of Liver Cancer. *Cancer* 122: 1312-1337, 2016.
3. Li L and Wang H: Heterogeneity of liver cancer and personalized therapy. *Cancer Lett* 379: 191-197, 2016.
4. Dawkins J and Webster RM: The hepatocellular carcinoma market. *Nat Rev Drug Discov* 18: 13-14, 2019.
5. Smith MD, Arake-Tacca L, Nitido A, Montabana E, Park A and Cate JH: Assembly of eIF3 mediated by mutually dependent subunit insertion. *Structure* 24: 886-896, 2016.
6. Zang Y, Zhang X, Yan L, Gu G, Li D, Zhang Y, Fang L, Fu S, Ren J and Xu Z: Eukaryotic translation initiation factor 3b is both a promising prognostic biomarker and a potential therapeutic target for patients with clear cell renal cell carcinoma. *J Cancer* 8: 3049-3061, 2017.
7. Xu F, Xu CZ, Gu J, Liu X, Liu R, Huang E, Yuan Y, Zhao G, Jiang J, Xu C, *et al*: Eukaryotic translation initiation factor 3B accelerates the progression of esophageal squamous cell carcinoma by activating β -catenin signaling pathway. *Oncotarget* 7: 43401-43411, 2016.
8. Liang H, Ding X, Zhou C, Zhang Y, Xu M, Zhang C and Xu L: Knockdown of eukaryotic translation initiation factors 3B (EIF3B) inhibits proliferation and promotes apoptosis in glioblastoma cells. *Neurol Sci* 33: 1057-1062, 2012.
9. Wang L and Ouyang L: Effects of EIF3B gene downregulation on apoptosis and proliferation of human ovarian cancer SKOV3 and HO-8910 cells. *Biomed Pharmacother* 109: 831-837, 2019.
10. Choi YJ, Lee YS, Lee HW, Shim DM and Seo SW: Silencing of translation initiation factor eIF3b promotes apoptosis in osteosarcoma cells. *Bone Joint Res* 6: 186-193, 2017.
11. Tian Y, Zhao K, Yuan L, Li J, Feng S, Feng Y, Fang Z, Li H and Deng R: EIF3B correlates with advanced disease stages and poor prognosis, and it promotes proliferation and inhibits apoptosis in non-small cell lung cancer. *Cancer Biomark* 23: 291-300, 2018.
12. Spilka R, Ernst C, Mehta AK and Haybaeck J: Eukaryotic translation initiation factors in cancer development and progression. *Cancer Lett* 340: 9-21, 2013.
13. Wang H, Ru Y, Sanchez-Carbayo M, Wang X, Kieft JS and Theodorescu D: Translation initiation factor eIF3b expression in human cancer and its role in tumor growth and lung colonization. *Clin Cancer Res* 19: 2850-2860, 2013.
14. Lin L, Holbro T, Alonso G, Gerosa D and Burger MM: Molecular interaction between human tumor marker protein p150, the largest subunit of eIF3, and intermediate filament protein K7. *J Cell Biochem* 80: 483-490, 2001.

15. Golob-Schwarzl N, Krassnig S, Toeglhofer AM, Park YN, Gogg-Kamerer M, Vierlinger K, Schröder F, Rhee H, Schicho R, Fickert P and Haybaeck J: New liver cancer biomarkers: PI3K/AKT/mTOR pathway members and eukaryotic translation initiation factors. *Eur J Cancer* 83: 56-70, 2017.
16. Team RDCJC: R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. 14: 12-21, 2009.
17. Wickham H: Ggplot2: Elegant graphics for data analysis. *J R Stat Soc* 174: 245-246, 2011.
18. Robin X, Turck N, Hainard A, Tiberti N, Lisacek F, Sanchez JC and Müller M: pROC: An open-source package for R and S+ to analyze and compare ROC curves. *BMC Bioinformatics* 12: 77, 2011.
19. Therneau TM and Grambsch PM: Modeling survival data: Extending the Cox model. Springer, New York, 2000.
20. Subramanian A, Tamayo P, Mootha VK, Mukherjee S, Ebert BL, Gillette MA, Paulovich A, Pomeroy SL, Golub TR, Lander ES and Mesirov JP: Gene set enrichment analysis: A knowledge-based approach for interpreting genome-wide expression profiles. *Proc Natl Acad Sci USA* 102: 15545-15550, 2005.
21. Cronin KA, Lake AJ, Scott S, Sherman RL, Noone AM, Howlander N, Henley SJ, Anderson RN, Firth AU, Ma J, *et al*: Annual report to the nation on the status of cancer, part I: National cancer statistics. *Cancer* 124: 2785-2800, 2018.
22. Xu N, Liu YN, Yin P, Wang LJ, Dou YS, Yang WJ and Zhou MG: Impact of liver cancer deaths on life expectancy in 14 counties (districts) from the Huai River Basin, 2013: Relationship between the water environment and liver cancer. *Zhonghua Yu Fang Yi Xue Za Zhi* 50: 629-633, 2016 (In Chinese).
23. Tu T, Bühler S and Bartenschlager R: Chronic viral hepatitis and its association with liver cancer. *Biol Chem* 398: 817-837, 2017.
24. Chen YJ, Wallig MA and Jeffery EH: Dietary broccoli lessens development of fatty liver and liver cancer in mice given diethylnitrosamine and fed a western or control diet. *J Nutr* 146: 542-550, 2016.
25. Global Burden of Disease Cancer Collaboration, Fitzmaurice C, Akinyemiju TF, Al Lami FH, Alam T, Alizadeh-Navaei R, Allen C, Alsharif U, Alvis-Guzman N, Amini E, *et al*: Global, regional, and national cancer incidence, mortality, years of life lost, years lived with disability, and disability-adjusted life-years for 29 cancer groups, 1990 to 2016: A systematic analysis for the global burden of disease study. *JAMA Oncol* 4: 1553-1568, 2018.
26. Jiao Y, Fu Z, Li Y, Meng L and Liu Y: High EIF2B5 mRNA expression and its prognostic significance in liver cancer: A study based on the TCGA and GEO database. *Cancer Manag Res* 10: 6003-6014, 2018.
27. Jiao Y, Fu Z, Li Y, Zhang W and Liu Y: Aberrant FAM64A mRNA expression is an independent predictor of poor survival in pancreatic cancer. *PLoS One* 14: e0211291, 2019.
28. Jiao Y, Li Y, Lu Z and Liu Y: High trophinin-associated protein expression is an independent predictor of poor survival in liver cancer. *Dig Dis Sci* 64: 137-143, 2019.
29. Jiao Y, Li Y, Fu Z, Hou L, Chen Q, Cai Y, Jiang P, He M and Yang Z: OGDHL expression as a prognostic biomarker for liver cancer patients. *Dis Markers* 2019: 9037131, 2019.
30. Jiao Y, Li Y, Jiang P, Han W and Liu Y: PGM5: A novel diagnostic and prognostic biomarker for liver cancer. *PeerJ* 7: e7070, 2019.
31. Jiao Y, Li Y, Liu S, Chen Q and Liu Y: ITGA3 serves as a diagnostic and prognostic biomarker for pancreatic cancer. *Oncotargets Ther* 12: 4141-4152, 2019.
32. Li Y, Jiao Y, Fu Z, Luo Z, Su J and Li Y: High miR-454-3p expression predicts poor prognosis in hepatocellular carcinoma. *Cancer Manag Res* 11: 2795-2802, 2019.
33. Li Y, Jiao Y, Li Y and Liu Y: Expression of La ribonucleoprotein domain family member 4B (LARP4B) in liver cancer and their clinical and prognostic significance. *Dis Markers* 2019: 1569049, 2019.
34. Li Y, Jiao Y, Luo Z, Li Y and Liu Y: High peroxidase-like expression is a potential and independent prognostic biomarker in breast cancer. *Medicine (Baltimore)* 98: e17703, 2019.
35. Zhang X, Cui Y, He M, Jiao Y and Yang Z: Lipocalin-I expression as a prognosticator marker of survival in breast cancer patients. *Breast Care*, 2019.
36. Cui Y, Jiao Y, Wang K, He M and Yang Z: A new prognostic factor of breast cancer: High carboxyl ester lipase expression related to poor survival. *Cancer Genet* 239: 54-61, 2019.
37. Hou L, Zhang X, Jiao Y, Li Y, Zhao Y, Guan Y and Liu Z: ATP binding cassette subfamily B member 9 (ABCB9) is a prognostic indicator of overall survival in ovarian cancer. *Medicine (Baltimore)* 98: e15698, 2019.
38. Cai H, Jiao Y, Li Y, Yang Z, He M and Liu Y: Low CYP24A1 mRNA expression and its role in prognosis of breast cancer. *Sci Rep* 9: 13714, 2019.
39. Ma F, Li X, Ren J, Guo R, Li Y, Liu J, Sun Y, Liu Z, Jia J and Li W: Downregulation of eukaryotic translation initiation factor 3b inhibited proliferation and metastasis of gastric cancer. *Cell Death Dis* 10: 623, 2019.
40. Ladu S, Calvisi DF, Conner EA, Farina M, Factor VM and Thorgeirsson SS: E2F1 inhibits c-Myc-driven apoptosis via p1K3CA/Akt/mTOR and COX-2 in a mouse model of human liver cancer. *Gastroenterology* 135: 1322-1332, 2008.
41. Xin B, Yamamoto M, Fujii K, Ooshio T, Chen X, Okada Y, Watanabe K, Miyokawa N, Furukawa H and Nishikawa Y: Critical role of Myc activation in mouse hepatocarcinogenesis induced by the activation of AKT and RAS pathways. *Oncogene* 36: 5087-5097, 2017.
42. Li X, Wu Q, Bu M, Hu L, Du WW, Jiao C, Pan H, Sdiri M, Wu N, Xie Y and Yang BB: Ergosterol peroxide activates Foxo3-mediated cell death signaling by inhibiting AKT and c-Myc in human hepatocellular carcinoma cells. *Oncotarget* 7: 33948-33959, 2016.
43. Barnes JL, Zubair M, John K, Poirier MC and Martin FL: Carcinogens and DNA damage. *Biochem Soc Trans* 46: 1213-1224, 2018.



This work is licensed under a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0) License.