



Effects of precipitation changes on soil bacterial community composition and diversity in the Junggar desert of Xinjiang, China

Ke Wu^{1,2,3}, Wenxuan Xu^{1,2} and Weikang Yang^{1,2}

¹CAS Key Laboratory of Biogeography and Bioresource in Arid Land, Xinjiang Institute of Ecology and Geography, Urumqi, China

²Mori Wildlife Ecological Monitoring and Experimentation Station, Xinjiang Institute of Ecology and Geography, Chinese Academy of Sciences, Mori, China

³University of Chinese Academy of Sciences, Beijing, China

ABSTRACT

Variation in precipitation can markedly affect the structure and function of soil microbial communities, especially in arid areas which are limited by water resources. Therefore, it is critical to understand how soil bacterial community composition and diversity will respond to variation in precipitation. In this study, we examined the soil bacterial community structure and diversity between five precipitation treatments (60% decrease, 30% decrease, control, 30% increase and 60% increase in precipitation) in the same arid site, in the Junggar desert of Xinjiang, northern China. The dominant bacterial phyla, present at similar frequencies in plots with different precipitation levels, were Actinobacteria, Proteobacteria, Bacteroidetes, Acidobacteria and Chloroflexi. The Shannon-Wiener and Chao1 indices of soil bacterial α -diversity were both positively correlated with plant diversity. Our results indicated that (1) extreme drought significantly decreased bacterial abundance and diversity compared with increased precipitation; (2) variation in precipitation did not change the dominant components of the bacterial communities; and (3) soil pH and total nitrogen concentration were the key factors affecting soil bacterial composition in the Junggar desert.

Submitted 27 August 2019
Accepted 19 December 2019
Published 27 January 2020

Corresponding authors
Wenxuan Xu, xwx@ms.xjb.ac.cn
Weikang Yang, yangwk@ms.xjb.ac.cn

Academic editor
Xavier Le Roux

Additional Information and
Declarations can be found on
page 17

DOI 10.7717/peerj.8433

© Copyright
2020 Wu et al.

Distributed under
Creative Commons CC-BY 4.0

OPEN ACCESS

Subjects Ecology, Microbiology, Climate Change Biology

Keywords Precipitation variation, Soil bacterial diversity, Drought, Junggar desert

INTRODUCTION

Global precipitation amounts and patterns will change in the context of global climate change (Cui, 2018). The changes in soil moisture caused by precipitation will directly affect water utilization by plants and soil microbes, and indirectly affect the functions and processes of terrestrial ecosystems. Soil microbes are one of the important groups of soil organisms, with almost all soil processes being directly or indirectly associated with soil microbes (Bardgett, 2005). The structure and diversity of soil microbial communities can reflect changes in processes affecting soil environmental quality, and are of great significance for maintaining soil quality and ecosystem stability (Witt & Setälä, 2010). Bacteria are responsible for diverse metabolic functions that affect soil and plant health.

All life forms rely on bacterial processes for their survival, and bacterial diversity is greater than the diversity of any other group of organisms ([Kennedy, 1999](#)).

Desert ecosystems are more sensitive than other regions to environmental changes, because of the characteristic stressful conditions of low water and nutrient availability ([Makhalanyane et al., 2015](#)). Moisture is believed to be one of the major constraints affecting microbial diversity, community structure and activity in arid ecosystems ([Li et al., 2016](#)). Variation in precipitation can affect soil bacterial communities directly by changing soil water availability ([Bagayoko et al., 2000](#)), and indirectly by altering soil nutrient availability ([Xu et al., 2014](#)) and plant community composition and productivity ([Gray et al., 2011](#)).

Over the past few decades, numerous studies have documented the response of soil bacterial communities to precipitation changes. However, as a result of extreme drought and temperature conditions, lack of soil organic matter, species-poor communities and community structure, and simple ecological functions in the desert, results from research into the effect of precipitation change on soil bacterial community in the desert are inconsistent. Most studies have shown that changes in precipitation can affect soil bacterial diversity and community composition ([Ochoa-Hueso et al., 2018](#); [She et al., 2018](#)). For example, [Zhang et al. \(2018\)](#) found that the relative abundances of Proteobacteria and Bacteroidetes did not change significantly with the addition of water, but that the frequency of Acidobacteria clearly decreased in response to water addition, whereas the relative abundance of Planctomycetes increased first and then decreased under two different water addition regimes. On the contrary, other studies have indicated that variation in precipitation did not change the diversity and the composition of the dominant bacterial communities ([Bachar et al., 2010](#); [Wang et al., 2014](#); [Zhang et al., 2019](#)). Collectively, the inconsistent responses of microbes to precipitation changes may result from ecosystem-specific responses, historical precipitation regimes, or indirect water-induced environmental factors ([She et al., 2018](#)).

Plant community composition and diversity is likely to affect the diversity of microbial species ([Johnson et al., 2003](#)). Some research has shown that an increase in plant diversity can increase the bacterial community diversity ([Stephan, Meyer & Schmid, 2000](#); [Loranger-Merciris et al., 2006](#)), because the diversity of bacteria is regulated by the composition of growth-limiting resources, which are mainly based on soil organic matter ([Lavelle et al., 1995](#)). However, some studies have shown inconsistent results. For instance, [Kuramae et al. \(2011\)](#) indicated that no significant changes in bacterial community structure and diversity occurred in response to an increase in plant diversity. Whether there is a relationship between plant diversity and soil bacterial diversity depends on the type of biological interactions between them and the temporal and spatial scale of organisms affected by ecological factors ([Hooper et al., 2000](#)).

The area of temperate desert in Xinjiang in northern China accounts for about 53% of the total desert area in China ([Cui, 2018](#)). Most researchers thought that the rainfall in Xinjiang tended to increase ([Qin, Ding & Su, 2006](#); [Ding et al., 2007](#)), but some researchers suggested that Xinjiang will enter a long-term period of low precipitation in the future ([Ren et al., 2015](#)). Changes in precipitation may alter the activities of soil microbes and thus,

affect the biogeochemical cycles in terrestrial ecosystems (Zhang *et al.*, 2013). Therefore, an accurate understanding of the response of soil microbes to precipitation changes will help us accurately predict how ecosystem functions will change in response to future climate conditions. The soil microbial community composition is influenced by a wide variety of factors, including soil type and texture, moisture, pH and temperature (Fierer & Jackson, 2006). Many of these factors interact with one another and have both direct and indirect effects on the soil microbial community (Buyer *et al.*, 2010). Identification of the factors that influence microbial community composition in the desert will have significant impact on our understanding of how precipitation change will affect soil bacterial community.

We took advantage of an on-going precipitation manipulation experiment in the Junggar desert of Xinjiang to investigate how increases and decreases in precipitation might alter soil bacterial diversity and bacterial community composition. Our main goals were (1) to test how soil bacterial diversity and community composition respond to increased precipitation and extreme drought; (2) to determine the relationship between plant diversity and bacterial diversity; and (3) to identify the key factors that determine soil bacterial community structure. The results from this current study should advance a potential mechanistic understanding of the responses of desert microbial communities to global environmental changes.

METHODS

Study site

The study was conducted in a desert site at the southeastern edge of the Junggar basin at Mori County, Xinjiang, China. The study area (43°59'N, 90°48'E) lies on the alluvial fan of the Tianshan Mountains, with an elevation ranging from 1,000 to 1,200 m above sea level (Fig. 1). The major soil type is the desert gray soil that is highly susceptible to wind erosion. The soil is highly alkaline ($\text{pH} = 9.08 \pm 0.05$) with low fertility. The climate of this area is arid and cold ($+20^\circ\text{C}$ on average in July, and -15°C on average in January). The mean annual precipitation is 150 mm, falling mainly in the spring and winter. The mean soil concentration of total nitrogen and organic matter were 0.49 g kg^{-1} and 6.13 g kg^{-1} , respectively.

The rainfall in the study area is concentrated mainly in the spring, from the beginning of April to the end of May, which is the period of the most vigorous growth of the ephemeral plants and also of the greatest species richness (Xu *et al.*, 2016).

The experimental plots, where the present study was conducted, were fenced as exclosures in 2014. The study area is typical desert, dominated by the shrub *Seriphidium borotalense*. The vegetation is typically sparse ($\sim 25\%$ coverage) and short ($\sim 20\text{ cm}$), with an admixture of scattered shrubs dominated by *Haloxylon ammodendron*, *Atraphaxis spinosa*, and *Reaumuria songarica*, among others. The herbaceous vegetation in the area flourishes in May and June. The most common herbaceous plants are *Ceratocarpus arenarius*, *Salsola paulsenii*, *Halogeton glomeratus*, and *strigosella* (*Malcolmia*) *Africana*.




Figure 1 Experimental site.

Full-size  DOI: [10.7717/peerj.8433/fig-1](https://doi.org/10.7717/peerj.8433/fig-1)

Experimental design

We analyzed the precipitation data in Mori County from 1960 to 2008 (data being obtained from local meteorological stations) and determined that the mean annual precipitation ranged from -30% to $+30\%$ of the average. Furthermore, precipitation is predicted to increase by 30% in northern China over the next 30 years (*Liu et al., 2017*), so the water addition treatments were adjusted by an extra $+30\%$ precipitation or by an extra $+60\%$ precipitation to represent an extreme situation; we also reduced precipitation by 30% and 60% to represent drought and extreme drought treatments, respectively.

In April 2016, fifteen plots were established, with each of them being $2\text{ m} \times 2\text{ m}$, with 3 m buffer zones between any two adjacent plots to avoid any ‘edge effect’. The fifteen plots were represented three replicates for each of five levels of precipitation: 60% decreased precipitation (-60%), 30% decreased precipitation (-30%), control (CK), 30% increased precipitation ($+30\%$), and 60% increased precipitation ($+60\%$).

For the drought treatments, rainout shelters had a metal frame supporting V-shaped clear acrylic bands, without a UV filter, covering an area of 9 m^2 ($3\text{ m} \times 3\text{ m}$). The roof had a 20° inclination, and on the lowest side, it had a gutter that channeled any intercepted water into a flexible storage tank, with a capacity of 170 L, made of white canvas covered with PVC (*Knapp et al., 2017*). The roof suspended over treatment plots only minimally affected other key environmental variables, and this approach is widely used in studies of many types of ecosystems (*Yahdjian & Sala, 2002*). Increased precipitation treatment plots received ambient precipitation collected from the roofs over the drought treatment plots immediately after rainfall. The ambient precipitation plots at each site served as controls (*Fig. 2*).

The number of plant species, percentage cover and plant height were determined in each sample plot in May 2018 (*Table 1*).




Figure 2 Precipitation decreased by 30 percent (A) and 60 percent (B).

Full-size DOI: [10.7717/peerj.8433/fig-2](https://doi.org/10.7717/peerj.8433/fig-2)

Table 1 The general vegetation of the study site.

Variable	Annuals	Perennials	Shrubs
Number of species	14.00 ± 1.26 ^a	2.00 ± 0.05	5.00 ± 0.61
Mean cover (%)	10.03 ± 0.04	5.14 ± 0.01	24.91 ± 0.25
Mean height (cm)	2.80 ± 0.25	14.20 ± 0.01	7.53 ± 1.37

Notes.

^amean ± standard error (SE).

Calculation of plant community diversity

Simpson's diversity index, richness index, Shannon-Wiener index and Pielou evenness index were used to measure community diversity (Lloyd & Ghelardi, 1964; Ma, 1994).

Richness index(R) : $R = S$

$$H' = - \sum_i^S (P_i) (\ln P_i)$$

Pielou index(E) : $E = H' / \ln S$

Where S is the number of species, and P_i is the proportion of individuals belonging to i species.

Soil sampling and analyses

Soil samples were carefully collected from each plot at the topsoil level (0–10 cm) in early August 2018 using an auger (3.5 cm diameter). The diagonal sampling method was adopted to set up four sampling points in each plot, with all soil samples from each replicate plot being mixed uniformly and passed through a 2.0 mm sieve. The fresh soil sample was divided into two subsamples, of which one was air-dried and used to measure physical and chemical properties and the other was stored at -80°C prior to DNA extraction (Petrosino *et al.*, 2009).

Soil water content was determined by taking soil cores in each plot and then oven-drying the cores for 48 h at 105°C . Soil pH was measured in a soil water suspension (1:2.5 w: v) with a digital pH meter (SevenEasy, METLER TOLEDO Corporation). Soil organic

matter (SOM, g kg^{-1}) was determined by oxidation with dichromate in the presence of H_2SO_4 without application of external heat, as described by [Rowell \(1994\)](#). Total nitrogen (TN, g kg^{-1}) was determined by the Kjeldahl method (8420, FOSS Analytical Corporation, Denmark); total phosphorus (TP, g kg^{-1}) was determined using the ignition method of [Saunders & Williams \(1955\)](#) with P detection by acid-molybdate colorimetry (Cary60, Agilent Technologies Inc.); and total K (TK, g kg^{-1}) was determined by Na_2CO_3 extraction ([Olsen & Sommers, 1982](#)) and measurement with an atomic absorption spectrometer (IRIS Advantage-ER, Thermo Jarrell Ash Corporation). Available N (NH_4^+ and NO_3^-) was extracted using KCl and determined calorimetrically in the soil extracts (8420, FOSS Analytical Corporation, Denmark). Available P (AP) was determined using the Kelowna method as described by [Van Lierop \(1988\)](#), using a solid: liquid extraction ratio of 1:5 and quantified by spectrophotometry (Cary60, Agilent Technologies Inc.). Available K (AK) were extracted with the Mehlich III solution ([John et al., 2007](#)) and measured using an inductively coupled plasma atomic-emission spectrometer (IRIS Advantage-ER, Thermo Jarrell Ash Corporation).

Soil DNA extraction and amplification of 16S rRNA genes

Extraction and purification of the total genomic DNA from soil was achieved using the Fast-DNA SPIN Kit for Soil (MP Biomedicals). The quality of the extracted genomic DNA was visualized by electrophoresis through a 1% agarose gel, and the DNA was stored at -80°C until use ([Caporaso et al., 2010](#)). We amplified a region of the 16S rRNA gene for bacteria. The primers used for sequencing were 341F (CCTAYGGGRBGCASCAG)/806R (GGACTACNNGGGTATCTAAT) flanking the V3 and V4 regions of the 16S rRNA gene ([Berg et al., 2012](#)). All PCR reactions were carried out in 30 μL reactions with 15 μL of Phusion® High-Fidelity PCR Master Mix (New England Biolabs); 0.2 μM of forward and reverse primers, and 10 ng template DNA. Thermal cycling consisted of initial denaturation at 98°C for 1 min, followed by 30 cycles of denaturation at 98°C for 10 s, annealing at 50°C for 30 s, and elongation at 72°C for 30 s, with a final incubation at 72°C for 5 min. The same volume of 1 \times loading buffer (contained SYB green) and the PCR products was subjected to electrophoresis on 2% agarose gel for detection. PCR products were mixed in equidensity ratios. Then, mixture PCR products was purified with GeneJET™ Gel Extraction Kit (Thermo Scientific). Samples with bright main strip between 400–450 bp were chosen for further experiments. PCR products was mixed in equimolar ratios. Then, mixture PCR products was purified with GeneJET Gel Extraction Kit (Thermo Scientific). The purified PCR amplicons products were sequenced on an Illumina MiSeq platform (Illumina Inc., San Diego, CA, USA) at Novogene Bioinformatics Technology Co. Ltd. (Beijing, China).

Sequence bioinformatics analysis

Firstly, using Cutadapt (v1.9.1, <http://cutadapt.readthedocs.io/en/stable/>) was used to shear reads low quality part, then, the data from each sample was split from the obtained reads based on Barcode, and raw reads were obtained by cutting off Barcode and primer sequence for preliminary quality control. Secondly, raw reads were compared with the

species annotation database through UCHIME. Algorithm to detect chimeric sequences and finally to remove the chimeric sequences, so as to obtain the clean reads. Clustering analysis was performed using UPARSE software (version 7.0.1001, <http://drive5.com/uparse/>). Sequences with $\geq 97\%$ similarity were assigned to the same operational taxonomic units (OTUs). We got 3,207 OTUs totally and we selected a representative sequence for each OTU and use the RDP classifier to annotate taxonomic information for each representative sequence. In order to compute alpha-diversity, we rarefied the OTU table and calculate three metrics: Chao1 estimates the species abundance; Observed Species estimates the number of unique OTUs found in each sample, and the Shannon-Wiener index estimates the abundance and evenness of the species present. Rarefaction curves were generated based on these three metrics.

Species taxa analysis was conducted using the Mothur method and the small subunit ribosomal RNA (SSU rRNA) database (*Wang et al., 2007*) of SILVA132 (<http://www.arb-silva.de/>) (threshold value was 0.8~1).

Bacterial community diversity

The data of each sample were normalized, where the least amount of data in the sample was taken as the standard for normalization, and the subsequent alpha-diversity analyses were based on data following homogenization. QIIME software (version 1.9.1, <http://qiime.org/>) was used to calculate the diversity values of the samples, including Chao1 index, Shannon-Wiener index, and Simpson's diversity index. R software (Version 2.15.3, <http://www.r-project.org/>) was used to analyze the differences between the groups of alpha-diversity indexes and to calculate a rarefaction curve (*Edgar, 2013*). Species richness and diversity of the bacterial community were characterized by the Chao1 and Shannon-Wiener indexes, respectively and the sequencing depth index was taken as the measure of sequencing coverage (*Wang et al., 2015*).

Chao1 index (*Chao, 1984*):

$$\text{Chao1} = S_{\text{obs}} + \frac{F_1(F_1 - 1)}{2(F_2 + 1)}$$

Where S_{obs} is the number of observed OTUs, F_1 is the number of OTUs with only one sequence, and F_2 is the number of OTUs with only two sequences.

Simpson's diversity index (*Magurran, 2014*):

$$C = 1 - \sum P_i^2$$

Where P_i is the proportion of the community represented by OTU i .

Rarefaction curve

Rarefaction curves can be used to understand the depth of sampling of a community, compared with its total diversity, and can reflect the rationality of sequencing data volume (*Olszewski, 2004*). The rarefaction curve showed a flat state, but one which had not reached a saturated state, indicating that sampling was reasonable and that the result represented the bacterial community structure in the soil environment (*Fig. 3*).




Figure 3 The rarefaction curve.

Full-size DOI: [10.7717/peerj.8433/fig-3](https://doi.org/10.7717/peerj.8433/fig-3)

Statistical analysis

All data were expressed as the mean \pm standard error (SE) of the mean. All data for a particular variable were first checked for normal distribution prior to analysis by using the Kolmogorov–Smirnov Z test. When the data were confirmed to be normally distributed, a parametric one-way measures analysis of variance (ANOVA) and multiple pairwise comparison analyse (Least Significant Difference, LSD) was carried out with SPSS 20.0 (IBM, Armonk, NY, USA), to measure the differences in soil properties and plant diversity among the five treatments. When the data were not normally distributed, the non-parametric Kruskal-Wallis test was used. The threshold probability level for significance in both cases was $P < 0.05$. A significance test was conducted with the pairwise PERMANOVA analysis method using the ‘adonis’ function in vegan package of R software (version 2.15.3, <http://www.r-project.org/>), based on the Bray-Curtis distance measure, to analyze differences in microbial community diversity. We used linear regression and Pearson’s correlation analysis to study the relationship between plant and bacterial diversity. Redundancy analysis (RDA) of the soil bacterial community was analyzed by CANOCO 4.5 (Microcomputer Power, Ithaca, NY, USA) to analyze the influence of environmental variables on the composition of the soil bacterial community. Structural equation modeling (SEM) was used to obtain a mechanistic understanding of how soil properties and plant diversity mediated changes in soil bacterial diversity under different precipitation regimes, by AMOS 17.0 (IBM, Armonk, NY, USA). SEM analysis was performed with the specification of a conceptual model of hypothetical relationships, assuming that precipitation altered soil pH, soil water content and plant diversity, each of which, in turn, affected bacterial community diversity (Zeng et al., 2016). In SEM analysis, we compared the model-derived variance–covariance matrix against the observed

Table 2 Physical and chemical properties of topsoil under different precipitation regimes.

Variable	Precipitation regime				
	−60%	−30%	CK	+30%	+60%
pH	9.19 ± 0.08a	9.44 ± 0.16a	9.24 ± 0.09a	9.56 ± 0.03a	9.45 ± 0.24a
SOM	4.56 ± 0.12c	5.00 ± 0.49bc	5.25 ± 0.20ab	5.29 ± 0.14ab	6.01 ± 0.10a
SWC (%)	2.20 ± 0.01b	3.03 ± 0.01a	3.25 ± 0.01a	4.29 ± 0.01a	4.08 ± 0.01a
TN	0.22 ± 0.03a	0.23 ± 0.05a	0.25 ± 0.05a	0.26 ± 0.06a	0.27 ± 0.06a
TP	0.56 ± 0.01a	0.56 ± 0.02a	0.53 ± 0.02a	0.54 ± 0.02a	0.55 ± 0.04a
TK	19.04 ± 0.86a	19.80 ± 0.44a	19.69 ± 0.48a	19.05 ± 0.17a	19.57 ± 0.26a
AN	13.23 ± 2.40a	13.49 ± 2.74a	15.36 ± 2.79a	15.98 ± 2.37a	13.5 ± 3.84a
AP	8.14 ± 0.32a	7.92 ± 0.83ab	8.17 ± 0.40a	6.19 ± 1.46b	7.43 ± 1.01ab
AK	241.40 ± 20.14a	269.50 ± 36.20a	210.90 ± 11.53a	247.80 ± 41.66a	193.10 ± 31.52a

Notes.

Data are presented as mean ± SE. CK, control (ambient precipitation). For a particular variable, any two samples with a common letter are not significantly different ($P > 0.05$). SOM (g kg^{-1}), soil organic matter; SWC, soil water content; TN (g kg^{-1}), total nitrogen; TP (g kg^{-1}), total phosphorus; TK (g kg^{-1}), total potassium; AN (mg kg^{-1}), available nitrogen; AP (mg kg^{-1}), available phosphorus; AK (mg kg^{-1}), available potassium.

variance–covariance matrix, and data were fitted to the models, using the maximum likelihood estimation. Adequate model fits were indicated by the χ^2 test ($df > 5$; $P > 0.05$) and a low root mean square error of approximation (RMSEA) ($P < 0.05$). The final model was improved by removing relationships between observed variables from prior models, based on these indices.

RESULTS

Soil properties and plant diversity index

After 2.5 years of the precipitation treatments, the physicochemical parameters of soil in the 15 plots were determined. The soil properties are summarized in Table 2. The pH values were similar (averaging 9.38 ± 0.14) in all plots. Soil water content (SWC) ranged from 4.56% to 6.01% across the plots, with soil pH and SWC being highest in the +30% plots. Compared with the control (CK), +30% and +60% plots, decreasing precipitation by 60% (−60%) significantly reduced the soil organic matter content (SOM), and there was a significantly lower SOM in −30% than in +60% (Table 2). The soil water content (SWC) under the −60% treatment was significantly lower than under the other four treatments, but any differences in SWC between the other four treatments were not significant. There was a general trend of increasing SOM, TN and SWC with increasing precipitation.

The plant species richness ranged from 5.80 to 8.00 across the plots, being highest in the +60% plot. The Shannon-Wiener index varied from 1.23 to 1.69, and the Pielou and Simpson indices were in the range 0.63–0.82 and 0.23–0.43, respectively (Table 3), with no significant differences in any of these variables between the treatments.

Soil bacterial diversity

A total of 392,089 optimized sequences were obtained. The number of optimized sequences from each soil sample over the different precipitation regimes ranged from 75,777 to 80,493, with a mean ± SE of 78,418 (± 892.61) sequences per sample. Soil bacterial diversity and

Table 3 Physical and chemical properties of topsoil under different precipitation regimes. Data are presented as mean \pm SE. CK, control (ambient precipitation). For a particular variable, any two samples with a common letter are not significantly different ($P > 0.05$).

	Precipitation regime				
	−60%	−30%	CK	+30%	+60%
Richness	5.70 \pm 0.72a	7.30 \pm 0.27a	6.60 \pm 0.27a	7.00 \pm 0.47a	8.00 \pm 0.82a
Shannon–Wiener index	1.23 \pm 0.03a	1.42 \pm 0.02a	1.40 \pm 0.02a	1.45 \pm 0.06a	1.69 \pm 0.09a
Pielou index	0.70 \pm 0.05a	0.82 \pm 0.04a	0.63 \pm 0.04a	0.81 \pm 0.07a	0.80 \pm 0.07a
Simpson index	0.39 \pm 0.04a	0.43 \pm 0.04a	0.29 \pm 0.00a	0.30 \pm 0.02a	0.23 \pm 0.05a

Table 4 Variance analysis of alpha-diversity and richness of soil bacterial communities.

Diversity index	<i>df</i>	<i>F</i>	<i>P</i>
Chao1	4	2.10	0.026
Shannon index	4	3.432	0.042
Simpson index	4	3.224	0.061

Table 5 Alpha-diversity and richness of soil bacterial communities under the different precipitation treatments. Data are presented as mean \pm SE. CK, control (ambient precipitation). For a particular variable, any two samples with a common letter are not significantly different ($P > 0.05$).

Diversity index	Precipitation regime				
	−60%	−30%	CK	+30%	+60%
Chao1	2156.47 \pm 75.81b	2390.46 \pm 96.93ab	2471.93 \pm 75.87ab	2463.31 \pm 79.83a	2503.69 \pm 74.88a
Shannon index	8.82 \pm 0.13b	9.09 \pm 0.09ab	9.30 \pm 0.08ab	9.21 \pm 0.13a	9.42 \pm 0.06a
Simpson index	0.991 \pm 0.001a	0.994 \pm 0.001a	0.995 \pm 0.000a	0.994 \pm 0.001a	0.996 \pm 0.000a

richness under the extreme drought treatment (−60%) were significantly lower than under treatments with higher precipitation. The bacterial diversity and richness both revealed responses to precipitation in the order: +60%>CK>+30%>−30%>−60% (Tables 4 and 5).

Figure 4 depicts a Venn diagram representing the unique and overlapping OTUs under the five treatments, indicating that most of the members comprising each of the soil communities were similar. The number of unique OTUs decreased with decreasing precipitation, and was significantly higher in the +60% precipitation plots than in the −60% and −30% plots (Fig. 4).

Relationship between bacterial diversity and plant diversity

Bacterial diversity was positively correlated with plant diversity. The Shannon-Wiener index and the Chao1 index of soil bacterial diversity were both positively correlated with plant diversity, but the relationship was significant only in the +30% plot for the Shannon-Wiener index and was significant in the −30% and +30% plots for the Chao1 index (Table 6). Bacterial richness was positively correlated with plant richness. There were significant differences in different treatments, except for the −30% plot (Table 7).

The response of bacterial diversity to changes in precipitation was investigated using the SEM model. We compared multiple candidate models of SEM and selected a best model (χ^2




Figure 4 Venn diagram.

Full-size DOI: 10.7717/peerj.8433/fig-4


Table 6 The correlation between bacterial diversity and plant diversity.

Plant Shannon index	Bacteria Shannon index		Bacteria Chao1 index	
	r	P	r	P
-60%	0.38	0.32	0.78	0.12
-30%	0.79	0.11	0.92	0.02
CK	0.63	0.25	0.75	0.14
+30%	0.94	0.02	0.98	<0.01
+60%	0.57	0.32	0.42	0.66

= 0.300, $P = 0.584$, RMSEA = 0.00), which accounted for 19% and 50% of the variation in pH and SWC, respectively, 47% in plant diversity, and 73% in bacterial diversity. Increased precipitation was associated with increased bacterial diversity both directly and indirectly, via increased soil pH, SWC and plant diversity. Changes in plant diversity resulted in associated shifts in bacterial diversity (Fig. 5). The relationships between the other variables were not significant, but they clearly improved the model when incorporated together.

Table 7 The correlation between bacterial richness and plant richness.

Plant Richness	Bacteria Richness	
	r	P
−60%	0.94	0.02
−30%	0.75	0.15
CK	0.92	0.03
+30%	0.92	0.03
+60%	0.96	0.01

**Figure 5** Structural equation model (SEM) of the effects of the precipitation on the bacterial diversity in the top soil (0–10 cm) soil depth.Full-size DOI: [10.7717/peerj.8433/fig-5](https://doi.org/10.7717/peerj.8433/fig-5)

Bacterial community structure

Soil bacterial community composition was consistent among the different precipitation treatments (Fig. 6). The bacterial communities from each treatment consisted of eight phyla, which were as follows: Actinobacteria (29% of the total amount of reads), Proteobacteria (22%), Bacteroidetes (20%), Acidobacteria (8%), Chloroflexi (7%), Gemmatimonadetes (3.47%), Oxyphotobacteria (2.89%) and Firmicutes (2.35%). The response in frequency of all phyla to variation in precipitation was inconsistent. The abundance of Gemmatimonadetes increased with increased precipitation, but the difference was not significant. The abundance of Proteobacteria and Bacteroidetes peaked at the +30% precipitation treatment, but there was no obvious trend with respect to how the abundance of the other groups in the bacterial community varied in response to precipitation. There




Figure 6 Bacterial composition of the different communities.

Full-size [DOI: 10.7717/peerj.8433/fig-6](https://doi.org/10.7717/peerj.8433/fig-6)

Table 8 Comparison between bacterial community composition at different precipitation regimes based on pairwise PERMANOVA. F represents the test value of F ; R^2 represents the ratio of group variance to total variance; $P < 0.05$.

Group comparison	F	R^2	P
-60%--30%	0.715	0.152	0.800
-60%--CK	0.701	0.149	0.900
-60%--+30%	0.825	0.171	0.700
-60%--+60%	0.736	0.155	0.801
-30%--CK	0.876	0.180	0.701
-30%--+30%	0.793	0.165	0.801
-30%--+60%	0.902	0.184	0.701
CK--+30%	1.100	0.216	0.200
CK--+60%	0.694	0.148	0.900
+30%--+60%	0.821	0.170	0.701

was no significant difference in community composition among the treatments ($P > 0.05$) (Table 8).

Relationships between soil properties and soil bacterial communities

The results of the RDA showed that the first two axes accounted for a total of 59.6% of the variance in bacterial community changes, with the first axis accounting for 41.2% and the second axis accounting for 18.4%. Soil pH and TN had stronger effects on the soil bacterial community, factors explain a total of 64.6% of the variation in community composition (Fig. 7).




Figure 7 Redundancy analysis to explore relationship between microorganism community (genus) and physiochemical characteristic.

Full-size DOI: [10.7717/peerj.8433/fig-7](https://doi.org/10.7717/peerj.8433/fig-7)

DISCUSSION

To reveal the bacterial diversity and community composition response to decreases and increases in precipitation in the Junggar desert, we examined five precipitation treatments. Using high throughput sequencing, we obtained 392,087 high quality sequences and examined the bacterial richness and diversity in each treatment. As expected, the results revealed significant differences in bacterial diversity between drought and increased precipitation treatment, but no significant differences in bacterial community composition among these treatments.

The effects of precipitation on bacterial diversity

The soil microbial community plays a central role in the plant-soil feedback system (Prestel, Salamitou & Dubow, 2008). Therefore, the nature of the relationship between plant diversity and soil microbial diversity has been widely reported (Zak et al., 2003; Wang et al., 2015). For example, research has found that, the generally positive correlation between plant diversity and soil microbial diversity (Prober et al., 2015), will turn into a negative correlation when water and high levels of N are added (Zhang et al., 2018). In arid and semi-arid regions, increasing precipitation alleviated water limitations, and increased the number and abundance of dormant microorganisms to varying degrees, leading to an increase in the Simpson's diversity index for soil bacteria (Tecon & Or, 2017). In our study, significant positive correlations of bacterial richness and diversity with plant

richness and diversity were found where water was added (Tables 6 and 7). Millard & Singh (2010) suggested that increased plant diversity can promote productivity, and that higher productivity leads to greater litter decomposition and more root secretion, which, in turn, increases the soil organic carbon content. In consequence, the increase in plant diversity can result in an increase in soil humus concentration and improved nutrient conditions, providing better conditions for microbial growth (Lange et al., 2015; Wang et al., 2015).

In the present study, soil bacterial diversity and richness increased with increasing precipitation, and there were significant differences between the extreme drought treatment (−60%) and precipitation enhancement treatments (+30% and +60%) (Table 5). The reasons why the difference between precipitation manipulation treatments and the control treatment was not significant in the short term were that the numbers and abundance of species in the desert plant communities were relatively low, the coverage was low, the community species composition was relatively similar between the five treatments (Table 3), and it takes a long time for changes in plant diversity to affect soil properties. Li et al. (2018) suggested that extreme drought can significantly reduce the number and biomass of soil bacteria, because extreme drought soil conditions can inhibit soil microbial growth, and reduce the activity of the microbial community. In the present study, for example, we found that SOM and SWC under the extreme drought treatment were significantly lower than those under higher precipitation levels (Table 2). Furthermore, drought can indirectly affect soil microbial community characteristics by affecting plant physiological processes. For example, drought can restrict the flow of nutrients from the plant to the soil by affecting photosynthesis, which subsequently leads to a decline in soil microbial abundance and diversity (Rousk, Smith & Jones, 2013). It has also been found that drought can lead to a decrease in plant aboveground net primary productivity (ANPP), which leads to a decrease in the amount of carbon and nitrogen entering the soil, and ultimately to a decrease in soil microbial abundance (Beier et al., 2012).

The effects of precipitation on bacterial community composition

Actinobacteria constitute one of the largest phyla among bacteria and represent Gram-positive bacteria, which can decompose considerable amounts of organic matter (Ventura et al., 2007). At the phylum level, Actinobacteria dominated changes in soil bacterial composition under extreme and moderate precipitation events (Zhang et al., 2019). Numerous studies have shown that Actinobacteria, Proteobacteria and Bacteroidetes are the main bacterial groups in desert soils, with the proportion of Actinomycetes reaching up to 50% (Bhatnagar & Bhatnagar, 2005; Bachar, Soares & Gillor, 2012). The abundance of the phylum Firmicutes, correlated negatively with soil moisture, being high in the deserts (Prestel, Salamitou & Dubow, 2008). Consistent with previous findings, at the phylum level, Actinobacteria, Proteobacteria and Bacteroidetes were dominated the composition of soil bacterial communities. In addition, Acidobacteria and Chloroflexi also account for a high proportion in the community (Fig. 6). Gram-positive Actinomycetes are tolerant of drought and can grow under dry conditions, and the relative abundance of Actinobacteria was higher in the less-watered treatments presumably because it has stronger cell walls, and can produce spores and grow in a filamentous pattern, which can effectively

reduce the damage caused by drought and high temperatures (Koyama *et al.*, 2018; Li *et al.*, 2018). Proteobacteria and Bacteroidetes are widespread in desert soils, due to the presence of a variety of nutritional-type bacteria in these phyla (An *et al.*, 2013).

There was little variation in the frequency patterns of these dominant components of soil bacterial communities under the different precipitation treatments in the present study, indicating that the dominant phyla of soil bacteria were stable under precipitation changes. The influence of precipitation change on soil bacterial community structure plays an important role in ecosystem function, such as decomposition of organic matter, transformation of nutrients, and maintenance of soil structure (Yuan *et al.*, 2015). As the dominant components of soil bacterial communities did not change, it can be speculated that soil bacterial community has a regulatory mechanism to influence natural factors on a small scale, to ensure the stability of its composition and maintain its ecosystem function.

The key factor driving the change of bacterial composition

Bacterial community composition can be correlated to an array of environmental factors (Fig. 7). Our studies indicated that indirect environmental factors, such as soil pH, that shift under precipitation changes, might play a larger role in determining microbial community composition than do direct changes to soil moisture. This finding is in accordance with earlier studies which showed that variation in bacterial community composition was strongly correlated with soil pH (Lauber *et al.*, 2008; Yao *et al.*, 2014). Fierer *et al.* (2009) revealed that soils in all biomes were dominated by the same soil bacterial phyla (Acidobacteria, Actinobacteria, Proteobacteria, and Bacteroidetes), with any variation in the composition of the bacterial communities being explained mostly by soil pH. The change in soil pH not only affects the diversity of soil bacteria directly (Fig. 5), but also affects it indirectly by affecting the supply of soil nutrients and the quantity and type of plant root exudates (Zhang *et al.*, 2018). In addition, TN has an important effect on the structure and metabolism of bacterial community (Srinandan *et al.*, 2012; Tu *et al.*, 2019), and has a significant positive correlation with both the richness and biomass of microbial communities (Cordova-kreylos *et al.*, 2006). In this research, we also found that soil pH and TN were the main factors affecting soil microbial community composition (Fig. 7), confirming that they were two of the most important factors involved in determining the composition of the soil microbial community. However, there were no significant differences in soil pH and TN among the five different precipitation treatments investigated in this study (Table 2), which may be the reason why there were no major differences in soil microbial community composition among the different precipitation treatments. Precipitation change is a long and gradual process, and many uncertainties remain to be resolved. Further field studies, with longer observation periods, will be necessary to disentangle the influences of precipitation changes on desert microbial communities.

CONCLUSIONS

In the current study, we manipulated precipitation levels under field conditions to assess the response of soil bacterial diversity and community structure to precipitation changes. The results indicated that extreme drought significantly decreased soil bacterial diversity

compared to increased precipitation, but the effects of increased precipitation on richness and diversity were not significant. There were no significant differences in bacterial community composition among the different precipitation treatments. The studies also indicated that bacterial diversity was positively correlated with plant diversity. Our findings demonstrated an inhibitory effect of extreme drought on soil microbial diversity in a short-term experiment and found the effect of increased precipitation on soil bacterial community was not significant. Precipitation change is a long and gradual process, and many uncertainties remain to be addressed, so that its cumulative effect on the components of the desert ecosystem will require long-term monitoring and research.

ACKNOWLEDGEMENTS

We thank International Science Editing for editing this manuscript. Thanks to the reviewers and issue editor of the journal for their valuable comments, suggestions, and revisions on this manuscript.

ADDITIONAL INFORMATION AND DECLARATIONS

Funding

This work was supported by the West Light Cross Team Project of CAS (2018-XBJCTD-001) and the National Natural Science Foundation of China (31872254). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Grant Disclosures

The following grant information was disclosed by the authors:

West Light Cross Team Project of CAS: 2018-XBJCTD-001.

National Natural Science Foundation of China: 31872254.

Competing Interests

The authors declare there are no competing interests.

Author Contributions

- Ke Wu conceived and designed the experiments, performed the experiments, analyzed the data, prepared figures and/or tables, authored or reviewed drafts of the paper, and approved the final draft.
- Wenxuan Xu conceived and designed the experiments, performed the experiments, authored or reviewed drafts of the paper, and approved the final draft.
- Weikang Yang conceived and designed the experiments, authored or reviewed drafts of the paper, and approved the final draft.

Data Availability

The following information was supplied regarding data availability:

Data is available at NCBI SRA: [SRR10020091–SRR10020105](https://www.ncbi.nlm.nih.gov/sra/SRR10020091-SRR10020105).

Supplemental Information

Supplemental information for this article can be found online at <http://dx.doi.org/10.7717/peerj.8433#supplemental-information>.

REFERENCES

- An S, Couteau C, Luo F, Neveu J, Dubow MS. 2013. Bacterial diversity of surface sand samples from the Gobi and Taklamaken deserts. *Microbial Ecology* **66**(4):850–860 DOI [10.1007/s00248-013-0276-2](https://doi.org/10.1007/s00248-013-0276-2).
- Bachar A, Al-Ashhab A, Soares MIM, Sklarz MY, Angel R, Ungar ED. 2010. Soil microbial abundance and diversity along a low precipitation gradient. *Microbial Ecology* **60**(2):453–461 DOI [10.1007/s00248-010-9727-1](https://doi.org/10.1007/s00248-010-9727-1).
- Bachar A, Soares MIM, Gillor O. 2012. The effect of resource islands on abundance and diversity of bacteria in arid soils. *Microbial Ecology* **63**(3):694–700 DOI [10.1007/s00248-011-9957-x](https://doi.org/10.1007/s00248-011-9957-x).
- Bagayoko M, Alvey S, Neumann G, Buerkert A. 2000. Root-induced increases in soil pH and nutrient availability to fieldgrown cereals and legumes on acid sandy soils of Sudano Sahelian West Africa. *Plant & Soil* **225**:117–127 DOI [10.1023/a:1026570406777](https://doi.org/10.1023/a:1026570406777).
- Bardgett RD. 2005. The biology of soil: a community and ecosystem approach. *Soil Use & Management* **22**(3):323–323 DOI [10.1093/acprof:oso/9780198525035.001.0001](https://doi.org/10.1093/acprof:oso/9780198525035.001.0001).
- Beier C, Beierkuhnlein C, Wothlgemuth T, Penuelas J, Emmett B, Korner C, Boeck H, Christensen JH, Leuzinger S, Janssens IA, Hansen K. 2012. Precipitation manipulation experiments challenges and recommendations for the future. *Ecology Letter* **15**:899–911 DOI [10.1111/j.1461-0248.2012.01793.x](https://doi.org/10.1111/j.1461-0248.2012.01793.x).
- Berg J, Brandt KK, Al-Soud WA, Holm PE, Hansen LH, Sorensen SJ, Nybroe O. 2012. Selection for Cu-tolerant bacterial communities with altered composition, but unaltered richness, via long-term Cu exposure. *Applied and Environmental Microbiology* **78**(20):7438–7446 DOI [10.1128/AEM.01071-12](https://doi.org/10.1128/AEM.01071-12).
- Bhatnagar A, Bhatnagar M. 2005. Microbial diversity in desert ecosystems. *Current Science India* **89**(1):91–100 DOI [10.1016/j.agee.2005.01.014](https://doi.org/10.1016/j.agee.2005.01.014).
- Buyer JS, Teasdale JR, Roberts DP, Zasada IA, Maul JE. 2010. Factors affecting soil microbial community structure in tomato cropping systems. *Soil Biology & Biochemistry* **42**(5):831–841 DOI [10.1016/j.soilbio.2010.01.020](https://doi.org/10.1016/j.soilbio.2010.01.020).
- Caporaso JG, Kuczynski J, Stombaugh J, Bittinger K, Bushman FD, Costello EK, Fierer N. 2010. QIIME allows analysis of high-throughput community sequencing data. *Nature Methods* **7**(5):335–336 DOI [10.1038/nmeth.f.303](https://doi.org/10.1038/nmeth.f.303).
- Chao A. 1984. Non-parametric estimation of the number of classes in a population. *Scandinavian Journal of Statistics* **11**:265–270 DOI [10.2307/4615964](https://doi.org/10.2307/4615964).
- Cordova-kreylos AL, Cao Y, Green PG, Hwang HM, Kuivila KM, LaMontagne MG, Werfhorst VD, Holden PA, Scow KM. 2006. Diversity, composition, and geographical distribution of microbial communities in California Salt

- marsh sediments. *Applied & Environmental Microbiology* 72(5):3357–3366
DOI 10.1128/AEM.72.5.3357–3366.2006.
- Cui XQ. 2018.** Effect of enhanced precipitation, temperature and nitrogen addition on nitrogen fate and plant stoichiometry in temperate desert ecosystem in Xinjiang. PhD thesis, China Agricultural University.
- Ding YH, Ren GY, Zhao ZC, Xu Y, Luo Y, Li QP, Zhang J. 2007.** Detection, attribution and projection of climate change over China. *Desert Oasis Meteo* 1:1–10
DOI 10.3969/j.issn.1002-0799.2007.01.001.
- Edgar RC. 2013.** UPARSE: highly accurate OTU sequences from microbial amplicon reads. *Nature Methods* 10:996–998 DOI 10.1038/NMETH.2604.
- Fierer N, Jackson RB. 2006.** The diversity and biogeography of soil bacterial community. *Proceedings of the Nation Academy of Sciences of the United States of America* 103(3):626–631 DOI 10.1073/pnas.0507535103.
- Fierer N, Strickland MS, Liptzin D, Bradford MA, Cleveland CC. 2009.** Global patterns in belowground communities. *Ecology Letter* 12:1238–1249
DOI 10.1111/j.1461-0248.2009.01360.x.
- Gray SB, Classen AT, Kardol P, Yermakov Z. 2011.** Multiple climate change factors interact to alter soil microbial community structure in an old-field ecosystem. *Soil Science Society of America Journal* 75:2217–2226 DOI 10.2136/sssaj2011.0135.
- Hooper DU, Bignell DE, Brown VK, Brussard L, Dangerfield JM, Wall DH. 2000.** Interactions between aboveground and belowground biodiversity in terrestrial ecosystems: Patterns, mechanisms, and feedbacks. *Bioscience* 50(12):1049–1061
DOI 10.1641/0006-3568(2000)050[1049:IBAABB]2.0.CO;2.
- John R, Dalling JW, Harms KE, Yavitt JB, Stallard RF, Mirabello M, Hubbell SP, Valencia R, Navarrete H, Vallejo M, Foster RB. 2007.** Soil nutrients influence spatial distributions of tropical tree species. *Proceedings of the Nation Academy of sciences of the United States of America* 104:864–869 DOI 10.1073/pnas.0604666104.
- Johnson D, Booth RE, Whiteley AS, Bailey MJ, Reada DJ. 2003.** Plant community composition affects the biomass, activity and diversity of microorganisms in limestone grassland soil. *European Journal of Soil Science* 54(4):671–677
DOI 10.1046/j.1351-0754.2003.0562.x.
- Kennedy AC. 1999.** Bacterial diversity in agroecosystems. *Agriculture, Ecosystems & Environment* 74:65–76 DOI 10.1016/S0167-8809(99)00030-4.
- Knapp AK, Avolio ML, Beier C, Carroll CJW, Collins S, Dukes JS, Fraser LH, Griffinnolan RJ, Hoover DL, Jentsch A, Loik ME, Phillips RP, Post AK, Sala OE, Slette IJ, Yahdjian L, Smith MD. 2017.** Pushing precipitation to the extremes in distributed experiments: recommendations for simulating wet and dry years. *Global Change Biology* 23(5):1774–1782 DOI 10.1111/gcb.13504.
- Koyama A, Steinweg JM, Haddix ML, Dukes JS, Wallenstein MD. 2018.** Soil bacterial community responses to altered precipitation and temperature regimes in an old field grassland are mediated by plants. *FEMS Microbiol Ecology* 94(1):fix156
DOI 10.1093/femsec/fix156.

- Kuramae E, Gamper H, Van Veen JV, Kowalchuk G. 2011.** Soil and plant factors driving the community of soil-borne microorganisms across chronosequences of secondary succession of chalk grasslands with a neutral pH. *FEMS Microbial Ecology* 77(2):285–294 DOI [10.1111/j.1574-6941.2011.01110](https://doi.org/10.1111/j.1574-6941.2011.01110).
- Lange M, Eisenhauer N, Sierra CA, Bessler H, Engels C, Griffiths RI, Mellado-vazquez PG, Malik AA, Roy J, Scheu S, Steinbeiss S, Thomson BC, Trumbore SE, Gleixner G. 2015.** Plant diversity increases soil microbial activity and soil carbon storage. *Nature Communication* 6:Article 6707 DOI [10.1038/ncomms7707](https://doi.org/10.1038/ncomms7707).
- Lauber CL, Strickland MS, Bradford MA, Fiener N. 2008.** The influence of soil properties on the structure of bacterial and fungi community across land-use types. *Soil Biology & Biochemistry* 40(9):2407–2415 DOI [10.1016/j.soilbio.2008.05.021](https://doi.org/10.1016/j.soilbio.2008.05.021).
- Lavelle P, Lattuada C, Trigo D, Barois I. 1995.** Mutualism and biodiversity in soils. *Plant & Soil* 170(1):23–33 DOI [10.1007/bf02183052](https://doi.org/10.1007/bf02183052).
- Li H, Xu Z, Yang S, Li X, Top EM, Wang R, Zhang YG, Cai JP, Yao F, Han XG, Jiang Y. 2016.** Responses of soil bacterial communities to nitrogen deposition and precipitation increment are closely linked with aboveground community variation. *Microbial Ecology* 71:974–989 DOI [10.1007/s00248-016-0730-z](https://doi.org/10.1007/s00248-016-0730-z).
- Li T, Zhang W, Liu GX, Chen T. 2018.** Advances in the study of microbial ecology in desert soil. *Journal of Desert Research* 38(2):329–338 DOI [10.7522/j.issn.1000-694X.2017.00113](https://doi.org/10.7522/j.issn.1000-694X.2017.00113).
- Liu YR, Delgado-Baquerizo M, Trivedi P, He JZ, Wang JT, Singh BK. 2017.** Identity of biocrust species and microbial communities drive the response of soil multifunctionality to simulated global change. *Soil Biology & Biochemistry* 107:208–217 DOI [10.1016/j.soilbio.2016.12.003](https://doi.org/10.1016/j.soilbio.2016.12.003).
- Lloyd M, Ghelardi RJ. 1964.** A table for calculating the ‘Equitability’ component of species diversity. *Journal of Animal Ecology* 33(2):217–225 DOI [10.2307/2628](https://doi.org/10.2307/2628).
- Loranger-Merciris G, Barthes L, Gastine A, Leadley P. 2006.** Rapid effects of plant species diversity and identity on soil microbial communities in experimental grassland ecosystems. *Soil Biology & Biochemistry* 38(8):2336–2343 DOI [10.1016/j.soilbio.2006.02.009](https://doi.org/10.1016/j.soilbio.2006.02.009).
- Ma P. 1994.** Measurement of biological community diversity- α diversity measurement method (I). *Chinese Biodiversity* 2(3):162–168 DOI [10.17520/biods.1994038](https://doi.org/10.17520/biods.1994038).
- Magurran AE. 2014.** *Measuring of biological diversity*. London: Blackwell Publishing.
- Makhalanyane TP, Valverde A, Gunnigle E, Frossard A, Ramond JB, Cowan DA. 2015.** Microbial ecology of hot desert edaphic systems. *FEMS Microbiol Reviews* 39:203–221 DOI [10.1093/femsre/fuu011](https://doi.org/10.1093/femsre/fuu011).
- Millard P, Singh BK. 2010.** Does grassland vegetation drive soil microbial diversity? *Nutrient Cycle in Agroecosystems* 88(2):147–158 DOI [10.1007/s10705-009-9314-3](https://doi.org/10.1007/s10705-009-9314-3).
- Ochoa-Hueso R, Collins SL, Delgado-Baquerizo M, Hamonts K, Power SA. 2018.** Drought consistently alters the composition of soil fungal and bacterial communities in grasslands from two continents. *Global Change Biology* 24(7):2818–2827 DOI [10.1111/gcb.14113](https://doi.org/10.1111/gcb.14113).

- Olsen SR, Sommers LE. 1982.** In: Miller RH, Keeney DR, eds. *Methods of soil analysis: part 2. Chemical and microbiological properties*. Madison: Monograph Number 9 Agronomy Monograph 9, ASA and SSSA 403–430.
- Olszewski TD. 2004.** A unified mathematical framework for the measurement of richness and evenness within and among multiple communities. *Oikos* **104**:377–387 DOI [10.2307/3547967](https://doi.org/10.2307/3547967).
- Petrosino JF, Highlander S, Luna RA, Gibbs RA. 2009.** Metagenomic pyrosequencing and microbial identification. *Clinical Chemistry* **55**:856–866 DOI [10.1373/clinchem.2008.107565](https://doi.org/10.1373/clinchem.2008.107565).
- Prestel E, Salamiou S, Dubow MS. 2008.** An examination of the bacteriophages and bacteria of the Namib Desert. *Journal of Microbiology* **46**:364–372 DOI [10.1007/s12275-008-0007-4](https://doi.org/10.1007/s12275-008-0007-4).
- Prober SM, Leff JW, Bates ST, Elizabeth TB, Jennifer F, Harpole WS, Lind EM, Seabillm EW, Adler PB, Bakker JD, Cleland EE, DeCrappeo NM, Delorenze E, Hagenah N, Hautier Y, Hofmockel KS, Kirkman KP, Knops JMH, Pierre KJL, MacDougall AS, McCulley RL, Mitchell CE, Risch CJ, Williams RJ, Fierer N. 2015.** Plant diversity predicts beta but not alpha diversity of soil microbes across grasslands worldwide. *Ecology Letter* **18**(1):85–95 DOI [10.1111/ele.12381](https://doi.org/10.1111/ele.12381).
- Qin DH, Ding YH, Su JL. 2006.** Assessment of climate and environment changes in China (I): climate and environment changes in china and their projection. *Advance Climate Change Research* **1**:4–9.
- Ren GY, Yuan YJ, Liu YJ, Ren YY. 2015.** Changes in precipitation over Northwest China. *Arid Zone Research* **33**(1):1–19 DOI [10.13866/j.azr.2016.01.01](https://doi.org/10.13866/j.azr.2016.01.01).
- Rousk J, Smith AR, Jones DL. 2013.** Investigating the long-term legacy of drought and warming on the soil microbial community across five European shrubland ecosystems. *Global Change Biology* **19**:3872–3884 DOI [10.1111/gcb.12338](https://doi.org/10.1111/gcb.12338).
- Rowell DL. 1994.** *Soil science: methods and applications*. Harlow: Longman Scientific and Technical Group, 86–87.
- Saunders WMH, Williams EG. 1955.** Observations on the determination of total organic phosphorus. *Journal of Soil Science* **6**:254–267 DOI [10.1111/j.1365-2389.1955.tb00849.x](https://doi.org/10.1111/j.1365-2389.1955.tb00849.x).
- She WW, Bai YX, Zhang YQ, Qin SG, Feng W, Sun YF, Zheng J, Wu B. 2018.** Resource availability drives responses of soil microbial communities to short-term precipitation and nitrogen addition in a Desert Shrubland. *Frontiers in Microbiology* **9**:Article 186 DOI [10.3389/fmicb.2018.00186](https://doi.org/10.3389/fmicb.2018.00186).
- Srinandan CS, D'souza G, Srivastava N, Nayak BB, Nerurkar AS. 2012.** Carbon sources influence the nitrate removal activity, community structure and biofilm architecture. *Bioresource Technology* **117**:292–299 DOI [10.1016/j.biortech.2012.04.079](https://doi.org/10.1016/j.biortech.2012.04.079).
- Stephan A, Meyer AH, Schmid B. 2000.** Plant diversity affects culturable soil bacteria in experimental grassland communities. *Journal of Ecology* **88**(6):988–998 DOI [10.1046/j.1365-2745.2000.00510.x](https://doi.org/10.1046/j.1365-2745.2000.00510.x).
- Tecon R, Or D. 2017.** Biophysical processes supporting the diversity of microbial life in soil. *FEMS Microbiol Reviews* **41**:599–623 DOI [10.1093/femsre/fux039](https://doi.org/10.1093/femsre/fux039).

- Tu Y, Li HX, Jiang L, Wang DQ. 2019.** Bacterial community structure and diversity in rhizosphere of different plants from Huixian wetland, Guangxi. *Ecology & Environmental Science* **28(02)**:252–261 DOI [10.16258/j.cnki.1674-5906.2019.02.005](https://doi.org/10.16258/j.cnki.1674-5906.2019.02.005).
- Van Lierop W. 1988.** Determination of available phosphorus in acid and calcareous soils with Kelowna multiple element extractant. *Soil Science* **146**:284–291 DOI [10.1097/00010694-198810000-00009](https://doi.org/10.1097/00010694-198810000-00009).
- Ventura M, Canchaya C, Tauch A, Chandra G, Fitzgerald GF, Chater KF. 2007.** Genomics of actinobacteria: tracing the evolutionary history of an ancient phylum. *Microbiology & Molecular Biology Reviews* **71(3)**:495–548 DOI [10.1128/MMBR.00005-07](https://doi.org/10.1128/MMBR.00005-07).
- Wang NN, Wang MJ, Li SL, Sui X, Han SJ, Feng FJ. 2014.** Effects of variation in precipitation on the distribution of soil bacterial diversity in the primitive Korean pine and broadleaved forests. *World Journal of Microbiology and Biotechnology* **30**:2975–2984 DOI [10.1007/s11274-014-1725-x](https://doi.org/10.1007/s11274-014-1725-x).
- Wang N, Wang MJ, Li SL, Wang NN, Feng FJ, Han SJ. 2015.** Effects of precipitation variation on growing seasonal dynamics of soil microbial biomass in broadleaved Korean pine mixed forest. *Chinese Journal of Applied Ecology* **26(05)**:1297–1305 DOI [10.13287/j.1001-9332.20150302.015](https://doi.org/10.13287/j.1001-9332.20150302.015).
- Wang Q, Garrity GM, Tiedje JM, Cole JR. 2007.** Naive Bayesian classifier for rapid assignment of rRNA sequences into the new bacterial taxonomy. *Applied & Environmental Microbiology* **73(16)**:5261–5267 DOI [10.1128/aem.00062-07](https://doi.org/10.1128/aem.00062-07).
- Witt C, Setàlà H. 2010.** Do plant species of different resource qualities form dissimilar energy channels below-ground. *Applied Soil Ecology* **44(3)**:270–278 DOI [10.1016/j.apsoil.2010.01.004](https://doi.org/10.1016/j.apsoil.2010.01.004).
- Xu WX, Wu K, Huang Y, Yang WK, David B. 2016.** Gradient variation of soil nutrients of a piosphere in Seriphidium desert of northern Xinjiang. *Acta Agrestia Sinica* **24(4)**:807–811 DOI [10.11733/j.issn.1007-0435.2016.04.014](https://doi.org/10.11733/j.issn.1007-0435.2016.04.014).
- Xu Y, Wang HT, Wang YP, Han YF, Jiang YZ, Zhu WR. 2014.** Correlation between soil nutrient availability and bacteria community succession in poplar plantations. *Chinese Journal of Applied and Environmental Biology* **20(3)**:491–498 DOI [10.3724/SP.J.1145.2014.11026](https://doi.org/10.3724/SP.J.1145.2014.11026).
- Yahdjian L, Sala OE. 2002.** A rainout shelter design for intercepting different amounts of rainfall. *Oecologia* **133**:95–101 DOI [10.1007/s00442-002-1024-3](https://doi.org/10.1007/s00442-002-1024-3).
- Yao M, Rui J, Li J, Dai Y. 2014.** Rate-specific responses of prokaryotic diversity and structure to nitrogen deposition in the *Leymus chinensis* steppe. *Soil Biology & Biochemistry* **79**:81–90 DOI [10.1016/j.soilbio.2014.09.009](https://doi.org/10.1016/j.soilbio.2014.09.009).
- Yuan HC, Wu H, Ge TD, Li KL, Wu JS, Wang JR. 2015.** Effects of long-term fertilization on bacterial and archaeal diversity and community structure within subtropical red paddy soils. *Chinese Journal of Applied Ecology* **26(6)**:1807–1813.
- Zak DR, Holmes WE, White DC, Peacocke AD, Tilman D. 2003.** Plant diversity, soil microbial communities, and ecosystem function: are there any Links? *Ecology* **84(8)**:2042–2050 DOI [10.1890/02-0433](https://doi.org/10.1890/02-0433).

- Zeng J, Liu XJ, Song L, Lin XG, Chu H. 2016.** Nitrogen fertilization directly affects soil bacterial diversity and indirectly affects bacterial community composition. *Soil Biology & Biochemistry* **92**:41–49 DOI [10.1016/j.soilbio.2015.09.018](https://doi.org/10.1016/j.soilbio.2015.09.018).
- Zhang HF, Liu HM, Zhao JN, Li G, Lai X, Li J. 2018.** Effects of simulated nitrogen deposition and precipitation change on soil bacterial community structure in a *Stipa baicalensis* steppe. *Acta Ecology Sinica* **38**(1):244–253 DOI [10.5846/stxb201612092539](https://doi.org/10.5846/stxb201612092539).
- Zhang H, Liu WJ, Kang XM, Cui XY, Wang YF, Zhao HT, Qian XQ, Hao YB. 2019.** Changes in soil microbial community response to precipitation events in a semi-arid steppe of the Xilin River Basin, China. *Journal of Arid Land* **11**(1):97–110 DOI [10.1007/s40333-018-0071-5](https://doi.org/10.1007/s40333-018-0071-5).
- Zhang N, Liu W, Yang H, Yu X, Jessica LMG. 2013.** Soil microbial responses to warming and increased precipitation and their implications for ecosystem C cycling. *Oecologia* **173**:1125–1142 DOI [10.1007/s00442-013-2685-9](https://doi.org/10.1007/s00442-013-2685-9).