

POSTER PRESENTATION

Open Access

Synchronizing and desynchronizing effects of nonlinear delayed feedback deep brain stimulation in Parkinson's disease

Andrey A Dovzhenok¹, Choongseok Park¹, Robert M Worth², Leonid L Rubchinsky^{1,3*}

From Twenty First Annual Computational Neuroscience Meeting: CNS*2012
Decatur, GA, USA. 21-26 July 2012

Hypokinetic symptoms of Parkinson's disease have been related to the excessive beta-band oscillations and synchrony in basal ganglia and other structures; suppression of these synchronized oscillations is believed to suppress the motor symptoms of Parkinson's disease [1]. This leads to a strong interest in a possibility of desynchronizing these pathological oscillations through electrical stimulation of affected subcortical areas. Deep brain stimulation (DBS) is an accepted clinical practice, however, no feedback-based DBS algorithm has been implemented, which would desynchronize neural activity by delivering small currents. One method received a lot of attention recently and has appeared to be very promising: nonlinear delayed feedback control. Numerical simulation showed that a delayed feedback through the mean field (being a proxy for easy-to-record local field potentials, LFP) may cancel the effect of coupling and desynchronize ensembles of coupled oscillators (see, for example, [2]). When the dynamics became desynchronized, the stimulation currents needed to support it were very small (vanishing in the large number of neurons limit). Subsequent studies provided further computational evidence for the ability of nonlinear delayed feedback to destabilize synchronized state and thus to desynchronize synchronous dynamics.

However, the dynamics of the neural activity in Parkinson's disease exhibits complex intermittent synchronous patterns [3], far from the idealized synchronous dynamics used to study the delayed feedback stimulation. To investigate the action of nonlinear delayed feedback stimulation on the parkinsonian

synchrony we employ the computational model of the basal ganglia networks [4], which successfully reproduces experimentally recorded neural activity, as ensured by the matching the temporal patterning of the synchronous dynamics [5]. The model [4] is based on the membrane properties of the basal ganglia cells and is can reproduce not only the average synchrony levels, but also the temporal patterns of the synchronous dynamics, recorded in experiment.

We study the effect of delayed feedback in numerical experiment. When the parameters of our model are such that the dynamics is strongly synchronous, the feedback, in line with earlier studies, exerts desynchronizing action. However, this strongly synchronous dynamics is not physiologically realistic. When the model is tuned to reproduce highly variable temporal patterns observed experimentally, the same delayed feedback tends to increase the synchrony in the system. "Desynchronizing" delayed feedback acting on realistic partially synchronous dynamics boosts rather than suppresses synchronization. The study also indicates that in general, delayed feedback stimulation may not necessarily exhibit desynchronization effect, when acting on a realistic partially synchronous dynamics (the kind of dynamics expected in neuroengineering applications), and sets an example of how to estimate the stimulation effect.

Acknowledgements

This study was supported by NIH grant R01NS067200 (NSF/NIH CRCNS program).

Author details

¹Department of Mathematical Sciences and Center for Mathematical Biosciences, Indiana University Purdue University Indianapolis, Indianapolis, IN 46202, USA. ²Department of Neurosurgery, Indiana University School of

* Correspondence: leo@math.iupui.edu

¹Department of Mathematical Sciences and Center for Mathematical Biosciences, Indiana University Purdue University Indianapolis, Indianapolis, IN 46202, USA

Full list of author information is available at the end of the article

Medicine, Indianapolis, IN 46202, USA. ³Stark Neurosciences Research Institute, Indiana University School of Medicine, Indianapolis, IN 46202, USA.

Published: 16 July 2012

References

1. Hammond C, Bergman H, Brown P: Pathological synchronization in Parkinson's disease: networks, models and treatments. *Trends Neurosci* 2007, 30:357-364.
2. Popovych OV, Hauptmann C, Tass PA: Impact of nonlinear delayed feedback on synchronized oscillators. *J Biol Phys* 2008, 34:267-279.
3. Park C, Worth RM, Rubchinsky LL: Fine temporal structure of beta oscillations synchronization in subthalamic nucleus in Parkinson's disease. *J Neurophysiol* 2010, 103:2707-2716.
4. Park C, Worth RM, Rubchinsky LL: Neural dynamics in parkinsonian brain: the boundary between synchronized and nonsynchronized dynamics. *Phys Rev E* 2011, 83:042901.
5. Ahn S, Park C, Rubchinsky LL: Detecting the temporal structure of intermittent phase locking. *Phys Rev E* 2011, 84:016201.

doi:10.1186/1471-2202-13-S1-P53

Cite this article as: Dovzhenok et al.: Synchronizing and desynchronizing effects of nonlinear delayed feedback deep brain stimulation in Parkinson's disease. *BMC Neuroscience* 2012 13(Suppl 1): P53.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

