

Emotional Arousal, Blood Glucose Levels, and Memory Modulation: Three Laboratory Exercises in Cognitive Neuroscience

Robert W. Flint, Jr.

Department of Psychology, The College of Saint Rose, Albany, NY 12203-1490

The relationships between emotional arousal and cognition in humans represent an important field in cognitive neuroscience. Studies examining the characteristics of emotion-induced memory enhancement and the mechanisms through which these effects occur are becoming increasingly common. This article describes three affordable laboratory exercises of relevance to the growing interest in this field. Specifically, Experiment one reviews a protocol for examining memory, hypermnnesia, reminiscence, and primacy/recency effects for emotional and neutral words. Experiments two and three provide opportunities to examine the relationships between blood glucose level and memory for either a list of pictures or the spatial location of pictures. Each laboratory exercise contains a certain amount of flexibility and is malleable to the specific needs of the instructor. For example, the use

of blood glucose monitoring may be of value to a variety of different exercises examining stress and/or emotional arousal and the stimuli used in each of the protocols may be varied, creating opportunities for a number of different novel exercises. A series of questions have been provided at the end of each exercise in order to help stimulate in-class discussion. The potential application of this line of research in cognitive neuroscience is conveyed through a list of references where glucose has been used to attenuate cognitive deficits in Alzheimer's disease, age-related cognitive decline, and other neuropsychological conditions.

Key words: glucose; emotional arousal; memory; memory modulation; blood sugar; neuroscience education; humans; teaching methods

The impact of acute emotional arousal on subsequent memory performance has been a topic of interest to cognitive neuroscientists for decades (Flint, 2002). The enhancing effects of arousal may be favorable, as in the flashbulb memory and Polyanna effects, or they may be detrimental, as in the case of post-traumatic stress disorder and learned helplessness. In this article I focus on the favorable characteristics of arousal, provide a foundation for the role of glucose in memory modulation of acute emotional arousal, and review three cost-effective laboratory exercises associated with the impact of emotional arousal on memory. Two of these exercises specifically examine the role that blood glucose may play in the memory enhancing effects associated with arousal and review some simple techniques that may be utilized in other laboratory exercises.

Acute emotional arousal, often a result of proximal environmental stimuli and frequently involving the to-be-remembered stimuli, activates a collection of responses known as the stress cascade. This stress cascade involves a series of parallel neuroendocrine responses (reviewed by Kalman & Grahn, 2004). To summarize, the hypothalamic-anterior pituitary-adrenocortical axis (HPA axis) is associated with the endogenous release of adrenocorticotropic hormone (ACTH) and glucocorticoids, whereas the sympatho-adrenomedullary axis (SAM axis) releases epinephrine and norepinephrine. All of these substances have been shown to have memory modulating effects (Gold & van Buskirk, 1975; Haroutunian & Riccio, 1977, 1979; Richardson et al., 1984; Ferry et al., 1999; McGaugh & Roozendaal, 2002).

Our understanding of the body's responsiveness under emotional stress and the need for immediate energy in the form of glucose allows us to modify and extend the




Figure 1. Extension of the Traditional Stress Cascade. The stress cascade, including the HPA-axis and SAM-axis, has been extended to reveal the pathways through which glucocorticoids and epinephrine elevate plasma glucose levels.

stress cascade. Glucocorticoid release produces increases in blood glucose by directly affecting glucose output from the liver, decreasing glucose absorption from peripheral tissue, and by releasing glucogenic amino acid (Figure 1). Similarly, epinephrine increases blood glucose levels directly via the liver.




Figure 2. Schematic Representation of the Dose- and Time-Dependent Effects of Glucose on Memory Consolidation. A) Small and large doses have little or no influence on performance administered immediately after training, but a moderate dose immediately following training provides optimal memory enhancement. B) The same moderate dose of glucose administered after a post-training delay has little or no influence on performance.

In 1984, Messier and White demonstrated that an injection of glucose enhanced subsequent memory performance in rats. Over the next few years research demonstrated both time- and dose-dependent effects of glucose on memory consolidation (Figure 2) that have now been mirrored in the human literature (Messier et al., 1998; Mohanty & Flint, 2001; Sunram-Lea et al., 2001, 2002). Glucose, of a specific dose, administered immediately following training enhances performance whereas an injection of the same dose that is delayed has little or no effect on performance (e.g., Flint & Riccio, 1995). Dose response studies have commonly revealed an inverted-U dose response curve, where small (e.g., 10 mg/kg) and large (e.g., 500 mg/kg) doses have little or no influence on performance, but moderate (e.g., 100 mg/kg) doses provide optimal memory enhancement.

Research examining the effects of glucose on cognition in humans and rodents may be designed in ways that allow scientists to assess different aspects of memory. Glucose administered prior to training (i.e., glucose is present during training) would allow the researcher to

examine memory acquisition (Pavone et al., 1998), although one could not rule out a memory consolidation effect. Glucose administered following the completion of training allows researchers to examine memory consolidation (Flint & Riccio, 1999), and glucose administered shortly prior to testing allows for the examination of memory retrieval (Flint & Riccio, 1997). Much of the research with human participants has involved administering glucose shortly prior to training in paradigms where testing is conducted right after training has been completed. In a protocol such as this, one is unable to determine whether an effect of glucose results from influences on memory acquisition, memory consolidation, or memory retrieval. In much of the non-human animal literature, glucose is administered immediately after the training episode is completed, and testing is delayed for at least 24 hours allowing the animal's system to return to baseline. In a paradigm such as this, any effect of glucose must result from alterations in processes associated with memory consolidation, as the glucose was not present during training or testing. Research with human participants has begun to use this consolidation paradigm more frequently since it is difficult to argue that glucose influenced the participant in some other way, giving the false appearance of memory facilitation (e.g., enhanced attention to task stimuli).

In the context of memory modulation resulting from acute emotional arousal, it is important to demonstrate that increases in blood glucose are associated with emotionally arousing stimuli. Research has shown, for example, that a strong footshock will produce a natural elevation in blood glucose that is comparable to the level in an animal that received a mild footshock plus an injection of 100 mg/kg of glucose (Hall & Gold, 1986). Importantly, animals in both conditions commonly display strong memory retention that is greater than animals that receive a mild footshock alone. This may actually represent an endogenous memory modulating mechanism where the body is capable of increasing the salience of memory in situations of high emotionality and/or stress.

Based on these results and similar studies in humans, one might hypothesize that memory for emotionally arousing material combined with systemic administration of a memory enhancing dose of glucose would actually impair performance in comparison to each treatment alone. We have, in fact, recently demonstrated this to be true (Mohanty & Flint, 2001) and argue that this result is similar to what one would find when systemically administering a large dose of glucose (i.e., glucose increase due to emotional arousal plus glucose increase from systemic administration equals higher than optimal glucose on inverted-U dose response curve).

EXPERIMENT 1 HYPERMNESIA AND REMINISCENCE FOR EMOTIONALLY AROUSING WORDS

The purpose of this exercise is to provide a demonstration of some basic characteristics of memory, specifically, enhanced memory for positive and negative emotionally arousing stimuli, hypermnesia, and

reminiscence. Hypermnnesia is described as an improvement in performance with repeated testing in the absence of additional study sessions, whereas reminiscence is the recall of previously forgotten or unrecalled items. Instructors may also utilize these data from this exercise to demonstrate other characteristics of memory including primacy and recency effects.

Participants The participants for all three of these exercises are typically students who volunteer to participate in the laboratory activity. If students outside the class are used it is recommended that the instructor consult the local Institutional Review Board (IRB) for research with human participants to determine whether or not IRB approval should be obtained.

For this exercise, a minimum of 20 participants works best. There are no specific requirements or restrictions, although it is recommended that participants refrain from eating or drinking for two hours and smokers refrain from smoking for at least 30 minutes prior to the exercise.

Neutral Words	Negative Words	Positive Words
Bench 655	Bomb 46	Cash 503
Banner 649	Demon 106	Orgasm 920
Cord 698	Rape 344	Thrill 438
Jelly 238	Slap 396	Sexy 530
Statue 995	Tumor 459	Kiss 248
Errand 150	Ambulance 15	Ecstasy 75
Habit 775	Cancer 60	Glory 189
Pamphlet 925	Tornado 444	Treasure 449
Hay 784	Crash 89	Fame 157
Prairie 325	Killer 244	Joke 826
Hairpin 776	Pervert 312	Flirt 754
Cow 554	Slave 398	Graduate 192
-----	-----	-----
Valence 5.10	2.15	8.05
Arousal 3.63	6.80	7.15
Frequency 14.33	14.4	14.08

Table 1. Neutral, Negative, and Positive Words. Words, word number, and mean ratings for valence, arousal, and word frequency from the *Affective Norms for English Words* (Bradley & Lang, 1999) used in Experiment 1.

Materials A list of 36 words, shown in Table 1, are needed for this exercise. These words have been selected from the *Affective Norms for English Words* (ANEW, Bradley & Lang, 1999, <http://www.phhp.ufl.edu/csea/>) which provides normative values for valence (positive/negative), arousal (high/low), and dominance (high/low) using a nine point Likert scale. Additionally, ANEW provides the word frequency for each word, as determined by Kucera and Francis (1987) in their analysis of American English. For this exercise, three lists of twelve words, neutral, negative arousing, and positive arousing, were selected. An attempt to match words across the three lists for word frequency was made to help assure that words differed primarily on the valence and arousal dimensions. These 36 words should be arranged in pseudo-random order into a single list, so that four words of each type appear in each third of the list and so that no more than three words of a particular type appear in a

sequence within a particular third of the list. This is done in order to distribute the types of words equally across the list for assessment of primacy and recency effects and to ensure that one particular type of word is not clustered together. Once the wordlist is arranged the words should be placed into a slideshow presentation program such as PowerPoint so that each word appears for five seconds followed by a three second blank screen. When using this as a group class exercise it is recommended that instructors use a computer projection system and projection screen to display the wordlist. In addition to these items instructors will need a stopwatch and will need to create a pleasantness rating sheet containing 36 Likert scales ranging from one (very unpleasant) to five (very pleasant) and three word recall forms with 36 blank spaces on each.

Methods Inform the participants that they will be viewing a slideshow that will present a total of 36 common words and that each word will appear on the screen for five seconds followed by a three second blank screen. At the end of the slideshow there will be a word recall test, so participants should pay close attention to the words when they are presented. In addition, participants will be rating the pleasantness, to ensure somewhat equal processing of each word, by circling the number on the Likert scale that best represents their gut feeling regarding the word. This rating should be done during the three second blank screen after each individual word appears.

Immediately following the slideshow presentation, collect the pleasantness rating forms and distribute one word recall form face down. Instruct the participants to write their name or some unique identifier on the back of the page and tell them that they will have two minutes to write down as many words as they can recall when told to turn the paper over. Do not tell them that there will be more recall tests. Once the first recall test is completed, collect the forms and distribute another blank recall sheet face down. Instruct the participants to write their name/identifier and the number two on the sheet and tell the participants that you want them to try to improve upon their previous recall performance. Give them another two minutes to recall as many words from the wordlist as possible. Once the second recall test is complete, repeat the procedure again, informing the participants that this is the last recall test, to write their name/identifier on the back along with test three, and that the participants should do their very best to recall more words from the wordlist than they did on the previous two trials.

Results & Discussion Once the final recall test is completed, ask each individual to score his/her own data. Distribute to each participant a list of the neutral, positive arousing, and negative arousing words and have them tabulate the following; total number of words recalled, number of neutral words recalled, number of positive arousing words recalled, and the number of negative arousing words recalled for each of the three tests. In addition, have the students calculate the number of each type of word recalled on test two that was not recalled on

test one and the number of words recalled on test three that were not recalled on test two. Finally, if interested in using this exercise to demonstrate primacy and recency effects, participants should be given three lists of words corresponding to each third of the list and asked to tabulate the number of words they recalled each time in each third of the list.

A variety of options exist for examining these data, although they may be unnecessary given that the purpose is to demonstrate simply some basic characteristics of memory in cognitive neuroscience. I recommend that students share their data and create a line graph of the class mean recall performance for neutral, negative arousing, and positive arousing words across the three trials. Figure 3 displays the combined results of a few similar exercises. The increase in performance across the three trials and the clear enhanced performance for emotionally arousing words should suffice to stimulate discussion. Some additional questions to stimulate discussion of these include: Why does performance improve across trials when no additional exposure to the test stimuli occurred? Why is performance greater for emotionally arousing stimuli? Did the class perform equally well with negative and positive arousing stimuli and if no, why not? Why is reminiscence necessary for hypermnesia to occur? Did any students find that they forgot some previously recalled items on tests two and/or three and if so, how does this impact reminiscence if hypermnesia is still demonstrated? How might word imagery have influenced recall performance? Did the emotional valence of the wordlist impact the traditional J-shaped primacy/recency curve? Finally, students may find it interesting to speculate about the arousing nature of the words and whether or not negative words were equally arousing as positive words.

EXPERIMENT 2 EMOTIONAL AROUSAL-INDUCED MEMORY ENHANCEMENT AND BLOOD GLUCOSE ELEVATION

The purpose of this exercise is to examine the relationship between blood glucose level and memory for negative emotionally arousing and neutral pictures. The exercise is based, in part, on studies by Blake et al. (2001) and Mohanty & Flint (2001).

Participants For this exercise it is best to have a minimum of ten participants per group which usually means a class size of 20 or more. Participants must fast (no food or drink except water) for a minimum of two hours prior to the lab exercise, and each student will need to provide a small blood sample. Thus, participants with blood sugar disorders (diabetes, hyper-/hypo-glycemia) or individuals with phenylketonuria, blood clotting disorders, or blood-based contagious infections must refrain from participating. Smokers are also encouraged to refrain from smoking for 30 minutes prior to the lab exercise.

Materials Instructors will need a copy of the *International Affective Picture System* (IAPS; Lang et al., 1999, see author notes for url) which may be obtained from Peter Lang at the Center for Research in Psychophysiology at the University of Florida. This database of pictures comes with normative values for level of arousal (high/low), valence (positive/negative), and dominance (high/low). A set of 25 highly arousing negative pictures and 25 neutral non-arousing pictures should be selected and placed into a computerized slideshow presentation system such as PowerPoint. If the exercise is conducted in a group, instructors may wish to use a room equipped with a projection system and projector screen. Table 1 displays the picture description and picture number for a pre-selected set of negative arousing and neutral pictures along with the mean valence and arousal ratings for each set (based on a nine point Likert scale). A generic pleasantness rating sheet should be developed with 25 Likert scales ranging from one (very unpleasant) to five (very pleasant). At least one, depending on class size, standard diabetic testing kit including glucometer, lancets, and enzyme strips will be needed. The One Touch Basic Glucometer from LifeScan has proven to be an inexpensive and durable glucometer for this purpose. These are available at most local drug stores and pharmacies. Instructors will also need gauze pads, rubbing alcohol, and band-aids. Finally, instructors will need a stopwatch and should prepare a single page of difficult multiplication and division problems to be distributed as a distracting task to prevent rehearsal.




Figure 3. Neutral, Negative Arousing, and Positive Arousing Word Recall on Trials 1, 2, and 3. The increasing performance on each subsequent recall test is evidence of hypermnesia and reminiscence. Negative and positive arousing words were clearly recalled at a higher rate than non-arousing words (Need & Flint, 2000; Need et al., 2001; Shatynski & Flint, 2000).

Procedure Review the participant requirements outlined in the participants section above and explain to the students that some of the pictures may include disturbing negative pictures and if they do not wish to participate in the exercise, they do not have to. Divide the participants into two groups and dismiss one group from the classroom for 30 minutes. Distribute glucometers, enzyme strips, lancets, gauze pads, rubbing alcohol, and band-aids to the participants. Explain that during this exercise participants

will be required to prick their finger twice to obtain two drops of blood, view and rate the pleasantness of 25 pictures, and complete a picture recall test.

Ask participants to select and clean a finger-tip on their non-dominant hand with gauze and rubbing alcohol. Once the fingertip has dried, prepare a lancet and enzyme strip, prick the fingertip, and place a drop of blood onto the enzyme strip in the glucometer to establish a baseline blood glucose level. Fasting blood glucose levels may be expected to range between 60 and 120 mg/dl, although results from a recent study in our lab indicated that blood glucose level ranged from 63 to 111 mg/dl with a mean of 83.92 (SD=8.35) mg/dl after eight hours of fasting (Flint & Zak, unpublished). Participants may wish to put pressure on the finger-prick with a gauze pad for a moment and apply a band-aid. Have each student record the blood glucose value on their datasheet.

Neutral Pictures		Negative Pictures		Positive Pictures	
Description	IAPS #	Description	IAPS #	Description	IAPS #
Fingerprint	2206	Mutilation	3150	Erotic	4290
Neutral Face	2210	Mutilation	3000	Female	4520
Boy	2280	Baby	2053	Erotic Male	4608
Elderly				Couple	4660
Woman	2516	Dying Man	3230	Erotic	4660
Shadow	2880	Boxer	8230	Couple	5260
Rocks	5130	Garbage	9373	Waterfall	5450
Mushrooms	5534	Roach	7380	Liftoff	5470
Plant	5740	Pizza	7380	Astronaut	5470
Bicyclist	5875	Jet	9622	Fireworks	5480
Outlet	6150	Native Boy	2730	Sky-divers	8185
Rolling Pin	7000	Ruins	9470	Windsurfer	5623
Bowl	7006	Knife	6300	Hang	5626
Woman	2271	Meat Slicer	7361	Gliders	5629
Girl	2381	Mutilation	3030	Hiker	5629
Shoes	7031	Abduction	6312	Ice Cream	7270
Hair Dryer	7050	HIV Tatoo	9006	Castle	7502
Car	7096	Aimed Gun	6230	Skier	8030
Trash Can	7060	Tank	6940	Sailing	8080
Lamp	7175	Dirty	9300	Rafters	8400
File Cabinets	7224	Skull	9480	Roller	8490
Building	7491	Injection	9592	Coaster	8501
Car Crash	7920	Bomb	9630	Money	8501
Runner	8010	Ship	9600	Sport Car	8531
Rain	9210	Mutilation	3051	Football	8116
		Tornado	5971	Erotic	4670
Mean Valence	4.78			Couple	4670
Mean Arousal	3.08			Nature	5270
				Lightning	5950

Table 2. Neutral, Negative, and Positive Pictures. Picture description, picture number, and mean valence and arousal ratings for 25 neutral, 25 negative emotionally arousing pictures, and 25 positive emotionally arousing pictures selected from the IAPS database (Lang et al., 1999) for Experiment 2.

Distribute a pleasantness rating sheet with 25 five point Likert scales to the participants. Instruct them that they will be viewing a picture slideshow where each picture will appear for five seconds, followed by a three second blank screen before the next picture appears. Ask

participants to study each picture while it is on the screen and to circle the number on the Likert scale that best represents their gut feeling of pleasantness when the blank screen appears.

Once the slideshow is completed (either negative arousing or neutral pictures), distribute the math worksheet and tell the participants to complete as many problems as possible in the next two minutes. After two minutes have passed, ask participants to turn over their math sheets and write down brief descriptions of as many of the pictures as they can recall. After five minutes, instruct the participants to stop writing, and to conduct another blood glucose test in the same manner as previously described.

This procedure should be repeated with the other group of participants using the remaining picture slideshow. Instructors should be careful to keep the datasheets from the two groups separate for data analysis.

Results & Discussion Once data collection is complete, have all students return to class and go through

score his/her answers. Create summary tables where students can indicate their group (neutral or negative arousing pictures), their blood glucose levels (pre and post slideshow), and the number of pictures correctly recalled. Once each student has a copy of the class data, instruct them to create a line graph for blood glucose levels for the two groups and to create a bar graph for recall performance. Results should indicate elevated blood glucose levels post-viewing for the group that saw negative emotionally arousing pictures as well as a higher rate of recall on the performance test. Engage the class in discussion of these results. Some specific questions instructors may wish to pose include: Why did blood glucose levels increase in one group and not the other and what mechanisms are likely responsible for this result? Can you trace the neurobiological pathways responsible for emotion-induced increases in plasma glucose? How might glucose have affected the brain to improve recall? What other neuroendocrine mechanisms might have contributed to an improvement in performance? Is this a memory effect, or could the results be confounded by attention? Do you think the same results would have occurred if positive emotionally arousing pictures had been used instead of negative emotionally arousing ones?

**EXPERIMENT 3
GLUCOSE-INDUCED ENHANCEMENT OF
SPATIAL MEMORY**

The purpose of this exercise is to demonstrate the memory-enhancing effects of glucose on acquisition of a spatial memory task. This exercise is originally based on a study by Mohanty & Flint (2001) where they demonstrated that 100 mg/kg of glucose administered prior to training enhanced learning (Figure 4).

Participants Participant requirements for this exercise are the same as those for Experiment 2.

Materials For this exercise instructors will need glucose (free samples are available from Arthur Daniels Midland Corn Processing), a non-nutritive artificial sweetener (e.g., saccharine solution Sweet-10 available at many grocery stores), and unsweetened lemon-flavored drink mix (e.g., Kool Aid). Two scales will be needed, one to weigh the participants and one to weigh 100 mg/kg of glucose to be mixed with eight oz of the lemon-flavored beverage. If using Sweet-10 as a control, two ml should be mixed with eight oz of the lemon-flavored beverage. Two 2-gal pitchers, weighing boats, stir sticks, paper towels, and large plastic cups will also be needed. Two blood samples will also be obtained during this exercise so glucometers, enzyme strips, and lancets will be needed as in Experiment 2. Finally, a stopwatch and a set of 16 neutral non-arousing pictures from the IAPS database (Lang et al., 1999, see author notes for url) will be needed. Once selected, the pictures should be printed on white paper, numbered on the back, and laminated. If the participants have not seen the IAPS pictures from Experiment 2, then instructors may simply select 16 of the neutral pictures in Table 2 for use in this exercise.




Figure 4. Mean Number of Errors on a Neutral Spatial Memory Task for Participants Receiving Either Saccharin or 100 mg/kg Glucose. (Results adapted from Mohanty & Flint, 2001).

Procedure Data for this exercise is collected from one student at a time and thus can be time consuming. In order to facilitate the data collection process, instructors may opt to create a second set of 16 neutral pictures and run the experiment twice, once with half the class acting as the researcher and then switching roles. The restrictions for participating in this exercise should be reviewed as described in Experiment 2 with the exception that no negative arousing pictures will be viewed. Participating students should be weighed and randomly assigned to either the saccharin or glucose group. Those in the glucose group should have the appropriate lemon-flavored beverage mixed to ensure a 100 mg/kg dose of glucose. Care should be taken to keep the participants blind to their condition, so the instructor should code the cups to ensure that conditions can be revealed following the exercise. Ask all participants to take a baseline blood glucose measure per the instructions provided in Experiment 2 and then consume the lemon-flavored beverage.

The pictures should be arranged on a large flat surface (e.g., tabletop) in a 4 x 4 grid where the picture labeled number 1 is in the upper left corner and the picture labeled 16 is in the lower right. The participant should be

facing away from the table when the pictures are arranged. Once the pictures have been arranged, the participant should be instructed to turn around and study the pictures paying particular attention to their position. Give the participant 20-sec and instruct them to turn back around and begin counting backward from 200 by 3 out loud (as a distraction to prevent rehearsal). While the participant is counting, shuffle the pictures well and place them into a randomly ordered pile. Tell the participant to turn around, hand him/her the pile of pictures with instructions to place them into a 4 x 4 grid as accurately as possible based on their original location. Once the participant has completed this task, ask him/her to turn back around, flip over all the pictures, and record the number of correct responses. Return the pictures to their correct location on the 4 x 4 grid and repeat this process two more times. After completing the last trial, ask the participant to repeat the blood glucose test.




Figure 5. Sample Baseline and 15-min Post-Consumption Blood Glucose Data. Participants consumed either saccharin or 100 mg/kg of glucose in an 8 oz lemon-flavored beverage and took blood samples before and 15-min after consumption (from Mohanty & Flint 2001).

Results & Discussion Analysis should begin with revealing the group assignment for each participant. Summary tables should be created including group assignment, baseline and post-consumption blood glucose levels, and recall performance on each spatial memory trial. Students should create a line graph, as in Figure 5, demonstrating an increase in plasma glucose levels for those participants that received 100 mg/kg of glucose in their beverage. A line graph for recall performance across the three trials in each group should also be drawn, followed by a class discussion of these results. If the glucose-induced facilitation of performance is difficult to see in a line graph, instruct the students to create a bar graph of these data collapsed across trials (as in Figure 4). Spatial memory has also been shown to be sensitive to gender differences. While the study by Mohanty & Flint (2001) lacked sufficient numbers to address potential gender differences, instructors with a large enough sample may find that an additional analysis of gender differences a useful exercise.

Instructors interested in expanding this exercise further may wish to add a verbal recall task that might allow them to do cross-task comparisons. For this it is recommend that the California Verbal Learning Test be utilized, as prior research has shown that it is sensitive to the memory enhancing effects of glucose in unimpaired adult participants (Sunram-Lea et al., 2002). Some focus questions for this lab include: Was performance in the glucose group greater than that in the saccharin group on each trial, or did the glucose group display facilitated performance on the 2nd and 3rd trials?, What is spatial memory and what sorts of strategies were utilized to complete this task?, If endogenous glucose release is one mechanism through which memories are enhanced under conditions of stress and/or high emotional arousal, how might the enhancement of spatial memory translate into real-world situations?, What areas of the brain and neurotransmitter systems are particularly important for spatial memory?, Given that acetylcholine in the hippocampus is important for spatial memory, how might glucose facilitate these processes (hint: glucose facilitates acetylcholine neurotransmission, Messier et al., 1990; Ragozzino et al., 1996)? (See also Results & Discussion for Experiment 2 for additional questions of relevance).

These exercises are based largely on primary empirical research which has been referenced with each activity.

The role of glucose in memory is also of interest because of its potential involvement in age-related cognitive decline, Alzheimer's disease, and other neuropsychological conditions. While the laboratory exercises presented here are not directly applicable to these conditions, instructors may wish to assign some additional reading before or after an exercise in order to stimulate discussion or provide sources for a written report of the laboratory exercise. Table 3 provides a list of a number of references where glucose has been effectively used to attenuate cognitive deficits for a variety of neuropsychological conditions in humans.

REFERENCES

Blake TM, Varnhagen CK, Parent MB (2001) Emotionally arousing pictures increase blood glucose levels and enhance recall. *Neurobio Learn Mem* 75:262-73.

Bradley MM, Lang PJ (1999) Affective norms for English words (ANEW): Instruction manual and affective ratings. Technical Report C-1, The Center for Research in Psychophysiology, Univeristy of Florida.

Craft S, Dagogo-Jack SE, Wiethop BV, Murphy C, Nevins RT, Fleischman S, Rice V, Newcomer JW, Cryer PE (1993) Effects of hyperglycemia on memory and hormone levels in dementia of the Alzheimer's type: A longitudinal study. *Behav Neurosci* 107:926-40.

Craft S, Zallen G, Baker LD (1992) Glucose and memory in mold senile dementia of the Alzheimer type. *J Clin Exp Neuropsych* 14:253-67.

Ferry B, Roozendaal B, McGaugh JL (1999) Role of norepinephrine in mediating stress hormone regulation of long-term memory storage: A critical involvement of the amygdale. *Bio Psychia* 46:1140-52.

Flint RW Jr (ed) (2002) Forget it? Sources, theories, & mechanisms of alterations in mnemonic function. North Chelmsford, MA: Courier Custom Publishing, Inc.

Flint RW Jr (2002) Glucose-induced memory modulation, In Forget it? Sources, theories, & mechanisms of alterations in mnemonic function (Flint RW Jr, ed) pp 197-216. North Chelmsford, MA: Courier Custom Publishing, Inc.

Flint RW Jr, Riccio DC (1996) Glucose administration attenuates hypothermia-induced retrograde amnesia in rats in a time- and dose-dependent manner. *Psychobio* 24:62-66.

Flint RW Jr, Riccio DC (1997) Pre-test administration of glucose attenuates infantile amnesia for passive avoidance in rats. *Dev Psychobio* 31:207-216.

Flint RW Jr, Riccio DC (1999) Post-training glucose administration attenuates forgetting of passive-avoidance conditioning in 18-day-old rats. *Neurobio Learn Mem* 72:62-67.

Gold PE, van Buskirk RB (1975) Facilitation of time-dependent memory processes with posttrial epinephrine injections. *Behav Bio* 13:145-53.

Gonder-Frederick L, Hall JL, Vogt J, Cox DJ, Green J, Gold PE (1987) Memory enhancement in elderly humans: Effects of glucose ingestion. *Physio Behav* 41:503-4

Hall JL, Gold PE (1986) The effects of training, epinephrine, and glucose injections on plasma glucose level in rats. *Behav Neural Bio* 46:156-76.

Hall JL, Gonder-Frederick LA, Chewing WW, Silveira J, Gold PE (1989) Glucose enhancement of performance on memory tests in young and aged humans. *Neuropsychologia* 27:1129-38.

Reference	Sample
Craft et al. (1992)	Alzheimer's Disease
Craft et al. (1993)	Alzheimer's Disease
Gonder-Frederick et al. (1987)	Elderly
Hall et al., (1989)	Elderly
Manning et al. (1990)	Elderly
Manning et al. (1992)	Elderly
Manning et al. (1993)	Alzheimer's Disease
Manning et al. (1998)	Down's Syndrome
Messier & Gagnon (1996)	Alzheimer's/Diabetes
Messier et al. (1997)	Elderly
Newcomer et al. (1999)	Schizophrenia
Pettersen & Skelton (2000)	Head Injury

Table 3. List of References for Further Reading on Glucose-Induced Memory Modulation in Humans. These references may provide additional context within which students can evaluate these laboratory exercises.

GENERAL DISCUSSION

This article reviews some of the literature regarding enhanced memory for emotionally arousing stimuli, provides a foundation from which to argue that increases in plasma glucose levels may be one of many endogenous mechanisms contributing to this effect, and describes three experiments associated with these effects in cognitive neuroscience. The materials necessary for these human laboratory exercises are quite affordable and the protocols are easily adapted to suit the needs of individual instructors. In particular, the techniques for monitoring blood glucose levels used in Experiments 2 and 3 are relatively simple, and may be easily applied to other experimental protocols where stress/emotional-arousal are of interest. For example, in non-human animals, a simple tail cut can be used to obtain a drop of blood for examination of blood glucose levels or assessment of plasma glucose levels could be added to another exercise such as the one described by Kalman & Grahn (2004).

- Haroutunian V, Riccio DC (1977) Effect of arousal conditions during reinstatement treatment upon learned fear in young rats. *Dev Psychobio* 10:25-32.
- Haroutunian V, Riccio DC (1979) Drug-induced "arousal" and the effectiveness of CS exposure in the reinstatement of memory. *Behav Neural Bio* 26:115-120.
- Kalman BA, Grahn RE (2004) Measuring salivary cortisol in the behavioral neuroscience laboratory. *J Undergrad Neurosci Edu* 2: A41-A49.
- Kucera H, Francis WN (1967) Computational analysis of present-day American English. Providence, RI: Brown Univ Press.
- Lang PJ, Bradley MM, Cuthbert BN (1999) The International Affective Picture System (IAPS). Technical report A-4. Gainesville: University of Florida Center for Research in Psychophysiology.
- Manning CA, Hall JL, Gold PE (1990) Glucose effects on memory and other neuropsychological tests in elderly humans. *Psych Sci* 1:307-11.
- Manning CA, Honn VJ, Stone WS, Jane JS, Gold PE (1998) Glucose effects on cognition in adults with Down's Syndrome. *Neuropsych* 12 479-84.
- Manning CA, Parsons MW, Gold PE (1992) Anterograde and retrograde enhancement of 24-h memory by glucose in elderly humans and young adults. *Behav Neural Bio* 58:125-30.
- Manning CA, Ragozzino ME, Gold PE (1993) Glucose enhancement of memory in patients with probable senile dementia of the Alzheimer's type. *Neurobio Aging* 14:523-28.
- McGaugh JL, Roozendaal B (2002) Role of adrenal stress hormones in forming lasting memories in the brain. *Curr Opin Neurobio* 12:205-10.
- Messier C, Durkin T, Mrabet O, Destrade C (1990) Memory-improving action of glucose: Indirect evidence for a facilitation of hippocampal acetylcholine synthesis. *Behav Brain Res* 39:135-43.
- Messier C, Gagnon M (1996) Glucose regulation and cognition functions: Relation to Alzheimer's Disease and diabetes. *Behav Brain Res* 75:1-11.
- Messier C, Gagnon M, Knott V (1997) Effect of glucose and peripheral glucose regulation on memory in the elderly. *Neurobio Aging* 18:297-304.
- Messier C, Pierre J, Desrochers A, Gravel M (1998) Dose-dependent action of glucose on memory processes in women: Effect on serial position and recall priority. *Cog Brain Res* 7:221-233.
- Messier C, White NM (1984) Contingent and non-contingent actions of sucrose and saccharin reinforcers: Effects on taste preferences and memory. *Physio Behav* 32:195-203.
- Mohanty A, Flint RW Jr (2001) Differential effects of glucose on modulation of emotional and nonemotional spatial memory tasks. *Cog Aff & Behav Neurosci* 1:90-95.
- Need C, Flint RW Jr (2000) Hypermnnesia, reminiscence, and forgetting for positive arousing and neutral words. Poster session presented at the annual meeting of the Eastern Psychological Assn, Baltimore, MD.
- Need C, Shatynski E, Gifford H, Flint RW Jr (2001) Hypermnnesia, reminiscence, and forgetting for negative and positive arousing words. Poster presented at the annual meeting of the Eastern Psychological Assn, Wash DC.
- Newcomer JW, Craft S, Fucetola R, Moldin SO, Selke G, Paras L, Miller R (1999) Glucose-induced increase in memory performance in patients with schizophrenia. *Schizo Bull* 25:321-35.
- Pavone F, Capone F, Battaglia M, Sansone M (1998) Shuttle-box avoidance learning in mice: Improvement by combined glucose and tacrine. *Neurobio Learn Mem* 69:204-210.
- Pettersen JA, Skelton RW (2000) Glucose enhances long-term declarative memory in mildly head-injured rugby players. *Psychobio* 28:81-89.
- Ragozzino ME, Unick KE, Gold PE (1996) Hippocampal acetylcholine release during memory testing in rats: Augmentation by glucose. *Proc Natl Acad Sci* 93:4693-98.
- Richardson R, Riccio DC, Devine L (1984) ACTH-induced recovery of extinguished avoidance responding. *Physio Psych* 12:1984-92.
- Shatynski E, Flint RW Jr (2000) Hypermnnesia, reminiscence, and forgetting for negative arousing and neutral words. Poster presented at the annual meeting of the Eastern Psychological Assn, Baltimore, MD.
- Sunram-Lea SI, Foster JK, Durlach P, Perez C (2001) Glucose facilitation of cognitive performance in healthy young adults: examination of the influence of fast-duration, time of day and pre-consumption plasma glucose levels. *Psychopharm* 157:46-54.
- Sunram-Lea SI, Foster JK, Durlach P, Perez C (2002) The effect of retrograde and anterograde glucose administration on memory performance in healthy young adults. *Behav Brain Res* 134:505-516.

Received August 16, 2004; revised September 20, 2004 accepted October 12, 2004.

The author would like to thank two anonymous reviewers for their feedback and helpful suggestions on an earlier version of this manuscript. ANEW and IAPS inquiries should be directed to Dr. Peter Lang at the Center for the Study of Emotion and Attention at the University of Florida, <http://www.php.ufl.edu/csea/>.

Address correspondence to: Dr. Robert W. Flint, Jr., 432 Western Avenue, The College of Saint Rose, Albany, NY 12203-1490. Email: flintr@strose.edu.