

Original Research Article

The Effects of Dual-Task Training on Patient Outcomes of Institutionalized Elderly Having Chronic Stroke

Yağmur Tetik Aydoğdu^a Onur Aydoğdu^b H. Serap İnal^c

^aDepartment of Physiotherapy and Rehabilitation, Haydarpasa Numune Research and Training Hospital, Istanbul, Turkey; ^bDepartment of Physiotherapy and Rehabilitation, Faculty of Health Sciences, Marmara University, Istanbul, Turkey; ^cDepartment of Physiotherapy and Rehabilitation, Faculty of Health Sciences, Bahcesehir University, Istanbul, Turkey

Keywords

Geriatrics · Rehabilitation · Stroke

Abstract

Background/Aims: The purpose of our study was to investigate the effects of dual-task training on balance, mobility, functional independence, and fear of falling in geriatrics with chronic stroke. **Methods:** Fifty-three geriatrics diagnosed with stroke were included in our study. The elderly were divided into 2 groups, i.e., those walking with (intervention group, $n = 25$) and those walking without (controls, $n = 28$) dual-task training. **Results:** We found statistically significant improvements in all parameters between pre- and posttreatment in both groups ($p < 0.05$). **Conclusion:** Based on our findings, walking with dual-task training can be used in addition to conventional stroke rehabilitation aiming to improve balance and mobility.

© 2018 The Author(s)
Published by S. Karger AG, Basel

Introduction

The restoration of gait is one of the main concerns of stroke rehabilitation to increase the independence of patients in many daily living activities and to increase their participation in society [1, 2] but also to decrease risks of falls, especially for those having dementia or cognitive impairments [3]. The stroke patients with limited cognitive status and attention may have difficulty performing 1 activity or at least 2 activities at the same time [4]. In other

Table 1. Physical characteristics of the patients

	Control group (<i>n</i> = 28)	Dual-task group (<i>n</i> = 25)	Z	<i>p</i>
Age, years	71.21±4.92 (65–86)	69.28±5.03 (65–85)	-1.783	0.075
Gender, female/male	9/19	5/20	-	-
Height, cm	166.78±7.92 (153–182)	167.48±8.21 (155–182)	-0.277	0.782
Body weight, kg	75.07±11.65 (55–102)	71.70±12.37 (50–106)	-0.990	0.322
Body mass index	27.00±1.51 (20.07–33.59)	27.48±2.08 (17.48–37.56)	-1.265	0.206

Values are means ± standard deviations (minimum–maximum).

words, they may have difficulty focusing their attention on different tasks at the same time [5] due to misallocation of their attentional resources [1]. Since gait is an automated function requiring high-level cognitive function [6], dual-task walking is also considered as a standard measure to understand the relationship between cognitive ability and gait [7]. Thus, patients with stroke may need special attention while walking and performing another activity. For instance, Yang et al. [1] reported that elderly with chronic stroke had disturbed balance and slowed walking speed while performing a motor task, such as carrying a tray with glasses. They may also have an increased risk of falling if they focus their attention on another activity, such as traffic lights while crossing a road [5]. In this regard, decreased gait velocity and increased double support as a percentage of stride time were reported for elderly stroke patients if they talked while walking [8]. Thus, the interest of the researchers in the dual-task training approach as a neurophysiologic procedure, and as an additional procedure to the conventional rehabilitation of stroke to restore patients' balance and gait, is increasing extensively [2]. Thus, in general, the literature focuses on the assessment of dual-task performance of patients with stroke. To our knowledge, there is no study reporting the effects of dual-task training as a treatment modality on the clinical findings of patients with chronic stroke. Therefore, the purpose of this study was to investigate the effects of dual-task training on balance, mobility, functional independence, and fear of falling in geriatrics with chronic stroke.

Methods

Study Participants

The 53 elderly (14 females and 39 males) with chronic stroke were selected according to the following inclusion criteria: being 65 years or older, having efficient cognitive function to be able to communicate (a Mini-Mental State Examination score of 23 points and above), walking independently 10 m (if necessary with a cane), and being able to stand on both feet for at least 90 s without assistance. The exclusion criteria were having a severe hearing loss and/or visual impairment, uncontrolled hypertension, and vertigo.

All participants were living independently in a government institution named Darulaceze. Their physical characteristics are given in Table 1. There were no statistical differences between the educational levels and the physical characteristics of the elderly in both groups. They were assigned randomly (according to their arrival) to 2 groups, i.e., those walking with dual-task training (intervention group, *n* = 25) and those walking without dual-task training (controls, *n* = 28). The intervention group included those who accepted the training with dual-task activities while walking (Table 2). Both groups had the same duration

Table 2. Verbal cognitive tasks given while patients were walking with dual-task training

Repeating numbers and letters	Give the patient a certain quantity of numbers and letters depending on the illness and let him/her repeat them from memory
Word spelling	Give the patient a word and let him/her spell the word given
Stroop test	Ask the patient to read color words printed in different colors, e.g. to read the word “yellow” in blue writing
Image description	Show the patient a picture and ask him/her to describe the picture in detail
Verbal communication	Give a letter of the alphabet to the patient and let him/her say words of a certain category, such as plant, animal, country, starting with that letter
Count the number	Ask the patient to count to 3 or from 3 backwards or forwards
Describing daily life activities	Ask the patient to describe something he/she has done on weekdays or on weekends and let him/her explain how to do it

Table 3. Differences between the groups after the treatment program

Outcomes	Dual-task group		Control group		Differences between the groups	
	pretreatment	posttreatment	pretreatment	posttreatment	Δ dual-task	p^2
Balance	42.8±10.0	46.4±10.0	41.5±10.4	43.1±9.1	3.6±3.2	0.020*
Mobility	11.5±3.0	13.2±2.5	8.3±3.2	9.0±3.3	1.7±1.4	0.027*
Functional independence	105.4±23.5	107.8±25.2	101.2±15.6	104.4±14.3	1.6±3.6	0.045*
Fear of falling	35.6±15.8	34.4±15.8	45.1±10.5	44.3±10.6	-1.2±2.3	0.036*

p^1 , Wilcoxon test. p^2 , Mann-Whitney U test. * $p < 0.05$.

of walking sessions, i.e., 5 days a week for 30 min each, and had undergone the same conventional rehabilitation programs (range of motion, strengthening, mat, and mobility exercises) with neurodevelopmental therapy (Bobath therapy) for 45 min 5 days a week. Their functional balance (Berg Balance Scale) [9], mobility (Rivermead Mobility Index) [10], functional independence (Functional Independence Measure) [11], and falls efficacy (Falls Efficacy Scale) [12] were assessed before and after the totally 8 weeks of the treatment program.

Statistical Analysis

All statistical analyses were performed using SPSS Statistics, version 15, software (IBM Corporation, USA). $p < 0.05$ was considered statistically significant. The demographic characteristics (age, height, and body mass index), postural balance, mobility levels, and fear of falling of the cases were expressed as means \pm standard deviations (minimum–maximum).

A Kolmogorov-Smirnov test was used to determine the distribution of the data. Since the groups were not evenly distributed, the data obtained from patients before and after treatment were assessed by the Wilcoxon signed-rank test. Additionally, the analysis of differences between the 2 groups was assessed by the Mann-Whitney U test.

Results

We found significant improvements in the outcomes of balance and mobility of the elderly in both groups ($p < 0.05$). However, the difference was significant in favor of the dual-task group ($p < 0.05$) (Table 3). Although the elderly in both groups had increased functional independence and decreased fear of falling ($p < 0.05$) after the treatment program, there was no significant difference between the groups.

We have found a positive correlation between balance and educational level of the elderly in the dual-task group ($r = -0.409$, $p = 0.043$) ($p < 0.05$). According to simple linear regression analyses, education had an effect of 15% on balance in the dual-task group ($R = 0.389$, $p < 0.05$).

Discussion

We found that balance was improved after the training sessions in the elderly with chronic stroke in both groups. However, the group which had dual-task training while walking showed a significantly better performance in balance than the group which only had a conventional rehabilitation program with neuromuscular therapy. This was also similar in their functional mobility activities. Supporting these outcomes, An et al. [13] reported that the motor and cognitive dual-task gait training was more effective in improving the balance of chronic stroke patients than if both trainings were performed conventionally as separate modalities. Kim et al. [14] also observed improved cognitive and walking abilities of patients with chronic stroke after dual-task training. However, Choi et al. [15] found that dual-task training was as effective as conventional balance training in improving balance and cognition in poststroke patients in a subacute phase. Although we have studied elderly with chronic stroke, we suggest including walking with dual-task training as a modality as early as possible in the rehabilitation program of stroke cases.

We have found a positive correlation between balance and educational level of the elderly in the dual-task group. Although this was a weak (15%) effect according to the linear regression analyses, we consider it as important. Thus, elderly with chronic stroke may derive more benefit from dual-task training for their balance performance if they are well educated.

Based on our findings, we conclude that walking with dual-task training is complementary to conventional stroke rehabilitation aiming at improving balance and mobility. However, for the best practice in physiotherapy and rehabilitation, the educational level of the elderly may be considered while planning the dual-task program.

Statement of Ethics

This study was approved by the Yeditepe University Ethics Committee for Clinical Research (Protocol No. 37068608-6100-15-1233).

Disclosure Statement

The authors acknowledge that there are no conflicts of interest pertaining to this article.

References

- 1 Yang YR, Chen YC, Lee CS, Cheng SJ, Wang RY. Dual-task-related gait changes in individuals with stroke. *Gait Posture*. 2007 Feb;25(2):185–90.
- 2 Kristensen HK, Ytterberg C, Jones DL, Lund H. Research-based evidence in stroke rehabilitation: an investigation of its implementation by physiotherapists and occupational therapists. *Disabil Rehabil*. 2016 Dec;38(26):2564–74.
- 3 Perrochon A, Tchalla AE, Bonis J, Perucaud F, Mandigout S. Effects of a Multicomponent Exercise Program on Spatiotemporal Gait Parameters, Risk of Falling and Physical Activity in Dementia Patients. *Dement Geriatr Cogn Disord Extra*. 2015 Sep;5(3):350–60.
- 4 Silsupadol P, Siu KC, Shumway-Cook A, Woollacott MH. Training of balance under single- and dual-task conditions in older adults with balance impairment. *Phys Ther*. 2006 Feb;86(2):269–81.
- 5 Yang L, He C, Pang MY. Reliability and Validity of Dual-Task Mobility Assessments in People with Chronic Stroke. *PLoS One*. 2016 Jan;11(1):e0147833.
- 6 Yogeve-Seligmann G, Hausdorff JM, Giladi N. The role of executive function and attention in gait. *Mov Disord*. 2008 Feb;23(3):329–42; quiz 472.
- 7 Martínez-Ramírez A, Martinikorena I, Lecumberri P, Gómez M, Millor N, Casas-Herrero A, et al. Dual Task Gait Performance in Frail Individuals with and without Mild Cognitive Impairment. *Dement Geriatr Cogn Disord*. 2016;42(1–2):7–16.
- 8 Bowen A, Wenman R, Mickelborough J, Foster J, Hill E, Tallis R. Dual-task effects of talking while walking on velocity and balance following a stroke. *Age Ageing*. 2001 Jul;30(4):319–23.
- 9 Sahin F, Yilmaz F, Ozmaden A, Kotevolu N, Sahin T, Kuran B. Reliability and validity of the Turkish version of the Berg Balance Scale. *J Geriatr Phys Ther*. 2008;31(1):32–7.
- 10 Rådman L, Forsberg A, Nilsagård Y. Modified Rivermead Mobility Index: a reliable measure in people within 14 days post-stroke. *Physiother Theory Pract*. 2015 Feb;31(2):126–9.
- 11 Ottenbacher KJ, Hsu Y, Granger CV, Fiedler RC. The reliability of the functional independence measure: a quantitative review. *Arch Phys Med Rehabil*. 1996 Dec;77(12):1226–32.
- 12 Yardley L, Beyer N, Hauer K, Kempen G, Piot-Ziegler C, Todd C. Development and initial validation of the Falls Efficacy Scale-International (FES-I). *Age Ageing*. 2005 Nov;34(6):614–9.
- 13 An HJ, Kim JI, Kim YR, Lee KB, Kim DJ, Yoo KT, et al. The effect of various dual task training methods with gait on the balance and gait of patients with chronic stroke. *J Phys Ther Sci*. 2014 Aug;26(8):1287–91.
- 14 Kim GY, Han MR, Lee HG. Effect of Dual-task Rehabilitative Training on Cognitive and Motor Function of Stroke Patients. *J Phys Ther Sci*. 2014 Jan;26(1):1–6.
- 15 Choi JH, Kim BR, Han EY, Kim SM. The effect of dual-task training on balance and cognition in patients with subacute post-stroke. *Ann Rehabil Med*. 2015 Feb;39(1):81–90.