

# BMJ Open Effect of home-based interventions on basic activities of daily living for patients who had a stroke: a systematic review with meta-analysis

Ping Qin,<sup>1,2</sup> Canxin Cai,<sup>1,2</sup> Xuan Chen,<sup>1,3</sup> Xijun Wei <sup>1,2</sup>

**To cite:** Qin P, Cai C, Chen X, *et al.* Effect of home-based interventions on basic activities of daily living for patients who had a stroke: a systematic review with meta-analysis. *BMJ Open* 2022;**12**:e056045. doi:10.1136/bmjopen-2021-056045

► Prepublication history and additional supplemental material for this paper are available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2021-056045>).

Received 09 August 2021  
Accepted 12 July 2022



© Author(s) (or their employer(s)) 2022. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

<sup>1</sup>Rehabilitation Lab of Mix Reality, Shenzhen Hospital, Southern Medical University, Shenzhen, Guangdong, China  
<sup>2</sup>Department of Rehabilitation Medicine, Shenzhen Hospital, Southern Medical University, Shenzhen, Guangdong, China  
<sup>3</sup>Department of Rehabilitation Medicine, Shenzhen Hospital, Peking University, Shenzhen, China

## Correspondence to

Dr Xijun Wei;  
[weixj2016@smu.edu.cn](mailto:weixj2016@smu.edu.cn)

## ABSTRACT

**Objectives** To investigate the effectiveness of home-based interventions in improving the ability to do basic activities of daily living in patients who had a stroke.

**Methods** Randomised controlled trials were searched through MEDLINE, Embase and CINAHL from their inception to 31 December 2021. We included studies involving home-based intervention prescribed by professionals and implemented at patients' homes. The characteristics of these studies were collected. Risk of bias of individual study was assessed by Physiotherapy Evidence Database scale. Meta-analyses were performed where studies reported comparable interventions and outcomes.

**Results** In total, 49 studies were included in the systematic review and 16 studies had sufficient data for meta-analyses. The short-term effect of home-based intervention showed no significant difference when compared with institution-based intervention (standardised mean difference (SMD)=0.24, 95% CI -0.15 to 0.62,  $I^2=0\%$ ). No significant difference was found between home-based intervention and usual care for long-term effect (SMD=0.02; 95% CI -0.17 to 0.22;  $I^2=0\%$ ). Home-based rehabilitation combined with usual care showed a significant short-term effect on the ability to do basic daily activities, compared with usual care alone (SMD=0.55; 95% CI 0.22 to 0.87;  $p=0.001$ ;  $I^2=3\%$ ).

**Conclusion** Home-based rehabilitation with usual care, which varied from no therapy to inpatient or outpatient therapy, may have a short-term effect on the ability to do basic activities of daily living for patients who had a stroke compared with usual care alone. However, the evidence quality is low because of the limited number of studies and participants included in the meta-analysis and the possible publication bias. Future research is needed to investigate the effectiveness of home-based rehabilitation in groups with stratification by stroke severity and time since stroke onset, with elaboration of details of the home-based and the control interventions. Moreover, more high-quality studies are required to prove the cost-effectiveness of newly developed strategies like caregiver-mediated rehabilitation and telerehabilitation.

**The primary source of funding** The Medical Research Fund of Guangdong Province (No: A2021041).

## STRENGTHS AND LIMITATIONS OF THIS STUDY

- ⇒ Investigated the effectiveness of home-based rehabilitation in improving the ability to do basic activities of daily living in patients who had a stroke, compared with institution-based intervention, usual care and no intervention, respectively.
- ⇒ Updated some newly developed home-based treatment strategies such as telerehabilitation and caregiver-mediated intervention, and investigated their effectiveness.
- ⇒ The number of studies included in the individual meta-analysis was limited because of the inadequate data in the individual studies.
- ⇒ The clinical heterogeneity between studies in terms of severity of stroke, onset time of stroke, interventions and manner of delivery also compromised the evidence strength of our meta-analyses.

## INTRODUCTION

Stroke is one of the major causes of death and is a leading cause of adult disability worldwide.<sup>1</sup> About half of patients who had a stroke are left with varying degrees of physical or cognitive impairments.<sup>2</sup> Previous studies have shown that 25%–74% of patients who had a stroke need assistance from caregivers for the activities of daily living,<sup>3</sup> and the quality of life of both patients and caregivers is heavily impacted.<sup>4</sup> Although the need for rehabilitation services for the patients who had a stroke after discharge from acute hospitals is widely recognised, outpatient and inpatient rehabilitation are often compromised for reasons such as lack of accessibility, increased costs and poor compliance.<sup>5 6</sup> On the other hand, the ability to perform activities of daily living in an institution-based environment may not be generalised to the home environment, which is the final discharge destination for most patients who had a stroke.<sup>7</sup> Moreover, the motor relearning of patients who had a stroke improves by context-specific training, and training in the patient's own

environment is preferred.<sup>8</sup> Early supported discharge from hospital with subsequent rehabilitation services at home has shown to be more cost-effective than usual care, with a lower caregiver burden and shorter length of stay in hospital.<sup>9 10</sup> Therefore, a home-based rehabilitation programme could be a viable alternative to institution-based stroke rehabilitation.

A Cochrane review of home-based therapy programmes for upper limb functional recovery following stroke found that there was insufficient good-quality evidence to determine the relative effect of home-based upper limb programmes on performance of basic activities of daily living (BADL), compared with placebo, no intervention or usual care.<sup>11</sup> The limited number of included studies and the heterogeneity in terms of the type of home-based therapy programmes limited the evidence strength. Apart from upper limb function, the ability to perform BADL in patients who had a stroke is influenced by much more factors such as mobility, cognition and communication,<sup>12</sup> environmental limitation<sup>13</sup> and psychological adaptation.<sup>14</sup> Moreover, upper limb function is not linearly related to the actual performance of daily activities, and the improved upper limb motor capacity does not translate into the increased upper limb performance in daily life.<sup>15</sup> Therefore, the effectiveness of home-based intervention including but not limited to upper extremity function training is needed to be investigated.

Another previous review found a significant short-term effect on functional independence in favour of home-based rehabilitation for community-dwelling people who had a stroke.<sup>16</sup> However, the evidence was weak because the control interventions mentioned in the previous review were mixed with usual care, centre-based intervention and no intervention. Moreover, as the development of home-based treatment strategy and also for the purpose of reservation of medical services, novel home-based intervention strategies such as telerehabilitation and caregiver-mediated intervention have emerged nowadays. An updated review is needed to investigate the effectiveness of home-based interventions in improving the ability to perform self-care activities in patients who had a stroke.

The objective of this systematic review was to evaluate the effectiveness of home-based interventions in performance of BADL, when comparing with institution-based intervention, usual care and no intervention, respectively, in patients who had a stroke.

## METHODS

The following items were reported according to the Preferred Reporting Items for Systematic reviews and Meta-Analyses.<sup>17</sup>

### Patient and public involvement

No patient involved.

### Search strategy

The MEDLINE, Embase and CINAHL databases were searched through PubMed, Embase and EBSCOhost

platforms, respectively, from inception to 31 December 2021. The search strategy is presented in online supplemental appendix 1.

### Inclusion and exclusion criteria

We only included participants in home-based intervention groups who were living in their own home. Studies that included participants in home-based intervention groups who were living in care homes and other forms of supported or sheltered accommodation were excluded. We defined the home-based interventions as (1) prescribed by professionals and (2) implemented in the patient's own home. Studies delivered solely in environmental modifications, ergonomic intervention, psychosocial interventions or medication were excluded. The comparison interventions included institution-based intervention, usual care and no intervention. We included studies which measured the ability to do BADL as one of the outcomes. Trials that studied solely instrumental activities of daily living were excluded. We only included peer-reviewed studies in English language. Grey literature like unpublished studies or conference abstracts were excluded.

### Selection of studies

Two reviewers independently extracted randomised controlled studies from the three databases. The duplicate articles were deleted, and the obviously irrelevant studies were eliminated by screening the titles and abstracts. If any one of the reviewers considered one reference as eligible, the full text was assessed and two reviewers evaluated the study separately based on the inclusion and exclusion criteria. The two reviewers also searched for relevant reviews reported on the similar topic from the three databases. Reference lists of those reviews were examined, and citation searching and full-text assessment were conducted to identify the additional eligible studies. Only the studies identified by both reviewers were included in the review. Any disagreements between the two reviewers were resolved through discussion with the third reviewer.

### Data extraction and management

Data were extracted from the included studies and recorded on a data extraction form by one reviewer and checked by another reviewer. The extracted information included the following items: (1) the total number of participants of each group; (2) characteristics of participants such as age, gender, disability level and time elapsed since stroke onset; (3) characteristics of home-based interventions and interventions in the control group, and details of home-based interventions; (4) outcome measures of performance of BADL and the time points of outcome measures; and (5) results of effectiveness.

### Assessment of methodological quality

Two reviewers independently assessed the methodological quality of the included studies using the Physiotherapy Evidence Database scale (PEDro scale).<sup>18</sup> The

PEDro scale is an 11-item scale assessing the following items of individual study: (1) specified eligibility criteria and source of participants; (2) random allocation; (3) concealed allocation; (4) similarity at baseline; (5) blindness of participants; (6) blindness of therapists; (7) blindness of assessors; (8) less than 15% dropouts; (9) intention-to-treat analysis; (10) between-group statistical comparisons; and (11) point and variability measures. Each satisfied item contributes 1 point to the total PEDro score, except for item 1 which pertains to external validity. Researchers have suggested that scores of <4 are considered poor quality, 4–5 are considered fair quality, 6–8 are considered good quality and 9–10 are considered excellent quality.<sup>19 20</sup> Any disagreements between the two reviewers were resolved through discussion with the third reviewer.

### Data analysis

The Cochrane Collaboration's Review Manager software (V.5.3) was used to carry out all statistical analyses. The overall estimate of the treatment effect was calculated using the means and SDs of outcome scores with continuous data in the home-based intervention group and control group. Short-term effect and long-term effect were analysed by comparing the statistical difference of outcome score between two groups at treatment endpoint and at the last follow-up, respectively. Those studies with no mean or SD of outcome measure reported were excluded from the meta-analysis. For the studies that used the same measurement tool, we calculated a pooled estimate of the mean differences (MDs) with 95% CIs. When different measurement tools were used, we used the standardised MDs (SMDs) instead of MDs.

Statistical heterogeneity was measured using the  $I^2$  statistic.  $I^2 > 50\%$  was considered to indicate substantial heterogeneity, which would result in the use of a random-effects model for the meta-analysis. When  $I^2$  was  $\leq 50\%$ , a fixed-effects model was used.<sup>11</sup> Inverse variance method was used to estimate the treatment effect.

We planned to perform several meta-analyses to evaluate (1) the effectiveness of home-based intervention compared with institution-based intervention at treatment endpoint and follow-up; (2) the effectiveness of home-based intervention compared with usual care at treatment endpoint and follow-up; (3) the effectiveness of home-based intervention combined with usual care compared with usual care at treatment endpoint and follow-up; and (4) the effectiveness of home-based intervention compared with no intervention at treatment endpoint and follow-up.

We planned to perform subgroup analyses to investigate the sources of heterogeneity, according to the (1) level of disability of the patients who had a stroke with stratification of mild, moderate and severe stroke; (2) the onset time of stroke; (3) type of home-based intervention; (4) self-mediated and/or caregiver mediated versus professional mediated.

We also planned to perform a sensitivity analysis to diminish the influence of studies with poor methodological quality on the effect size estimate. The studies with poor PEDro score ( $\leq 3$ ) were deleted from the meta-analysis. All of the statistical tests were two tailed, and  $p < 0.05$  represented statistical significance.

We planned to test for funnel plot asymmetry to assess the publication bias if there were more than 10 studies included in the meta-analysis.<sup>21</sup>

### Assessment of certainty of the evidence

Two reviewers independently assessed the quality of the evidence using the Grading of Recommendations, Assessment, Development and Evaluation (GRADE) approach.<sup>22</sup> Five factors result in rating down the quality of evidence including study limitations, inconsistency of results, indirectness of evidence, imprecision and publication bias. We rated the overall quality of evidence as high, moderate, low or very low for each outcome. We justified all decisions to downgrade the quality of evidence.

## RESULTS

### Study identification


The search of the electronic bibliographical databases identified 466 articles (MEDLINE=221, Embase=231, CINAHL=14). Five additional studies were identified through the reference lists of relevant articles and reviews. In total, 49 studies met the eligibility criteria and were included into this systematic review (figure 1).

### Study characteristics

Twenty-one randomised controlled trials compared home-based rehabilitation with institution-based rehabilitation on an inpatient or outpatient basis.<sup>23–43</sup> Fifteen randomised controlled trials compared home-based rehabilitation with usual care, which was provided according to routine practice without the involvement of the research team and might include no therapy, home care, instructions for home rehabilitation, inpatient therapy and outpatient therapy.<sup>32 38 44–56</sup> Five randomised controlled trials evaluated the effect of specific home-based interventions by comparing with blank control or sham control.<sup>57–61</sup> Ten randomised controlled trials compared home-based interventions combined with usual care with usual care alone.<sup>62–71</sup> The main characteristics of the included studies are shown in online supplemental table 1. The summary of details of the home-based intervention in each included study according to the Template for Intervention Description and Replication<sup>72</sup> is shown in online supplemental table 2.

There were various home-based interventions performed in the included studies. Twelve studies did not describe the details of treatment strategy.<sup>28 29 32 33 37 43 48 50 52 56 68 70</sup> Among 37 studies which reported the details of treatment strategy, 19 studies provided physical exercise practice,<sup>23 26 27 31 34–36 45–47 49 53 57–59 61 63 64 71</sup> 20 studies provided

PRISMA 2020 flow diagram for new systematic reviews which included searches of databases, registers and other sources



**Figure 1** PRISMA 2020 flow diagram for new systematic reviews which included searches of databases, registers and other sources. Process of study selection. BADL, basic activities of daily living; PRISMA, Preferred Reporting Items for Systematic reviews and Meta-Analyses.

training of daily activities,<sup>23–27 30 36 38 45 46 53–55 58–61 65 66 71</sup> 8 studies provided task/functional-specific training<sup>35 39–42 51 61 69</sup> and 2 studies provided constraint-induced movement treatment.<sup>44 62</sup> Other treatments like speech and communication therapy,<sup>42</sup> psychosocial intervention,<sup>67</sup> emotion management,<sup>71</sup> electromyography-triggered neuromuscular stimulation,<sup>26 27</sup> environmental modification,<sup>47 64</sup> application of leisure activities,<sup>66 67 71</sup> providing adaptive aids and equipment,<sup>59 66</sup> providing fall prevention strategies<sup>63</sup> and providing splint or orthoses<sup>34</sup> were also used as part of home-based interventions for patients who had a stroke.

The delivery strategy of home-based intervention has transformed in recent years. Before 2009, 24 out of 25 included studies reported that home-based interventions were provided by professionals during home visits. Among 24 studies published from 2009, 11 studies involved self/caregiver-mediated intervention,<sup>23 26 27 31 35 36 44 51 61 62 64</sup> in which 4 studies provided telerehabilitation supervised by professionals.<sup>26 27 35 64</sup>

### Risk of bias in included studies

The methodological quality of the included studies is presented in [table 1](#). Thirty-three out of the included 49 studies were of good methodological quality (PEDro score=6–8). Twelve studies were of fair quality (PEDro score=4–5) and four were of poor quality (PEDro score=1–3).

### Effectiveness of home-based intervention compared with institution-based intervention

Twenty-one studies investigated the effectiveness of home-based intervention compared with institution-based intervention. Three studies did not specify the between-group statistical comparison.<sup>33 35 38</sup> Fifteen studies found no significant difference between two groups in the ability to do daily living no matter at treatment endpoints or at follow-ups.<sup>23–32 37 39–42</sup> While one study<sup>43</sup> demonstrated the superiority of home-based intervention compared with institution-based intervention, one study<sup>34</sup> showed an opposite result. Seven studies reported improvement in ability to do daily living in both groups.<sup>23 26 31 32 36 37 43</sup>

**Table 1** Physiotherapy Evidence Database scores of included studies

Studies	Items											Score (0–10)
	1	2	3	4	5	6	7	8	9	10	11	
Asano <i>et al</i> <sup>23</sup>	√	√					√		√	√	√	5
Baskett <i>et al</i> <sup>24</sup>	√	√	√	√			√	√		√	√	7
Björkdahl <i>et al</i> <sup>25</sup>	√	√	√	√			√	√	√		√	7
Chen <i>et al</i> <sup>26</sup>	√	√	√	√			√	√	√	√	√	8
Chen <i>et al</i> <sup>27</sup>	√	√	√	√			√	√	√	√	√	8
Gladman and Lincoln <sup>28</sup>	√	√		√			√			√	√	5
Gladman <i>et al</i> <sup>29</sup>	√	√	√				√	√		√	√	6
Han <i>et al</i> <sup>30</sup>	√	√		√			√	√	√	√	√	7
Hesse <i>et al</i> <sup>31</sup>	√	√		√			√	√	√	√	√	7
Hofstad <i>et al</i> <sup>32</sup>	√	√	√	√	√		√		√	√	√	8
Kalra <i>et al</i> <sup>33</sup>	√	√	√	√			√	√	√	√	√	8
Özdemir <i>et al</i> <sup>34</sup>	√	√		√				√	√		√	5
Pandian <i>et al</i> <sup>35</sup>	√	√					√					2
Redzuan <i>et al</i> <sup>36</sup>	√	√								√	√	3
Roderick <i>et al</i> <sup>37</sup>	√	√	√	√						√	√	5
Taule <i>et al</i> <sup>38</sup>	√	√	√				√			√	√	5
Thorsén <i>et al</i> <sup>39</sup>	√	√	√				√			√	√	5
von Koch <i>et al</i> <sup>40</sup>	√	√	√				√	√		√	√	6
von Koch <i>et al</i> <sup>41</sup>	√	√	√				√	√		√	√	6
Widén Holmqvist <i>et al</i> <sup>42</sup>	√	√	√				√	√		√	√	6
Young and Forster <sup>43</sup>	√	√		√			√	√		√	√	6
Barzel <i>et al</i> <sup>44</sup>	√	√	√	√			√	√	√	√	√	8
Chaiyawat and Kulkantrakorn <sup>45</sup>	√	√	√	√				√	√	√	√	7
Chaiyawat <i>et al</i> <sup>46</sup>	√	√	√	√				√	√	√	√	7
Chen <i>et al</i> <sup>47</sup>	√	√		√	√			√	√	√	√	7
Deng <i>et al</i> <sup>48</sup>	√	√	√	√			√	√	√	√	√	8
Duncan <i>et al</i> <sup>49</sup>	√	√	√	√				√	√	√		6
Lincoln <i>et al</i> <sup>50</sup>	√	√					√			√	√	4
Lindley <i>et al</i> <sup>51</sup>	√	√		√			√	√	√	√	√	7
Mayo <i>et al</i> <sup>52</sup>	√	√	√	√				√		√	√	6
Rasmussen <i>et al</i> <sup>53</sup>	√	√	√	√						√	√	5
Santana <i>et al</i> <sup>54</sup>	√	√	√	√			√			√	√	6
Walker <i>et al</i> <sup>55</sup>	√	√	√	√			√	√		√	√	7
Wolfe <i>et al</i> <sup>56</sup>	√	√	√						√	√	√	5
Azab <i>et al</i> <sup>62</sup>	√						√					1
Batchelor <i>et al</i> <sup>63</sup>	√	√	√	√			√		√	√	√	7
Chumbler <i>et al</i> <sup>64</sup>	√	√	√	√			√		√	√	√	7
Corrand Bayer <sup>65</sup>	√	√	√	√						√	√	5
Goldberg <i>et al</i> <sup>67</sup>	√	√		√								2
Gilbertson <i>et al</i> <sup>66</sup>	√	√	√				√	√	√	√	√	7
Mandigout <i>et al</i> <sup>68</sup>	√	√		√			√		√	√	√	6
Ricauda <i>et al</i> <sup>69</sup>	√	√		√			√		√	√	√	6
Rudd <i>et al</i> <sup>70</sup>	√	√	√	√						√	√	5

Continued

Table 1 Continued

Studies	Items											Score (0–10)
	1	2	3	4	5	6	7	8	9	10	11	
Wong and Yeung <sup>71</sup>	√	√	√	√			√	√	√	√	√	8
Koç <sup>57</sup>	√	√	√	√						√	√	5
Lin <i>et al</i> <sup>58</sup>	√	√		√		√	√	√	√	√	√	8
Wade <i>et al</i> <sup>59</sup>	√	√		√			√	√		√	√	6
Walker <i>et al</i> <sup>60</sup>	√	√		√			√	√		√	√	6
Wang <i>et al</i> <sup>61</sup>	√	√					√	√	√	√	√	6

Rating items: 1—eligibility criteria and source of participants; 2—random allocation; 3—concealed allocation; 4—baseline comparability; 5—blinded participants; 6—blinded therapists; 7—blind assessors; 8—adequate follow-up; 9—intention-to-treat analysis; 10—between-group comparisons; 11—point estimates and variability. Item 1 evaluates external validity, does not contribute to the total score.

Three studies with adequate data were included in the meta-analysis to evaluate the effect of home-based intervention compared with institution-based intervention.<sup>26 31 36</sup> Two studies measured the performance of BADL at treatment endpoint,<sup>26 31</sup> and one study<sup>26</sup> measured at the follow-up after intervention.<sup>73</sup> One study did the follow-up assessment during the treatment period.<sup>36</sup> For the endpoint analysis, a fixed-effects analysis produced an insignificant result (SMD=0.24; 95% CI=-0.15 to 0.62; n=104; I<sup>2</sup>=0%) (figure 2).

At treatment endpoint, very low-quality evidence indicates the uncertainty of the effect of home-based intervention compared with institution-based intervention. The evidence was downgraded one level for publication bias and two levels for imprecision as the sample size is small and the 95% CI estimated includes both null effect and appreciable benefit or harm.

### Effectiveness of home-based intervention compared with usual care

Fifteen studies investigated the effectiveness of home-based intervention compared with usual care. Ten studies found no significant difference between two groups in the ability to do daily living no matter at treatment endpoints or at follow-ups.<sup>32 38 44 49–54 56</sup> Five studies showed significantly better improvement in the home-based intervention than in the usual care group.<sup>45–48 55</sup>

Six studies were pooled in the meta-analysis to evaluate the effect of home-based intervention compared with usual care.<sup>44 47 48 51 52 54</sup> Four studies measured the performance of BADL at treatment endpoint.<sup>44 47 48 52</sup> For the

endpoint analysis, a random-effects analysis produced an insignificant result with high heterogeneity between studies (SMD=0.62; 95% CI=-0.07 to 1.31; n=475; I<sup>2</sup>=92%) (figure was eliminated). Three studies measured outcome at the follow-up after intervention.<sup>44 52 54</sup> For the follow-up analysis (after intervention), a fixed-effects analysis produced an insignificant result (SMD=0.02; 95% CI=-0.17 to 0.22; n=399; I<sup>2</sup>=0%) (figure 3). Three studies measured outcome during the period of treatment.<sup>47 48 51</sup> For the follow-up analysis (during the treatment period), a random-effects analysis produced an insignificant result with high heterogeneity between studies (SMD=1.03; 95% CI=-0.21 to 2.27; n=1264; I<sup>2</sup>=98%) (figure was eliminated).

At treatment endpoint, very low-quality evidence indicates the uncertainty of the effect of home-based intervention compared with usual care. The evidence was downgraded one level for publication bias, one level for inconsistency because of the heterogeneity between results, and one level for imprecision as the 95% CI estimated includes both null effect and appreciable benefit or harm.

At follow-up after intervention, very low-quality evidence indicates the uncertainty of the effect of home-based intervention compared with usual care. The evidence was downgraded one level for publication bias and two levels for imprecision as the sample size is small and the 95% CI estimated includes both null effect and appreciable benefit or harm.





Figure 2 Forest plot comparing the effectiveness of home-based intervention with institution-based intervention at treatment endpoint.



**Figure 3** Forest plot comparing the effectiveness of home-based intervention with usual care at the follow-up (after treatment).

At follow-up during the intervention period, very low-quality evidence indicates the uncertainty of the effect of home-based intervention compared with usual care. The evidence was downgraded one level for publication bias, one level for inconsistency because of the heterogeneity between results, and one level for imprecision as the 95% CI estimated includes both null effect and appreciable benefit or harm.

### Effectiveness of home-based intervention compared with no intervention

Five studies investigated the effectiveness of home-based intervention compared with no intervention. Four out of five included studies showed significantly greater improvements of BADL in home-based intervention group than in the control group.<sup>57 58 60 61</sup> Three studies demonstrated significant improvements on the BADL in home-based intervention but not in the control group.<sup>57 60 61</sup>

Four studies were pooled in the meta-analysis to evaluate the effect of home-based intervention compared with no intervention.<sup>57-59 61</sup> All of them measured the performance of BADL at treatment endpoint, and a random-effects analysis produced an insignificant result with high heterogeneity between studies ( $\text{SMD}=0.84$ ; 95% CI=-0.38 to 2.05;  $n=231$ ;  $I^2=94\%$ ) (figure was eliminated).

At treatment endpoint, very low-quality evidence indicates the uncertainty of the effect of home-based intervention compared with no intervention. The evidence was downgraded one level for publication bias, one level for inconsistency because of the heterogeneity between results, and two levels for imprecision as the sample size is small and the 95% CI estimated includes both null effect and appreciable benefit or harm.

### Effectiveness of home-based intervention addition to usual care compared with usual care


Ten studies investigated the effectiveness of home-based intervention compared with no intervention. One study

did not specify the between-group statistical comparison.<sup>67</sup> Seven studies found no significant difference between two groups in the ability to do daily living no matter at treatment endpoints or at follow-ups.<sup>63-66 68-70</sup> Two studies demonstrated significantly greater improvement in home-based intervention group than control group.<sup>62 71</sup> Wong and Yeung<sup>71</sup> and Ricauda *et al*<sup>69</sup> found significant improvement in both groups, while Batchelor *et al*<sup>63</sup> showed there was no significant improvement either in home-based intervention group or in the control group.

Four studies were pooled in the meta-analysis to evaluate the effect of home-based intervention addition to usual care compared with usual care.<sup>63 64 70 71</sup> Two studies<sup>64 71</sup> measured the performance of BADL at treatment endpoint, and a fixed-effects analysis produced a significant result ( $\text{SMD}=0.55$ ; 95% CI=0.22 to 0.87;  $n=152$ ;  $I^2=3\%$ ) (figure 4). All of them measured at the follow-up, and a random-effects analysis produced an insignificant result with high heterogeneity between studies ( $\text{SMD}=0.23$ ; 95% CI=-0.16 to 0.62;  $n=545$ ;  $I^2=77\%$ ) (figure was eliminated).

At treatment endpoint, low-quality evidence indicates the home-based intervention addition to usual care may have little or no effect on BADL compared with usual care alone. The evidence was downgraded one level for publication bias and one level for imprecision as the sample size is small.

At follow-up after intervention, very low-quality evidence indicates the uncertainty of the effect of home-based intervention addition to usual care compared with usual care alone. The evidence was downgraded one level for publication bias, one level for inconsistency because of the heterogeneity between results, and one level for imprecision as the 95% CI estimated includes both null effect and appreciable benefit or harm.



**Figure 4** Forest plot comparing the effectiveness of home-based intervention addition to usual care with usual care at the treatment endpoint.

### Subgroup analysis

We were unable to conduct subgroup analyses according to the level of disability of the patients who had a stroke, the onset time of stroke, type of intervention or manner of delivery of the treatment because of the clinical diversity between studies, the insufficient information in the individual study and the limited number of included studies.

### Sensitivity analysis

We were unable to conduct sensitivity analyses because there was no low-quality study included in the meta-analyses.

### Assessment of reporting bias

We were unable to conduct the funnel plot to assess the reporting biases because of the limited number of included studies in each meta-analysis.

## DISCUSSION

Our review found that home-based intervention combined with usual care may have short-term benefits for patients who had a stroke compared with usual care alone. However, the evidence was weak because of the limited number of studies and participants included in the meta-analysis and the possible publication bias. We speculated the intensive dosage of intervention attributes to the effect of home-based intervention combined with usual care. Most included studies conducted the professional-mediated home-based intervention to participants and the usual care was also provided, which ensured the high intensity of therapy to improve the recovery of patients who had a stroke.

There was insufficient evidence to determine the short-term effect of home-based intervention compared with the institution-based intervention, or to determine the long-term effect of home-based intervention compared with the usual care, because of the limited number of studies and participants included in the meta-analyses. There was insufficient evidence to prove the short-term effect of home-based intervention on the performance of activities of daily living in patients who had a stroke, when compared with no intervention. There was insufficient evidence to suggest the short-term effect of home-based intervention compared with usual care, or to suggest the long-term effect of home-based intervention addition to usual care compared with usual care. The heterogeneity between the studies limited the conclusions that could be drawn.

Many current rehabilitation interventions are developed in clinical setting, and some are translated to home. Home-based intervention strategies vary in type, duration, intensity, frequency and delivery manner. This systematic review revealed that exercise physiology practice and training of activity of daily living were commonly performed as home-based interventions which are supported by current evidence. American

Heart Association/American Stroke Association (AHA/ASA) guideline suggested that lower extremity strengthening exercise and cardiovascular exercise are beneficial to improve gait capacity of patients who had a stroke and can also improve their ability to perform gait-related mobility tasks.<sup>74</sup> Moreover, training of activities of daily living is strongly recommended for community-dwelling patients who had a stroke.<sup>75</sup> Some newly developed home-based interventions like caregiver-mediated rehabilitation and telerehabilitation have emerged for the past two decades to replace the traditional home visits by professionals. A Cochrane systematic review found that the caregiver-mediated rehabilitation did not increase the caregiver burden but the effectiveness in the ability to perform BADL in patients who had a stroke was uncertain.<sup>73</sup> One large study found that the lower dose of caregiver-guided rehabilitation training and non-multidisciplinary coordination might decrease the efficacy of caregiver-mediated home-based interventions.<sup>51</sup> Telerehabilitation seemed to be a good alternative to traditional rehabilitation. Chen *et al* delivered the same treatment strategy to home-based telerehabilitation group and institution-based rehabilitation group.<sup>26</sup> At the end of intervention and at follow-up, both groups showed significant improvement in the ability to do activities of daily living, and there was no significant difference between two groups throughout the time. Similarly, when comparing with the traditional face-to-face way of home-based intervention performed by professionals, home-based telerehabilitation showed equal positive effect on enhancing the ability to do activities of daily living of patients who had a stroke.<sup>76</sup> More high-quality studies and practice are required to prove the effectiveness of those new strategies.

### Limitations

There is no sufficient study to determine the effectiveness of home-based intervention compared with other approaches. Although 49 studies were included in the review, many of them did not report adequate data so they were not included in the meta-analyses; therefore, only two to four studies were included in each meta-analysis. Moreover, among the 49 included studies, several studies of the original study and their follow-up study shared the same study population, including two studies conducted by Chaiyawat *et al*,<sup>45 46</sup> four studies conducted by von Koch *et al*<sup>39–42</sup> and two studies conducted by Gladman *et al*.<sup>28 29</sup> Therefore, only 44 trials were included. Nearly half of studies included in the meta-analyses had sample size smaller than 30. The clinical heterogeneity between studies in terms of severity of stroke, onset time of stroke, interventions and manner of delivery also compromised the evidence strength of our meta-analyses. Estimation of publication bias using funnel plots failed because of the insufficient number of included studies in each meta-analysis. Publication bias was possibly increased as we have not searched for the grey literature.



## CONCLUSION

Our finding reveals that the home-based intervention combined with usual care may be more effective than usual care alone for the short-term effect. However, the evidence is weak. Future research with larger sample size is needed to investigate the effectiveness of home-based rehabilitation, including (1) on groups with stratification of stroke severity defined by Brunnstrom stage; (2) on groups with stratification of acute, subacute and chronic stroke; (3) with elaboration of the details of the home-based interventions and the control interventions. Moreover, more high-quality studies are required to prove the cost-effectiveness of newly developed strategies like caregiver-mediated rehabilitation and telerehabilitation.

**Contributors** XJW and PQ designed the review. PQ and CXC screened and selected the study. PQ and CC rated the study quality and extracted the data. PQ analysed the data and drafted the paper. XJW revised the paper. All authors acknowledged and agreed with the format and content of the paper before submission for publication. XJW is the guarantor and responsible for the overall contents of this study.

**Funding** This work is supported by the Medical Research Fund of Guangdong Province (No: A2021041), The Science and Technology Project of Shenzhen Municipality (No: JCYJ20190814112607443) and Seedling Program of Shenzhen Hospital of Southern Medical University (No: 2018MM08).

**Competing interests** None declared.

**Patient and public involvement** Patients and/or the public were not involved in the design, or conduct, or reporting, or dissemination plans of this research.

**Patient consent for publication** Not required.

**Ethics approval** Not applicable.

**Provenance and peer review** Not commissioned; externally peer reviewed.

**Data availability statement** No data are available. Not applicable.

**Supplemental material** This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

**Open access** This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

## ORCID iD

Xijun Wei <http://orcid.org/0000-0002-7643-0475>

## REFERENCES

- Bonita R, Mendis S, Truelsen T, *et al*. The global stroke initiative. *Lancet Neurol* 2004;3:391–3.
- Leys D, Hénon H, Mackowiak-Cordoliani M-A, *et al*. Poststroke dementia. *Lancet Neurol* 2005;4:752–9.
- Veerbeek JM, Kwakkel G, van Wegen EEH, *et al*. Early prediction of outcome of activities of daily living after stroke: a systematic review. *Stroke* 2011;42:1482–8.
- Di Carlo A. Human and economic burden of stroke. *Age Ageing* 2009;38:4–5.
- Allen L, John-Baptiste A, Meyer M, *et al*. Assessing the impact of a home-based stroke rehabilitation programme: a cost-effectiveness study. *Disabil Rehabil* 2019;41:2060–5.
- Dodakian L, McKenzie AL, Le V, *et al*. A home-based Telerehabilitation program for patients with stroke. *Neurorehabil Neural Repair* 2017;31:923–33.
- Siemonsma P, Döpp C, Alpay L, *et al*. Determinants influencing the implementation of home-based stroke rehabilitation: a systematic review. *Disabil Rehabil* 2014;36:2019–30.
- Langhorne P, Bernhardt J, Kwakkel G. Stroke rehabilitation. *Lancet* 2011;377:1693–702.
- Teng J, Mayo NE, Latimer E, *et al*. Costs and caregiver consequences of early supported discharge for stroke patients. *Stroke* 2003;34:528–36 <https://www.ahajournals.org/doi/>
- Langhorne P, Baylan S, Early Supported Discharge Trialists. Early supported discharge services for people with acute stroke. *Cochrane Database Syst Rev* 2017;7:CD000443.
- Coupar F, Pollock A, Legg LA, *et al*. Home-based therapy programmes for upper limb functional recovery following stroke. *Cochrane Database Syst Rev* 2012;5:CD006755.
- Wallace GL. Profile of life participation after stroke and aphasia. *Top Stroke Rehabil* 2010;17:432–50.
- Nishioka S, Wakabayashi H, Nishioka E, *et al*. Nutritional improvement correlates with recovery of activities of daily living among malnourished elderly stroke patients in the convalescent stage: a cross-sectional study. *J Acad Nutr Diet* 2016;116:837–43.
- López-Espuela F, Pedrera-Zamorano JD, Jiménez-Caballero PE, *et al*. Functional status and disability in patients after acute stroke: a longitudinal study. *Am J Crit Care* 2016;25:144–51.
- Doman CA, Waddell KJ, Bailey RR, *et al*. Changes in upper-extremity functional capacity and daily performance during outpatient occupational therapy for people with stroke. *Am J Occup Ther* 2016;70:7003290040p1–7003290040.
- Hillier S, Inglis-Jassiem G. Rehabilitation for community-dwelling people with stroke: home or centre based? A systematic review. *Int J Stroke* 2010;5:178–86.
- Page MJ, Moher D, Bossuyt PM, *et al*. PRISMA 2020 explanation and elaboration: updated guidance and exemplars for reporting systematic reviews. *BMJ* 2021;372:n160.
- Maher CG, Sherrington C, Herbert RD, *et al*. Reliability of the PEDro scale for rating quality of randomized controlled trials. *Phys Ther* 2003;83:713–21.
- Foley NC, Teasell RW, Bhogal SK, *et al*. Stroke rehabilitation evidence-based review: methodology. *Top Stroke Rehabil* 2003;10:1–7.
- Gonzalez GZ, Moseley AM, Maher CG, *et al*. Methodologic quality and statistical reporting of physical therapy randomized controlled trials relevant to musculoskeletal conditions. *Arch Phys Med Rehabil* 2018;99:129–36.
- Higgins JP, Thomas J, Chandler J. Cochrane Handbook for systematic reviews of interventions version 6.3 (updated February 2022), 2022. Cochrane. Available: [www.training.cochrane.org/handbook](http://www.training.cochrane.org/handbook)
- The GRADE Working Group. Grade handbook for grading quality of evidence and strength of recommendations, 2019. Available: <https://gdt.gradepro.org/app/handbook/handbook.html>
- Asano M, Tai BC, Yeo FY, *et al*. Home-based tele-rehabilitation presents comparable positive impact on self-reported functional outcomes as usual care: the Singapore Tele-technology aided rehabilitation in stroke (stars) randomised controlled trial. *J Telemed Telecare* 2021;27:231–8.
- Baskett JJ, Broad JB, Reekie G, *et al*. Shared responsibility for ongoing rehabilitation: a new approach to home-based therapy after stroke. *Clin Rehabil* 1999;13:23–33.
- Björkdahl A, Nilsson AL, Grimby G, *et al*. Does a short period of rehabilitation in the home setting facilitate functioning after stroke? A randomized controlled trial. *Clin Rehabil* 2006;20:1038–49.
- Chen J, Jin W, Dong WS, *et al*. Effects of home-based Telesupervising rehabilitation on physical function for stroke survivors with hemiplegia: a randomized controlled trial. *Am J Phys Med Rehabil* 2017;96:152–60.
- Chen J, Sun D, Zhang S, *et al*. Effects of home-based telerehabilitation in patients with stroke: a randomized controlled trial. *Neurology* 2020;95:e2318–30.
- Gladman JR, Lincoln NB. Follow-up of a controlled trial of domiciliary stroke rehabilitation (domino study). *Age Ageing* 1994;23:9–13.
- Gladman JR, Lincoln NB, Barer DH. A randomised controlled trial of domiciliary and hospital-based rehabilitation for stroke patients after discharge from hospital. *J Neurol Neurosurg Psychiatry* 1993;56:960–6.
- Han D-S, Chuang P-W, Chiu E-C. Effect of home-based reablement program on improving activities of daily living for patients with stroke: a pilot study. *Medicine* 2020;99:e23512.



- 31 Hesse S, Welz A, Werner C, *et al.* Comparison of an intermittent high-intensity vs continuous low-intensity physiotherapy service over 12 months in community-dwelling people with stroke: a randomized trial. *Clin Rehabil* 2011;25:146–56.
- 32 Hofstad H, Gjelsvik BEB, Næss H, *et al.* Early supported discharge after stroke in Bergen (ESD stroke Bergen): three and six months results of a randomised controlled trial comparing two early supported discharge schemes with treatment as usual. *BMC Neurol* 2014;14:239.
- 33 Kalra L, Evans A, Perez I, *et al.* Alternative strategies for stroke care: a prospective randomised controlled trial. *Lancet* 2000;356:894–9.
- 34 Özdemir F, Birtane M, Tabatabaei R, *et al.* Comparing stroke rehabilitation outcomes between acute inpatient and nonintense home settings. *Arch Phys Med Rehabil* 2001;82:1375–9.
- 35 Pandian JD, Felix C, Kaur P, *et al.* Family-Led rehabilitation after stroke in India: the attend pilot study. *Int J Stroke* 2015;10:609–14.
- 36 Redzuan NS, Engkasan JP, Mazlan M, *et al.* Effectiveness of a video-based therapy program at home after acute stroke: a randomized controlled trial. *Arch Phys Med Rehabil* 2012;93:2177–83.
- 37 Roderick P, Low J, Day R, *et al.* Stroke rehabilitation after hospital discharge: a randomized trial comparing domiciliary and day-hospital care. *Age Ageing* 2001;30:303–10.
- 38 Taule T, Strand LI, Assmus J, *et al.* Ability in daily activities after early supported discharge models of stroke rehabilitation. *Scand J Occup Ther* 2015;22:355–65.
- 39 Thorsén A-M, Holmqvist LW, de Pedro-Cuesta J, *et al.* A randomized controlled trial of early supported discharge and continued rehabilitation at home after stroke: five-year follow-up of patient outcome. *Stroke* 2005;36:297–303.
- 40 von Koch L, de Pedro-Cuesta J, Kostulas V, *et al.* Randomized controlled trial of rehabilitation at home after stroke: one-year follow-up of patient outcome, resource use and cost. *Cerebrovasc Dis* 2001;12:131–8.
- 41 von Koch L, Widén Holmqvist L, Kostulas V, *et al.* A randomized controlled trial of rehabilitation at home after stroke in Southwest Stockholm: outcome at six months. *Scand J Rehabil Med* 2000;32:80–6.
- 42 Widén Holmqvist L, von Koch L, Kostulas V, *et al.* A randomized controlled trial of rehabilitation at home after stroke in Southwest Stockholm. *Stroke* 1998;29:591–7.
- 43 Young JB, Forster A. The Bradford community stroke trial: results at six months. *BMJ* 1992;304:1085–9.
- 44 Barzel A, Ketels G, Stark A, *et al.* Home-based constraint-induced movement therapy for patients with upper limb dysfunction after stroke (HOMECIMT): a cluster-randomised, controlled trial. *Lancet Neurol* 2015;14:893–902.
- 45 Chaiyawat P, Kulkantrakorn K. Effectiveness of home rehabilitation program for ischemic stroke upon disability and quality of life: a randomized controlled trial. *Clin Neurol Neurosurg* 2012;114:866–70.
- 46 Chaiyawat P, Kulkantrakorn K, Sritipsukho P. Effectiveness of home rehabilitation for ischemic stroke. *Neurol Int* 2009;1:e10.
- 47 Chen S, Lv C, Wu J, *et al.* Effectiveness of a home-based exercise program among patients with lower limb spasticity post-stroke: a randomized controlled trial. *Asian Nurs Res* 2021;15:1–7.
- 48 Deng A, Yang S, Xiong R. Effects of an integrated transitional care program for stroke survivors living in a rural community: a randomized controlled trial. *Clin Rehabil* 2020;34:524–32.
- 49 Duncan P, Richards L, Wallace D, *et al.* A randomized, controlled pilot study of a home-based exercise program for individuals with mild and moderate stroke. *Stroke* 1998;29:2055–60.
- 50 Lincoln NB, Walker MF, Dixon A, *et al.* Evaluation of a multiprofessional community stroke team: a randomized controlled trial. *Clin Rehabil* 2004;18:40–7.
- 51 Lindley RI, Anderson CS, Billot L, *et al.* Family-led rehabilitation after stroke in India (attend): a randomised controlled trial. *Lancet* 2017;390:588–99.
- 52 Mayo NE, Wood-Dauphinee S, Côté R, *et al.* There's no place like home: an evaluation of early supported discharge for stroke. *Stroke* 2000;31:1016–23.
- 53 Rasmussen RS, Østergaard A, Kjær P, *et al.* Stroke rehabilitation at home before and after discharge reduced disability and improved quality of life: a randomised controlled trial. *Clin Rehabil* 2016;30:225–36.
- 54 Santana S, Rente J, Neves C, *et al.* Early home-supported discharge for patients with stroke in Portugal: a randomised controlled trial. *Clin Rehabil* 2017;31:197–206.
- 55 Walker MF, Gladman JR, Lincoln NB, *et al.* Occupational therapy for stroke patients not admitted to hospital: a randomised controlled trial. *Lancet* 1999;354:278–80.
- 56 Wolfe CD, Tilling K, Rudd AG. The effectiveness of community-based rehabilitation for stroke patients who remain at home: a pilot randomized trial. *Clin Rehabil* 2000;14:563–9.
- 57 Koç A. Exercise in patients with subacute stroke: a randomized, controlled pilot study of home-based exercise in subacute stroke. *Work* 2015;52:541–7.
- 58 Lin J-H, Hsieh C-L, Lo SK, *et al.* Preliminary study of the effect of low-intensity home-based physical therapy in chronic stroke patients. *Kaohsiung J Med Sci* 2004;20:18–22.
- 59 Wade DT, Collen FM, Robb GF, *et al.* Physiotherapy intervention late after stroke and mobility. *BMJ* 1992;304:609–13.
- 60 Walker MF, Drummond A, Lincoln NB. Evaluation of dressing practice for stroke patients after discharge from Hospital: a crossover design study. *Clin Rehabil* 1996;10:23–31.
- 61 Wang T-C, Tsai AC, Wang J-Y, *et al.* Caregiver-mediated intervention can improve physical functional recovery of patients with chronic stroke: a randomized controlled trial. *Neurorehabil Neural Repair* 2015;29:3–12.
- 62 Azab M, Al-Jarrah M, Nazzal M, *et al.* Effectiveness of constraint-induced movement therapy (CIMT) as home-based therapy on Barthel index in patients with chronic stroke. *Top Stroke Rehabil* 2009;16:207–11.
- 63 Batchelor FA, Hill KD, Mackintosh SF, *et al.* Effects of a multifactorial falls prevention program for people with stroke returning home after rehabilitation: a randomized controlled trial. *Arch Phys Med Rehabil* 2012;93:1648–55.
- 64 Chumbler NR, Quigley P, Li X, *et al.* Effects of telerehabilitation on physical function and disability for stroke patients: a randomized, controlled trial. *Stroke* 2012;43:2168–74.
- 65 Corr S, Bayer A. Occupational therapy for stroke patients after hospital discharge — a randomized controlled trial. *Clin Rehabil* 1995;9:291–6.
- 66 Gilbertson L, Langhorne P, Walker A, *et al.* Domiciliary occupational therapy for patients with stroke discharged from Hospital: randomised controlled trial. *BMJ* 2000;320:603–6.
- 67 Goldberg G, Segal ME, Berk SN, *et al.* Stroke transition after inpatient rehabilitation. *Top Stroke Rehabil* 1997;4:64–79.
- 68 Mandigout S, Chaparro D, Borel B, *et al.* Effect of individualized coaching at home on walking capacity in subacute stroke patients: a randomized controlled trial (Ticac' dom). *Ann Phys Rehabil Med* 2021;64:101453.
- 69 Ricauda NA, Bo M, Molaschi M, *et al.* Home hospitalization service for acute uncomplicated first ischemic stroke in elderly patients: a randomized trial. *J Am Geriatr Soc* 2004;52:278–83.
- 70 Rudd AG, Wolfe CD, Tilling K, *et al.* Randomised controlled trial to evaluate early discharge scheme for patients with stroke. *BMJ* 1997;315:1039–44.
- 71 Wong FKY, Yeung SM. Effects of a 4-week transitional care programme for discharged stroke survivors in Hong Kong: a randomised controlled trial. *Health Soc Care Community* 2015;23:619–31.
- 72 Hoffmann TC, Glasziou PP, Boutron I, *et al.* Better reporting of interventions: template for intervention description and replication (TIDieR) checklist and guide. *BMJ* 2014;348:g1687.
- 73 Vloothuis JD, Mulder M, Veerbeek JM, *et al.* Caregiver-mediated exercises for improving outcomes after stroke. *Cochrane Database Syst Rev* 2016;12:CD011058.
- 74 Winstein CJ, Stein J, Arena R, *et al.* Guidelines for adult stroke rehabilitation and recovery: a guideline for healthcare professionals from the American heart Association/American stroke association. *Stroke* 2016;47:e98–169.
- 75 Stroke Foundation. *Clinical Guidelines for Stroke Management 2017*. Melbourne Australia, 2017.
- 76 Fordeucey PG, Glueckauf RL, Bergquist TF, *et al.* Telehealth for persons with severe functional disabilities and their caregivers: facilitating self-care management in the home setting. *Psychol Serv* 2012;9:144–62.