

CHEMISTRY

Direct transformation of dinitrogen: synthesis of *N*-containing organic compounds via N–C bond formation

Ze-Jie Lv¹, Junnian Wei¹, Wen-Xiong Zhang¹, Ping Chen², Dehui Deng², Zhang-Jie Shi³ and Zhenfeng Xi ^{1,*}

ABSTRACT

N-containing organic compounds are of vital importance to lives. Practical synthesis of valuable *N*-containing organic compounds directly from dinitrogen (N₂), not through ammonia (NH₃), is a holy-grail in chemistry and chemical industry. An essential step for this transformation is the functionalization of the activated N₂ units/ligands to generate N–C bonds. Pioneering works of transition metal-mediated direct conversion of N₂ into organic compounds via N–C bond formation at metal-dinitrogen [N₂-M] complexes have generated diversified coordination modes and laid the foundation of understanding for the N–C bond formation mechanism. This review summarizes those major achievements and is organized by the coordination modes of the [N₂-M] complexes (end-on, side-on, end-on-side-on, etc.) that are involved in the N–C bond formation steps, and each part is arranged in terms of reaction types (*N*-alkylation, *N*-acylation, cycloaddition, insertion, etc.) between [N₂-M] complexes and carbon-based substrates. Additionally, earlier works on one-pot synthesis of organic compounds from N₂ via ill-defined intermediates are also briefed. Although almost all of the syntheses of *N*-containing organic compounds via direct transformation of N₂ so far in the literature are realized in homogeneous stoichiometric thermochemical reaction systems and are discussed here in detail, the sporadically reported syntheses involving photochemical, electrochemical, heterogeneous thermo-catalytic reactions, if any, are also mentioned. This review aims to provide readers with an in-depth understanding of the state-of-the-art and perspectives of future research particularly in direct catalytic and efficient conversion of N₂ into *N*-containing organic compounds under mild conditions, and to stimulate more research efforts to tackle this long-standing and grand scientific challenge.

Keywords: dinitrogen transformation, metal-dinitrogen complex, N–C bond formation, *N*-containing organic compounds

INTRODUCTION

As the most abundant constituent in Earth's atmosphere (atm), dinitrogen (N₂) is the main nitrogen source of *N*-containing compounds on the Earth. Therefore, N₂ fixation and activation are essential both for nature and humans. Nevertheless, the high bond dissociation energy (942 kJ/mol) and large highest occupied molecular orbital (HOMO)—lowest unoccupied molecular orbital (LUMO) gap (10.82 eV) make N₂ exhibit extremely low reactivity and be regarded as an inert gas. Currently, the N₂ fixation and conversion in nature and industry mainly

rely on two pathways, in which ammonia (NH₃) is the product [1]. In nature, nitrogenase metalloenzymes employ iron-sulfur clusters as the key cofactor (FeMo, FeV or FeFe cofactor) and water as the proton source to transfer N₂ into NH₃ at ambient temperature and pressure [2]. This biosynthetic NH₃ is a versatile precursor for the synthesis of *N*-containing organic compounds, such as amino acids and nucleic acids. Although the precise biological N₂ reduction mechanism is still controversial, spectroscopic and computational studies suggested the presence of an interstitial carbon atom at the center of the FeMo and FeV cofactors [3–5].

¹Beijing National Laboratory for Molecular Sciences (BNLMS), Key Laboratory of Bioorganic Chemistry and Molecular Engineering of Ministry of Education, College of Chemistry, Peking University, Beijing 100871, China;

²Dalian Institute of Chemical Physics, Chinese Academy of Sciences, Dalian 116023, China and

³Department of Chemistry, Fudan University, Shanghai 200433, China

*Corresponding author. E-mail: zfxi@pku.edu.cn

Received 28 May 2020; Revised 21 June 2020; Accepted 21 June 2020

In industry, more than 170 million metric tons of NH_3 is produced from the Haber-Bosch process annually, in which N_2 reacts with dihydrogen (H_2) under the harsh condition in the presence of iron or ruthenium catalysts. This NH_3 synthesis process consumes 1–2% of the world's annual energy supply along with the huge CO_2 emission, due to the drastic reaction condition and the energy requirement for H_2 production from fossil fuels and water [6]. As the main route of N_2 fixation and transformation in industry, ~20% of NH_3 produced from the Haber-Bosch process is used as the feedstock to produce *N*-containing chemicals, including higher-value *N*-containing organic compounds, like amines, nitriles, nitro and so on. To better understand the reaction mechanism of biological and industrial reduction of N_2 into NH_3 , several catalytic systems including homogeneous molecular systems, electrochemical systems and heterogeneous systems have been studied for decades, and there are comprehensive reviews that readers may refer to [7–13].

Compared to NH_3 -based N_2 fixation process, an alternative route of N_2 fixation is the direct conversion of N_2 into *N*-containing organic compounds under mild condition. This approach is always targeted because it provides the potential solution to developing a sustainable system with reduced fossil fuel requirements. The earliest study towards this goal began in the 1960s, when Vol'pin *et al.* discovered that the titanium species, for example, Cp_2TiCl_2 could react with PhLi under N_2 to give aniline after hydrolysis [14]. However, further application of this reaction was hindered by the low yields and the lack of reaction details. During the same period, the first metal-dinitrogen (N_2 -*M*) complex $[\text{Ru}(\text{NH}_3)_5(\text{N}_2)]^{2+}$ was reported in 1965 [15]. After that, thousands of N_2 -*M* complexes have been documented [16]. The reactivity exploration reveals that the functionalization of the N_2 ligands can also be fulfilled for some N_2 -*M* complexes [17]. Making *N*-C bonds from the reactions of transition metal N_2 complexes with carbon-based reagents has received much attention in recent decades, although the catalysis system has not been realized [18,19].

This review will focus on the previous works regarding the transformation of N_2 into organic compounds. In almost all of these works, the *N*-C bond formation steps are fulfilled upon the well-defined N_2 -*M* complexes with diversified coordination modes. This review is organized by the coordination modes of the N_2 -*M* complexes (to clarify, the N_2 -derived metal nitrides are also considered as a coordination mode of N_2 -*M* complexes) that





Figure 1. The classification of this review. The *N*-C bond formation is reported (✓) or not reported (✗).

are involved in the *N*-C bond formation steps, and each part is arranged in terms of the type of reactions between N_2 -*M* complexes and carbon-based substrates. The earlier works about one-pot synthesis of organic compounds from N_2 via ill-defined intermediates are also introduced briefly in this review (Fig. 1).

N-C BOND FORMATION VIA END-ON N_2 -*M* COMPLEXES

End-on bond is the most prevalent bonding mode for N_2 -*M* complexes and the N_2 -*M* complexes with this binding mode have been known to assemble *N*-C bond for a long time. Main works were achieved via the reaction of end-on terminal N_2 -*M* complexes with alkyl or acyl halides and their analogues. *N*-C bond formations from the cycloaddition and insertion reactions of end-on-bridged N_2 -*M* complexes with imido-like N_2 ligands have also been reported.



Scheme 1. *N*-alkylation of end-on terminal N₂-Mo complexes by electrophiles. (a) *N*-methylation of N₂-Mo complex by MeOTs to afford methyldiazenido complex. (b) *N*-methylation of N₂-Mo complexes by MeOTs and MeOTf to afford methyldiazenido and *N,N*-dimethylhydrazido complexes.

N-alkylation


N-alkylation by electrophiles

The strong electrophilic alkyl triflates and their analogues are often employed to functionalize the end-on N₂-M complexes because the N₂ ligands in these complexes feature a nucleophilic character by electron donation from the electron-rich metal centers. Peters *et al.* [20] and Greco and Schrock [21] reported that the methylation reaction occurs when the anionic end-on N₂-Mo complexes **1** and **3** are treated with methyl tosylate (MeOTs) to provide methyldiazenido complexes **2** and **4** (Scheme 1a and b). Additionally, **4** could further undergo N–C bond formation to furnish *N,N*-dimethylhydrazido complexes **5** by reaction with excess methyl triflate (MeOTf) or MeOTs (Scheme 1b).

Although many late-transition metal complexes with end-on N₂ ligands have been documented, reports on their reactivity toward electrophiles to make N–C bond are very rare. Peters *et al.* described that the anionic end-on terminal N₂ complexes of Fe **6** and Co **8** react with MeOTs to give *N*-methylation species **7** and **9** (Scheme 2a) [22]. In 2016, the same group found that the modified N₂-Fe complex **10** bearing monoanionic tetradentate trisphosphinosilyl ligand can also be alkylated to afford *N,N*-dimethylated product **11** (Scheme 2b) [23].


N-alkylation by *in situ* formed radicals

There are only a few examples of N–C bond formation at N₂-M complexes by radicals. One ex-



Scheme 2. *N*-alkylation of end-on terminal N₂-Fe, Co complexes by MeOTf or MeOTs. (a) *N*-methylation of N₂-Fe, Co complexes by MeOTs to afford methyldiazenido complexes. (b) *N*-methylation of N₂-Fe complexes by MeOTf to afford *N,N*-dimethylhydrazido complex.

ample is the reactions of terminal end-on N₂-Mo, W complexes **12** with alkyl halides, driven by light (*vide infra*). Mechanism investigation reveals that the radicals in these reactions are generated *in situ* by the homolysis of the alkyl halides within the coordination sphere. The attacking of these alkyl radicals at the N₂ ligands provides **13**. Furthermore, dialkylhydrazido complexes **14**, **15** and **16** can also be obtained via alkylation of **13** or one-pot dialkylation of **12** (Scheme 3a) [24,25]. It is noteworthy that if the diphosphine ligands in **12** are replaced by the monophosphine ligands, the corresponding N₂-Mo, W complexes fail to react with alkyl halides to assemble N–C bond. Another example that involves the radical mechanism is the *N*-functionalization of the terminal end-on N₂-Mo complex **17**, which possesses higher reactivity than **12** (Scheme 3b) [26]. For instance, the treatment of **17** with BnBr or aryl iodide gives the *N,N*-dibenzyl product **18** or *N*-arylation complex **19**, the latter of which can also be converted to the organo-hydrazido species **20** by further reaction with MeI. More intriguingly, when **17** is treated with MeI in toluene, the prospective product **21** is formed together with isolation of an unexpected product **22**. A plausible mechanism is raised for the generation of **22** (Scheme 3c). The initial reaction between **17** and MeI results in iodine atom abstracting to afford intermediate **A** and the methyl radical, which would abstract an *H*-atom from toluene to yield benzyl radical. The latter reaction between **A** and benzyl radical gives the *N*-benzylation intermediate **B**, which can further react with MeI to afford the final product. The formation of **22** confirms




Scheme 3. *N*-alkylation of end-on terminal $\text{N}_2\text{-Mo}$, W complexes by *in situ* formed radicals. (a) *N*-alkylation of $\text{N}_2\text{-Mo}$, W complexes supported by diphosphine ligands. (b) *N*-alkylation of $\text{N}_2\text{-Mo}$ complex supported by tetra-thioether ligand. (c) A plausible mechanism for the generation of **22**.


the radical process of these *N*-alkylation reactions again.

N-alkylation by nucleophiles

For end-on terminal $\text{N}_2\text{-M}$ complexes, simple Lewis formulas could be used to depict their structures. As shown in Scheme 4a, the *N* atom adjacent to the metal atom (N_α) features positive charge and could be attacked by nucleophiles in theory. Surprisingly, there is only one example of this reactivity for $\text{N}_2\text{-M}$ complexes [27,28]. Sellman *et al.* found that an end-on terminal $\text{N}_2\text{-Mn}$ complex **23** reacts with methyl or phenyl lithium reagent at low temperature to give the N_α -functionalized products **24**, which could subsequently react with Meerwein reagent Me_3OBF_4 upon N_β atom to afford **25**. This



Scheme 4. Manganese-promoted direct conversion of N_2 into azomethane via the reaction between nucleophiles and $\text{N}_2\text{-Mn}$ complex. (a) Simple Lewis formulas for end-on terminal $\text{N}_2\text{-M}$ complexes. (b) A synthetic cycle for synthesis of azo-compound from N_2 .



Scheme 5. *N*-acylation of end-on terminal $\text{N}_2\text{-Mo}$, $\text{N}_2\text{-W}$ complexes.


azomethane complex would ultimately liberate free azomethane by pressuring with 100 bar of N_2 along with reforming $\text{N}_2\text{-Mn}$ complex **23**. Thus, a synthetic cycle was raised for synthesis of azo-compound from N_2 .

N-acylation

Besides alkyl halides, acyl chlorides are also used to functionalize end-on $\text{N}_2\text{-M}$ complexes. Chatt *et al.* found that the $\text{N}_2\text{-Mo}$, W complexes **12** supported by bidentate phosphines ligands react with acyl chloride to afford acyldiazenido complexes (Scheme 5) [29,30]. These *N*-acylation reactions possibly proceed through nucleophilic attacking of the N_2 ligands on the acyl carbons.

Cycloaddition and insertion


For some end-on-bridged $\text{N}_2\text{-M}$ complexes with strongly activated N_2 ligands, the imido-like structures make them able to undergo cycloaddition or insertion reactions with carbon-based



Scheme 6. N–C bond formation from cycloaddition of end-on-bridged N_2 -Ti complex with phenylallene, $tBuNCO$ and CO_2 .

unsaturated substrates to assemble N–C bond. In 2017, the reaction of an end-on bridging binuclear N_2 -Ti complex **27** with phenylallene, *tert*-butyl isocyanate ($tBuNCO$) and CO_2 was investigated by Kawaguchi *et al.*, to provide N–C bond formation products (Scheme 6) [31]. Treatment of **27** with an excess of phenylallene results in the formation of dititanium hydrazido complexes **28** and **29** as a mixture of isomers. The formation of **28** and **29** can be rationalized in terms of an initial [2+2] cycloaddition of phenylallene with Ti = N bond in **27** to give the 4-membered titanacycle intermediates (two isomers), and the subsequent protonolysis of the Ti–C bonds in these intermediates to give the final products. Further studies indicate that the proton source in this reaction could be a second equivalent of phenylallene, the ancillary ligands, or even adventitious impurities present in the reaction mixture. The reaction of **27** with $tBuNCO$ also proceeds through a formal [2+2] cycloaddition reaction to afford **30**. However, when **27** is introduced with an atm of CO_2 , the insertion of three molecules of CO_2 into Ti = N bonds in **27** is achieved to furnish **31**. By adding an excess amount of $TMSCl$, **31** could be converted to organic compound $N_2(TMS)(CO_2TMS)_3$, which is unstable under the reaction condition and decomposes to two hydrazine derivatives $[TMS(CO_2TMS)N]_2$ and $(TMS)_2NN(CO_2TMS)_2$ via decarboxylation.


The cycloaddition reactions between group 5 end-on N_2 -M complexes and carbon-based unsaturated bonds have also been observed. For example, N_2 -Nb complex **32** and N_2 -Ta complex **33** with di-imido bridging N_2 ligands are known to react with aldehyde and acetone to afford the corresponding ketazines (Scheme 7) [32,33].



Scheme 7. N–C bond formation from the reactions of end-on-bridged N_2 -Nb, Ta complexes with aldehyde or acetone. (a) The reaction of N_2 -Nb complex with benzaldehyde. (b) The reaction of N_2 -Ta complex with acetone.

N-protonation/carbonylation

An alternative route for making N–C bond is the treatment of carbon-based substrates with the *N*-hydrogenated complexes derived from N_2 because in some cases *N*-hydrogenation are more accessible than *N*-alkylation for end-on N_2 -M complexes. Seminal works about these transformations were finished by Hidai and others [25,34]. They reported that the N_2 -Mo, W complexes **34** and **12** supported by monophosphine or diphosphine ligand react with HX ($X = Cl, Br$ and I) or HBF_4 to afford the hydrazido complexes **35** and **36**, which can act as the versatile precursors to construct N–C bond [25,35] (Scheme 8a). For instance, **35** could react with diphenylketene and phthalaldehyde to provide **37** and **38**, while the reaction between **36** and succinyl chloride gives rise to **39**. More intriguingly, these hydrazido complexes **35** and **36** are also reported to undergo a condensation reaction with ketones and aldehydes in the presence of catalytic amounts of acid to afford all kinds of diazoalkane




Scheme 8. N-C bond formation from the reactions of hydrazido Mo, W complexes with carbon-based reagents. (a) The reaction of N_2 -Mo, W complexes with HX (X = Cl, Br and I) or HBF_4 to afford the hydrazido complexes **35** and **36** to assemble N-C bond. (b) Carbonylation of hydrazido complexes **35** and **36** to assemble N-C bond. (c) A synthetic cycle for synthesis of 1H-pyrrole from N_2 .

complexes **40** and **41** (Scheme 8b). The liberation of the N-containing organic compounds from these N-functionalized complexes has also been explored [36]. For example, when the cyclic hydrazido complex **42**, produced from the reaction of **36** and the cyclic acetal of succinaldehyde, is treated with $LiAlH_4$, the reductive destruction of **42** is observed to release 1H-pyrrole accompanied by the generation of the tetrahydride complex **43**. Furthermore, this tetrahydride tungsten complex could be converted to the initial N_2 -W complex **12** under photolytic conditions to achieve a cycle (Scheme 8c) [37].

Involvement of photochemistry

Photochemistry is an emerging approach for the transfer of N_2 . The earliest observation of photocatalyzed N-C bond formation of N_2 -M complexes



Scheme 9. An electrochemical cycle for synthesis of piperidine direct from N_2 via end-on terminal N_2 -Mo, W complexes.

is of the reactions between end-on terminal N_2 -M complexes **12** and alkyl halides (Scheme 3a) [25]. In the case of N_2 -W complex, visible light or a tungsten lamp is often necessary for these N-alkylation reactions. However, for the N_2 -Mo complex, it could react with alkyl bromide slowly in the dark. It is also reported that the N_2 ligands in **12** are not evolved in the absence of the alkyl halides since irradiation of the N_2 -M complexes without organic halide caused no change. These results indicate the possibility of photo engaging in the assistance of alkyl radicals formation in these reactions [24].

Involvement of electrochemistry

Besides photochemistry, electrochemistry is another versatile method in the N_2 conversion process. Although the direct involvement of electrochemistry in the N-C bond formation step has not been discovered, the electrochemical reduction of the N-alkylated complexes to release the final organic products has been developed. For example, the organohydrazido complexes **16**, which is synthesized from the reaction of N_2 -M complexes **12** with 1,5-dibromopentane, undergoes electrochemical reduction at a Pt electrode in tetrahydrofuran (THF) under N_2 by using $[NBu_4][BF_4]$ as the electrolyte to liberate piperidine accompanied by the regeneration of N_2 -M complexes **12**. According to the control experiment under the atmosphere of Ar or CO, a M(II) hydrazido intermediate is proposed in this piperidine releasing process. Based on these results, an electrochemical cycle to synthesize dialkylhydrazine from N_2 was reported by Leigh *et al.* (Scheme 9) [38].

N-C BOND FORMATION VIA SIDE-ON N_2 -M COMPLEXES


The side-on bonding modes are often observed at group 3 and group 4 transition metal N_2 -M complexes. The N_2 ligands in these side-on N_2 -M

complexes can be divided into neutral N_2 , $(N_2)^{2-}$, $(N_2)^{3-}$ and $(N_2)^{4-}$, and the N–C bond formation usually takes place at side-on-bridged binuclear N_2 -M complexes with $(N_2)^{4-}$ moiety. Very recently, the N-functionalization of a $(N_2)^{3-}$ -Sc complex was also fulfilled.


N-alkylation

There are two reports about the reaction of group 4 side-on N_2 -M complexes with alkyl halides or their analogues to make N–C bond. One example was reported by Hirotsu *et al.* in 2007, in which the side-on-bridged N_2 -Hf complexes **44** with extremely activated $(N_2)^{4-}$ ligands can react with ethyl bromide (EtBr) to provide the N-ethylated products **45** (Scheme 10a) [39]. Controlled experiments indicate that this reaction is remarkably sensitive to the steric effects of the ancillary ligands. For example, when the R' group in **44b** is changed from Et to ⁱPr, the corresponding N-ethylation product could not be obtained. Besides, **45a** and **45b** fail to undergo further N-ethylation, even in the presence of excess EtBr. The other work is reported by the reaction of methyl triflate (MeOTf) with a hafnocene complex **46** that also bears side-on bridging $(N_2)^{4-}$ ligand (Scheme 10b) [40]. This reaction offers a mixture of products and one of them is the N_2 ligand monomethylated product **47**, which could be converted to the final organic compound N-methylhydrazine by treating with excess HCl. Besides, an unprecedented triflate hafnocene hydrazonato complex **48** is generated via a second N–C bond formation when additional MeOTf is added to **47**.

Compared with the group 4 transition metals, rare-earth metal promoted direct conversion of N_2 into organic compounds attracts less attention. The only example of this topic was reported by Xi, Zhang *et al.* in 2019 (Scheme 11) [41]. Treatment of the $(N_2)^{3-}$ -bridged discandium complex **49** with MeOTf leads to the formation of N,N'-dimethylation discandium complex **50** in 43% yield. The yield of **50** can be improved via the treatment of **49** with MeOTf and potassium several times. Transformation of the $(N_2Me_2)^{2-}$ ligand into organic compounds could be accomplished by treatment of **50** with I_2 , HCl, BnBr and acyl chloride to afford azomethane, 1,2-dimethylhydrazine and a series of tetra-substituted hydrazine derivatives, concomitant with the regeneration of the precursors of the N_2 -Sc complexes. Hence, a three-step synthetic cycle for scandium-mediated direct conversion of N_2 and carbon-based electrophiles to multi-substituted hydrazine derivatives could be realized. The insertion of a CO molecule into the Sc–N bond of **50**



Scheme 10. N-alkylation of side-on-bridged N_2 -Hf complexes by EtBr or MeOTf. (a) N-ethylation of N_2 -Hf complex by EtBr. (b) N-methylation of N_2 -Hf complex by MeOTf.




Scheme 11. Scandium-promoted direct conversion of N_2 into hydrazine derivatives via the reaction between MeOTf and N_2 -Sc complex.

with further N–C bond formation is also observed to provide **51**.

Cycloaddition and insertion


The group 4 side-on bridging N_2 -M complexes are known to undergo cycloaddition and insertion reactions with carbon-based reagents that contain C = X (X = N, O) bonds or C≡C bond, such as carbon dioxide (CO_2), isocyanates (RNCO) or alkynes, owing to their imido-like reactivity. Compared with N-alkylation and acylation reactions, these cycloaddition and insertion reactions are more atom-efficient for N_2 functionalization,



Scheme 12. N–C bond formation of side-on N₂-Zr complex by reaction with alkynes.


because the formation of transition metal halides and inorganic salts as the by-products is avoided in these reactions. The earliest study of the cycloaddition reactions between group 4 N₂-M complexes and unsaturated bond to assemble N–C bond was finished by Fryzuk *et al.* via the reaction of side-on bridging N₂-Zr complex **53** with arylacetylene (RC≡CH; R = Ph, 4-Me-C₆H₄ and 4-t-Bu-C₆H₄) (Scheme 12) [42]. The N-functionalization products **54** may result from a sequence of two successive steps: cycloaddition of alkyne across a Zr–N bond in **53** leading to the zircona-aza-cyclobutene intermediate **D**, which subsequently encounters Zr–C bond cleavage by protonation with another molecular terminal alkyne to yield **54**.

By elegant modulation of the substitutions on the multi-substituted Cp ligands, Chirik *et al.* accomplished a series of reactions of dinuclear N₂-Zr, Hf complexes with isocyanates or CO₂ to assemble N–C bond. N₂-Hf complex **46** bearing tetramethylcyclopentadienyl (Cp^{4Me}) ligand is reported to react with PhNCO to provide the initial product **56** via a possible intermediate **55**. In the solution, **56** also reacts quickly with another molecule of ArNCO (Ar = Ph and *p*-MeC₆H₄) to afford **57**, which could also be prepared directly from **46** (Scheme 13a) [43]. Besides, further studies indicate that another N-functionalization product **58**, in which the same nitrogen atom is di-carboxylated, would be formed predominately when CO₂ is bubbled into a solution of **46** (Scheme 13b) [44]. Subsequent reaction of **58** with TMSI gives rise to the generation of **59**, which is known to liberate the corresponding hydrazine derivative (TMS)₂NN(CO₂TMS)₂ by further reacting with excess TMSI. Unfortunately, the similar N-functionalization reactions of PhNCO and CO₂ with zirconium congener of **46** are unsuccessful, which is believed to be caused by the deleterious ligand-induced side-on, end-on isomerization of the (N₂)⁴⁻ ligand. Hence, a [Me₂Si]-bridged *ansa*-zirconocenes N₂ complex **60** with higher energy barrier for the side-on, end-on isomerization was designed and prepared to investigate the reactivity toward CO₂ [45]. The treatment of **60** with



Scheme 13. N–C bond formation from the reactions of the side-on N₂-Zr, Hf complexes with isocyanates and CO₂. (a) The reaction of N₂-Hf complex with PhNCO. (b) The reaction of N₂-Hf complex with CO₂. (c) The reaction of N₂-Zr complex with CO₂.

CO₂ leads to the immediate generation of **61**, where N₂-functionalization takes place at each N-atom. Organic compound N,N'-dicarboxylated hydrazine can be released from **61** by reacting with TMSI. Furthermore, a second N–C bond formation occurs when **61** is treated with MeOTf to provide **62**, which is known to liberate region-specific hydrazine



Scheme 14. CO-induced N_2 scission and functionalization at side-on N_2 -Zr and Hf complexes. (a) The reaction of N_2 -Zr, Hf complexes and CO. (b) A plausible mechanism for this CO-induced N_2 scission and functionalization reaction.

$X(COOMe)NN(COOMe)Me$ ($X = H, TMS$) by further reacting with H_2O or TMSI (Scheme 13c). These results indicate that small modifications of the ligands will change the reactivity of the N_2 -M complexes dramatically.

CO-induced N_2 functionalization and cleavage


Being isoelectronic with N_2 , CO is an abundant and cheap diatomic molecule with BDE of 1 079 kJ/mol. Hence, the transformation of CO and N_2 into N—C bond is a challenge but a significant process. Until now, only two systems of CO-induced N_2 ligand scission and functionalization at N_2 -M complexes have been developed, in which all of the N_2 ligands adopt side-on bridging coordination mode.

Following their earlier work on N—C bond formation from N_2 -Zr and Hf complexes, Chirik *et al.* reported in 2010 the first example that treatment of the *ansa*-zirconocene and hafnocene N_2 complexes **60** and **63** with 4 atm or 1 atm of CO leads to the generation of the dinuclear oxamide complexes **64** and **65** as two isomers [46,47]. Besides, when **63** is treated with less CO (1.5 equiv), a new product of imido-bridged dihafnium complex **66** could be isolated, in which the *H*-atom on the bridging imido is derived from the cyclometallation of the *t*Bu group (Scheme 14a). Protonolysis enables these products to release the corresponding *N*-containing organic compounds of free oxamide and isocyanic acid.

Density functional theory (DFT) calculations [48] and experimental results [49] reveal that the formation of **64** and **65** is assumed to be initiated by CO insertion into an Hf—N bond and followed by the retro [2+2] cycloaddition to provide the presumptive μ -nitride species **67**. The coordination and insertion of CO to the μ -nitrido intermediate **67** results in the formation of **68**, which was characterized by multinuclear nuclear magnetic resonance (NMR) spectroscopy at low temperature. **68** undergoes C—C bond formation via coupling of the terminal and bridging isocyanate units along with the loss of the terminal carbonyl ligand to give the final products (Scheme 14b).


More studies indicate that these CO-induced N_2 cleavage and functionalization reactions are also compatible with other zirconocene and hafnocene N_2 complexes. Therefore, a tetrametallic hafnocene oxamide complex **71** could be obtained via a dimeric hafnium intermediate **70** when the N_2 -Hf complex **69** is treated with CO [50]. The transformations of these oxamide complexes were also elaborated. Thermolysis of **71** at 110°C provides a μ -oxo hafnocene complex **72** with both terminal cyanide and isocyanate ligands that undergoes preferential group transfer of the cyanide unit to liberate organonitriles of TMSCN or MeCN along with the generation of **73** by reacting with TMSI or MeOTf (Scheme 15a) [50]. Oxamide complex **64** reacts with CO_2 and *t*BuNCO to give the formal [2+2] cycloaddition products **74** and **75** [51] (Scheme 15b). Additionally, various free *N,N'*-dialkyloxamide could be formed via stepwise *N*-alkylation of the oxamide complex **76** and following protonolysis with HCl or ethanol (Scheme 15c).

The characterization and reactivity studies of the μ -nitride intermediates were also developed (Scheme 16). Rapid bubbling of CO into N_2 -Hf complex **69** at a low temperature produces a metastable dihafnocene nitride complex **79**, which is characterized by IR and multinuclear NMR spectroscopy. This base-free μ -nitride can react with various substrates [52–55]. For instance, the treatment of **79** with TMSI affords silylureate complex **80**. This reaction is involved in the initial iodide ion abstraction to give a transient silyl cation and a formally anionic bridging nitride intermediate, whose nucleophilicity is increased by weakening the Hf—N multiple bonding. Hence, this intermediate could undergo nucleophilic attacking of the nitride group to the terminal isocyanate moiety to form the ureate core, which is then trapped by the silyl cation to yield **80**. Besides, when **79** is treated with cyclooctyne, monosubstituted allenes and isocyanates, the formal [2+2] cycloaddition reactions occur to



Scheme 15. N–C bond formation of CO- and N₂-derived oxamidate complexes. (a) Thermolysis of oxamidate complex **71**. (b) The reaction of oxamidate complex **64** with CO₂ and *t*BuNCO. (c) *N*-alkylation of the oxamidate complex **76** to afford *N,N*-dialkyloxamidates.


afford **81**, **84** and **85**, respectively. The alkyne and isocyanates products are kinetically unstable at elevated temperature and engage in additional N–C bond formations to give **82** and **86**. Complex




Scheme 16. N–C bond formation of CO- and N₂-derived Hf-nitride complexes.

82 could be converted to a binuclear complex **83** with two bridging carbodiimidyl ligands by reacting with TMSCl. The μ -oxo complex **86** can liberate *N*-containing organic compound of carbodiimide along with the generation of dihafnium oxo complex **73** by reacting with MeOTf. In contrast, exposure of the nitride complex **79** to another heterocumulene of CO₂ provides μ -oxo bis(isocyanate) complex **87**, resulting from deoxygenation of CO₂ accompanied by N–C bond formation. Furthermore, the Hf–nitride bond in **79** also engages in the insertion of cyclohexylnitrile (CyCN) to provide **88**, which can continue reacting with another molecule of CyCN to afford bridging ureate-type complex **89** via additional N–C bond formation (Scheme 16).

The other system of CO-induced N₂ scission and functionalization was discovered by Mazzanti *et al.* using uranium complexes. A side-on-bridged binuclear N₂-U complex **90** with μ -nitride ligands reacts with CO to provide the oxo/cyanate diuranium complex **91** accompanied by releasing of potassium cyanate (KCN), which is formed from the reaction of nitride unit with CO [56] (Scheme 17a). To understand the role of the bridging nitride in these transformations, a similar N₂-U complex **92** with bridged μ -oxo ligand was synthesized and its reactivity toward CO was also investigated [57]. The reaction between **92** and CO immediately results in the generation of cyanamido bridged complex **93** with retaining the μ -oxo moiety via both cleavages of N–N single bond and C \equiv O triple bond (Scheme 17b). DFT calculation indicates that the different reactivity of **90** and **92** is attributed to the different bonding nature of the N₂ ligands, in which the μ -nitride is involved in the binding and resultant activation of RT, but the μ -oxo is not.



Scheme 17. CO-induced N_2 scission and functionalization at side-on N_2 -U complexes. (a) The reaction between CO and N_2 -U complex **90** with μ -nitride ligand. (b) The reaction between CO and N_2 -U complex **92** with μ -oxo ligand.



Scheme 18. *N*-alkylation of side-on-end-on N_2 -Ta complex by BnBr.

N—C BOND FORMATION VIA SIDE-ON-END-ON N_2 -M COMPLEXES


The side-on-end-on bound mode is much less common relative to the aforementioned two coordination modes in N_2 -M complexes. All of the work regarding the making of N—C bond from N_2 -M complex with this bonding mode were finished by Fryzuk *et al.* by employing a binuclear N_2 -Ta complex.

N-alkylation

In 2001, the *N*-alkylation of the side-on-end-on bridging binuclear N_2 -Ta complex **94** was developed to afford *N*-benzilation product **95** in high yield by reaction with benzyl bromide (BnBr) (Scheme 18) [58]. This reaction was similar to the *N*-alkylation reaction of the side-on N_2 -Zr complexes **44** (Scheme 10a).

Cycloaddition and insertion

Besides, this side-on-end-on N_2 -Ta complex **94** was also reported to undergo [2+2] cycloaddition reaction by treating with heteroatom 1,2-cumulenes (Scheme 19a) [59]. For example, the reaction between **94** and *N,N'*-diphenyl carbodiimide results in the formation of **96**. However, when carbon disulfite or isothiocyanates are added, the *N*-functionalization product **97** is generated




Scheme 19. N—C bond formation from the cycloaddition reactions of side-on-end-on N_2 -Ta complexes with heteroatom 1,2-cumulenes. (a) The reaction of N_2 -Ta complex **94** with carbodiimide, carbon disulfite and isothiocyanates. (b) A plausible mechanism for the generation of **97** and **98**.


concomitant with the N—N bond scission. In the case of *tert*-butyl isothiocyanate (*t*BuNCS), the generated intermediate **98** would further undergo N—Si bond formation at elevated temperature to give **99**. As depicted in Scheme 19b, the formation of **97** and **98** can be rationalized by the following mechanism. The initial [2+2] cycloaddition reactions between **94** and the C=S bond of the substrates give intermediate E, followed by reductive elimination of H_2 to provide a transient intermediate F that contains a Ta—Ta bond. The Ta—Ta bond in F would trigger the N—N bond cleavage to afford the final products.

N—C BOND FORMATION VIA METAL NITRIDES

The complete reduction of N_2 -M complexes might cleave the N—N bond of the N_2 ligands to give metal nitrides. In the most terminal metal nitrides, the strong metal—nitrogen bonds result in these nitrides often exhibiting weak nucleophilicity and just reacting with high-energy species such as alkyl triflates and acyl chlorides to assemble N—C bond. However, some bridging nitrides derived from N_2 can also react with other carbon-based substrates, like MeI and CO, to form N—C bond.



Scheme 20. *N*-methylation of Nb-nitride by reaction with MeI.




Scheme 21. *N*-alkylation of Mo-nitride by reaction with MeI.

N-alkylation


In 2007, Kawaguchi *et al.* reported the reaction between MeI and bis(μ -nitrido) diniobium complex **101**, which is prepared from the tetra(μ -hydride) diniobium precursor **100** (Scheme 20) [60]. Stepwise methylation of **101** by MeI yields mono-imido **102** and bi-imido **103**, the latter of which could also react with excess pyridine to give terminal imido **104**, which reacts with CO₂ to generate **105** and **106** through further N–C bond formation. A plausible mechanism for this process was raised by the authors. A [2+2] cycloaddition of **104** with CO₂ followed by extrusion of methyl isocyanate (MeNCO) results in the formation of a terminal oxo species that dimerizes to give **106**. Meanwhile, the generated MeNCO would also undergo [2+2] cycloaddition with another molecule of **104** to form **105** [61].

Cummins *et al.* found that the terminal molybdenum nitride **108** synthesized from the three coordination Mo(III) complex **107**, undergoes *N*-alkylation by reacting with MeI to provide **109** (Scheme 21) [62].

Another example of making N–C bond from metal nitrides was fulfilled by Schneider *et al.* in 2016. The reactions between ROTf (R = Me, Et and Bn) and a terminal rhenium nitride **111**, which is prepared from the reduction of the dichloride precursor **110** with sodium amalgam or CoCp*₂, give the *N*-alkylation complexes **112** (Scheme 22) [63–65]. Further studies suggest that the nitriles (RCN) can be liberated by deprotonation and oxidation of **112** via the ketimido intermediates **113**, accompanied by the generation of trichloride complex **114**, which could also be reduced by sodium amal-



Scheme 22. Re-promoted conversion of N₂ into nitriles via *N*-alkylation of Re-nitride.




Scheme 23. *N*-methylation of Fe-nitride by reaction with MeOTs.

gam to afford the starting material **111**. According to these results, a full synthetic cycle for synthesis of nitriles from N₂ with moderate isolated yields was established (Scheme 22).

Because the group 8–10 N₂-derived nitrides are rare, the *N*-functionalization of these nitrides is hardly observed. One exception was reported by Holland *et al.*, who employed an unprecedented trinuclear iron nitride **116** to achieve this transformation. This nitride complex **116**, obtained from the reduction of the chloride precursor **115** with precisely equivalent KC₈ under N₂ atmosphere, can react with MeOTs and 18-crown-6 (18-C-6) to give the methylimido complex **118** via a presumptive two-coordinate nitride **117** with higher reactivity (Scheme 23) [66].

N-acylation

The *N*-acylation of N₂-derived nitride has also been investigated. For example, the *N*-acylation




Scheme 24. Conversion of N₂ into nitriles via *N*-acylation of Mo-nitride.


products **119** are obtained when **108** is treated with acyl chlorides in the presence of additives, such as [TMS(py)]⁺[OTf]⁻ and ^tPr₃SiOTf. Furthermore, when the *N*-acylated products **119** reacts with magnesium anthracene (MgC₁₄H₁₀) and trimethylsilyl triflate (TMSOTf) in one pot, it would be converted to the trimethylsiloxy-substituted ketimide **121** via the intermediates of **120**. Further reaction of **121** with SnCl₂ or ZnCl₂ affords the corresponding organic nitriles commitment with the generation of molybdenum chloride complex **122**, a precursor of the trisamide molybdenum complex **107**. In consequence, an efficient synthetic cycle that can directly convert N₂ to nitrile was accomplished (Scheme 24) [67].

N-acylation/elimination

In addition to simple *N*-acylated products, the reactions between N₂-derived metal nitrides and acyl chlorides also afford nitriles proceeded through *N*-acylation and subsequent elimination in formal. Hou *et al.* discovered that the reaction of titanium trialkyl complex **123** with N₂ and H₂ results in a novel diimide/tetrahydride complex **124**. This complex can react with N₂ at elevated temperature to provide a tetranuclear diimide/dinitride complex **125** that can further react with a series of acyl chloride to afford the corresponding nitriles in high yield (Scheme 25) [68]. Based on the experimental and computational results, the authors think that the functionalization of the imide ligands is prior to the




Scheme 25. Titanium-promoted direct conversion of N₂ into nitriles via the reaction between acyl chlorides and Ti-nitride.



Scheme 26. Niobium-promoted direct conversion of N₂ into nitriles via the reaction between acyl chlorides and Nb-nitride.

nitride groups in these reactions. Furthermore, by treatment of the crude reaction mixture with HCl, the titanium trichloride complex **127** is isolated, which could be easily converted to **123** by reacting with TMSCH₂Li. Hence, a synthetic cycle of titanium-promoted synthesis of nitriles direct from N₂ was proposed (Scheme 25).

Another synthetic cycle for providing organic nitriles from N₂ was developed by Cummins *et al.* via a niobium nitride intermediate (Scheme 26) [69]. An end-on bridging heterodinuclear N₂-M complex **129**, prepared from the reaction of the niobium triflate complex **128** and the aforementioned N₂-Mo




Scheme 27. Vanadium-promoted direct conversion of N_2 into potassium cyanate via the reaction between CO and N_2 -V complex.

complex **1**, could undergo N_2 ligand scission to form anionic niobium nitride **130** along with the formation of molybdenum nitride **108**, when **129** is treated with sodium amalgam. Treatment of **130** with acyl chloride results in releasing of nitriles accompanied by the generation of the niobium oxo complex **131**. By treating with triflic anhydride, this oxo complex **131** can be converted to a bistriflate complex **132** that could be reduced to the initial compound **128** to finish the cycle.

Redox-coupled *N*-atom transfer

In comparison with *N*-alkylation or acylation and subsequent reduction, a more efficient route to transfer the nitride into organic compounds is the transformation of nitride-*N* atom into an incoming substrate with concurrent metal reduction. In 2014, Kawaguchi *et al.* reported the redox-coupled *N*-atom transformation of a V-nitride (Scheme 27) [70]. Reduction of the V(III) complex **133** by KH under N_2 results in a split of the N_2 to provide the μ -nitride V(IV) complex **134**, which could be oxidized to V(V) nitride compound **135** via reacting with *p*-benzoquinone. When **135** is treated with CO or 2,6-xylylisocyanide in the presence of [2.2.2]-cryptand, the *N*-atom transformation of the substrates is observed concomitant with the formation of **136**. The contact-ion-pair complex **137** could also be isolated from the reaction of **135** and CO. Although the extrusion of the cyanate or carbodiimide ligand in **136** are not easy, the contact-ion-pair **137** readily undergoes ligand exchange with 2-butyne to liberate potassium cyanate (KNCO) with the



Scheme 28. *N*-C bond formation via the reaction of silylimido complexes with CO_2 .


formation of the alkyne adduct **138**. Additionally, **138** is facilely converted into the starting complex **133** upon dissolving in THF. Hence, a synthetic cycle for direct conversion of N_2 and CO into KNCO was completed. However, achieving the catalytic process of this synthetic cycle remains elusive due to the incompatibility of the individual steps in this cycle, such as the requirement of solvents in the *N*-C bond formation step and KNCO releasing step being different.

N-silylation/imido transfer

Besides *N*-hydrogenated intermediates, N_2 -derived *N*-silylated complexes are also good precursors to make *N*-C bond. For instance, a cycle of Mo- and W-promoted synthesis of isocyanates from N_2 via a silyl-imido intermediate was established (Scheme 28) [71]. Photolysis of the end-on bridging N_2 -Mo, W complexes **140** leads to the generation of nitride intermediates via *N*-*N* bond cleavage (*vide infra*), which would be trapped *in situ* by TMSCl to afford silyl-imido complexes **141**. When TMSCl is replaced by Ph_3SiCl , Me_3CCl or Me_3GeCl , the similar reaction could also take place to provide the corresponding imido complexes. Besides, the organic compound TMSNCO could be obtained concomitant with the formation of the mono-nuclear oxo complexes **142** by treatment of **141** with CO_2 . These oxo complexes **142** are known to react with additional TMSCl to regenerate the dichloride complexes **139** that are the precursors of the N_2 -M complexes **140**.

Metal-ligand cooperative *N*-atom transfer

Metal-ligand cooperative *N*-atom transfer is also an efficient strategy because of the avoiding of extra protons and electrons. Recently, a



Scheme 29. Metal-ligand cooperative *N*-atom transfer of a Re-nitride.


metal-ligand cooperative *N*-atom transfer of a Re-nitride was reported by a cooperative $2 H^+ / 2 e^-$ transfer of the pincer ligand (Scheme 29) [72]. This Re-nitride complex **144**, which is generated from photo-promoted cleaving of the end-on bridging binuclear N_2 -Re complex **143** (*vide infra*), could react with benzoyl chloride to afford benzamide (PhCONH₂), benzonitrile (PhCN) and benzoic acid (PhCOOH) along with the formation of trichloride Re complexes **145**, in which the pincer ligand is oxidized to an imine-type ligand. The producing of PhCN and PhCOOH is caused from the reaction of the initially formed benzamide with excess benzoylchloride in the crude.

Involvement of photochemistry


Besides photo-promoted radical generation, another pathway of the photochemistry participating in N_2 transformation is the direct photolytic splitting of N_2 ligands into nitrides, which could further engage in *N*-atom transfer. Two examples of this method have been reported. In the first example, irradiation of the above-mentioned end-on bridging N_2 -Mo, W complexes **140** over several days by medium-pressure Hg lamps leads to the generation of two metal nitrides **146** and **147** (Scheme 30a). Furthermore, when **140** are photolyzed in the presence of excess TMSCl, the terminal silylimido complexes **141** are obtained in moderate yield with the formation of dichloride complexes **139** (Scheme 29) [71]. The other example involves the photolysis of an end-on-bridged N_2 -Re complex **143** that has abnormal thermal-stability, to provide the aforementioned Re-nitride **144** (Scheme 30b) [72]. It is noteworthy that the photo source of this reaction could be Xe(Hg) lamp ($\lambda > 305$ nm) or a 390 nm LED lamp.

Involvement of electrochemistry

Electrochemical N_2 reduction is an alternative to chemical N_2 reduction for the synthesis of N_2 -M complexes. This approach has been utilized to regain the N_2 -Re complex **143** to achieve a cycle (Scheme 31) [72]. Schneider *et al.* found that in the controlled potential electrolysis experiment, the



Scheme 30. Photolytic cleavage of end-on bridging N_2 -Mo, W and Re complexes into nitrides. (a) Photolytic cleavage of end-on bridging N_2 -Mo, W complexes **140**. (b) Photolytic cleavage of end-on bridging N_2 -Re complexes **143**.




Scheme 31. Electrochemical reduction involved synthetic cycle of direct conversion of N_2 into benzamide and benzonitrile.

trichloride Re complexes **145** formed from the reaction of Re-nitride with PhCOCl, could be converted to the N_2 -Re complex **143** via electrolyzing at $E = -1.65$ V for 8 h in the presence of proton source of 2,6-dichlorophenol (DCP) and subsequently electrolyzing at $E = -1.85$ V for 5 h under N_2 . Thus, a three-step cycle for the synthesis of PhCONH₂/PhCN from N_2 was established, in which the creative approaches of metal-ligand cooperation and photo- and electrochemistry were all used.

N—C BOND FORMATION VIA UNCHARACTERIZED N_2 -M INTERMEDIATES

Compared to the above works, the earlier reports about the conversion of N_2 into organic compounds were achieved by one-pot reactions of ill-defined




Scheme 32. Ti-promoted transformation of N_2 into amines via ill-defined intermediates. (a) The reaction of Cp_2TiCl_2 with aryllithium reagents under N_2 to afford aromatic amines. (b) The reaction of diaryltitanocenes Cp_2TiAr_2 with alkali or alkaline metal under N_2 to afford aromatic amines. (c) Synthesis of organic amines from the reaction between ketones and a supposed titanium nitride species.

N_2 -complexes or their derivatives with carbon-based substrates and followed by hydrolysis. In this section, some examples of this method were introduced briefly.

The initial works of transition metal promoted direct conversion of N_2 into organic compounds were reported more than 50 years ago, when Vol'pin and Shur *et al.* developed two systems for transformation of N_2 into aromatic amines mediated by titanium species [14,73]. In the first system, several aromatic amines are obtained when Cp_2TiCl_2 is treated with excess of aryllithium (aryl = Ph, *m*- and *p*-MeC₆H₄) reagents under N_2 pressure of 80–100 atm and followed by hydrolysis (Scheme 32a). When the aryllithium in these reactions is replaced by alkyl-lithium reagents, the corresponding alkylamines could not be obtained. The other system, which also gives arylamines by subsequent hydrolysis, involves the reaction of diaryltitanocenes Cp_2TiAr_2 (Ar = Ph, *m*- and *p*-MeC₆H₄) with alkali or alkaline metal (Li, Na and Mg) and N_2 (100 atm) (Scheme 32b). Although plenty of effort has been made, the detailed mechanisms of these works are yet unclear.

Meanwhile, a related work was reported in 1970 by van Tamelen and Rudler, who succeeded in the synthesis of organic amines from the reaction between ketones and a supposed titanium nitride species prepared through the reaction of Cp_2TiCl_2 with magnesium under N_2 (Scheme 32c) [74].

Additionally, Mori *et al.* achieved incorporation of N_2 into organic compounds via the *N*-silylation titanium complexes **148**, which were prepared from the one-pot reaction of titanium species ($TiCl_4$ or $Ti(O^iPr)_4$), Li and TMSCl under N_2 or dry air (1 atm) [75]. Although the precise components and structures of **148** have not been determined so far, they are considered to contain $XTi = NTMS$, $X_2Ti-N(TMS)_2$ ($X = Cl, O^iPr$) and $N(TMS)_3$



Scheme 33. Ti-promoted N–C bond formation via ill-defined *N*-silylation titanium species. (a) Preparation of *N*-silylation titanium complexes **148** from one-pot reaction of $TiCl_4$ or $Ti(O^iPr)_4$ with Li and TMSCl under N_2 or dry air. (b) The reaction between **148** and keto-carbonyl compounds to afford nitric heterocycles. (c) Palladium-catalyzed synthesis of aryl- or allyl- amines and amide derivatives from **148** and aryl or allyl halides in the absence or the presence of CO.

[76–78]. **148** could serve as a nitrogenation reagent to react with a series of keto-carbonyl compounds to provide kinds of nitric heterocycles, such as indole, quinoline, pyrrole, pyrrolizine, lactams and indolizine derivatives. Besides, when **148** is treated with palladium complexes, the transmetalation of *N*-atom from **148** to the palladium center occurs. Hence, palladium-catalyzed synthesis of aryl- or allyl- amines and amide derivatives from aryl or allyl halides and **148** could be fulfilled, in the absence or the presence of CO (Scheme 33).

CONCLUSION AND OUTLOOK

Direct transformation of N_2 into *N*-containing organic compounds is of fundamental and practical significance. In the past 60 years, the area of direct incorporation of N_2 into N–C bond effectuated

many great achievements. Relative to the traditional methods of assembling N–C bond via *N*-alkylation of N₂-M complexes, more atom-efficient approaches, such as cycloaddition, insertion and redox-coupled *N*-atom transfer for making N–C bond have been developed and received more attention in recent years. By the delicate design, some synthetic cycles about direct conversion of N₂ into organic compounds have also been developed. In these cycles, photo- and electrochemistry are sometimes used to prepare the N₂-M complexes, cleave the N–N bond or release the final products. However, all of these reactions are stoichiometric and the catalytic system for the direct introduction of N₂ into organic compounds has not been realized yet. The main factors that prevent these complete synthetic cycles from becoming a catalytic process are the rigorous reaction conditions of the N–C bond formation and *N*-containing organic compounds releasing steps in these cycles, which are incompatible with the preparation steps for N₂-M complexes. Hence, developing milder systems are imperative. Besides, new reaction types also need to be explored. In this context, we think the following fields can be considered in the future.

New reaction systems

So far, most N–C bond formation occurs at N₂-M complexes of group 4–6 transition metals. Exploring other metal promoted N–C bond formation is an attractive topic. Besides, the multi-metal synergistically promoted N₂ activation and functionalization also need to be studied. Toward this end, the design of new types of ligands should be considered.

New reaction types

Reductive elimination, an essential step in catalytic amination reactions, has not been found to take place at N₂-derived *N*-containing transition metal complexes. This process should be explored in future because it provides an efficient approach to assemble N–C bond concomitant with regaining the N₂-M complexes or their precursors. Additionally, other intriguing reaction modes, such as the [4+2] cycloadditions of N₂ ligands and insertion of N₂ into metal–carbon bond should also be investigated.

Polynuclear metal species cooperative N₂ scission and functionalization

Stimulated by the previous works on multinuclear Ti, Fe complexes-promoted N₂ cleavage and N–C

bond formation [66,68], the strategy to realize synergistic N₂-splitting and subsequent functionalization using polynuclear metal complexes should be further explored. Additionally, recent reports on gas-phase polynuclear metal clusters-mediated N₂ scission and subsequent N–C bond formation deserve further attention [79,80].

Main group elements promoted N–C bond formation

The recent report about N₂ reduction by borylenes from Braunschweig *et al.* suggests the potential of boron mediated formation of N–C bond from N₂ [81]. Besides, some calculation results indicate that the direct reactions of boron or carbene with N₂ are also permitted in some cases. For example, Li and Schaefer *et al.* designed a new molecular system for nitrogen reduction, involving a 2,3'-bipyridine-anchored, end-on-bridging dinitrogen complex of the Me₂B–BMe₂ intermediate by theoretical calculations [82]. Zhu *et al.* designed a metal-free dinitrogen activation system based on the boron and NHC carbene system [83]. These results offer inspiration for future work on p-block elements promoting or catalytic conversion of N₂ into organic compounds.

Analogue of PCET: lessons from N₂-to-NH₃ catalysis system

Very recently, Nishibayashi *et al.* achieved a remarkable N₂-to-NH₃ process via a molybdenum-catalysis system. By using the samarium diiodide (SmI₂) as the reductant and alcohol or water as the proton sources, the total turnover number (TON) of this reaction reaches up to 4350 with 91% yield of NH₃. Further studies reveal that a proton-coupled electron-transfer (PCET) process, in which O–H bonds in water or alcohols are weakened by coordination to SmI₂, is the key to this high reactivity [84]. Inspired by this, a similar process of C–X (X = O, Cl, Br, I) bonds coordinated to a relevant reductant to weaken the C–X bonds, which could be named as carbocation-coupled electron-transfer (CCET) should also be studied. This CCET process may offer a path forward for developing catalysis systems that incorporate N₂ into amines via successive alkylation of a N₂-derived nitride.

Photochemistry

Limited complexes capable of N₂ photoactivation are currently known, and the underlying photophysical and photochemical processes after light absorption are largely unresolved. Light can induce the

split of $\text{N}\equiv\text{N}$ bond in the M-N_2 complexes and the resulting nitride complexes are typically reactive. Hence, the following N-C formation reaction should be possible. Besides, the excitation into N-N π^* orbitals is also possible, which can lead to a weakened π -bond, and hence a following N-C formation directly from M-N_2^* and carbon-based substrates including CO , CO_2 might be possible.

Electrochemistry

Previous work indicates that electrochemical reduction could release final organic compounds along with regeneration of $\text{N}_2\text{-M}$ complexes. This stimulates us to create an electrochemical reduction system, in which the electrolyzation step is compatible with the N-C bond formation step. Besides, the recent report of N_2 and CO_2 coupling to produce urea, which was conducted by an electrocatalyst consisting of PdCu alloy nanoparticles on TiO_2 nanosheets, suggests that designing a new solid catalyst to incorporate N_2 into high-value N -containing product beyond NH_3 should also be attractive [85].

Heterogeneous catalysis systems

Although the industrial Haber-Bosch process produces NH_3 over the surface of heterogeneous solid-state catalysts, a similar process, in which a heterogeneous catalyst catalyzes or promotes direct transformation of N_2 into organic compounds, has not been reported in literature. Development of the new systems, where the merits of the homogeneous molecular systems and the heterogeneous systems are rationally combined, is a promising approach toward the goal.

Via N -protonation or silylation intermediates

The strategy of conversion of N_2 into organic compounds via the N -silylation and N -protonation complexes can be further extended. As an example, converting the less active M-N_2 complexes into M-N-Si/H species *in situ* followed by catalytic reaction with carbon-based substrates, might result in various valuable organic compounds being synthesized.

In a word, with combined efforts from cross-disciplines the dream of direct catalytic and efficient conversion of N_2 into N -containing organic compounds under mild conditions is believed to be attainable in the future.

ACKNOWLEDGEMENTS

The authors thank Dr. Mingdong Zhong, Dr. Chunhai Wang, Mr. Gaoxiang Wang and Mr. Zhubao Yin of Peking University for their discussion during the preparation of this manuscript.

FUNDING

Prof. Ping Chen, Dehui Deng, Zhang-Jie Shi and Zhenfeng Xi are working together as a team in the National Natural Science Foundation of China Center of 'Transformation Chemistry of Key Components of Air' (21988101). Activation and conversion of dinitrogen directly to nitrogenous organic compounds is the major aim of this Center.

Conflict of interest statement. None declared.

REFERENCES

1. Walter MD. Recent advances in transition metal-catalyzed dinitrogen activation. In: Pedro JP (ed.). *Advances in Organometallic Chemistry*. New York: Academic Press, 2016, vol. **65**, Chapter 5: 261–377.
2. Hoffman BM, Dean DR and Seefeldt LC. Climbing nitrogenase: toward a mechanism of enzymatic nitrogen fixation. *Acc Chem Res* 2009; **42**: 609–19.
3. Spatzal T, Aksoyoglu M and Zhang L *et al.* Evidence for interstitial carbon in nitrogenase FeMo cofactor. *Science* 2011; **334**: 940.
4. Lancaster KM, Romelt M and Ettenhuber P *et al.* X-ray emission spectroscopy evidences a central carbon in the nitrogenase iron-molybdenum cofactor. *Science* 2011; **334**: 974–7.
5. Stappen CV, Decamps L and Cutsail GE, III *et al.* The spectroscopy of nitrogenases. *Chem Rev* 2020; **120**: 5005–81.
6. Rafiqul I, Weber C and Lehmann B *et al.* Energy efficiency improvements in ammonia production-perspectives and uncertainties. *Energy* 2005; **30**: 2487–504.
7. Jia HP and Quadrelli EA. Mechanistic aspects of dinitrogen cleavage and hydrogenation to production ammonia in catalysis and organometallic chemistry: relevance of metal hydride bonds and dihydrogen. *Chem Soc Rev* 2014; **43**: 547–64.
8. Nishibayashi Y (ed.). *Nitrogen Fixation, Topics in Organometallic Chemistry*, vol. **60**. Cham: Springer International Publishing, 2017.
9. Foster SL, Bakovic SIP and Duda RD *et al.* Catalysts for nitrogen reduction to ammonia. *Nat Catal* 2018; **1**: 490–500.
10. Chalkley MJ, Drover MW and Peters JC. Catalytic N_2 -to- NH_3 (or $-\text{N}_2\text{H}_4$) conversion by well-defined molecular coordination complexes. *Chem Rev* 2020; **120**: 5582–636.
11. Tang C and Qiao SZ. How to explore ambient electrocatalytic nitrogen reduction reliably and insightfully. *Chem Soc Rev* 2019; **48**: 3166–80.
12. Jennings JR (ed.). *Catalytic Ammonia Synthesis: Fundamentals and Practice*. New York: Plenum, 1991, 451.
13. Aika K and Tamaru K. *Ammonia: Catalysis and Manufacture*. Nielsen A (ed.). Berlin: Springer-Verlag, 1995, 103–48.

14. Vol'pin ME, Shur VB and Kudryavtsev RV *et al.* Amine formation in molecular nitrogen fixation: nitrogen insertion into transition metal-carbon bonds. *Chem Commun* 1968; 1038–40.
15. Allen AD and Senoff CV. Nitrogenopentammineruthenium(II) complexes. *Chem Commun* 1965; 621–2.
16. Tanabe Y and Nishibayashi Y. Overviews of the Preparation and Reactivity of Transition Metal–Dinitrogen Complexes. In: Nishibayashi Y (ed.). *Transition Metal–Dinitrogen Complexes: Preparation and Reactivity*. Weinhei: Wiley–VCH, 2019, 1–77.
17. Burford RJ and Fryzuk MD. Examining the relationship between coordination mode and reactivity of dinitrogen. *Nat Rev Chem* 2017; **1**: 0026.
18. Li J, Yin J and Yu C *et al.* Direct transformation of N₂ to N-containing organic compounds. *Acta Chim Sinica* 2017; **75**: 733–43 (in Chinese).
19. Kim S, Loose F and Chirik PJ. Beyond ammonia: nitrogen–element bond forming reactions with coordinated dinitrogen. *Chem Re* 2020; **120**: 5637–81.
20. Peters JC, Cherry JPF and Thomas C *et al.* Redox-catalyzed binding of dinitrogen by molybdenum N-*tert*-hydrocarbylanilide complexes: implications for dinitrogen functionalization and reductive cleavage. *J. Am Chem Soc* 1999; **121**: 10053–67.
21. Greco GE and Schrock RR. Synthesis, structure, and electrochemical studies of molybdenum and tungsten dinitrogen, diazenido, and hydrazido complexes that contain aryl-substituted triamidoamine ligands. *Inorg Chem* 2001; **40**: 3861–78.
22. Betley TA and Peters JC. Dinitrogen chemistry from trigonally coordinated iron and cobalt platforms. *J Am Chem Soc* 2003; **125**: 10782–3.
23. Rittle J and Peters JC. An Fe–N₂ complex that generates hydrazine and ammonia via Fe = NNH₂: demonstrating a hybrid distal-to-alternating pathway for N₂ reduction. *J Am Chem Soc* 2016; **138**: 4243–8.
24. Chatt J, Diamantis AA and Heath GA *et al.* Reactions of ligating dinitrogen to form carbon–nitrogen bonds. *Dalton Trans* 1977; 688–97.
25. Hidaï M and Mizobe Y. Recent advances in the chemistry of dinitrogen complexes. *Chem Rev* 1995; **95**: 1115–33.
26. Yoshida T, Adachi T and Ueda T *et al.* N-arylation and N,N-dibenzylolation of coordinated N₂ with organic halides; differences in the reactivity of *trans*-[Mo(N₂)₂(Me₆[16]aneS₄)] and its phosphane analogues. *Angew Chem Int Ed Engl* 1989; **28**: 1040–2.
27. Sellmann D and Weiss W. The first reaction of the N₂ ligand with bases; reduction of coordinated dinitrogen by nucleophilic attack. *Angew Chem Int Ed Engl* 1977; **16**: 880–1.
28. Sellmann D and Weiss W. Consecutive nucleophilic and electrophilic attack on N₂ ligands: synthesis of Azo compounds from molecular nitrogen. *Angew Chem Int Ed Engl* 1978; **17**: 269–70.
29. Chatt J, Heath GA and Leigh GJ. The formation of a nitrogen to carbon bond in a reaction of a dinitrogen complex. *Chem Commun* 1972; 444–5.
30. Sato M, Kodama T and Hidaï M *et al.* Preparation and properties of molybdenum- and tungsten-dinitrogen complexes 7. The molecular structure of MoCl(N₂COPh)(Ph₂PCH₂CH₂PPh₂)₂. *J Organomet Chem* 1978; **152**: 239–54.
31. Nakanishi Y, Ishida Y and Kawaguchi H. Nitrogen-carbon bond formation by reactions of a titanium-potassium dinitrogen complex with carbon dioxide, *tert*-butyl isocyanate, and phenylallene. *Angew Chem Int Ed* 2017; **56**: 9193–7.
32. Caselli A, Solari E and Scopelliti R *et al.* Dinitrogen rearranging over a metal-oxo surface and cleaving to nitride: from the end-on to the side-on bonding mode, to the stepwise cleavage of the N≡N bonds assisted by Nb^{III}-calix[4]arene. *J Am Chem Soc* 2000; **122**: 3652–70.
33. Turner HW, Fellmann JD and Rocklage MS *et al.* Tantalum complexes containing diimido bridging dinitrogen ligands. *J Am Chem Soc* 1980; **102**: 7809–11.
34. Chatt J, Dilworth JR and Richards RL. Recent advances in the chemistry of nitrogen fixation. *Chem Rev* 1978; **78**: 589–625.
35. Hidaï M. Chemical nitrogen fixation by molybdenum and tungsten complexes. *Coord Chem Rev* 1999; **185–6**: 99–108.
36. Hussain W, Leigh JG and Pickett CJ. Stepwise conversion of dinitrogen coordinated to molybdenum into an amine and an imido-complex. Relevance to the reactions of nitrogenase. *Chem Commun* 1982; 747–8.
37. Seino H, Ishii Y and Sasagawa T *et al.* Synthesis and reactivities of pyrrolylimido complexes of molybdenum and tungsten: formation of pyrrole and N-aminopyrrole from molecular nitrogen. *J Am Chem Soc* 1995; **117**: 12181–93.
38. Pickett CJ and Leigh GJ. Towards a nitrogen-fixing cycle: electrochemical reduction of a hydrazido-complex of molybdenum(IV) under N₂ to yield the dialkylhydrazine and a molybdenum(0) dinitrogen complex. *Chem Commun* 1981; 1033–5.
39. Hirotsu M, Fontaine PP and Zavalij PY *et al.* Extreme N≡N bond elongation and facile N-atom functionalization reactions within two structurally versatile new families of group 4 bimetallic 'side-on-bridged' dinitrogen complexes for zirconium and hafnium. *J Am Chem Soc* 2007; **129**: 12690–2.
40. Knobloch DJ, Benito-Garagorri D and Bernskoetter WH *et al.* Addition of methyl triflate to a hafnocene dinitrogen complex: stepwise N₂ methylation and conversion to a hafnocene hydrazonato compound. *J Am Chem Soc* 2009; **131**: 14903–12.
41. Lv ZJ, Huang Z and Zhang WX *et al.* Scandium-promoted direct conversion of dinitrogen into hydrazine derivatives via N–C bond formation. *J Am Chem Soc* 2019; **141**: 8773–7.
42. Morello L, Love JB and Patrick BO *et al.* Carbon–nitrogen bond formation via the reaction of terminal alkynes with a dinuclear side-on dinitrogen complex. *J Am Chem Soc* 2004; **126**: 9480–1.
43. Bernskoetter WH, Olmos AV and Pool JA *et al.* N–C bond formation promoted by a hafnocene dinitrogen complex: comparison of zirconium and hafnium congeners. *J Am Chem Soc* 2006; **128**: 10696–7.
44. Bernskoetter WH, Lobkovsky E and Chirik PJ. Nitrogen–carbon bond formation from N₂ and CO₂ promoted by a hafnocene dinitrogen complex yields a substituted hydrazine. *Angew Chem Int Ed* 2007; **46**: 2858–61.
45. Knobloch DJ, Toomey HE and Chirik PJ. Carboxylation of an *ansa*-zirconocene dinitrogen complex: regiospecific hydrazine synthesis from N₂ and CO₂. *J Am Chem Soc* 2008; **130**: 4248–9.
46. Knobloch DJ, Lobkovsky E and Chirik PJ. Dinitrogen cleavage and functionalization by carbon monoxide promoted by a hafnium complex. *Nat Chem* 2010; **2**: 30–5.
47. Knobloch DJ, Lobkovsky E and Chirik PJ. Carbon monoxide-induced dinitrogen cleavage with group 4 metallocenes: reaction scope and coupling to N–H bond formation and C–O deoxygenation. *J Am Chem Soc* 2010; **132**: 10553–64.
48. Zhang X, Butschke B and Schwarz H. N₂ activation by a hafnium complex: a DFT study on CO-assisted dinitrogen cleavage and functionalization. *Chem Eur J* 2010; **16**: 12564–9.
49. Knobloch DJ, Semproni SP and Lobkovsky E *et al.* Studies into the mechanism of CO-induced N₂ cleavage and C–C bond formation promoted by an *ansa*-hafnocene complex. *J Am Chem Soc* 2012; **134**: 3377–86.
50. Semproni SP, Margulieux GW and Chirik PJ. Di- and tetrametallic hafnocene oxamidides prepared from CO-induced N₂ bond cleavage and thermal rearrangement to hafnocene cyanide derivatives. *Organometallics* 2012; **31**: 6278–87.

51. Knobloch DJ, Lobkovsky E and Chirik PJ. Functionalization of hafnium oxamide complexes prepared from CO-induced N₂ cleavage. *J Am Chem Soc* 2010; **132**: 15340–50.
52. Semproni SP, Milsmann C and Chirik PJ. Structure and reactivity of a hafnocene μ -nitrido prepared from dinitrogen cleavage. *Angew Chem Int Ed* 2012; **51**: 5213–6.
53. Semproni SP and Chirik PJ. Synthesis of a base-free hafnium nitride from N₂ cleavage: a versatile platform for dinitrogen functionalization. *J Am Chem Soc* 2013; **135**: 11373–83.
54. Semproni SP and Chirik PJ. Activation of dinitrogen-derived hafnium nitrides for nucleophilic N–C bond formation with a terminal isocyanate. *Angew Chem Int Ed* 2013; **52**: 13203–7.
55. Semproni SP and Chirik PJ. N–H and N–C bond formation with an N₂-derived dihafnium μ -nitrido complex. *Organometallics* 2014; **33**: 3727–37.
56. Falcone M, Chatelain L and Scopelliti R *et al.* Nitrogen reduction and functionalization by a multimetallic uranium nitride complex. *Nature* 2017; **547**: 332–5.
57. Falcone M, Barluzzi L and Andrez J *et al.* The role of bridging ligands in dinitrogen reduction and functionalization by uranium multimetallic complexes. *Nat Chem* 2019; **11**: 154–60.
58. Fryzuk MD, Johnson SA and Patrick BO *et al.* New mode of coordination for the dinitrogen ligand: formation, bonding, and reactivity of a tantalum complex with a bridging N₂ unit that is both side-on and end-on. *J Am Chem Soc* 2001; **123**: 3960–73.
59. Ballmann J, Yeo A and Patrick BO *et al.* Carbon–nitrogen bond formation by the reaction of 1,2-cumulenes with a ditantalum complex containing side-on- and end-on-bound dinitrogen. *Angew Chem Int Ed* 2011; **50**: 507–10.
60. Akagi F, Matsuo T and Kawaguchi H. Dinitrogen cleavage by a diniobium tetrahydride complex: formation of a nitride and its conversion into imide species. *Angew Chem Int Ed* 2007; **46**: 8778–81.
61. Akagi F, Suzuki S and Ishida Y *et al.* Reactions of a niobium nitride complex prepared from dinitrogen: synthesis of imide and ureate complexes and ammonia formation. *Eur J Inorg Chem* 2013; **2013**: 3930–6.
62. Sceats EL, Figueroa JS and Cummins CC *et al.* Complexes obtained by electrophilic attack on a dinitrogen-derived terminal molybdenum nitride: electronic structure analysis by solid state CP/MAS ¹⁵N NMR in combination with DFT calculations. *Polyhedron* 2004; **23**: 2751–68.
63. Klopsch I, Finger M and Würtele C *et al.* Dinitrogen splitting and functionalization in the coordination sphere of rhenium. *J Am Chem Soc* 2014; **136**: 6881–3.
64. Klopsch I, Kinauer M and Finger M *et al.* Conversion of dinitrogen into acetonitrile under ambient conditions. *Angew Chem Int Ed* 2016; **55**: 4786–9.
65. Klopsch I, Schendzielorz F and Volkmann C *et al.* Synthesis of benzonitrile from dinitrogen. *Z Anorg Allg Chem* 2018; **644**: 916–9.
66. MacLeod KC, Menges FS and McWilliams SF *et al.* Alkali-controlled C–H cleavage or N–C bond formation by N₂-derived iron nitrides and imides. *J Am Chem Soc* 2016; **138**: 11185–91.
67. Curley JJ, Sceats EL and Cummins CC. A cycle for organic nitrile synthesis via dinitrogen cleavage. *J Am Chem Soc* 2006; **128**: 14036–7.
68. Guru MM, Shima T and Hou Z. Conversion of dinitrogen to nitriles at a multinuclear titanium framework. *Angew Chem Int Ed* 2016; **55**: 12316–20.
69. Figueroa JS, Piro NA and Clough CR *et al.* A nitridoniobium(V) reagent that effects acid chloride to organic nitrile conversion: synthesis via heterodinuclear (Nb/Mo) dinitrogen cleavage, mechanistic insights, and recycling. *J Am Chem Soc* 2006; **128**: 940–50.
70. Ishida Y and Kawaguchi H. Nitrogen atom transfer from a dinitrogen-derived vanadium nitride complex to carbon monoxide and isocyanide. *J Am Chem Soc* 2014; **136**: 16990–3.
71. Keane AJ, Farrell WS and Yonke BL *et al.* Metal-mediated production of isocyanates, R₃EN=C=O from dinitrogen, carbon dioxide, and R₃ECl. *Angew Chem Int Ed* 2015; **54**: 10220–4.
72. Schendzielorz F, Finger M and Abbeneth J *et al.* Metal-ligand cooperative synthesis of benzonitrile via electrochemical reduction and photolytic splitting of dinitrogen. *Angew Chem Int Ed* 2019; **58**: 830–4.
73. Vol'pin ME, Shur VB and Berkovich EG. Transformations of molecular nitrogen into aromatic amines under the action of titanium compounds. *Inorg Chim Acta* 1998; **280**: 264–74.
74. van Tamelen EE and Rudler H. Utilization of molecular nitrogen in the synthesis of organic amines and nitriles. *J Am Chem Soc* 1970; **92**: 5253–4.
75. Kawaguchi M, Hamaoka S and Mori M. Incorporation of molecular nitrogen into organic compounds. Titanium catalyzed nitrogenation. *Tetrahedron Lett* 1993; **34**: 6907–10.
76. Mori M. Activation of nitrogen for organic synthesis. *J Organomet Chem* 2004; **689**: 4210–27.
77. Mori M, Akashi M and Hori M *et al.* Nitrogen fixation: synthesis of heterocycles using molecular nitrogen as a nitrogen source. *Bull Chem Soc Jpn* 2004; **77**: 1655–70.
78. Tanabe Y and Nishibayashi Y. Recent advances in catalytic silylation of dinitrogen using transition metal complexes. *Chem Soc Rev* 2019; **389**: 73–93.
79. Li ZY, Mou LH and Wei GP *et al.* Coupling in N₂ fixation by the ditantalum carbide cluster anions Ta₂C₄[−]. *Inorg Chem* 2019; **58**: 4701–5.
80. Li ZY, Li Y and Mou LH *et al.* A facile N≡N bond cleavage by the trinuclear metal center in vanadium carbide cluster anions V₃C₄[−]. *J Am Chem Soc* 2020; **142**: 10747–54.
81. Légaré MA, Bélanger-Chabot G and Dewhurst RD *et al.* Nitrogen fixation and reduction at boron. *Science* 2018; **359**: 896–900.
82. Li L, Wu Z and Zhu H *et al.* Reduction of dinitrogen via 2,3'-bipyridine-mediated tetraboration. *J Am Chem Soc* 2020; **142**: 6244–50.
83. Zhu J. Rational design of a carbon-boron frustrated lewis pair for metal-free dinitrogen activation. *Chem Asian J* 2019; **14**: 1413–7.
84. Ashida Y, Arashiba K and Nakajima K *et al.* Molybdenum-catalysed ammonia production with samarium diiodide and alcohols or water. *Nature* 2019; **568**: 536–40.
85. Chen C, Zhu X and Wen X *et al.* Coupling N₂ and CO₂ in H₂O to synthesize urea under ambient conditions. *Nat Chem* 2020; doi: 10.1038/s41557-020-0481-9.