




Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.



## Review

## Mammalian phosphatidylinositol 4-kinases as modulators of membrane trafficking and lipid signaling networks

Emma L. Clayton, Shane Minogue, Mark G. Waugh\*

UCL Institute for Liver &amp; Digestive Health, UCL Royal Free Campus, Rowland Hill Street, London NW3 2PF, United Kingdom

## ARTICLE INFO

## Article history:

Received 31 December 2012

Accepted 8 April 2013

Available online 19 April 2013

## ABSTRACT

The four mammalian phosphatidylinositol 4-kinases modulate inter-organelle lipid trafficking, phosphoinositide signalling and intracellular vesicle trafficking. In addition to catalytic domains required for the synthesis of PI4P, the phosphatidylinositol 4-kinases also contain isoform-specific structural motifs that mediate interactions with proteins such as AP-3 and the E3 ubiquitin ligase Itch, and such structural differences determine isoform-specific roles in membrane trafficking. Moreover, different permutations of phosphatidylinositol 4-kinase isozymes may be required for a single cellular function such as occurs during distinct stages of GPCR signalling and in Golgi to lysosome trafficking. Phosphatidylinositol 4-kinases have recently been implicated in human disease. Emerging paradigms include increased phosphatidylinositol 4-kinase expression in some cancers, impaired functioning associated with neurological pathologies, the subversion of PI4P trafficking functions in bacterial infection and the activation of lipid kinase activity in viral disease. We discuss how the diverse and sometimes overlapping functions of the phosphatidylinositol 4-kinases present challenges for the design of isoform-specific inhibitors in a therapeutic context.

© 2013 Elsevier Ltd. Open access under [CC BY license](http://creativecommons.org/licenses/by/3.0/).

## Contents

1. Introduction	294
2. PI 4-kinases and the intracellular compartmentation of PI4P synthesis	295
3. Cell signalling	296
4. Interactions with other lipid pathways	297
4.1. Sterol regulation of PI4KII $\alpha$	297
4.2. PI 4-kinases & sphingomyelin synthesis	297
4.3. PI 4-kinases & glycosphingolipid synthesis	298
5. Emerging roles for PI 4-kinases in disease	298
5.1. Cancer	299
5.2. Neurological disease	299
5.3. Bacterial infection	299
5.4. Viral disease	300
6. Pharmacological targeting of the PI 4-kinases and future perspectives	300
Acknowledgement	301
References	301

**Abbreviations:** ARF, ADP-ribosylation factor; DAG, diacylglycerol; EGF, epidermal growth factor; EGFR, epidermal growth factor receptor; ER, endoplasmic reticulum; FAPP2, Four-phosphate adaptor protein 2; FAAT motif, two phenylalanines in an acidic tract motif; GPCR, G-protein-coupled receptor; HCV, hepatitis C virus; NCS-1, neuronal calcium sensor-1; OSBP, oxysterol binding protein; OSH, oxysterol binding protein homologue; PKD, protein kinase D; PH, pleckstrin homology; PI, phosphatidylinositol; PI4P, phosphatidylinositol 4-phosphate; PI(4,5)P<sub>2</sub>, phosphatidylinositol (4,5)-bisphosphate; PI(3,4,5)P<sub>3</sub>, phosphatidylinositol (3,4,5)-trisphosphate; PLC, phospholipase C; SARS, Severe acute respiratory syndrome; TGN, trans Golgi network; TRPV, transient receptor potential vanilloid.

\* Corresponding author. Tel.: +44 2074332822.

E-mail address: [m.waugh@ucl.ac.uk](mailto:m.waugh@ucl.ac.uk) (M.G. Waugh).

## 1. Introduction


The four enzymes that make up the mammalian phosphatidylinositol (PI) 4-kinase family can be divided into two groups based on primary sequence and biochemical properties: consisting of type II (PI4KII $\alpha$  and PI4KII $\beta$ ) and the type III (PI4KIII $\alpha$  and PI4KIII $\beta$ ) isozymes [1,2]. All members catalyse the phosphorylation of phosphatidylinositol (PI) at the D4 position of the inositol head-group

to synthesize phosphatidylinositol 4-phosphate (PI4P) which is an essential precursor in the enzymatic pathways that produce PI(4,5)P<sub>2</sub> and PI(3,4,5)P<sub>3</sub> for receptor-activated phospholipase C (PLC) and phosphoinositide 3-kinase signalling [3]. In addition to the generation of phosphoinositides for receptor signalling, PI 4-kinase activity underlies the recruitment a number of PI4P-specific binding proteins such as the pleckstrin homology (PH) domain containing proteins CERT [4] and FAPP2 [5] which regulate lipid synthesis on Golgi membranes, and also the recruitment of the clathrin adaptors AP-1 [6] and AP-3 [7] during Golgi-endosomal trafficking. However, PI 4-kinases do not just produce PI4P. Another level of cellular regulation exists at the non-catalytic, protein-interaction level, where structural binding motifs mediate isoform-specific interactions with molecules such as Rab11 [8], NCS-1 [9,10], AP-3 [7,11,12] and the E3 ubiquitin ligase itch [13]. Recent work has also established that different PI 4-kinase isoform permutations are required during receptor-activated PLC signalling [14–16] and in Golgi-endosomal trafficking [16,17] all of which suggests that there is still much still to be discovered about this, the least well studied mammalian PI kinase family.

## 2. PI 4-kinases and the intracellular compartmentation of PI4P synthesis

For over 40 years there has been intense interest in mapping out the cellular compartments where PI 4-kinases are active [18–24]. Pioneering work from Michell, Harwood and Hawthorne established an early precedent for PI4P synthesis in plasma membrane enriched fractions [20,25] and the presence of a biochemically distinct PI 4-kinase activity associated with the ER which could be activated by the non-ionic detergent Cutscum [19]. More recently, in the post-cloning era, these original observations have been built on and there has been substantial progress in mapping the subcellular distributions of the four mammalian PI 4-kinase isoforms [26–29] (Fig. 1). Subcellular fractionation and immunocytochemical experiments have established that PI4KII $\alpha$  (55 kDa) localises to the *trans*-Golgi network (TGN) [6,28–31] where its lipid kinase activity has been implicated in the recruitment of AP-1 [6] and GGA clathrin adaptors [32]. PI4KII $\alpha$  also localises to various membranes of the endosomal system [12,26,29,33,34] as well as specialised trafficking intermediates such as synaptic vesicles [35] and Glut4 transport vesicles [36]. In addition, there is biochemical evidence for a minor but highly active pool of PI4KII $\alpha$  at the endoplasmic reticulum (ER) [29] where it may be activated by interactions with the c-FOS transcription factor [37]. On endosomal membranes both PI4KII $\alpha$  catalysed PI4P generation and the presence of a dileucine sorting motif are required for optimal binding of AP-3 [7] – a key step in cargo selection and trafficking to late endosomes/lysosomes. Despite its high degree of structural homology to the PI4KII $\alpha$  isoform, PI4KII $\beta$  has a different subcellular distribution profile and has been observed in both in the cytosol [38,39] and in association with trafficking vesicles [26,40].

As for the two wortmannin-sensitive PI4KIIIs [41–43], PI4KIII $\alpha$  [42,44–46] (230 kDa) has been localised by immunostaining with isoform specific-antibodies to cytoplasmic membranes [47–49], and more specifically to early *cis*-Golgi compartments [50] and the nucleolus [47]. PI4KIII $\alpha$  has also been identified the main isozyme responsible for PI4P generation at the plasma membrane [14,27,51]. Recently, Nakatsu et al. [52] demonstrated that PI4KIII $\alpha$  visits the plasma membrane, in a dynamic process mediated by interactions with the palmitoylated, membrane-resident protein EFR3B and TTC7B – a protein recruited from the cytosol [52]. Most strikingly, the generation of PI4P hotspots at the plasma membrane in yeast requires similar molecular components indicating a very



**Fig. 1.** Organisation and subcellular compartmentation of PI4P synthesis. The four mammalian PI 4-kinases have different subcellular localisations. The main pool of plasma membrane PI4P is maintained by PI4KIII $\alpha$  which has been imaged in separate studies either in association with the ER or recruited to the plasma membrane by EFR3B and TTC7B. PI4KIII $\beta$  and PI4KII $\alpha$  generate PI4P at the *trans*-Golgi network and PI 4-kinase activity on these membranes is required for the formation of constitutive plasma membrane vesicular carriers, secretory vesicles and in trafficking to late endosomes. Additionally, PI4P at the TGN targets lipid transfer proteins such as FAPP2 and CERT which effect non-vesicular transfer of glucosylceramide and ceramide to late Golgi compartments. PI4KII $\beta$  has been observed in the cytosol and in association with membranes, here we show PI4KII $\beta$  in association with intracellular endosomal-like vesicles. PI4KII $\alpha$  also localises to late endosomes where it modulates the trafficking and degradation of internalised ligand-activated receptors and cargo sorting during TGN to late endosome trafficking.

high degree of evolutionary conservation for this process [53]. These recent insights also revealed a major function for mammalian PI4KIII $\alpha$  in maintaining the normal proteomic and lipid composition of the plasma membrane as evidenced by the mistrafficking of GPCRs and cholesterol to intracellular compartments in PI4KIII $\alpha$  knock-out cells [52]. In a separate study, Hammond and colleagues have established multiple, non-precursor, roles for PI4P at the plasma membrane [54,55]. The first such function could be considered a general one, whereby PI4P in combination with other negatively charged phosphoinositide species, electrostatically defines the cytoplasmic-facing plasma membrane layer as a negatively charged lipid landscape with the capacity to recruit a range of proteins containing polybasic lipid binding domains such as those found in MARCKS and K-Ras. Secondly they observed that PI4P regulates particular ion channels such as the transient receptor potential vanilloid 1 (TRPV1) cation channel, the activity of which was found to be inhibited through either selective depletion of PI4P or PI(4,5)P<sub>2</sub> [54]. Additionally, PI4P, at physiologically relevant concentrations as low as 2 mol%, can induce curvature in model membranes [56]. Therefore, PI4P can contribute to the biochemical and biophysical identity of the plasma membrane, and this multifunctional pool of PI4P is generated principally by PI4KIII $\alpha$ . The other wortmannin-sensitive isoform, PI4KIII $\beta$  [57–60] (92 kDa) does not have a major role in the generation of plasma membrane PI4P but instead seems to mainly function in the generation of Golgi-derived carriers [8,31,61–63]. In line with these trafficking functions, PI4KIII $\beta$  can interact with Arf1 [63–65], neuronal calcium sensor-1 (NCS-1) [9,10,65–67] and the Rab11 GTPase [8]. In addition to the Golgi apparatus, PI4KIII $\beta$  has also been visualised on lysosomes [68] where it functions to maintain lysosomal membrane integrity and in a PKD-phosphorylated form in the nucleus [69] where its physiological function has yet to be determined.

In summary, the markedly different subcellular distributions of the four PI 4-kinases give rise to highly compartmentalised PI4P synthesis and thus organelle-specific functions for this phosphoinositide species [5,70,71].

### 3. Cell signalling

The requirement for PI4P synthesis and resupply during receptor-activated phosphoinositide signalling was one of the principal spurs for early studies aimed at identifying and purifying the mammalian PI 4-kinases. Early work demonstrated the presence of an EGF-activated PI 4-kinase activity in membrane fractions that was associated with a decreased apparent  $K_m$  for PI substrate but with no increase in the  $V_{max}$  for PI phosphorylation activated EGFR [72,73]. Subsequent investigations demonstrated that activated EGFR could be co-immunoprecipitated in complex with a PI4KII activity, PLC $\gamma$  and phosphatidylinositol transfer protein [74,75] (which was hypothesized to transfer PI substrate to the lipid kinases during ligand-activated signalling). Further detailed analyses of phosphoinositide turnover kinetics during receptor-stimulated PLC signalling, demonstrated a requirement for a wortmannin-inhibited PI4KIII activity in the generation of a signaling pool of PI4P [76,77]. A subsequent study that used RNA interference to abrogate the expression of individual PI 4-kinase isozymes discovered that wortmannin-sensitive PI4KIII $\alpha$  was primarily responsible for maintenance of the plasma membrane phosphoinositide pool which is turned over rapidly during GPCR signalling [14,27].

A possible disconnection between sites of PI4P synthesis and utilisation was revealed in elegant studies that used organelle-targeted, recombinant phosphoinositide phosphatases such as Sac1, to deplete PI4P levels at particular subcellular membranes [16,54]. Using this strategy Golgi-compartmentalised PI4P synthesis was shown to be required during the substrate replenishment phase of the angiotensin-stimulated PLC response [16]. At first glance it seems counterintuitive that PI4P synthesis at distal sites can supply the plasma membrane during ligand-activated signalling. However, the recent discovery of stable inter-organelle membrane contacts which facilitate lipid transfer between different organelles [78,79] is worth considering in this context. Most strikingly, a precedent for such inter-organelle regulation of phosphoinositide concentrations has been demonstrated in yeast where the ER-associated PI4P phosphatase Sac1, can act *in trans* at inter-membrane contact sites to dephosphorylate PI4P at the plasma membrane [80]. This process requires the PI4P-binding Osh3 (oxysterol homology 3) protein that appears to simultaneously function both as a plasma membrane PI4P sensor and activator of the Sac1 PI4P phosphatase activity at intermembrane contact sites. Additionally, 6 proteins have been identified that mediate ER-plasma membrane tethering and regulate plasma membrane PI4P levels. These proteins are the vesicle-associated membrane protein-associated proteins (VAP) Scs2 and Scs22 which are similar to mammalian VAP proteins that target FFAT (two phenylalanines in an Acidic Tract [81,82]) motif-containing proteins to the ER; 3 tricalbin protein isoforms, Tcb1, Tcb2 and Tcb3 that contain membrane spanning, lipid binding and multiple C2 domains and are related to the mammalian synaptotagmins [83,84]; and Ist2, which contains multiple transmembrane spanning domains and is orthologous to the mammalian TMEM16 family of Ca<sup>2+</sup>-activated chloride channels [85–88]. Interestingly, Ist2 has been shown in an unconnected study to mediate interactions between the plasma membrane and ER [89].

While mammalian cells express orthologues of the yeast proteins required for the formation of ER-plasma membrane contacts and the control of plasma membrane PI4P concentrations, it is not yet known if an exactly analogous regulatory system exists in higher eukaryotes. Nonetheless, it is noteworthy that in mammalian cells, transient ER plasma membrane contacts are known to form during agonist-stimulated PLC and Ca<sup>2+</sup> signalling in the capacitive re-entry phase when extracellular Ca<sup>2+</sup> replenishes depleted ER stores [79,90–92]. This store refilling event is mediated

by direct interactions between an ER protein STIM1 and a plasma membrane Ca<sup>2+</sup> channel Orai1 [91,92]. Therefore, this Ca<sup>2+</sup>-triggered membrane contact event demonstrates that it may be feasible for an ER-associated PI 4-kinase to supply phosphoinositide substrate to receptors during signalling. Of relevance to this proposal are the findings that PI4KIII $\alpha$  inhibition results in decreased store-operated Ca<sup>2+</sup> entry [90] and that PI4KIII $\alpha$  expression is required for the formation of STIM1-Orai1 intermembrane contacts [52]. Intriguingly, we have recently discovered that CDP-DAG synthase and phosphatidylinositol synthase – two enzymes required to produce the PI 4-kinase substrate PI, also localise to ER membrane microdomains that are in close contact with the plasma membrane [93]. Hence, it seems more and more likely that inter-organelle membrane contact sites play a pivotal role in the cellular organization of receptor-evoked phosphoinositide signalling (Fig. 1).

Since non-vesicular lipid transfer between contacting membrane sites tends to be high flux and energy independent [94,95] such a mechanism would be particularly well suited to supporting large scale and rapid turnover of PI4P pools following GPCR activation. These insights may mean that apparent spatial restrictions on PI4P generation due to the differential organelle targeting of the PI 4-kinases may not be as restrictive for signalling as previously envisaged. Furthermore, the ability for lipids to be transferred between closely apposed membranes suggests that the dynamics of organelle contact site formation and dissolution may impact on the spatiotemporal control of cellular PI4P metabolism.

While PI4KIII $\alpha$  has a prominent role in GPCR stimulated PLC signalling which tends to be a very dramatic and robust response mediated by G-protein activated PLC $\beta$ , the situation differs with the two PI4KII isoforms which seem to have at best minor roles in the agonist-stimulated Ca<sup>2+</sup> response [14,90]. However there are specific instances of PI4KII $\alpha$  being important for the regulation of receptor tyrosine kinase [15,33] and Wnt signalling [13,96,97]. In particular, TGN/endosome localised PI4KII $\alpha$  regulates the endosomal traffic of activated EGFR [33] and knockdown of this enzyme retards the degradation of internalised EGFR. Further evidence for this isozyme modulating receptor signalling has emerged from a study which demonstrated a role for PI4KII $\alpha$  in regulating Wnt3a signalling through Frizzled seven transmembrane receptors [96,97]. This particular mode of canonical Wnt signalling is associated with  $\beta$ -catenin and T cell factor-induced gene transcription rather than Wnt-activated PLC and Ca<sup>2+</sup> signalling. These studies found that PI4KII $\alpha$  is required for PI(4,5)P<sub>2</sub> formation and Wnt3a-dependent phosphorylation of the low-density lipoprotein receptor-related protein 6 (LRP6) [96]. Furthermore, Dishevelled (Dvl) – a cytosolic protein recruited to Wnt3a-bound Frizzled receptors, forms a complex with PI4KII $\alpha$  resulting in a doubling of lipid kinase activity [97]. Conversely it was later discovered that Itch, a HECT-type E3 ubiquitin ligase that regulates the endosomal trafficking of Wnt-activated Frizzled receptors, inhibits PI4KII $\alpha$  activity and that this effect is mediated by direct association of the Itch WW domains with a PPXY motif of PI4KII $\alpha$  (residues 15PPDY<sub>18</sub> in the amino terminus of PI4KII $\alpha$ ) [13]. Whilst PI4P synthesis is inhibited by the PI4KII $\alpha$  Itch complex formation, the ubiquitin ligase activity of Itch is enhanced [13]. Hence, the opposing effects of Dvl and Itch are consistent with activated PI4P synthesis being important in the early events of Wnt signalling, particularly for phosphorylation of the LRP6 co-receptor, but with a non-catalytic role for PI4KII $\alpha$  becoming predominant during the later stages of Wnt-receptor trafficking and sorting via the ubiquitin pathway. Interestingly while RNAi studies have not implicated the PI4KIIIs in Wnt signalling, there is evidence that both PI4KIII $\alpha$  and PI4KIII $\beta$  are required for activation of Hedgehog signalling in mammalian cells – a process which in *Drosophila* is thought to involve PI4P upregulating



vesicular trafficking of the Smoothed receptor to the plasma membrane [98].

Unexpectedly, non-catalytic functions of the PI 4-kinases may also be important in regulating receptor-stimulated phosphoinositide signalling. This can particularly be seen in the case of the PI4KII $\alpha$  isoform where recombinant overexpression of a lipid kinase-inactive mutant impairs EGF-stimulated Akt activation [15]. Overexpression of kinase-inactive PI4KII $\alpha$  does not inhibit endogenous PI4P synthesis but induces defective TGN-endosomal trafficking as evidenced by enlarged late endosomes and reduced transferrin receptor trafficking to recycling endosomes [7,26,33]. Therefore, the effect on Akt activation can be attributed to defective intracellular vesicle trafficking rather than reduced PI4P substrate supply and subsequent PI(3,4,5)P<sub>3</sub> generation [15]. PI4KII $\alpha$  expression levels determine the time course of EGFR trafficking [33] to the lysosome and are important for the recruitment of clathrin adaptors during Golgi-endosomal trafficking [6,7,11,32,34]. Moreover, these trafficking functions are mediated through both lipid kinase and modular protein binding properties of the enzyme. Therefore, non-lipid kinase functions may regulate phosphoinositide signalling pathways, albeit indirectly through effects on the intracellular trafficking dynamics of signalling proteins.

Although it only accounts for a small fraction of overall cellular PI4P production, the PI4KII $\beta$  isoform has been reported to undergo membrane recruitment following receptor activation [38]. The mechanism ascribed for this regulatory event is unique amongst the PI kinases and it involves in the absence of agonist, PI4KII $\beta$  being sequestered in a cytosolic complex with Hsp90 where it is catalytically inactive but at the same time protected from ubiquitination and consequent proteolytic degradation. Following receptor tyrosine kinase activation, the Hsp90 interaction is interrupted and PI4KII $\beta$  undergoes membrane translocation [38] and possible activation by the Rac GTPase. [99] PI4KII $\beta$  palmitoylation is associated with membrane recruitment and catalytic activity but unlike the PI4KII $\alpha$  isoform it is not constitutively palmitoylated [38,100]. Membrane-associated PI4KII $\beta$  is subject to agonist-dependent phosphorylation in its structurally unique amino terminal domain but this post-translational modification does not alter its lipid kinase activity [38]. Biochemical evidence suggests that this isozyme can also associate with crosslinked T cell receptor CD3 zeta chains following tyrosine phosphorylation of the receptor [101]. PI4KII activity has been observed by several groups in complex with tetraspanins (transmembrane-4 superfamily proteins) [40,102–107] which are cell surface, plasma membrane spanning proteins with important roles in adhesion and signalling [108–110]. In the case of the tetraspanin CD81, PI4KII $\beta$  has been shown to be the co-immunoprecipitated isoform [40]. However, the molecular mechanisms that determine PI4KII $\beta$  interactions with either tetraspanins or tetraspanin-enriched membrane domains have yet to be elucidated.

#### 4. Interactions with other lipid pathways

There are now multiple reports that PI4P generation on Golgi membranes is intrinsically linked with the synthesis of other lipids specifically glycosphingolipids and sphingomyelin, and that this is mediated by lipid transfer proteins that associate with Golgi membranes via PI4P-binding PH domains [5]. In addition to its role in promoting non-phosphoinositide lipid production there is also evidence that PI 4-kinase activity is itself modulated by membrane composition [111–113] and thus PI4P synthesis plays a key role in sensing and controlling the concentrations of a variety of lipid classes on late Golgi membranes.

#### 4.1. Sterol regulation of PI4KII $\alpha$

PI4KII $\alpha$  localises mainly to the TGN and endosomes and is only PI kinase enzyme that constitutively associates with membranes [28,29,100,114–116]. Different to all the other PI kinases, the biochemistry and enzymology of PI4KII $\alpha$  is largely defined by its targeting to cholesterol and glycosphingolipid-enriched microdomains of the TGN and the modulation of its lipid kinase activity by membrane sterol and oxysterol concentrations [28,100,111–113,116,117]. Work from our laboratory has shown that manipulating the sterol concentration of intracellular membranes with methyl- $\beta$ -cyclodextrin results in changes to the morphology and size of these membranes which affects both the diffusion rate and mobile fraction of the enzyme [112]. There is also evidence that enhanced sterol concentrations augment the catalytic activity of PI4KII $\alpha$  [111–113]. Targeting of PI4KII $\alpha$  to TGN rafts is achieved through dual-palmitoylation of cysteines within a CCPC motif located in the catalytic region of the protein by Golgi-associated palmitoyl transferases [100,115–117]. Interestingly, tight membrane association of PI4KII $\alpha$  does not necessitate its prior palmitoylation since palmitoyl-mutants remain tightly membrane bound. However, palmitoylation is indispensable for PI phosphorylation and for the correct subcellular targeting of the enzyme [100,115]. Most recent work has revealed that PI4KII $\alpha$  is in fact palmitoylated by DHH3 and DHH7 which are two Golgi-associated palmitoyl acyltransferases [117]. PI4KII $\alpha$  can be co-immunoprecipitated from cholesterol-rich lipid raft-like domains of the TGN in complex with these palmitoyl acyltransferases and these interactions are lost following sterol depletion with methyl- $\beta$ -cyclodextrin [117]. In this way reversible palmitoylation represents a unique cholesterol-sensitive regulatory mechanism that simultaneously links the lipid kinase activity of PI4KII $\alpha$  with its targeting to raft-like microdomains of TGN-endosomal membranes [117]. Another way of viewing these biochemical relationships is to consider PI4KII $\alpha$  localisation and activity at this subcellular locus as reporters for the sterol content and lipid microdomain organization of the tubulovesicular TGN compartment [111,112].

#### 4.2. PI 4-kinases & sphingomyelin synthesis

Sphingomyelin is generated from phosphatidylcholine and ceramide at the TGN in a reaction catalysed by sphingomyelin synthase 1 which also produces diacylglycerol (DAG). However, ceramide is synthesized in the ER and therefore must be transported to the TGN in order for sphingomyelin synthesis to take place. The transfer of ceramide to the Golgi is effected by CERT – a lipid transfer protein [118–120]. Structurally, in addition to carboxy-terminal StART lipid-binding domain, CERT contains a FAAT motif [81,82,121–123] which can bind to ER-localised VAP proteins and in its amino terminus a PH domain which specifically binds PI4P at the Golgi. The positioning of a Golgi-interacting PH domain and ER interacting FFAT motif at opposite ends of CERT facilitates the simultaneous binding of two compositionally distinct membranes and the non-vesicular transfer of ceramide across this protein-mediated inter-organelle contact site [118,120]. Through pharmacological inhibition and RNA interference studies the Balla laboratory established that the Golgi/TGN-localised PI4KIII $\beta$  isoform is required for CERT-mediated non-vesicular trafficking of ceramide to the TGN [4]. In this way, PI4KIII $\beta$  controls the supply of substrate to sphingomyelin synthase and thereby the rate of sphingomyelin production at the TGN. More recently, a study from Banerji and colleagues has detailed how OSBP – another PH domain containing protein, regulates sterol levels at the TGN which affects PI4KII $\alpha$  activity, PI4P-dependent CERT recruitment and consequently sphingomyelin production [111]. This places PI4KII $\alpha$  downstream of OSBP and of critical importance in

integrating changes to cholesterol concentration with sphingomyelin synthesis. Since cholesterol and sphingomyelin are proposed to be concentrated within lipid rafts, one inference from these studies is that PI4KII $\alpha$  is a key enzyme in driving lipid raft formation at the TGN. While this is an attractive model it should be noted that the Balla laboratory only found a minor role for PI4KII $\alpha$  in CERT recruitment to the TGN [4], hence there is some debate over which PI 4-kinase isoform is most important for maintaining sphingomyelin levels.

The reaction catalyzed by sphingomyelin synthase also produces DAG [124] which is required for activation [125] and targeting Protein Kinase D (PKD) to TGN membranes [126,127]. Furthermore, PI4KIII $\beta$  catalytic activity is activated through PKD catalysed phosphorylation of serine 294 [62,128]. This suggests the existence of an integrated lipid-based regulatory mechanism where the generation of DAG by sphingomyelin synthase leads to increased PI4P production through the DAG-PKD-PI4KIII $\beta$  axis [129]. PKD-induced phosphorylation of PI4KIII $\beta$  upregulates the formation of plasma-membrane destined vesicular carriers in a process that involves the formation of a complex between PI4KIII $\beta$ , 14-3-3 $\gamma$  adaptin dimers and CtBP1-S/BARS [130]. Whilst membrane trafficking to the plasma membrane driven by PI4KIII $\beta$  is increased, ceramide delivery to the TGN is decreased as PKD phosphorylation of CERT on serine 132 inhibits interaction of the CERT PH-domain with PI4P [131]. PKD also serine phosphorylates oxysterol-binding protein (OSBP) another protein implicated in PI4P-dependent sphingomyelin synthesis at the TGN resulting in impaired sterol and oxysterol-dependent recruitment of OSBP [132]. Together these recent findings illustrate the central importance of PKD in the homeostatic control of the TGN lipid composition.

#### 4.3. PI 4-kinases & glycosphingolipid synthesis


The synthesis of complex glycosphingolipids at the Golgi is critical for maintaining the characteristic lipid compositions of the both the TGN and plasma membrane, and depends absolutely on the activity of PI4KIII $\beta$  and to a lesser extent PI4KIII $\alpha$  [133–135]. Four-phosphate adaptor protein 2 (FAPP2) [119,120,136] like CERT,

is a member of the family of lipid transfer proteins which contains an N-terminal PI4P-binding PH domain [119,120,137]. However, unlike CERT, FAPP2 contains a C-terminal glycolipid transfer protein homology domain which is required for efficient transfer of glucosylceramide from *cis*- to *trans*-Golgi compartments, or as laid out in alternative model, the retrograde transfer of glucosylceramide to the ER followed by subsequent transport back to the Golgi for additional enzymatic glycosylation [138]. CERT also contains a FFAT-like motif which may mediate interaction with ER associated VAP proteins [139]. PI4P production by PI4KIII $\beta$  and to a lesser extent PI4KII $\alpha$  has been implicated in the recruitment of FAPP2 to late Golgi membranes [134]. Furthermore, FAPP2 is a component of the recently described PKD- and PI4KIII $\beta$ -regulated molecular complex that facilitates the formation of plasma membrane-destined vesicular carriers [5,130,140].

Since PI4KIII $\beta$  activity is under the control of DAG activated PKD and PI4KII $\alpha$  is responsive to membrane sterol concentrations, it could be said that the Golgi/TGN PI 4-kinases co-ordinate a membrane composition and sensing function that can transduce changes in membrane environment into altered PI4P generation thereby modulating the synthesis of the lipid raft-defining sphingomyelin and glycosphingolipids (Fig. 2). As both PI 4-kinases also mediate vesicle trafficking from the TGN, augmented PI4P synthesis has the potential to modify the lipid composition of distal membranes such as the plasma membrane. Therefore, PI 4-kinases occupy a key role in co-ordinating and integrating the lipid composition of the various post-Golgi membranes with enormous consequences for signalling, trafficking and membrane organization at extra-Golgi loci [5,141,142].

#### 5. Emerging roles for PI 4-kinases in disease

In this section we explore the emerging roles for the PI 4-kinases across a wide range of human disease but particularly in cancer, neurological disease and infections caused by both bacteria and viruses.



**Fig. 2.** Redistribution of PI4P from the plasma membrane to a membranous web during HCV infection. PI4P (green) was imaged by immunostaining with anti-PI4P antibody and the plasma membrane and intracellular pools of this lipid were visualised by microscopy. Nuclei are stained blue with the Hoescht dye and the Golgi protein giantin appears in red. In control Huh7.5 hepatoma cells, there is a sizeable plasma membrane-associated pool of PI4P. However, in cells replicating subgenomic replicons of HCV (genotype 2a) there is a reduction in PI4P at the plasma membrane and a concomitant large increase in intracellular PI4P in the HCV-induced membranous web which originates from the ER. PI4KIII $\alpha$  is the PI 4-kinase implicated in producing both the plasma membrane and membranous web pools of PI4P. This figure is adapted and reproduced from the work of Bianco and co-workers [51].

### 5.1. Cancer

The indications so far from cell based studies are that alterations to individual PI 4-kinase isozyme expression levels can modulate receptor tyrosine kinase, Wnt, integrin and tetraspanin signalling – all of which when deregulated can contribute towards the development of neoplastic disease. However, in the absence of disease causing mutations in the PI 4-kinases, it is not yet established whether alterations to catalytic or non-catalytic functions of the enzymes are most important in cancer (reviewed in [143]). So far, increased expression of PI4KII $\alpha$  and PI4KII $\beta$  has been reported for a range of cancers with increased PI4KII $\alpha$  levels associated with augmented activation of the HER2 receptor kinase pathway, HIF production and angiogenesis [144]. In a separate study, a potential anti-metastatic role for PI4KII $\beta$  has emerged which involves this isozyme promoting the trafficking of CD81 tetraspanin proteins away from the plasma membrane into a population of intracellular trafficking vesicles that also contain actinin thereby inducing anti-migratory remodelling of the actin cytoskeleton [40]. Meanwhile PI4KIII has been identified in non-biased screens as contributing towards a more aggressive metastatic phenotype of pancreatic ductal carcinoma cells [145] and was identified as one of several proteins that mediate resistance to the chemotherapeutic agents gemcitabine [146] and cisplatin [147]. However, unlike the well established case of constitutively active phosphoinositide 3-kinase mutations as found in PI3KCA or phosphatase PTEN deletions which can both elevate PI(3,4,5)P<sub>3</sub> levels and drive oncogenic signaling [148,149], it is significant that neither activating mutations of the PI 4-kinases nor deletions of PI4P phosphatases have yet been discovered in cancer. While there is some evidence that stimulation of PI4KIII $\beta$  activity by the oncogenic eukaryotic protein elongation factor eEF1A2 may have a role in the development of metastatic breast cancer [150–152], it is not yet clear whether increased catalytic or non-catalytic functionality of the PI 4-kinases, or even a combination of both properties, are important for oncogenesis. However, the balance of evidence so far suggests that defective vesicular trafficking, an important parameter in mediating the spatiotemporal control of receptor signalling, may be an important contributory factor in malignancies associated with augmented PI 4-kinase expression.

### 5.2. Neurological disease

Decreases in PI 4-kinase expression levels have also been associated with neuronal dysfunction and in particular the impaired survival of specific cell populations within the CNS. Reduced PI4KIII $\alpha$  expression in the CA1 region of the hippocampus following surgically induced-transient ischemia [153], while PI4KIII $\beta$  expression is reduced in the brain of rodents subject to chronic ethanol consumption [154]. There is also some evidence that polymorphisms in the PIK4CA gene which encodes for PI4KIII $\alpha$  may be associated with genetic predisposition to schizophrenia [155] and psychiatric disorders associated with chromosome 22q.11 [156,157], although this may be confined to particular, restricted populations [158].

While cell-based studies have revealed important roles for the PI 4-kinases in neuronal vesicular trafficking [11] it is not yet clear which PI 4-kinase dependent functions lead to aberrant neuronal survival. However, Genetrap mice which do not express PI4KII $\alpha$  are subject to selective cerebellar cell loss and progression to cerebellar spinal degeneration which is ultimately fatal [159]. Interestingly, transgenic animals with knocked-out PI4KIII $\alpha$  expression or those with a conditional knock-in of a lipid kinase inactive version of this isozyme do not exhibit any neurological abnormalities [160]. This may be an indication that the non-catalytic functions of PI4KII $\alpha$  may be important for neuronal survival

since unlike PI4P generation, these isoform-specific functions are less likely to be compensated for by other isozymes. Cell based studies have shown that PI4KII $\alpha$  is recruited to clathrin-coated vesicles through interactions with the clathrin adaptor AP-3 and is part of the protein complex that transports proteins such as dysbindin [11] and calcyon [161]. These observations intimate that at least for PI4KII $\alpha$ , alterations to clathrin-dependent intracellular trafficking could possibly be important for maintaining neuronal viability.

### 5.3. Bacterial infection

All of the mammalian PI 4-kinases have roles in either bacterial entry or replication. In the case of *Listeria monocytogenes*, which enters the cell through initial binding of the bacterial protein InIB to the Met hepatocyte growth factor receptor in membrane sites that also contain the tetraspanin CD81 [162]. Additionally both PI4KII $\alpha$  and PI4KII $\beta$  are required for bacterial internalisation and this may be in part mediated by these isozymes regulating cell surface levels of CD81 through affects on the intracellular trafficking dynamics of this protein [40]. This is an interesting parallel with the proposed role for PI4KII $\beta$  in hepatocellular carcinoma where alterations to cell adhesion and motility are due to this isoform regulating cell surface levels of CD81 [40]. PI4KII $\alpha$  has also been localised along with the PI4P producing PI(4,5)P<sub>2</sub> 5-phosphatase OCRL to the surface of *Chlamydia trachomatis* inclusions [163]. These bacterial inclusions are non-acidified vacuoles that fuse with the Golgi and multivesicular bodies during the cytosolic remodeling phase of their development. Protected within inclusions, *Chlamydiae* can survive and replicate whilst avoiding trafficking to the lysosomes. Further evidence for PI 4-kinase involvement in this disease mechanism stems from the observation that the PI4P-binding and ceramide transporting protein CERT is recruited to the inclusions at sites of interorganelle contact with the ER [164,165]. Therefore, PI4P synthesis on *Chlamydial* inclusions sustains the lipid identity of this replicative membrane compartment and thus the trafficking of this structure away from lysosomes but towards the perinuclear region of the cell.

The PI4KIII $\beta$  isoform has been implicated in the replication of *Legionella pneumophila* [166], a process which occurs inside intracellular vacuoles and is concomitant with the subversion and interference of host cell lipid signalling and trafficking pathways [167]. Amongst the bacterial proteins translocated into the host cytosol are two proteins called SidC and SidM/DrrA [167–169] that compete for a limited pool of PI4KIII $\beta$ -synthesised PI4P on the cytosolic-facing vacuolar membrane. DrrA is of particular note since the carboxy terminal helical domain of this protein mediates the highest yet recorded binding affinity for PI4P [169]. These PI4P-anchored proteins can redirect membrane trafficking by an unusual mechanism involving the recruitment and activation of Rab1b by DrrA-catalyzed adenosine monophosphorylation (AMPylation) of the small GTPase on tyrosine 77 [170]. This PI4P-dependent, covalent modification of Rab1b causes ER-derived vesicles to be recruited to the replicative vacuole resulting in a major diversion of intracellular membrane trafficking pathways [170–172]. Therefore, bacterial entry is dependent on PI 4-kinases to transport the bacterium into the cell in association with endogenous cell surface binding partners while replication depends on silencing and masking of phagosomal PI4P functions which would otherwise result in host-mediated elimination.

Finally, PI4KIII $\alpha$  has been identified in an RNA interference screen as an endogenous host factor along with an ubiquitin hydrolase USP22, and the ubiquitin ligase CDC27, which are required for the intracellular replication of *Francisella tularensis* [173]. Unlike the scenarios with *Chlamydia* and *Legionella* where PI4P production maintains a pro-replicative membrane environ-

ment through subversion of host membrane trafficking, PI4KIII $\alpha$  is required during the later cytosolic phase of *Francisella* proliferation. Hence these recent findings demonstrate that PI 4-kinases can regulate multiple steps in different phases of infection by pathogenic bacteria and with the possible exception of PI4KIII $\alpha$  this occurs by molecular exploitation and modulation of host endosomal trafficking pathways that are dependent on PI4P generation.

#### 5.4. Viral disease

In the past three years there has been huge progress in identifying PI 4-kinases as essential host factors required for the propagation of a number of RNA viruses including those responsible for human diseases such as Hepatitis C [49,160,174–184], polio [185] and severe acute respiratory syndrome (SARS) [186]. Similar to the emerging functions for PI 4-kinases in bacterial infection, PI4P production has been implicated in the generation of intracellular membranous web-like structures that provide an environment supportive of pathogen replication. This could be viewed as further evidence for the role of PI 4-kinases and particularly the PI4KIIs in promoting organelle biogenesis at the level of lipid synthesis. One important difference with all other pathologies involving the PI 4-kinases is that viral proteins actively stimulate PI4P synthesis through direct binding [49]. To date, PI4KIII $\beta$  has been implicated in the intracellular RNA replication of several enteroviruses [185,187,188] and in the endosomal fusion phase of SARS coronavirus [186] infection. While in the case of the Hepatitis C virus (HCV), PI4KIII $\alpha$  has emerged from RNA interference screens and pharmacological studies as a host protein that is bound and activated by the non-structural viral protein NS5A [49,181] to produce large amounts of PI4P and thus an ER-derived membranous web that promotes viral replication [51,174,189] (Fig. 2). This represents the first example of activated PI 4-kinase activity having a pathological role. Unlike the scenario with PI4P on the surface of bacterial vacuoles which is exploited by bacterial proteins to redirect intracellular trafficking there is as yet little evidence that formation of the large ER-derived PI4P rich web structure impairs vesicular trafficking along the secretory pathway. Indeed the presence of the Golgi-localised PI4KIII $\beta$  isoform is required for the propagation of Hepatitis C virus [182,183] and enteroviruses [185,188].

There is a possibility that the PI4P enriched surface of the membranous web may act as a docking surface for host, as opposed to viral proteins that contain apposite binding domains. In concordance with this hypothesis both OSBP [190] and CERT [191] – proteins with key roles in non-phosphoinositide lipid transfer and sensing at the TGN, have been identified as additional host factors required for HCV replication. This raises another issue as to whether such a massive enrichment of PI4P molecularly defines the identity of this virally induced organelle in terms of its surface electrostatics and also in the recruitment and homeostasis of other components of this membrane compartment [192]. Of relevance to this idea is the observation that host sphingolipid synthesis, a process known to be under the control of PI 4-kinases, is known to be greatly upregulated during HCV infection [193,194]. In addition, the PI4P-binding protein OSBP, which may control CERT-mediated ceramide transfer to sphingomyelin synthase, interacts with the amino terminal domain I of the HCV protein NS5A protein at the TGN and is required for the process of HCV maturation [190,191].

Another possible outcome of virally-induced PI4KIII $\beta$  activation is greatly enhanced signalling due to elevated PI4P substrate levels. Consistent with this proposal, a number of reports have demonstrated HCV-induced alterations to pro-oncogenic signalling pathways [195–199]. It remains to be established though, if elevated PI4P at the virally-induced compartment constitutes a signalling competent pool capable of supplying substrate to upstream phos-

phoinositide kinases and phospholipases at the plasma membrane and endosomes. Another proviso is that alterations to signalling and trafficking could be accounted for by the interactions of viral proteins with multiple host proteins. As an example, in addition to PI4KIII $\beta$ , NS5A can bind other host proteins including the cellular retinoic acid binding protein 1, centaurin $\delta$ 2 [174] and other lipid metabolizing enzymes such as calcium-dependent, group IVA phospholipase A2 [200]. In terms of PI signalling, Centaurin $\delta$ 2, also known as ARAP1, is especially interesting as it is a multi-domain protein that contains phosphoinositide binding PH domains, ankyrin repeats as well as GAP domains specific for Arf and Rho GTPases [201,202]. Furthermore, the related protein ARAP2 has been implicated in the internalization of *Listeria monocytogenes* which suggests the engagement of similar host factors for bacterial and viral infections [203]. Therefore it seems likely that alterations to PI4KIII $\alpha$  activity may contribute to an overall systems level change in signalling outputs during HCV infection [192,200].

#### 6. Pharmacological targeting of the PI 4-kinases and future perspectives

Due to their emerging and important role in several diseases there has been some interest in developing small molecule, isoform-selective inhibitors of the mammalian PI 4-kinases. There has been some progress in identifying reasonably specific inhibitors of the PI4KIIs. Particularly noteworthy in this regard are PIK93, the anti-enteroviral T-00127-HEV1 and some aminoimidazole inhibitors developed by Novartis [183] all of which selectively inhibit PI4KIII $\beta$ . Although there is now one report demonstrating that some coxsackievirus mutants can overcome their dependence on PI4P for their intracellular replication [187], suggesting that the acquisition of resistance may become a problem in targeting host PI4KIII $\beta$  in viral disease. As regards PI4KIII $\alpha$  the anti-viral molecules AL-9 – a 4-anilino quinazoline molecule [51], and the Boehringer Ingelheim compounds A and B [160] are all newly identified, selective inhibitors of this isoform. This contrasts with the current situation with the PI4KIIs where with the exception of less specific molecules such as resveratrol [204,205], and epigallocatechin gallate [206] there has been little progress. Nevertheless, the pharmacological targeting of individual isoforms may not be straightforward and this may in part due to the multiplicity of cell functions controlled by the PI 4-kinases coupled with compensation particularly at the level of PI4P synthesis – a common function shared by all isoforms. Hence to effectively silence PI 4-kinase-dependent trafficking, it may also be necessary to inhibit their modular protein-binding functions. A good example of such a scenario is the role of PI4KII $\alpha$  in the recruitment of AP-3 on late endosomal membranes as this function requires both PI4P synthesis and a dileucine AP-3 interaction motif [7]. Therefore, co-incident targeting of more than one aspect of PI 4-kinase structure may be required to silence a single biological function. To extend this idea, there are now documented instances where more than one PI 4-kinase controls distinct steps along the same trafficking pathway. This has been observed in yeast in the sequential recruitment of AP-1 and clathrin adaptors at the TGN [207], and in mammalian cells in the trafficking of  $\beta$ -glucocerebrosidase enzyme from the TGN to lysosomes in distinct steps requiring PI4KIII $\beta$  and PI4KII $\alpha$  consecutively [17]. These new insights suggest that it may be necessary to inhibit the activity of more than one PI 4-kinase isoform to comprehensively suppress a particular biological pathway.

A further challenge in targeting the PI 4-kinases has emerged from more recent studies which have revealed differential cell and tissue dependencies on the PI 4-kinase isoforms that are not always predictable. As an example, animal studies have shown that



genetic knockdown of PI4KIII $\alpha$  leads to severe changes in the gastrointestinal mucosal epithelium [160] whereas loss of PI4KIII $\alpha$  leads to selective loss of specific neuronal cell populations such as cerebellar Purkinje cells [159]. Therefore, it is not yet clear if inhibiting individual PI 4-kinase isoforms in a therapeutic setting represents a feasible strategy. However, a more comprehensive knowledge of the roles of different PI 4-kinase permutations in modulating lipid metabolism, signalling and trafficking, may illuminate which PI4P pathways to target in human disease.

## Acknowledgement

The authors acknowledge support from the BBSRC (grant BB/G021163/1) and the Royal Free Charity.

## References

- [1] Balla A, Balla T. Phosphatidylinositol 4-kinases: old enzymes with emerging functions. *Trends Cell Biol* 2006;16:351–61.
- [2] Minogue S, Waugh MG. The phosphatidylinositol 4-kinases: don't call it a comeback. *Subcell Biochem* 2012;58:1–24.
- [3] Bunney TD, Katan M. Phosphoinositide signalling in cancer: beyond PI3K and PTEN. *Nat Rev Cancer* 2010;10:342–52.
- [4] Toth B, Balla A, Ma H, Knight ZA, Shokat KM, Balla T. Phosphatidylinositol 4-kinase IIIbeta regulates the transport of ceramide between the endoplasmic reticulum and Golgi. *J Biol Chem* 2006;281:36369–77.
- [5] Godi A, Di Campli A, Konstantakopoulos A, Di Tullio G, Alessi DR, Kular GS, et al. FAPPs control Golgi-to-cell-surface membrane traffic by binding to ARF and PtdIns(4)P. *Nat Cell Biol* 2004;6:393–404.
- [6] Wang YJ, Wang J, Sun HQ, Martinez M, Sun YX, Macia E, et al. Phosphatidylinositol 4 phosphate regulates targeting of clathrin adaptor AP-1 complexes to the Golgi. *Cell* 2003;114:299–310.
- [7] Craige B, Salazar G, Faundez V. Phosphatidylinositol-4-kinase type II alpha contains an AP-3 sorting motif and a kinase domain that are both required for endosome traffic. *Mol Biol Cell* 2008.
- [8] de Graaf P, Zwart WT, van Dijken RA, Deneka M, Schulz TK, Geijssen N, et al. Phosphatidylinositol 4-kinasebeta is critical for functional association of rab11 with the Golgi complex. *Mol Biol Cell* 2004;15:2038–47.
- [9] Taverna E, Francolini M, Jeromin A, Hilfiker S, Roder J, Rosa P. Neuronal calcium sensor 1 and phosphatidylinositol 4-OH kinase beta interact in neuronal cells and are translocated to membranes during nucleotide-evoked exocytosis. *J Cell Sci* 2002;115:3909–22.
- [10] Zhao X, Varnai P, Tuymetova G, Balla A, Toth ZE, Oker-Blom C, et al. Interaction of neuronal calcium sensor-1 (NCS-1) with phosphatidylinositol 4-kinase beta stimulates lipid kinase activity and affects membrane trafficking in COS-7 cells. *J Biol Chem* 2001;276:40183–9.
- [11] Larimore J, Tornieri K, Ryder PV, Gokhale A, Zlatic SA, Craige B, et al. The schizophrenia susceptibility factor dysbindin and its associated complex sort cargoes from cell bodies to the synapse. *Mol Biol Cell* 2011;22:4854–67.
- [12] Salazar G, Craige B, Wainer BH, Guo J, De Camilli P, Faundez V. Phosphatidylinositol-4-kinase type II alpha is a component of adaptor protein-3-derived vesicles. *Mol Biol Cell* 2005;16:3692–704.
- [13] Mossinger J, Wiewfer M, Krause E, Freund C, Gerth F, Krauss M, et al. Phosphatidylinositol 4-kinase IIalpha function at endosomes is regulated by the ubiquitin ligase Itch. *EMBO Rep* 2012.
- [14] Balla A, Kim YJ, Varnai P, Szentpetery Z, Knight Z, Shokat KM, et al. Maintenance of hormone-sensitive phosphoinositide pools in the plasma membrane requires phosphatidylinositol 4-kinase III(alpha). *Mol Biol Cell* 2008;19:711–21.
- [15] Chu K, Minogue S, Hsuan J, Waugh M. Differential effects of the phosphatidylinositol 4-kinases, PI4KIIalpha and PI4KIIIbeta, on Akt activation and apoptosis. *Cell Death Dis* 2010;1:e106.
- [16] Szentpetery Z, Varnai P, Balla T. Acute manipulation of Golgi phosphoinositides to assess their importance in cellular trafficking and signaling. *Proc Natl Acad Sci USA* 2010;107:8225–30.
- [17] Jovic M, Kean MJ, Szentpetery Z, Polevoy G, Gingras AC, Brill JA, et al. Two phosphatidylinositol 4-kinases control lysosomal delivery of the Gaucher disease enzyme, beta-glucocerebrosidase. *Mol Biol Cell* 2012;23:1533–45.
- [18] Wong K, Meyers dd R, Cantley LC. Subcellular locations of phosphatidylinositol 4-kinase isoforms. *J Biol Chem* 1997;272:13236–41.
- [19] Harwood JL, Hawthorne JN. The properties and subcellular distribution of phosphatidylinositol kinase in mammalian tissues. *Biochim Biophys Acta* 1969;171:75–88.
- [20] Michell RH, Harwood JL, Coleman R, Hawthorne JN. Characteristics of rat liver phosphatidylinositol kinase and its presence in the plasma membrane. *Biochim Biophys Acta* 1967;144:649–58.
- [21] Cockcroft S, Taylor JA, Judah JD. Subcellular localisation of inositol lipid kinases in rat liver. *Biochim Biophys Acta* 1985;845:163–70.
- [22] Waugh MG, Lawson D, Tan SK, Hsuan JJ. Phosphatidylinositol 4-phosphate synthesis in immunisolated caveolae-like vesicles and low buoyant density non-caveolar membranes. *J Biol Chem* 1998;273:17115–21.
- [23] Jelsema CL, Morre DJ. Distribution of phospholipid biosynthetic enzymes among cell components of rat liver. *J Biol Chem* 1978;253:7960–71.
- [24] Michell RH, Hawthorne JN. The site of diphosphoinositide synthesis in rat liver. *Biochim Biophys Res Commun* 1965;21:333–8.
- [25] Galliard T, Michell RH, Hawthorne JN. Incorporation of phosphate into diphosphoinositide by subcellular fractions from liver. *Biochim Biophys Acta* 1965;106:551–63.
- [26] Balla A, Tuymetova G, Barshishat M, Geiszt M, Balla T. Characterization of type II phosphatidylinositol 4-kinase isoforms reveals association of the enzymes with endosomal vesicular compartments. *J Biol Chem* 2002;277:20041–50.
- [27] Balla A, Tuymetova G, Tsiomenko A, Varnai P, Balla T. A plasma membrane pool of phosphatidylinositol 4-phosphate is generated by phosphatidylinositol 4-kinase type-III alpha: studies with the PH domains of the oxysterol binding protein and FAPP1. *Mol Biol Cell* 2005;16:1282–95.
- [28] Waugh MG, Minogue S, Blumenkrantz D, Anderson JS, Hsuan JJ. Identification and characterization of differentially active pools of type IIalpha phosphatidylinositol 4-kinase activity in unstimulated A431 cells. *Biochem J* 2003;376:497–503.
- [29] Waugh MG, Minogue S, Anderson JS, Balinger A, Blumenkrantz D, Calnan DP, et al. Localization of a highly active pool of type II phosphatidylinositol 4-kinase in a p97/valosin-containing-protein-rich fraction of the endoplasmic reticulum. *Biochem J* 2003;373:57–63.
- [30] Waugh MG, Chu KM, Clayton EL, Minogue S, Hsuan JJ. Detergent-free isolation and characterization of cholesterol-rich membrane domains from trans-Golgi network vesicles. *J Lipid Res* 2011;52:582–9.
- [31] Weixel KM, Blumental-Perry A, Watkins SC, Aridor M, Weisz OA. Distinct Golgi populations of phosphatidylinositol 4-phosphate regulated by phosphatidylinositol 4-kinases. *J Biol Chem* 2005;280:10501–8.
- [32] Wang J, Sun HQ, Macia E, Kirchhausen T, Watson H, Bonifacio JS, et al. PI4P promotes the recruitment of the GGA adaptor proteins to the trans-Golgi network and regulates their recognition of the ubiquitin sorting signal. *Mol Biol Cell* 2007;18:2646–55.
- [33] Minogue S, Waugh MG, De Matteis MA, Stephens DJ, Berditchevski F, Hsuan JJ. Phosphatidylinositol 4-kinase is required for endosomal trafficking and degradation of the EGF receptor. *J Cell Sci* 2006;119:571–81.
- [34] Salazar G, Zlatic S, Craige B, Peden AA, Pohl J, Faundez V. Hermansky-Pudlak syndrome protein complexes associate with phosphatidylinositol 4-kinase type II alpha in neuronal and non-neuronal cells. *J Biol Chem* 2009;284:1790–802.
- [35] Guo J, Wenk MR, Pellegrini L, Onofri F, Benfenati F, De Camilli P. Phosphatidylinositol 4-kinase type IIalpha is responsible for the phosphatidylinositol 4-kinase activity associated with synaptic vesicles. *Proc Natl Acad Sci USA* 2003;100:3995–4000.
- [36] Xu Z, Huang G, Kandror KV. Phosphatidylinositol 4-kinase type IIalpha is targeted specifically to cellugyrin-positive glucose transporter 4 vesicles. *Mol Endocrinol* 2006;20:2890–7.
- [37] Alfonso Pecchio AR, Cardozo Gizzi AM, Renner ML, Molina-Calavita M, Caputto BL. C-Fos activates and physically interacts with specific enzymes of the pathway of synthesis of polyphosphoinositides. *Mol Biol Cell* 2011;22:4716–25.
- [38] Jung G, Wang J, Wlodarski P, Barylko B, Binns DD, Shu H, et al. Molecular determinants of activation and membrane targeting of phosphoinositol 4-kinase IIbeta. *Biochem J* 2008;409:501–9.
- [39] Jung G, Barylko B, Lu D, Shu H, Yin H, Albanesi JP. Stabilization of phosphatidylinositol 4-kinase type IIbeta by interaction with Hsp90. *J Biol Chem* 2011;286:12775–84.
- [40] Mazzocca A, Liotta F, Carloni V. Tetraspanin CD81-regulated cell motility plays a critical role in intrahepatic metastasis of hepatocellular carcinoma. *Gastroenterology* 2008;135 [244–256 e241].
- [41] Balla T, Downing GJ, Jaffe H, Kim S, Zolyomi A, Catt KJ. Isolation and molecular cloning of wortmannin-sensitive bovine type III phosphatidylinositol 4-kinases. *J Biol Chem* 1997;272:18358–66.
- [42] Nakagawa T, Goto K, Kondo H. Cloning, expression, and localization of 230-kDa phosphatidylinositol 4-kinase. *J Biol Chem* 1996;271:12088–94.
- [43] Wong K, Cantley LC. Cloning and characterization of a human phosphatidylinositol 4-kinase. *J Biol Chem* 1994;269:28878–84.
- [44] Gehrman T, Gulkan H, Suer S, Herberg FW, Balla A, Vereb G, et al. Functional expression and characterisation of a new human phosphatidylinositol 4-kinase PI4K230. *Biochim Biophys Acta* 1999;1437:341–56.
- [45] Szentpetery Z, Szakacs G, Bojireddy N, Tai AW, Balla T. Genetic and functional studies of phosphatidylinositol 4-kinase type IIIalpha. *Biochim Biophys Acta* 1811;2011:476–83.
- [46] Vereb G, Balla A, Gergely P, Wymann MP, Gulkan H, Suer S, et al. The ATP-binding site of brain phosphatidylinositol 4-kinase PI4K230 as revealed by 5'-p-fluorosulfonylbenzoyladenine. *Int J Biochem Cell Biol* 2001;33:249–59.
- [47] Kakuk A, Friedlander E, Vereb Jr G, Kasa A, Balla A, Balla T, et al. Nucleolar localization of phosphatidylinositol 4-kinase PI4K230 in various mammalian cells. *Cytometry A* 2006;69:1174–83.
- [48] Balla A, Vereb G, Gulkan H, Gehrman T, Gergely P, Heilmeyer Jr LM, et al. Immunohistochemical localisation of two phosphatidylinositol 4-kinase isoforms, PI4K230 and PI4K92, in the central nervous system of rats. *Exp Brain Res* 2000;134:279–88.
- [49] Reiss S, Rebhan I, Backes P, Romero-Brey I, Erfle H, Matula P, et al. Recruitment and activation of a lipid kinase by hepatitis C virus NS5A is

- essential for integrity of the membranous replication compartment. *Cell Host Microbe* 2011;9:32–45.
- [50] Dumaresq-Doiron K, Savard MF, Akam S, Costantino S, Lefrancois S. The phosphatidylinositol 4-kinase PI4KIIIalpha is required for the recruitment of GBF1 to Golgi membranes. *J Cell Sci* 2010;123:2273–80.
- [51] Bianco A, Reghellin V, Donnici L, Fenu S, Alvarez R, Baruffa C, et al. Metabolism of phosphatidylinositol 4-kinase IIIalpha-dependent PI4P is subverted by HCV and is targeted by a 4-anilino quinazoline with antiviral activity. *PLoS Pathog* 2012;8:e1002576.
- [52] Nakatsu F, Baskin JM, Chung J, Tanner LB, Shui G, Lee SY, et al. PtdIns4P synthesis by PI4KIIIalpha at the plasma membrane and its impact on plasma membrane identity. *J Cell Biol* 2012;199:1003–16.
- [53] Baird D, Stefan C, Audhya A, Weys S, Emr SD. Assembly of the PtdIns 4-kinase Stt4 complex at the plasma membrane requires Ypp 1 and Efr3. *J Cell Biol* 2008;183:1061–74.
- [54] Hammond GR, Fischer MJ, Anderson KE, Holdich J, Koteci A, Balla T, et al. PI4P and PI(4,5)P2 are essential but independent lipid determinants of membrane identity. *Science* 2012;337:727–30.
- [55] Hammond GR, Schiavo G, Irvine RF. Immunocytochemical techniques reveal multiple, distinct cellular pools of PtdIns4P and PtdIns(4,5)P(2). *Biochem J* 2009;422:23–35.
- [56] Furse S, Brooks NJ, Seddon AM, Woscholski R, Templer RH, Tate EW, et al. Lipid membrane curvature induced by distearoyl phosphatidylinositol 4-phosphate. *Soft Matter* 2012;8:3090–3.
- [57] Meyers R, Cantley LC. Cloning and characterization of a wortmannin-sensitive human phosphatidylinositol 4-kinase. *J Biol Chem* 1997;272:4384–90.
- [58] Nakagawa T, Goto K, Kondo H. Cloning and characterization of a 92 kDa soluble phosphatidylinositol 4-kinase. *Biochem J* 1996;320(Pt 2):643–9.
- [59] Suer S, Sickmann A, Meyer HE, Herberg JW, Heilmeyer Jr LM. Human phosphatidylinositol 4-kinase isoform PI4K92. Expression of the recombinant enzyme and determination of multiple phosphorylation sites. *Eur J Biochem* 2001;268:2099–106.
- [60] Zhao XH, Bondeva T, Balla T. Characterization of recombinant phosphatidylinositol 4-kinase beta reveals auto- and heterophosphorylation of the enzyme. *J Biol Chem* 2000;275:14642–8.
- [61] Bruns JR, Ellis MA, Jeromin A, Weisz OA. Multiple roles for phosphatidylinositol 4-kinase in biosynthetic transport in polarized Madin-Darby canine kidney cells. *J Biol Chem* 2002;277:2012–8.
- [62] Hausser A, Storz P, Martens S, Link G, Tokar A, Pfizenmaier K. Protein kinase D regulates vesicular transport by phosphorylating and activating phosphatidylinositol-4 kinase IIIbeta at the Golgi complex. *Nat Cell Biol* 2005;7:880–6.
- [63] Godi A, Pertile P, Meyers R, Marra P, Di Tullio G, Iurisci C, et al. ARF mediates recruitment of PtdIns-4-OH kinase-beta and stimulates synthesis of PtdIns(4,5)P2 on the Golgi complex. *Nat Cell Biol* 1999;1:280–7.
- [64] Haynes LP, Sherwood MW, Dolman NJ, Burgoyne RD. Specificity, promiscuity and localization of ARF protein interactions with NCS-1 and phosphatidylinositol-4 kinase-III beta. *Traffic* 2007;8:1080–92.
- [65] Haynes LP, Thomas GM, Burgoyne RD. Interaction of neuronal calcium sensor-1 and ADP-ribosylation factor 1 allows bidirectional control of phosphatidylinositol 4-kinase beta and trans-Golgi network-plasma membrane traffic. *J Biol Chem* 2005;280:6047–54.
- [66] Rajebhosale M, Greenwood S, Vidugiriene J, Jeromin A, Hilfiker S. Phosphatidylinositol 4-OH kinase is a downstream target of neuronal calcium sensor-1 in enhancing exocytosis in neuroendocrine cells. *J Biol Chem* 2003;278:6075–84.
- [67] Mikhaylova M, Reddy PP, Munsch T, Landgraf P, Suman SK, Smalla KH, et al. Calneurons provide a calcium threshold for trans-Golgi network to plasma membrane trafficking. *Proc Natl Acad Sci USA* 2009;106:9093–8.
- [68] Sridhar S, Patel B, Aphkhasava D, Macian F, Santambrogio L, Shields D, et al. The lipid kinase PI4KIIIbeta preserves lysosomal identity. *EMBO J* 2012;32:324–39.
- [69] Szivak I, Lamb N, Heilmeyer LM. Subcellular localization and structural function of endogenous phosphorylated phosphatidylinositol 4-kinase [PI4K92]. *J Biol Chem* 2006;281:16740–9.
- [70] De Matteis MA, Di Campli A, Godi A. The role of the phosphoinositides at the Golgi complex. *Biochim Biophys Acta* 2005;1744:396–405.
- [71] De Matteis MA, Luini A. Exiting the Golgi complex. *Nat Rev Mol Cell Biol* 2008;9:273–84.
- [72] Walker DH, Pike LJ. Phosphatidylinositol kinase is activated in membranes derived from cells treated with epidermal growth factor. *Proc Natl Acad Sci USA* 1987;84:7513–7.
- [73] Pike LJ, Eakes AT. Epidermal growth factor stimulates the production of phosphatidylinositol monophosphate and the breakdown of polyphosphoinositides in A431 cells. *J Biol Chem* 1987;262:1644–51.
- [74] Kauffmann-Zeh A, Klinger R, Endemann G, Waterfield MD, Wetzker R, Hsuan JJ. Regulation of human type II phosphatidylinositol kinase activity by epidermal growth factor-dependent phosphorylation and receptor association. *J Biol Chem* 1994;269:31243–51.
- [75] Kauffmann-Zeh A, Thomas GM, Ball A, Prosser S, Cunningham E, Cockcroft S, et al. Requirement for phosphatidylinositol transfer protein in epidermal growth factor signaling. *Science* 1995;268:1188–90.
- [76] Nakanishi S, Catt KJ, Balla T. A wortmannin-sensitive phosphatidylinositol 4-kinase that regulates hormone-sensitive pools of inositolphospholipids. *Proc Natl Acad Sci USA* 1995;92:5317–21.
- [77] Willars GB, Nahorski SR, Challiss RA. Differential regulation of muscarinic acetylcholine receptor-sensitive polyphosphoinositide pools and consequences for signaling in human neuroblastoma cells. *J Biol Chem* 1998;273:5037–46.
- [78] Lebedzinska M, Szabadkai G, Jones AW, Duszynski J, Wieckowski MR. Interactions between the endoplasmic reticulum, mitochondria, plasma membrane and other subcellular organelles. *Int J Biochem Cell Biol* 2009;41:1805–16.
- [79] Varnai P, Toth B, Toth DJ, Hunyady L, Balla T. Visualization and manipulation of plasma membrane-endoplasmic reticulum contact sites indicates the presence of additional molecular components within the STIM1-Orai1 Complex. *J Biol Chem* 2007;282:29678–90.
- [80] Stefan CJ, Manford AG, Baird D, Yamada-Hanff J, Mao Y, Emr SD. Osh proteins regulate phosphoinositide metabolism at ER-plasma membrane contact sites. *Cell* 2011;144:389–401.
- [81] Kaiser SE, Brickner JH, Reilein AR, Fenn TD, Walter P, Brunger AT. Structural basis of FFAT motif-mediated ER targeting. *Structure* 2005;13:1035–45.
- [82] Loewen CJ, Levine TP. A highly conserved binding site in vesicle-associated membrane protein-associated protein (VAP) for the FFAT motif of lipid-binding proteins. *J Biol Chem* 2005;280:14097–104.
- [83] Creutz CE, Snyder SL, Schulz TA. Characterization of the yeast tricalbins: membrane-bound multi-C2-domain proteins that form complexes involved in membrane trafficking. *Cell Mol Life Sci* 2004;61:1208–20.
- [84] Schulz TA, Creutz CE. The tricalbin C2 domains: lipid-binding properties of a novel, synaptotagmin-like yeast protein family. *Biochemistry* 2004;43:3987–95.
- [85] Yang YD, Cho H, Koo JY, Tak MH, Cho Y, Shim WS, et al. TMEM16A confers receptor-activated calcium-dependent chloride conductance. *Nature* 2008;455:1210–5.
- [86] Caputo A, Caci E, Ferrera L, Pedemonte N, Barsanti C, Sondo E, et al. TMEM16A, a membrane protein associated with calcium-dependent chloride channel activity. *Science* 2008;322:590–4.
- [87] Schroeder BC, Cheng T, Jan YN, Jan LY. Expression cloning of TMEM16A as a calcium-activated chloride channel subunit. *Cell* 2008;134:1019–29.
- [88] Almaca J, Tian Y, Aldehni F, Ousingawatt J, Kongsuphol P, Rock JR, et al. TMEM16 proteins produce volume-regulated chloride currents that are reduced in mice lacking TMEM16A. *J Biol Chem* 2009;284:28571–8.
- [89] Wolf W, Kilic A, Schrul B, Lorenz H, Schwappach B, Seedorf M. Yeast Ict2 recruits the endoplasmic reticulum to the plasma membrane and creates a ribosome-free membrane microcompartment. *PLoS One* 2012;7:e39703.
- [90] Korzeniowski MK, Popovic MA, Szentpetery Z, Varnai P, Stojilkovic SS, Balla T. Dependence of STIM1/Orai1-mediated calcium entry on plasma membrane phosphoinositides. *J Biol Chem* 2009;284:21027–35.
- [91] Mullins FM, Park CY, Dolmetsch RE, Lewis RS. STIM1 and calmodulin interact with Orai1 to induce Ca<sup>2+</sup>-dependent inactivation of CRAC channels. *Proc Natl Acad Sci USA* 2009;106:15495–500.
- [92] Park CY, Hoover PJ, Mullins FM, Bachhawat P, Covington ED, Raunser S, et al. STIM1 clusters and activates CRAC channels via direct binding of a cytosolic domain to Orai1. *Cell* 2009;136:876–90.
- [93] Waugh MG, Minogue S, Clayton EL, Hsuan JJ. CDP-diacylglycerol phospholipid synthesis in detergent-soluble, non-raft, membrane microdomains of the endoplasmic reticulum. *J Lipid Res* 2011;52:2148–58.
- [94] Hao M, Lin SX, Karylowski OJ, Wustner D, McGraw TE, Maxfield FR. Vesicular and non-vesicular sterol transport in living cells. The endocytic recycling compartment is a major sterol storage organelle. *J Biol Chem* 2002;277:609–17.
- [95] Prinz WA. Lipid trafficking sans vesicles: where, why, how? *Cell* 2010;143:870–4.
- [96] Pan W, Choi SC, Wang H, Qin Y, Volpicelli-Daley L, Swan L, et al. Wnt3a-mediated formation of phosphatidylinositol 4,5-bisphosphate regulates LRP6 phosphorylation. *Science* 2008;321:1350–3.
- [97] Qin Y, Li L, Pan W, Wu D. Regulation of phosphatidylinositol kinases and metabolism by Wnt3a and Dvl. *J Biol Chem* 2009;284:22544–8.
- [98] Yavari A, Nagaraj R, Owusu-Ansah E, Follick A, Ngo K, Hillman T, et al. Role of lipid metabolism in smoothed derepression in hedgehog signaling. *Dev Cell* 2010;19:54–65.
- [99] Wei YJ, Sun HQ, Yamamoto M, Wlodarski P, Kunii K, Martinez M, et al. Type II phosphatidylinositol 4-kinase beta is a cytosolic and peripheral membrane protein that is recruited to the plasma membrane and activated by Rac-GTP. *J Biol Chem* 2002;277:46586–93.
- [100] Barylko B, Mao YS, Wlodarski P, Jung G, Binns DD, Sun HQ, et al. Palmitoylation controls the catalytic activity and subcellular distribution of phosphatidylinositol 4-kinase II[alpha]. *J Biol Chem* 2009;284:9994–10003.
- [101] Srivastava R, Sinha RK, Subrahmanyam G. Type II phosphatidylinositol 4-kinase beta associates with TCR-CD3 zeta chain in Jurkat cells. *Mol Immunol* 2006;43:454–63.
- [102] Carloni V, Mazzocca A, Ravichandran KS. Tetraspanin CD81 is linked to ERK/MAPK signaling by Shc in liver tumor cells. *Oncogene* 2004;23:1566–74.
- [103] Yauch RL, Hemler ME. Specific interactions among transmembrane 4 superfamily (TM4SF) proteins and phosphoinositide 4-kinase. *Biochem J* 2000;351(Pt 3):629–37.
- [104] Claas C, Wahl J, Orlicky DJ, Karaduman H, Schnolzer M, Kempf T, et al. The tetraspanin D6.1A and its molecular partners on rat carcinoma cells. *Biochem J* 2005;389:99–110.
- [105] Yauch RL, Berditchevski F, Harler MB, Reichner J, Hemler ME. Highly stoichiometric, stable, and specific association of integrin alpha3beta1 with

- CD151 provides a major link to phosphatidylinositol 4-kinase, and may regulate cell migration. *Mol Biol Cell* 1998;9:2751–65.
- [106] Claas C, Stipp CS, Hemler ME. Evaluation of prototype transmembrane 4 superfamily protein complexes and their relation to lipid rafts. *J Biol Chem* 2001;276:7974–84.
- [107] Berditchevski F, Toliass KF, Wong K, Carpenter CL, Hemler ME. A novel link between integrins, transmembrane-4 superfamily proteins (CD63 and CD81), and phosphatidylinositol 4-kinase. *J Biol Chem* 1997;272:2595–8.
- [108] Berditchevski F. Complexes of tetraspanins with integrins: more than meets the eye. *J Cell Sci* 2001;114:4143–51.
- [109] Berditchevski F, Odintsova E, Sawada S, Gilbert E. Expression of the palmitoylation-deficient CD151 weakens the association of alpha 3 beta 1 integrin with the tetraspanin-enriched microdomains and affects integrin-dependent signaling. *J Biol Chem* 2002;277:36991–7000.
- [110] Hemler ME. Tetraspanin functions and associated microdomains. *Nat Rev Mol Cell Biol* 2005;6:801–11.
- [111] Banerji S, Ngo M, Lane CF, Robinson CA, Minogue S, Ridgway ND. Oxysterol binding protein-dependent activation of sphingomyelin synthesis in the golgi apparatus requires phosphatidylinositol 4-kinase IIalpha. *Mol Biol Cell* 2010;21:4141–50.
- [112] Minogue S, Chu KM, Westover EJ, Covey DF, Hsuan JJ, Waugh MG. Relationship between phosphatidylinositol 4-phosphate synthesis, membrane organization, and lateral diffusion of PI4KIIalpha at the trans-Golgi network. *J Lipid Res* 2010;51:2314–24.
- [113] Waugh MG, Minogue S, Chotai D, Berditchevski F, Hsuan JJ. Lipid and peptide control of phosphatidylinositol 4-kinase IIalpha activity on Golgi-endosomal rafts. *J Biol Chem* 2006;281:3757–63.
- [114] Waugh MG, Minogue S, Anderson JS, dos Santos M, Hsuan JJ. Signalling and non-caveolar rafts. *Biochem Soc Trans* 2001;29:509–11.
- [115] Barylko B, Gerber SH, Binns DD, Grichine N, Khvotchev M, Sudhof TC, et al. A novel family of phosphatidylinositol 4-kinases conserved from yeast to humans. *J Biol Chem* 2001;276:7705–8.
- [116] Barylko B, Wlodarski P, Binns DD, Gerber SH, Earnest S, Sudhof TC, et al. Analysis of the catalytic domain of phosphatidylinositol 4-kinase type II. *J Biol Chem* 2002;277:44366–75.
- [117] Lu D, Sun HQ, Wang H, Barylko B, Fukata Y, Fukata M, et al. Phosphatidylinositol 4-kinase IIalpha is palmitoylated by Golgi-localized palmitoyltransferases in cholesterol-dependent manner. *J Biol Chem* 2012;287:21856–65.
- [118] Hanada K, Kumagai K, Yasuda S, Miura Y, Kawano M, Fukasawa M, et al. Molecular machinery for non-vesicular trafficking of ceramide. *Nature* 2003;426:803–9.
- [119] Yamaji T, Kumagai K, Tomishige N, Hanada K. Two sphingolipid transfer proteins, CERT and FAPP2: their roles in sphingolipid metabolism. *IUBMB Life* 2008;60:511–8.
- [120] Hanada K, Kumagai K, Tomishige N, Yamaji T. CERT-mediated trafficking of ceramide. *Biochim Biophys Acta* 2009;1791:684–91.
- [121] Perry RJ, Ridgway ND. Molecular mechanisms and regulation of ceramide transport. *Biochim Biophys Acta* 2005;1734:220–34.
- [122] Amarilio R, Ramachandran S, Sabanay H, Lev S. Differential regulation of endoplasmic reticulum structure through VAP-Nir protein interaction. *J Biol Chem* 2005;280:5934–44.
- [123] Wyles JP, Ridgway ND. VAMP-associated protein-A regulates partitioning of oxysterol-binding protein-related protein-9 between the endoplasmic reticulum and Golgi apparatus. *Exp Cell Res* 2004;297:533–47.
- [124] Villani M, Subathra M, Im YB, Choi Y, Signorelli P, Del Poeta M, et al. Sphingomyelin synthases regulate production of diacylglycerol at the Golgi. *Biochem J* 2008;414:31–41.
- [125] Van Lint J, Ni Y, Valius M, Merlevede W, Vandenhede JR. Platelet-derived growth factor stimulates protein kinase D through the activation of phospholipase Cgamma and protein kinase C. *J Biol Chem* 1998;273:7038–43.
- [126] Baron CL, Malhotra V. Role of diacylglycerol in PKD recruitment to the TGN and protein transport to the plasma membrane. *Science* 2002;295:325–8.
- [127] Subathra M, Qureshi A, Luberto C. Sphingomyelin synthases regulate protein trafficking and secretion. *PLoS One* 2011;6:e23644.
- [128] Hausser A, Link G, Hoene M, Russo C, Selchow O, Pfizenmaier K. Phospho-specific binding of 14–3-3 proteins to phosphatidylinositol 4-kinase III beta protects from dephosphorylation and stabilizes lipid kinase activity. *J Cell Sci* 2006;119:3613–21.
- [129] Malhotra V, Campelo F. PKD regulates membrane fission to generate TGN to cell surface transport carriers. *Cold Spring Harb Perspect Biol* 2011;3:1.
- [130] Valente C, Turacchio G, Mariggio S, Pagliuso A, Gaibisso R, Di Tullio G, et al. A 14–3-3gamma dimer-based scaffold bridges CtBP1-S/BARS to PI(4)KIIbeta to regulate post-Golgi carrier formation. *Nat Cell Biol* 2012;14:343–54.
- [131] Fugmann T, Hausser A, Schoffler P, Schmid S, Pfizenmaier K, Olayioye MA. Regulation of secretory transport by protein kinase D-mediated phosphorylation of the ceramide transfer protein. *J Cell Biol* 2007;178:15–22.
- [132] Nhek S, Ngo M, Yang X, Ng MM, Field SJ, Asara JM, et al. Regulation of oxysterol-binding protein Golgi localization through protein kinase D-mediated phosphorylation. *Mol Biol Cell* 2010;21:2327–37.
- [133] D'Angelo G, Vicinanza M, De Matteis MA. Lipid-transfer proteins in biosynthetic pathways. *Curr Opin Cell Biol* 2008;20:360–70.
- [134] D'Angelo G, Polishchuk E, Di Tullio G, Santoro M, Di Campli A, Godi A, et al. Glycosphingolipid synthesis requires FAPP2 transfer of glucosylceramide. *Nature* 2007;449:62–7.
- [135] De Matteis MA, Di Campli A, D'Angelo G. Lipid-transfer proteins in membrane trafficking at the Golgi complex. *Biochim Biophys Acta* 2007;1771:761–8.
- [136] D'Angelo G, Rega LR, De Matteis MA. Connecting vesicular transport with lipid synthesis: FAPP2. *Biochim Biophys Acta* 1821;2012:1089–95.
- [137] Lenoir M, Coskun U, Grzybek M, Cao X, Buschhorn SB, James J, et al. Structural basis of wedging the Golgi membrane by FAPP pleckstrin homology domains. *EMBO Rep* 2010;11:279–84.
- [138] Halter D, Neumann S, van Dijk SM, Wolthoorn J, de Maziere AM, Vieira OV, et al. Pre- and post-Golgi translocation of glucosylceramide in glycosphingolipid synthesis. *J Cell Biol* 2007;179:101–15.
- [139] Mikitova V, Levine TP. Analysis of the key elements of FFAT-like motifs identifies new proteins that potentially bind VAP on the ER, including two AKAPs and FAPP2. *PLoS One* 2012;7:e30455.
- [140] Vieira OV, Verkade P, Manninen A, Simons K. FAPP2 is involved in the transport of apical cargo in polarized MDCK cells. *J Cell Biol* 2005;170:521–6.
- [141] Vieira OV, Gaus K, Verkade P, Fullekrug J, Vaz WL, Simons K. FAPP2, cilium formation, and compartmentalization of the apical membrane in polarized Madin-Darby canine kidney (MDCK) cells. *Proc Natl Acad Sci USA* 2006;103:18556–61.
- [142] Olsen HL, Hoy M, Zhang W, Bertorello AM, Bokvist K, Capito K, et al. Phosphatidylinositol 4-kinase serves as a metabolic sensor and regulates priming of secretory granules in pancreatic beta cells. *Proc Natl Acad Sci USA* 2003;100:5187–92.
- [143] Waugh MG. Phosphatidylinositol 4-kinases, phosphatidylinositol 4-phosphate and cancer. *Cancer Lett* 2012;325:125–31.
- [144] Li J, Lu Y, Zhang J, Kang H, Qin Z, Chen C. PI4KIIalpha is a novel regulator of tumor growth by its action on angiogenesis and HIF-1alpha regulation. *Oncogene* 2010;29:2550–9.
- [145] Ishikawa S, Egami H, Kurizaki T, Akagi J, Tamori Y, Yoshida N, et al. Identification of genes related to invasion and metastasis in pancreatic cancer by cDNA representational difference analysis. *J Exp Clin Cancer Res* 2003;22:299–306.
- [146] Giroux V, Iovanna J, Dagorn JC. Probing the human kinome for kinases involved in pancreatic cancer cell survival and gemcitabine resistance. *FASEB J* 2006;20:1982–91.
- [147] Guerreiro AS, Fattet S, Kulesza DW, Atamer A, Elsing AN, Shalaby T, et al. A sensitized RNA interference screen identifies a novel role for the PI3K p110gamma isoform in medulloblastoma cell proliferation and chemoresistance. *Mol Cancer Res* 2011;9:925–35.
- [148] Altomare DA, Testa JR. Perturbations of the AKT signaling pathway in human cancer. *Oncogene* 2005;24:7455–64.
- [149] Yuan TL, Cantley LC. PI3K pathway alterations in cancer: variations on a theme. *Oncogene* 2008;27:5497–510.
- [150] Amiri A, Noei F, Jeganathan S, Kulkarni G, Pinke DE, Lee JM. EEF1A2 activates Akt and stimulates Akt-dependent actin remodeling, invasion and migration. *Oncogene* 2007;26:3027–40.
- [151] Jeganathan S, Morrow A, Amiri A, Lee JM. Eukaryotic elongation factor 1A2 cooperates with phosphatidylinositol-4 kinase III beta to stimulate production of filopodia through increased phosphatidylinositol-4,5 bisphosphate generation. *Mol Cell Biol* 2008;28:4549–61.
- [152] Pinke DE, Lee JM. The lipid kinase PI4KIIbeta and the eEF1A2 oncogene cooperate to disrupt three-dimensional in vitro acinar morphogenesis. *Exp Cell Res* 2011;317:2503–11.
- [153] Furuta Y, Uehara T, Nomura Y. Correlation between delayed neuronal cell death and selective decrease in phosphatidylinositol 4-kinase expression in the CA1 subfield of the hippocampus after transient forebrain ischemia. *J Cereb Blood Flow Metab* 2003;23:962–71.
- [154] Saito M, Smiley J, Toth R, Vadasz C. Microarray analysis of gene expression in rat hippocampus after chronic ethanol treatment. *Neurochem Res* 2002;27:1221–9.
- [155] Jungerius BJ, Hoogendoorn ML, Bakker SC, Van't Slot R, Bardoel AF, Ophoff RA, et al. An association screen of myelin-related genes implicates the chromosome 22q11 PIK4CA gene in schizophrenia. *Mol Psychiatry* 2008;13:1060–8.
- [156] Saito T, Stopkova P, Diaz L, Papolos DF, Boussemart L, Lachman HM. Polymorphism screening of PIK4CA: possible candidate gene for chromosome 22q11-linked psychiatric disorders. *Am J Med Genet B Neuropsychiatr Genet* 2003;116:77–83.
- [157] Vorstman JA, Chow EW, Ophoff RA, van Engeland H, Beemer FA, Kahn RS, et al. Association of the PIK4CA schizophrenia-susceptibility gene in adults with the 22q11.2 deletion syndrome. *Am J Med Genet B Neuropsychiatr Genet* 2009;150B:430–3.
- [158] Kanahara N, Iyo M, Hashimoto K. Failure to confirm the association between the PIK4CA gene and schizophrenia in a Japanese population. *Am J Med Genet B Neuropsychiatr Genet* 2009;150B:450–2.
- [159] Simons JP, Al-Shawi R, Minogue S, Waugh MG, Wiedemann C, Evangelou S, et al. Loss of phosphatidylinositol 4-kinase 2alpha activity causes late onset degeneration of spinal cord axons. *Proc Natl Acad Sci USA* 2009;106:11535–9.
- [160] Vaillancourt FH, Brault M, Pilote L, Uyttersprot N, Gaillard ET, Stoltz JH, et al. Evaluation of Phosphatidylinositol-4-Kinase IIIalpha as a Hepatitis C Virus Drug Target. *J Virol* 2012;86:11595–607.
- [161] Muthusamy N, Faundez V, Bergson C. Calcyon, a mammalian specific NEEP21 family member, interacts with adaptor protein complex 3 (AP-3) and regulates targeting of AP-3 cargoes. *J Neurochem* 2012;123:60–72.



- [162] Pizarro-Cerda J, Payrastra B, Wang YJ, Veiga E, Yin HL, Cossart P. Type II phosphatidylinositol 4-kinases promote *Listeria monocytogenes* entry into target cells. *Cell Microbiol* 2007;9:2381–90.
- [163] Moorhead AM, Jung JY, Smirnov A, Kaufer S, Scidmore MA. Multiple host proteins that function in phosphatidylinositol-4-phosphate metabolism are recruited to the chlamydial inclusion. *Infect Immun* 2010;78:1990–2007.
- [164] Derre I, Swiss R, Agaisse H. The lipid transfer protein CERT interacts with the Chlamydia inclusion protein IncD and participates to ER–Chlamydia inclusion membrane contact sites. *PLoS Pathog* 2011;7:e1002092.
- [165] Elwell CA, Jiang S, Kim JH, Lee A, Wittmann T, Hanada K, et al. Chlamydia trachomatis co-opts GBF1 and CERT to acquire host sphingomyelin for distinct roles during intracellular development. *PLoS Pathog* 2011;7:e1002198.
- [166] Weber SS, Ragaz C, Reus K, Nyfeler Y, Hilbi H. *Legionella pneumophila* exploits PI(4)P to anchor secreted effector proteins to the replicative vacuole. *PLoS Pathog* 2006;2:e46.
- [167] Goody RS, Muller MP, Schoebel S, Oesterlin LK, Blumer J, Peters H, et al. The versatile *Legionella* effector protein DrrA. *Commun Integr Biol* 2011;4:72–4.
- [168] Brombacher E, Urwyler S, Ragaz C, Weber SS, Kami K, Overduin M, et al. Rab1 guanine nucleotide exchange factor SidM is a major phosphatidylinositol 4-phosphate-binding effector protein of *Legionella pneumophila*. *J Biol Chem* 2009;284:4846–56.
- [169] Schoebel S, Blankenfeldt W, Goody RS, Itzen A. High-affinity binding of phosphatidylinositol 4-phosphate by *Legionella pneumophila* DrrA. *EMBO Rep* 2010;11:598–604.
- [170] Muller MP, Peters H, Blumer J, Blankenfeldt W, Goody RS, Itzen A. The *Legionella* effector protein DrrA AMPylates the membrane traffic regulator Rab1b. *Science* 2010;329:946–9.
- [171] Urwyler S, Brombacher E, Hilbi H. Endosomal and secretory markers of the *Legionella*-containing vacuole. *Commun Integr Biol* 2009;2:107–9.
- [172] Urwyler S, Nyfeler Y, Ragaz C, Lee H, Mueller LN, Aebbersold R, et al. Proteome analysis of *Legionella* vacuoles purified by magnetic immunoseparation reveals secretory and endosomal GTPases. *Traffic* 2009;10:76–87.
- [173] Akimana C, Al-Khodor S, Abu Kwaik Y. Host factors required for modulation of phagosome biogenesis and proliferation of *Francisella tularensis* within the cytosol. *PLoS One* 2010;5:e11025.
- [174] Ahn J, Chung KS, Kim DU, Won M, Kim L, Kim KS, et al. Systematic identification of hepatocellular proteins interacting with NS5A of the hepatitis C virus. *J Biochem Mol Biol* 2004;37:741–8.
- [175] Alvisi G, Madan V, Bartenschlager R. Hepatitis C virus and host cell lipids: an intimate connection. *RNA Biol* 2011;8:258–69.
- [176] Tai AW, Benita Y, Peng LF, Kim SS, Sakamoto N, Xavier RJ, et al. A functional genomic screen identifies cellular cofactors of hepatitis C virus replication. *Cell Host Microbe* 2009;5:298–307.
- [177] Berger KL, Randall G. Potential roles for cellular cofactors in hepatitis C virus replication complex formation. *Commun Integr Biol* 2009;2:471–3.
- [178] Trotard M, Lepere-Douard C, Regeard M, Piquet-Pellorce C, Lavillette D, Cosset FL, et al. Kinases required in hepatitis C virus entry and replication highlighted by small interference RNA screening. *FASEB J* 2009;23:3780–9.
- [179] Vaillancourt FH, Pilote L, Cartier M, Lippens J, Liuzzi M, Bethell RC, et al. Identification of a lipid kinase as a host factor involved in hepatitis C virus RNA replication. *Virology* 2009;387:5–10.
- [180] Berger KL, Kelly SM, Jordan TX, Tartell MA, Randall G. Hepatitis C virus stimulates the phosphatidylinositol 4-kinase III alpha-dependent phosphatidylinositol 4-phosphate production that is essential for its replication. *J Virol* 2011;85:8870–83.
- [181] Lim YS, Hwang SB. Hepatitis C virus NS5A protein interacts with phosphatidylinositol 4-kinase type IIIalpha and regulates viral propagation. *J Biol Chem* 2011;286:11290–8.
- [182] Borawski J, Troke P, Puyang X, Gibaja V, Zhao S, Mickanin C, et al. Class III phosphatidylinositol 4-kinase alpha and beta are novel host factor regulators of hepatitis C virus replication. *J Virol* 2009;83:10058–74.
- [183] Lamarche MJ, Borawski J, Bose A, Capacci-Daniel C, Colvin R, Dennehy M, et al. Anti-hepatitis C virus activity and toxicity of type III phosphatidylinositol-4-kinase beta inhibitors. *Antimicrob Agents Chemother* 2012;56:5149–56.
- [184] Berger KL, Cooper JD, Heaton NS, Yoon R, Oakland TE, Jordan TX, et al. Roles for endocytic trafficking and phosphatidylinositol 4-kinase III alpha in hepatitis C virus replication. *Proc Natl Acad Sci USA* 2009;106:7577–82.
- [185] Arita M, Kojima H, Nagano T, Okabe T, Wakita T, Shimizu H. Phosphatidylinositol 4-kinase III beta is a target of enviroxime-like compounds for antipoliiovirus activity. *J Virol* 2011;85:2364–72.
- [186] Yang N, Ma P, Lang J, Zhang Y, Deng J, Ju X, et al. Phosphatidylinositol 4-kinase IIIbeta is required for severe acute respiratory syndrome coronavirus spike-mediated cell entry. *J Biol Chem* 2012;287:8457–67.
- [187] van der Schaar HM, van der Linden L, Lanke KH, Strating JR, Purstinger G, de Vries E, et al. Coxsackievirus mutants that can bypass host factor PI4KIIIbeta and the need for high levels of PI4P lipids for replication. *Cell Res* 2012;22:1576–92.
- [188] Sasaki J, Ishikawa K, Arita M, Taniguchi K. ACBD3-mediated recruitment of PI4KB to picornavirus RNA replication sites. *EMBO J* 2011;31:754–66.
- [189] Tai AW, Salloum S. The role of the phosphatidylinositol 4-kinase PI4KA in hepatitis C virus-induced host membrane rearrangement. *PLoS One* 2011;6:e26300.
- [190] Amako Y, Sarkeshik A, Hotta H, Yates 3rd J, Siddiqui A. Role of oxysterol binding protein in hepatitis C virus infection. *J Virol* 2009;83:9237–46.
- [191] Amako Y, Syed GH, Siddiqui A. Protein kinase D negatively regulates hepatitis C virus secretion through phosphorylation of oxysterol-binding protein and ceramide transfer protein. *J Biol Chem* 2011;286:11265–74.
- [192] Diamond DL, Syder AJ, Jacobs JM, Sorensen CM, Walters KA, Proll SC, et al. Temporal proteome and lipidome profiles reveal hepatitis C virus-associated reprogramming of hepatocellular metabolism and bioenergetics. *PLoS Pathog* 2010;6:e1000719.
- [193] Hirata Y, Ikeda K, Sudoh M, Tokunaga Y, Suzuki A, Weng L, et al. Self-enhancement of hepatitis C virus replication by promotion of specific sphingolipid biosynthesis. *PLoS Pathog* 2012;8:e1002860.
- [194] Sakamoto H, Okamoto K, Aoki M, Kato H, Katsume A, Ohta A, et al. Host sphingolipid biosynthesis as a target for hepatitis C virus therapy. *Nat Chem Biol* 2005;1:333–7.
- [195] Zhao LJ, Wang L, Ren H, Cao J, Li L, Ke JS, et al. Hepatitis C virus E2 protein promotes human hepatoma cell proliferation through the MAPK/ERK signaling pathway via cellular receptors. *Exp Cell Res* 2005;305:23–32.
- [196] Zhang Q, Gong R, Qu J, Zhou Y, Liu W, Chen M, et al. Activation of the Ras/Raf/MEK pathway facilitates hepatitis C virus replication via attenuation of the interferon-JAK-STAT pathway. *J Virol* 2012;86:1544–54.
- [197] Jiang YF, He B, Li NP, Ma J, Gong GZ, Zhang M. The oncogenic role of NS5A of hepatitis C virus is mediated by up-regulation of survivin gene expression in the hepatocellular cell through p53 and NF-kappaB pathways. *Cell Biol Int* 2011;35:1225–32.
- [198] Mankouri J, Griffin S, Harris M. The hepatitis C virus non-structural protein NS5A alters the trafficking profile of the epidermal growth factor receptor. *Traffic* 2008;9:1497–509.
- [199] Schmitz KJ, Wohlschlaeger J, Lang H, Sotiropoulos GC, Malago M, Steveling K, et al. Activation of the ERK and AKT signalling pathway predicts poor prognosis in hepatocellular carcinoma and ERK activation in cancer tissue is associated with hepatitis C virus infection. *J Hepatol* 2008;48:83–90.
- [200] Menzel N, Fischl W, Hueging K, Bankwitz D, Frentzen A, Haid S, et al. MAP-kinase regulated cytosolic phospholipase A2 activity is essential for production of infectious hepatitis C virus particles. *PLoS Pathog* 2012;8:e1002829.
- [201] Miura K, Jacques KM, Stauffer S, Kubosaki A, Zhu K, Hirsch DS, et al. ARAP1: a point of convergence for Arf and Rho signaling. *Mol Cell* 2002;9:109–19.
- [202] Campa F, Yoon HY, Ha VL, Szentpetery Z, Balla T, Randazzo PA. A PH domain in the Arf GTPase-activating protein (GAP) ARAP1 binds phosphatidylinositol 3,4,5-trisphosphate and regulates Arf GAP activity independently of recruitment to the plasma membranes. *J Biol Chem* 2009;284:28069–83.
- [203] Gavicherla B, Ritchey L, Gianfelice A, Kolokoltsov AA, Davey RA, Ireton K. Critical role for the host GTPase-activating protein ARAP2 in InlB-mediated entry of *Listeria monocytogenes*. *Infect Immun* 2010;78:4532–41.
- [204] Srivastava R, Ratheesh A, Gude RK, Rao KV, Panda D, Subrahmanyam G. Resveratrol inhibits type II phosphatidylinositol 4-kinase: a key component in pathways of phosphoinositide turn over. *Biochem Pharmacol* 2005;70:1048–55.
- [205] Naveen B, Shankar BS, Subrahmanyam G. FcepsilonRI cross-linking activates a type II phosphatidylinositol 4-kinase in RBL 2H3 cells. *Mol Immunol* 2005;42:1541–9.
- [206] Sinha RK, Patel RY, Bojjireddy N, Datta A, Subrahmanyam G. Epigallocatechin gallate (EGCG) inhibits type II phosphatidylinositol 4-kinases: a key component in pathways of phosphoinositide turnover. *Arch Biochem Biophys* 2011;516:45–51.
- [207] Daboussi L, Costaguta G, Payne GS. Phosphoinositide-mediated clathrin adaptor progression at the trans-Golgi network. *Nat Cell Biol* 2012;14:239–48.