

MEETING ABSTRACT

Open Access

Biological, pharmacological and immunological activities of novel 6-amino-acid-substituted 14-alkoxy-N-methylmorphinans

Elena Guerrieri, Valeria Follia, Dorota Garczarczyk, Silvia B Rief, Monika Fink, Muhammad F Asim, Mariana Spetea, Helmut Schmidhammer*

From 17th Scientific Symposium of the Austrian Pharmacological Society (APHAR). Joint meeting with the Hungarian Society of Experimental and Clinical Pharmacology (MFT)
Innsbruck, Austria. 29-30 September 2011

Background

Of the three opioid receptors, μ (MOP), δ (DOP) and κ (KOP), the MOP type is the most involved in the action of opioids in the gut. Experimental studies on models of intestinal inflammation and inflammatory bowel disease (IBD) support an anti-inflammatory role of peripheral MOP receptors in the gut, besides their involvement in pain control (analgesia) and gastrointestinal motility (anti-diarrheal effects). Research focuses increasingly on exploring the therapeutic potential of peripheral MOP receptors aiming for identification of peripheral ligands as improved treatment for debilitating conditions associated with bowel functions. One strategy to increase peripheral selectivity includes chemical modifications that enhance hydrophilicity [1,2]. Our work in the field of peripherally acting opioids has led to a series of opioids with zwitterionic moieties (i.e. amino acid residues) attached to the C-6 position of 14-O-methyloxymorphone, which may represent novel therapeutic molecules for IBD. These 14-alkoxymorphinans were pharmacologically and immunologically characterized.

Methods

Synthesis of novel zwitterionic 14-alkoxymorphinans was accomplished by multi-step syntheses. Binding, functional and immunomodulatory activities were determined *in vitro*. Antinociceptive activities were assessed using acetic acid-induced writhing and tail-flick tests.

Physicochemical properties ($\log P$ and $\log D$) were determined using the MarvinSketch software.

Results

In vitro, the new 6-amino-acid-substituted 14-alkoxymorphinans bound with high affinity and showed agonist activity towards the MOP receptor. They significantly inhibited the nuclear transcription factor kappaB (NF- κ B) activation in tumor-necrosis factor- α (TNF- α) and lipopolysaccharide (LPS)-stimulated human monocytic THP-1 cells. *In vivo*, they produced dose-dependent antinociceptive effects in mice after subcutaneous administration, being several-fold more potent than morphine. Based on the calculated $\log P$ and $\log D$ values, an increase of hydrophilicity, and thus peripheral selectivity can be achieved by attachment of amino acid residues to the morphinan skeleton.

Conclusions

Novel MOP agonists acting in the periphery with combined immunosuppressive and analgesic properties may provide a new approach for the treatment of IBD.

Acknowledgements

Supported by the Austrian Science Fund (FWF: P21350 and TRP 19-B18) and Tyrolean Research Fund (TWF-UNI-0404/949).

Published: 5 September 2011

References

1. Schmidhammer H, Spetea M: *Synthesis of 14-alkoxymorphinan derivatives and their pharmacological actions*. *Top Curr Chem* 2011, 299:63-91.
2. Spetea M, Windisch P, Guo Y, Bileviciute-Ljungar I, Schütz J, Asim MF, Riba P, Kiraly K, Fürst S, Al-Khasani M, Schmidhammer H: *Synthesis and*

* Correspondence: helmut.schmidhammer@uibk.ac.at

Department of Pharmaceutical Chemistry, Institute of Pharmacy and Center for Molecular Biosciences, University of Innsbruck, 6020 Innsbruck, Austria

pharmacological activities of 6-glycine substituted 14-phenylpropoxymorphinans, a novel class of opioids with high opioid receptor affinities and antinociceptive potencies. *J Med Chem* 2011, 54:980-988.

doi:10.1186/1471-2210-11-S2-A5

Cite this article as: Guerrieri et al.: Biological, pharmacological and immunological activities of novel 6-amino-acid-substituted 14-alkoxy-N-methylmorphinans. *BMC Pharmacology* 2011 11(Suppl 2):A5.

Submit your next manuscript to BioMed Central
and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

