

RESEARCH

Open Access



What matters: non-pharmaceutical interventions for COVID-19 in Europe

Yan Liu¹, Qiuyan Yu², Haoyu Wen¹, Fang Shi¹, Fang Wang³, Yudi Zhao¹, Qiumian Hong¹ and Chuanhua Yu^{1*} 

Abstract

Objectives: The purpose of this study is to describe the situation of COVID-19 in European countries and to identify important factors related to prevention and control.

Methods: We obtained data from World Health Statistics 2020 and the Institute for Health Metrics and Evaluation (IHME). We calculated the R_t values of 51 countries in Europe under different prevention and control measures. We used lasso regression to screen factors associated with morbidity and mortality. For the selected variables, we used quantile regression to analyse the relevant influencing factors in countries with different levels of morbidity or mortality.

Results: The government has a great influence on the change in R_t value through prevention and control measures. The most important factors for personal and group prevention and control are the mobility index, testing, the closure of educational facilities, restrictions on large-scale gatherings, and commercial restrictions. The number of ICU beds and doctors in medical resources are also key factors. Basic sanitation facilities, such as the proportion of safe drinking water, also have an impact on the COVID-19 epidemic.

Conclusions: We described the current status of COVID-19 in European countries. Our findings demonstrated key factors in individual and group prevention measures.

Keywords: Non-pharmaceutical interventions, COVID-19, Europe, Lasso regression, Quantile regression

Background

Since 7 December 2020, there have been 66,243,918 cumulative cases and 1,528,984 cumulative deaths worldwide according to data from the World Health Organization (WHO) on coronavirus disease (COVID-19). The WHO declared COVID-19 to be a pandemic on 12 March 2020; a pandemic is a global public health emergency of high international concern. COVID-19 is an acute respiratory infectious disease transmitted mainly through aerosols and droplets [1], and is characterised by higher infectivity but lower mortality than SARS, so we should strengthen its prevention and control [2].

The chief non-pharmacological prevention and control measures included individual and group prevention. Personal prevention primarily included wearing masks, washing one's hands, proper social distancing, and reducing participation in outdoor activities. Group prevention mostly encompassed travel restrictions, home isolation, the closure of educational facilities, prohibitions on public gatherings, all commercial activities, and non-essential commercial activities. The effect of prevention and control may also be related to medical resources (e.g. doctors, nurses, pharmacists, ventilators, ordinary beds, ICU beds), as well as the country's economic situation and ability to respond to public health incidents (e.g. gross domestic product [GDP] and global health security [GHS]).

Switzerland regards testing as the central method to control COVID-19 [3]. A meta-analysis showed that

*Correspondence: yuchua@whu.edu.cn

¹ Department of Epidemiology and Biostatistics, School of Public Health,

Wuhan University, Hubei, China

Full list of author information is available at the end of the article



© The Author(s) 2021. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

medical masks and N95 masks can protect against viral respiratory infection [4]. Community-wide mask wearing can control COVID-19 by preventing contact with large amounts of saliva and respiratory droplets [5]. Hand hygiene is the most important factor in control activities and infection prevention [6]. The use of antiseptic hand soaps leads to a greater reduction in the number of microorganisms compared with regular soaps [7]. However, the toxicity and stability of surface disinfectants are issues that should be further investigated [8]. The findings of a systematic review and meta-analysis support the idea that physical distancing of 1 m, wearing face masks, and eye protection prevent the person-to-person transmission of COVID-19 [9].

The first country in Europe where the COVID-19 outbreak occurred was Italy. On 9 March 2020, the Italian government implemented a comprehensive 'blockade' policy, mostly consisting of travel restrictions, mandatory stay-at-home orders (except for health problems, emergencies, or regulated shopping for necessities), and the temporary closure of non-essential businesses and shops, which lasted until 3 May 2020. In the following months, many European states adopted similar measures [10]. In March 2020, 47 countries did not allow commercial flights to land [11]. Moreover, it became necessary to develop an international framework to outline the method, time, and scale of travel restrictions according to the stage of the epidemic [12]. A cross-sectional study suggests that issuing a stay-at-home ban may help limit the spread of COVID-19 cases [13]. On 12 March, Norway issued stricter measures and instituted quarantines for those who entered the country. That same day, the government closed all schools and kindergartens; training centres and offices for psychologists and physiotherapists; hair salons; and swimming pools, and forbade cultural and sporting events. Sweden chose a different strategy: Kindergartens, elementary schools, other businesses, and training facilities remained open, and children's sports continued [14]. Many sporting events were restricted or cancelled to limit the spread of disease [15]. Some non-essential businesses, such as vape shops, were closed [16].

The disease incidence and mortality of COVID-19 were related to country healthcare resources and economic status [17]. In Italy, 16% of hospitalised COVID-19 patients required intensive care. Compared with Germany, the medical resources in northern Italy became overwhelmed by the increase in patients. In contrast, Germany had a wider distribution of cases and was able to make better use of its resources. The high mortality rate in Italy may reflect the relationship between the availability of medical resources and outcomes [18]. In

addition, the prevention and control effects of COVID-19 may be tied to indicators such as GDP and GHS [19].

Materials and methods

Research objective and data sources

We obtained data from World Health Statistics 2020 and the Institute for Health Metrics and Evaluation (IHME). The IHME is an independent population health research centre at the University of Washington Medicine. We selected 51 countries in Europe as our research objective. There are 53 countries in Europe according to WHO regional groupings. The data from Monaco and Turkmenistan were incomplete, so we discarded those two countries.

Where does the IHME obtain its data? These forecasts include data from local and national governments, hospital networks and associations, the WHO, third-party aggregators, and a range of other sources.

For testing data, the IHME relies primarily on data reported by Our World in Data. However, Cyprus, Italy, and Spain used government data. The IHME obtains hospital resource data from sources such as government websites, hospital associations, the Organisation for Economic Co-operation and Development (OECD), the WHO, and published studies. For population density, they used gridded population count estimates for 2020 at the 1×1 km (km) level from World-Pop. For mobility index data, the IHME used anonymised, aggregated data from Google, Facebook, and Apple. Their data on mask use come from the Premise, Facebook Global Symptom Survey (research based on survey results from the University of Maryland Social Data Science Center), the Kaiser Family Foundation (KFF), and the YouGov COVID-19 Behaviour Tracker survey.

Statistical analysis method

Changes in R_t under different prevention and control measures

For this study, we used the time-dependent basic reproduction number method. The time-dependent reproduction number (R_t) is the average number of secondary cases of a single infected person during t day of infection. R_t is usually applied to describe the transmission characteristics of pathogens during an epidemic, and can also evaluate the effectiveness of interventions. When R_t is greater than 1, it indicates that the number of infections is rising rapidly; when R_t is less than 1 and close to 0, it suggests that the epidemic has been effectively controlled [20].

Lasso regression

Traditional linear regression belongs to the subset selection method, and the chief approach for screening

variables is the ordinary least squares (OLS) technique. However, OLS has some defects: Variable selection is separated from model parameter estimation, so the model error increases; small changes in variables have a great influence on variable selection. Variable selection is not suitable for high-dimensional data [21, 22]. The least absolute shrinkage and selection operator regression (lasso regression) is the representative regularisation method that can effectively optimise the OLS estimation and treatment overfitting problem [23]. By introducing a penalty term into model estimation, lasso regression can obtain higher prediction accuracy and model generalisation ability. It can also effectively address overfitting and multicollinearity problems [24–26]. The specific formula can be expressed as:

$$\hat{\beta}(\text{lasso}) = \arg \min_{\beta} \|y - \sum_{j=1}^p X_j \beta_j\|^2 + \lambda \sum_{j=1}^p |\beta_j|$$

The first part represents the standard OLS loss function, while the second part denotes the penalty function. It represents the tuning parameter for controlling the degree of regression coefficient compression. When λ is ≥ 0 , the greater the value, the stronger the penalty. When $\lambda=0$, the loss function does not penalise the model. Lasso regression shrinks some coefficients and sets others to 0, and tries to retain the good features of both subset selection and ridge regression [27]. Researchers adopt different forms of penalty functions according to the different characteristics of independent variables in regression analysis on the basis of lasso regression. Many kinds of regularisation models have been established and developed, such as relaxed lasso [28], adaptive lasso [26], Bayesian lasso [29], fused lasso [24], group lasso [30], and elastic net [25].

Quantile regression

Quantile regression (QR) estimates the linear relationship between the different quantiles of the dependent variable and the independent variable. Both QR and ordinary least squares (OLS) can solve specific minimisation problems. The estimation of OLS regression is grounded in the smallest residual square, and the estimation of QR is rooted in the smallest weighted absolute value residual. The minimum weighted absolute deviation of quantile regression is as follows:

$$\min \{w_t |y_t - \alpha|\} = - \sum_{i:y_i < \alpha} (1 - \tau)(y_t - \alpha) + \sum_{i:y_i \geq \alpha} \tau(y_t - \alpha)$$

The purpose of quantile regression is to calculate the regression coefficients of different quantile values of the

dependent variable. It can comprehensively display all data information to a certain extent. Thus, it has unique advantages over traditional linear regression models. Especially for the condition of a non-normal distribution, quantile regression is more comprehensive and accurate than traditional linear regression coefficient estimation [31]. We used R software 4.0.2 and Stata 15.0 for all of our statistical analyses.

Results


Changes in Rt under different prevention and control measures

We plotted the changes in Rt values of 51 countries in Europe. After an outbreak period ranging from 2–3 months, the Rt of France, Denmark, Belgium, Armenia, Germany, the Netherlands, Spain, Portugal, Moldova, Sweden, Tajikistan, Turkey, and Uzbekistan were basically stable, and the value fluctuated at approximately 1. Russia and Ukraine had a longer outbreak period, but the Rt afterwards also stabilised at approximately 1. The Rt of other countries fluctuated to varying degrees. The general trend of changes in Rt in most European countries showed an increasing trend after October and exceeded 1.


Let us take Finland and Switzerland as examples. In Fig. 1a, we describe the changes in the Rt value in Finland under different prevention and control measures. The first case of COVID-19 occurred on 28 January in Finland, and on 18 March, any business activities were banned and educational facilities were closed. Travel restrictions were implemented on 25 March, and unnecessary business activities were prohibited on 4 April. Prevention and control effects were achieved, and the Rt value dropped below 1 under the above four powerful group interventions. On 29 May, the government ended travel restrictions. On 1 June, the government ended the ban on restricting any business activities and non-essential commercial activities. Due to the relaxation of the intervention policy, the Rt value rebounded and rose. On 24 July, the Rt value reached a small peak of 3.296. On 13 August, when the Rt value dropped to 1.398, the restriction on school facility closure ended. Then, on 11 October, the Rt value fluctuated repeatedly, and on 11 October, it reached 1.966.

In Fig. 1b, we describe the changes in Rt values in Switzerland under different prevention and control measures.

Cases first appeared in Switzerland on 23 February, and large gatherings were banned on 28 February. On 13



A Changes of Rt in Finland under different prevention and control measures



B Changes of Rt in Switzerland under different prevention and control measures

Fig. 1 a Changes of Rt in Finland under different prevention and control measures b Changes of Rt in Switzerland under different prevention and control measures

March, education facilities were closed. On 16 March, any commercial activities and non-essential commercial activities were prohibited. Under the above group intervention, a certain prevention and control effect was achieved, and the Rt value dropped below 1. Thus, on 27 April, the government ended the restrictions on

non-essential commercial activities. On 6 June, the government ended the ban on any commercial activities. On 8 June, the government decided to end the closure of school education facilities. Due to the relaxation of the intervention policy, the Rt value rebounded and rose. On 2 July, the Rt value exhibited a small peak of 2.88.

Subsequently, the R_t value dropped to approximately 1. On 16 October, there was a small peak of 2.51. The graphs of R_t changes in other countries can be seen in the supplemental materials.

Results of lasso regression

In this study, we explored the relationship between two dependent variables and 17 independent variables, including cumulative morbidity (Y1) and cumulative mortality (Y2). The independent variables were the density of medical doctors/per 10,000 people (X1), the density of medical nursing and midwifery personnel/per 10,000 people (X2), the density of medical pharmacists/per 10,000 people (X3), GDP per capita (US dollars per capita) (X4), the proportion of the population using safely managed drinking-water services (X5), total tests (X6), the mask use rate (X7), the mobility composite (X8), excess bed capacity for COVID-19 (X9), the ICU excess bed capacity for COVID-19 (X10), travel restrictions (X11), stay-at-home orders (X12), the closure of educational facilities (X13), restrictions on gathering (X14), business closures (X15), non-essential businesses being ordered to close (X16), and the GHS index (X17).

We finally screened five out of 17 variables related to cumulative morbidity (Y1) and seven variables related to cumulative mortality (Y2) after lasso regression selection. The specific content is shown in Fig. 2. We can see that when the average mean-squared error was the smallest, lasso regression screened out five variables in Fig. 2a; at this time, $\lambda=0.09559$. Among them, the selected variables were X1, X5, X8, X14, and X16. When the average mean-squared error was the smallest, lasso regression screened out seven variables in Fig. 2b; at this time, $\lambda=0.0822$. Among them, the selected variables were X5, X6, X8, X10, X13, X14, and X16. The selected indicators were closely related to cumulative morbidity and mortality.

Results of quantile regression

The results of quantile regression were different from the results of lasso regression, and quantile regression provided more comprehensive information. The specific content is presented in Table 1. The overall finding is that the mobility index, the ratio of safe drinking water, and the closure of non-essential businesses are related to the cumulative incidence. The closure of educational facilities, restrictions on gathering, and the closure of non-essential businesses are tied to cumulative mortality. In low- and medium-incidence countries, the mobility composite is connected with the cumulative incidence. In high-incidence countries, the closure of educational

facilities and restrictions on gathering are linked to cumulative mortality.


Discussion

We found that the population prevention and control measures implemented by the government had an impact on the change in the R_t value. In most countries, the R_t value had a clear upwards trend in October. The most important factor in personal prevention and control is the mobility composite. Group prevention of the total testing, the closure of educational facilities, restrictions on large-scale gatherings, and commercial restrictions are very important for prevention and control. Moreover, the number of ICU beds and the average number of doctors in medical resources are key elements. Basic sanitation, such as the proportion of safe drinking water, has also had a certain impact on the COVID-19 epidemic.


The rate of masks used in individual prevention does not seem to be related to cumulative mortality or morbidity, which does not mean that the use of masks has no effect on prevention and control. Research on the effectiveness of masks for prevention and control has been confirmed [4]. The lasso regression identifies variables with a very large degree of correlation, so the mobility index generated by personal behaviour in prevention and control may be more important than wearing a mask. Second, one possible reason could be that the data on the rate of mask use come from social surveys, and there may be large errors.

Some studies have reached conclusions consistent with ours. The outcomes of the dynamic SEIR model show that the lockdown control measures implemented by China on 23 January 2020 were essential to ultimately reducing the scale of the COVID-19 epidemic [32]. Another study found that the early detection and isolation of cases prevented more infections than restrictions on travel and reduced contact. Our research also reveals a certain relationship between cumulative mortality and testing [33]. Some scholars have even proposed that the best strategy is to use both robot recognition and migration restriction strategies. European countries can also take this approach to reduce exposure to infection and provide help for the prevention and control of diseases [34]. Once the initial pandemic is under control, we must turn our attention to how to improve the adverse effects of the lockdown [35].

Studies have also indicated that medical resources are related to the mortality rate of COVID-19, which proves our research conclusions. Our study demonstrates that the number of doctors per capita and the number of hospital beds per capita are linked to the incidence or death of COVID-19 [36]. The experience in Wuhan implies that if medical resources become scarce, the government



A The process of selecting variables in Lasso regression



B The process of selecting variables in Lasso regression

Fig. 2 **a** The process of selecting variables in Lasso regression and **b** the process of selection variables in Lasso regression

should establish temporary hospitals, and medical staff will be deployed from areas where the epidemic is relatively mild to ease the pressure in severely affected zones. Effective quarantine via quick detection prevents a larger

outbreak [37, 38]. It is necessary to establish medically necessary, time-sensitive procedure scoring systems during the COVID-19 pandemic [39]. Within days or weeks, the health system is reorganised. We must optimise

Table 1 Results of the quantile regression

	Quantiles								
	0.1	0.2	0.3	0.4	0.5	0.6	0.7	0.8	0.9
Y1									
(Intercept)	-3.88	-1.35	-14.06	-8.92	-11.81	-27.89	-25.56	-27.45	-3.62
X1	0.12	0.05	0.11	0.01	0.18	0.10	0.01	-0.01	0.46
X5	0.04	0.07*	0.16*	0.17*	0.10*	0.31*	0.32*	0.35	0.31
X8	-0.21	-0.09	-0.44	-0.42*	-0.52*	-0.52	-0.69	-0.68	-0.51
X14	-2.73	-3.36	-4.67	-5.39	-4.26	2.12	-0.39	-0.20	-28.64
X16	4.32	6.44*	5.12*	5.15*	8.30*	6.51*	8.81*	12.12	17.33
Y2									
(Intercept)	-8.54	-14.20	0.79	-8.84	35.08	15.56	-22.43	-17.87	-111.83
X5	0.08	0.01	-0.04	0.09	0.19	0.23	0.54	0.74	1.44
X6	0.00	0.00	0.01	0.01	0.00	0.00	0.00	-0.01	0.00
X8	-0.30	-0.69	-1.09*	-0.86	-0.82*	-1.28*	-1.74*	-1.47*	-2.60*
X10	7.46	5.64	1.78	0.15	4.52	8.27	-4.70	-15.30	-33.26
X13	3.83	-7.50	-57.16	-51.27	-48.17	-44.13	-31.18	-20.35*	8.17
X14	-6.59	7.77	39.84	41.69	-10.86	-6.85	1.25	-14.87*	5.63
X16	3.87	10.08*	13.73*	12.66*	16.70*	21.59*	19.06*	25.98	1.94

*The coefficient is statistically significant

health resources. The fight against the disease occurs via a joint medical team composed of doctors, nurses, pharmacists, and respiratory therapists [40].

The ratio of safe drinking water in basic health resources has a relationship with the cumulative morbidity and mortality of COVID-19, which may be related to the fact that hand washing can reduce the number of hand viruses and achieve a certain prevention and control effect. Basic cleaning services are a prerequisite for infection prevention and control [41].

We did not include certain indicators—such as nurses per capita, pharmacists per capita, GDP, and GHS—in the regression model, which may be due to the relatively low degree of correlation. After the Ebola outbreak in 2014, the GHS index was developed to measure countries' ability to respond to infectious disease outbreaks. Six core elements were evaluated: prevention, detection and reporting, response, the health system, compliance with norms, and the risk of infectious disease outbreaks. The higher the GHS score, the better the preparation.

The GHS index has low predictive value for the death outcomes of COVID-19, and we have reached the same conclusion [42, 43]. For example, the UK, which ranks second in terms of GHS index score, also bears a huge burden of disease [44]. However, the GHS index has a predictive effect on the burden of COVID-19, but in the opposite direction [45].

The outcomes of quantile regression indicated that low- and medium-incidence countries should pay more

attention to personal prevention (mobility composite), and that high-incidence countries should close educational facilities and impose restrictions on gatherings.

Our research has some advantages. First, we described the prevalence and control of COVID-19 in European countries. Second, we included many independent variables to analyse their relationship with dependent variables. The independent variables mostly include individual and group prevention, medical resources, basic health facilities, and comprehensive indicators. Third, we used lasso regression to screen variables with a smaller error than traditional regression, and the results are more accurate. We further performed quantile regression to quantify the specific situation of each divided point, thus providing more information than traditional regression. Of course, our article also has some shortcomings. For example, A small number of proven effective prevention and control measures did not enter into our regression model, which may result from the accuracy of the data and the impact of the variables.

Conclusions

We comprehensively described the status of COVID-19 prevention and control in European countries. We attempted to identify key factors in individual and group prevention measures, which can provide a policy basis for the prevention and control of epidemics in European countries.

Acknowledgements

Thanks to IHME and WHO for providing open data resources.

Authors' contributions

YL acquired the data, performed the analysis of data, and wrote the manuscript. QY and HW acquired the data and contributed to the analysis of data. FS, FW, YZ and QH contributed to the coding of the statistical analysis. CY designed and evaluated the whole work. All authors read and approved the final manuscript.

Funding

This research was funded by the National Natural Science Foundation of China (Grant No. 81773552), the National Key Research and Development Program of China (Grant No. 2018YFC1315302) and National College Student Innovation Training Project (Grant No. 202110486088).

Availability of data and materials

The datasets used and/or analysed during the current study are available from the corresponding author on reasonable request.

Declarations**Ethics approval and consent to participate**

Not applicable.

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Author details

¹Department of Epidemiology and Biostatistics, School of Public Health, Wuhan University, Hubei, China. ²Department of Epidemiology and Medicine Statistics, Public Health and Management School, Wenzhou Medical University, Zhejiang, China. ³School of Public Health, Xuzhou Medical University, Jiangsu, China.

Received: 6 March 2021 Accepted: 6 December 2021

Published online: 09 January 2022

References

- Jayaaweera M, Perera H, Gunawardana B, Manatunge J. Transmission of COVID-19 virus by droplets and aerosols: A critical review on the unresolved dichotomy. *Environ Res.* 2020;188:1098–119.
- Peeri NC, Shrestha N, Rahman MS, et al. The SARS, MERS and novel coronavirus (COVID-19) epidemics, the newest and biggest global health threats: what lessons have we learned? *Int J Epidemiol.* 2020;49(3):717–26.
- Salathé M, Althaus CL, Neher R, et al. COVID-19 epidemic in Switzerland: on the importance of testing, contact tracing and isolation. *Swiss Med Wkly.* 2020;150:202–5.
- Bartoszko JJ, Farooqi MAM, Alhazzani W, Loeb M. Medical masks vs N95 respirators for preventing COVID-19 in healthcare workers: a systematic review and meta-analysis of randomized trials. *Influenza Other Respir Viruses.* 2020;14(4):365–73.
- Cheng VC, Wong SC, Chuang VW, et al. The role of community-wide wearing of face mask for control of coronavirus disease 2019 (COVID-19) epidemic due to SARS-CoV-2. *J Infect.* 2020;81(1):107–14.
- Boyce JM, Pittet D. Guideline for hand hygiene in health-care settings. *MMWR Recommend Rep.* 2002;51(16):1–45.
- Embil JM, Dyck B, Plourde P. Prevention and control of infections in the home. *CMAJ Can Med Assoc J.* 2009;180(11):E82–86.
- Pradhan D, Biswasroy P, Kumar Naik P, Ghosh G, Rath G. A review of current interventions for COVID-19 prevention. *Arch Med Res.* 2020;51(5):363–74.
- Chu DK, Akl EA, Duda S, Solo K, Yaacoub S, Schünemann HJ. Physical distancing, face masks, and eye protection to prevent person-to-person transmission of SARS-CoV-2 and COVID-19: a systematic review and meta-analysis. *Lancet (London, England).* 2020;395(10242):1973–87.
- Lavezzo E, Franchin E, Ciavarella C, et al. Suppression of a SARS-CoV-2 outbreak in the Italian municipality of Vo'. *Nature.* 2020;584(7821):425–9.
- Devi S. Travel restrictions hampering COVID-19 response. *Lancet (London, England).* 2020;395(10233):1331–2.
- Xue W, Lam C, Yeung HH, Wong CS, Chan VLY, Wong YS. Travel restrictions in the rising COVID-19 pandemic. *Hong Kong Med J Xianggang yi xue za zhi.* 2020;26(3):255–7.
- Lyu W, Wehby GL. Comparison of estimated rates of coronavirus disease 2019 (COVID-19) in border counties in Iowa without a stay-at-home order and border counties in Illinois with a stay-at-home order. *JAMA Netw Open.* 2020;3(5):e2011102.
- Helsing LM, Refsum E, Gjøstein DK, et al. The COVID-19 pandemic in Norway and Sweden—threats, trust, and impact on daily life: a comparative survey. *BMC Public Health.* 2020;20(1):1597.
- McCloskey B, Zumla A, Ippolito G, et al. Mass gathering events and reducing further global spread of COVID-19: a political and public health dilemma. *Lancet (London, England).* 2020;395(10230):1096–9.
- Berg CJ, Callanan R, Johnson TO, et al. Vape shop and consumer activity during COVID-19 non-essential business closures in the USA. *Tobacco Control.* 2021;30(e1):e41–e44.
- Lai CC, Wang CY, Wang YH, Hsueh SC, Ko WC, Hsueh PR. Global epidemiology of coronavirus disease 2019 (COVID-19): disease incidence, daily cumulative index, mortality, and their association with country healthcare resources and economic status. *Int J Antimicrob Agents.* 2020;55(4):105946.
- Karagiannidis C, Mostert C, Hentschker C, et al. Case characteristics, resource use, and outcomes of 10 021 patients with COVID-19 admitted to 920 German hospitals: an observational study. *Lancet Respir Med.* 2020;8(9):853–62.
- Ravi SJ, Warmbrod KL, Mullen L, et al. The value proposition of the Global Health Security Index. *BMJ Glob Health.* 2020;5(10):e003648.
- Cori A, Ferguson NM, Fraser C, Cauchemez S. A new framework and software to estimate time-varying reproduction numbers during epidemics. *Am J Epidemiol.* 2013;178(9):1505–12.
- Breiman L. Better subset regression using the nonnegative garrote. *Technometrics.* 1995;37(4):373–84.
- Tibshirani RJ. Regression shrinkage and selection via the lasso: a retrospective. *R Stat Soc.* 2011;73(3):267–88.
- Candes E, Tao TJ. The Dantzig selector: statistical estimation when p is much larger than n. *Ann Stat.* 2005;35(6):2313–51.
- Tibshirani R, Saunders M, Rosset S, Zhu J, Knight K. Sparsity and smoothness via the fused lasso. *J R Stat.* 2005;67(1):91–108.
- Zou Hui T, et al. Addendum: regularization and variable selection via the elastic net. *J R Stat Soc.* 2005;67(5):768.
- Hui Z. The adaptive Lasso and its oracle properties. *Publ Am Stat Assoc.* 2006;101(476):1418–29.
- Hesterberg T, Choi NH, Meier L, Fraley C. Least angle and l1 penalized regression: a review. *Stat Surv.* 2008;2:61–93.
- Meinshausen N. Relaxed lasso. *Comput Stat Data Anal.* 2007;52(1):374–93.
- Park TC, George R. The Bayesian lasso. *Publ Am Stat Assoc.* 2008;103(482):681–6.
- Lin MYY. Model selection and estimation in regression with grouped variables. *J R Stat Soc.* 2006;68(1):49–67.
- Hallock KF, Koenker RW. Quantile regression. *J Econ Perspect.* 2001;15(4):143–56.
- Yang Z, Zeng Z, Wang K, et al. Modified SEIR and AI prediction of the epidemics trend of COVID-19 in China under public health interventions. *J Thorac Dis.* 2020;12(3):165–74.
- Lai S, Ruktanonchai NW, Zhou L, et al. Effect of non-pharmaceutical interventions to contain COVID-19 in China. *Nature.* 2020;585(7825):410–3.
- Baba IA, Baba BA, Esmaili P. A mathematical model to study the effectiveness of some of the strategies adopted in curtailing the spread of COVID-19. *Comput Math Methods Med.* 2020;2020:5248569.
- McEachan RRC, Dickerson J, Bridges S, et al. The Born in Bradford COVID-19 Research Study: protocol for an adaptive mixed methods research study to gather actionable intelligence on the impact of COVID-19 on health inequalities amongst families living in Bradford. *Wellcome Open Res.* 2020;5:191.

36. Ji Y, Ma Z, Peppelenbosch MP, Pan Q. Potential association between COVID-19 mortality and health-care resource availability. *Lancet Global Health*. 2020;8(4):e480.
37. Wang LP, Wang J, Zhao HY, et al. Modelling and assessing the effects of medical resources on transmission of novel coronavirus (COVID-19) in Wuhan, China. *Math Biosci Eng MBE*. 2020;17(4):2936–49.
38. Yuan Y, Qiu T, Wang T, et al. The application of Temporary Ark Hospitals in controlling COVID-19 spread: The experiences of one Temporary Ark Hospital, Wuhan, China. *J Med Virol*. 2020;92(10):2019–26.
39. Prachand VN, Milner R, Angelos P, et al. Medically necessary, time-sensitive procedures: scoring system to ethically and efficiently manage resource scarcity and provider risk during the COVID-19 pandemic. *J Am Coll Surg*. 2020;231(2):281–8.
40. Lee CCM, Thampi S, Lewin B, et al. Battling COVID-19: critical care and peri-operative healthcare resource management strategies in a tertiary academic medical centre in Singapore. *Anaesthesia*. 2020;75(7):861–71.
41. McGriff JA, Denny L. What COVID-19 Reveals about the neglect of WASH within infection prevention in low-resource healthcare facilities. *Am J Trop Med Hyg*. 2020;103(5):1762–4.
42. Haider N, Yavlinsky A, Chang YM, et al. The Global Health Security index and Joint External Evaluation score for health preparedness are not correlated with countries' COVID-19 detection response time and mortality outcome. *Epidemiol Infect*. 2020;148:e210.
43. Abbey EJ, Khalifa BAA, Oduwole MO, et al. The Global Health Security Index is not predictive of coronavirus pandemic responses among Organization for Economic Cooperation and Development countries. *PLoS ONE*. 2020;15(10):e0239398.
44. Aitken T, Chin KL, Liew D, Ofori-Asenso R. Rethinking pandemic preparation: Global Health Security Index (GHSI) is predictive of COVID-19 burden, but in the opposite direction. *J Infect*. 2020;81(2):318–56.
45. Boyd MJ, Wilson N, Nelson C. Validation analysis of Global Health Security Index (GHSI) scores 2019. *BMJ Glob Health*. 2020;5(10):e003276.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

