

feedback forces HL's output to be correlated with the sensed tilt angles. Thus, activity modulated Hebbian learning re-sharpens VA's efferent tuning and the modeled organism relearns to balance in an upright position – even without biofeedback. Phenomenologically this effect is also observed in human patients.

Acknowledgements

The authors would like to thank ETH Research Grant ETH-23 08-1.

Published: 20 July 2010

References

1. Tyler M, Danilov YP, Bach-Y-Rita P: **Closing an open-loop control system: vestibular substitution through the tongue.** *J Integr Neurosci* 2003, **2**(2):159-164.
2. Danilov YP, Tyler ME, Skinner KL, Hogle RA, Bach-y-Rita P: **Efficacy of electrotactile vestibular substitution in patients with peripheral and central vestibular loss.** *J Vestib Res* 2007, **17**(23):119-130.

doi:10.1186/1471-2202-11-S1-P111

Cite this article as: Jug *et al.*: A model of functional recovery after significant loss of neural tissue: biofeedback based healing of vestibular dysfunction. *BMC Neuroscience* 2010 **11**(Suppl 1):P111.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

