

Functional analysis of a dihydroflavonol 4-reductase gene in *Ophiorrhiza japonica* (OjDFR1) reveals its role in the regulation of anthocyanin

Wei Sun^{1,*}, Nana Zhou^{1,*}, Cai Feng¹, Shiyu Sun¹, Ming Tang¹, Xiaoxin Tang¹, Zhigang Ju² and Yin Yi^{1,3}

¹ Key Laboratory of State Forestry Administration on Biodiversity Conservation in Karst Mountain Area of Southwest of China, School of Life Science, Guizhou Normal University, Guiyang, China

² Pharmacy College, Guizhou University of Traditional Chinese Medicine, Guiyang, China

³ Key Laboratory of Plant Physiology and Development Regulation, School of Life Science, Guizhou Normal University, Guiyang, China

* These authors contributed equally to this work.

ABSTRACT

Dihydroflavonol 4-reductase (DFR), a key regulatory enzyme, participated in the biosynthesis of anthocyanins, proanthocyanidins and other flavonoids that essential for plant survival and human health. However, the role of this enzyme in *Ophiorrhiza japonica* is still unknown. Here, three putative *DFR-like* genes were firstly isolated from *O. japonica*. Phylogenetic analysis indicated that *OjDFR1* was classified into DFR subgroup, while the rest two were clustered into other NADPH-dependent reductases. Then, functions of the three genes were further characterized. Expression analysis showed that *OjDFR1* transcripts had strong correlations with the accumulation pattern of anthocyanin during the flower developmental, whereas other two were not, this suggested the potential roles of *OjDFR1* in anthocyanin biosynthesis. Subsequently, all three clones were functionally expressed in *Escherichia coli*, but confirming that only *OjDFR1* encode active DFR proteins that catalyzed the reduction of dihydroflavonols to leucoanthocyanidin. Consistent with the biochemical assay results, overexpressing *OjDFR1* in Arabidopsis *tt3-1* mutant successfully restored the deficiency of anthocyanin and proanthocyanidin, hinting its function as DFR in planta. Additionally, heterologous expression of *OjDFR1* in transgenic tobacco contributed to darker flower color *via* up-regulating the expressions of endogenous *NtANS* and *NtUGT*, which suggested that *OjDFR1* was involved in flower color development. In summary, this study validates the functions of *OjDFR1* and expands our understanding of anthocyanin biosynthesis in *O. japonica*.

Subjects Biochemistry, Molecular Biology, Plant Science

Keywords Anthocyanin biosynthesis, Dihydroflavonol 4-reductase, Biochemical characterization, Flower color, *Ophiorrhiza japonica*

INTRODUCTION

Anthocyanins, a kind of flavonoids, were first used to denote the blue pigments of flowers in 1835 (*Shibata, Shibata & Kasiwagi, 1919*). They are water-soluble and occur widely in

Submitted 29 July 2021

Accepted 26 September 2021

Published 20 October 2021

Corresponding authors

Zhigang Ju, juzhigangz@163.com

Yin Yi, yiyin@gznu.edu.cn

Academic editor

Pedro Silva

Additional Information and
Declarations can be found on
page 14

DOI 10.7717/peerj.12323

© Copyright

2021 Sun et al.

Distributed under

Creative Commons CC-BY 4.0

OPEN ACCESS




Figure 1 Diagrammatic representation of the lignins, flavonols, proanthocyanidins and anthocyanins biosynthetic pathways in plants. CHS, chalcone synthase; CCR, cinnamoyl CoA reductase; CHI, chalcone isomerase; F3H, flavanone 3-hydroxylase; F3'H, flavonoid 3'-hydroxylase; F3'5'H, flavonoid 3',5'-hydroxylase; FLS, flavonol synthase; DFR, dihydroflavonol 4-reductase; LAR, leucoanthocyanidin reductase; ANS, anthocyanidin synthase; ANR, anthocyanidin reductase; UFGT, anthocyanidin 3-O-glucosyltransferase; AT, anthocyanin aromatic acyltransferase.

Full-size DOI: 10.7717/peerj.12323/fig-1

the plant kingdom. Anthocyanins play a significant role during plant diverse development processes. Firstly, they are vital coloring pigments that impart wide range colors (pale yellow, orange, pink, red, purple, and blue) to different organs of plants. Simultaneously, these bright colors let plants more outstanding, thus attracting pollinators and seed dispersers for the purpose of breeding (Sara, Pierdomenico & Silvia, 2020; Tanaka, Sasaki & Ohmiya, 2008; Zhao & Tao, 2015). In addition, anthocyanins are also produced for protecting plants from various biotic and abiotic stresses such as fungal infection, insect attacks, high/low temperatures, sunlight exposure, drought, salty soils, as well as heavy metals (Ferreyra, Rius & Casati, 2012). During the last decade, great many researches have proved that anthocyanins are beneficial not only for plants, but also for human health because of their antioxidant, anti-viral, anti-microbial, anti-inflammatory and anticancer actions (Sonia, Diego & Cristina, 2010; Dharmawansa, David & Vasantha, 2020).

Anthocyanin biosynthesis involves many enzymes, among them, dihydroflavonol 4-reductase (DFR) is a rate-limited enzyme that controls the carbon flux direction of anthocyanin pathway, thereby leading to strikingly different anthocyanin profiles (Xie et al., 2004; Miyagawa et al., 2015). DFR catalyzes the conversion of three colorless dihydroflavonols (DHM, DHK, DHQ) to the corresponding leucoanthocyanidin and is recognized as a pivotal regulatory point during the biosynthesis of anthocyanins (Fig. 1) (Ni et al., 2020). In general, DFR can be classified into three types because of the difference at amino acid residue 134 (Johnson et al., 2001; Xie et al., 2018). First, there are the asparagine-type (Asn-type) DFRs, which have an Asn at position 134 and are diffusely

distributed in plants. Second, there are the aspartic acid-type (Asp-type) DFRs, which contain an Asp at this position and cannot reduce DHK efficiently. And the third type is called non-Asn/Asp-type DFRs, which possess neither Asn nor Asp. Meanwhile, previous studies have also reported that DFR is commonly found as a single gene in several plants, such as *Arabidopsis thaliana*, *Lycopersicon esculentum*, *Vitis vinifera*, *Antirrhinum majus* and *Oryza sativa* (Shirley, Hanley & Goodman, 1992; Bongue-Bartelsman et al., 1994; Sparvoli et al., 1994; Holton & Cornish, 1995; Chen, SanMiguel & Bennetzen, 1998). Although multiple *DFR* genes are existed, only one of them is catalytically active. For example, there are three tandem *DFR* genes in common morning glory (*Ipomoea purpurea*), but only mutation in *DFR-B* will stop the production of anthocyanin (Hoshino, Johzuka-Hisatomi & Iida, 2001). Similarly, multiple *DFR* genes are also found in *Gerbera hybrid*, but only *GDFR1* was expressed in flowers and catalytically active (Helariutta et al., 1993). Additionally, further biochemical analyses reveal that DFR proteins in certain species have substrate specificities, and this can be the main reason for affecting the content and ratios of anthocyanins, thus determine the final colors of plants (Liu et al., 2019). For example, *Petunia (Petunia hybrida)* and *Cymbidium hybrid* are devoid of pelargonidin-type anthocyanins as well as orange-red flowers, the reason is their DFRs cannot efficiently accept monohydroxylated DHK (Meyer et al., 1987; Johnson et al., 1999). So DFR is crucial in anthocyanin pathway, and controlling its expression levels is often regard as most effective for modifying plant colors. The details of DFRs have been extensively studied in many plants (Feng et al., 2021), but very few information is known in *O. japonica*. Our previous analyses showed that pelargonidin-type anthocyanins were not detected in *O. japonica*, therefore, activity study of its DFRs is necessary for interpreting anthocyanin biosynthesis of *O. japonica*.

Ophiorrhiza japonica, a precious Chinese medicinal plant, contains many kinds of bioactive compounds and is commonly used to cure ulcers, leprosy, rheumatism and so on (Sun et al., 2019). Anthocyanins as one of effective medicinal constituents in *O. japonica*, information on its metabolism is very limited. Characterization of key enzymes participated in anthocyanin biosynthesis will aid *O. japonica* medicinal properties improvement. In this current study, three putative *DFR*-like genes were isolated from flowers of *O. japonica*, and only one of them was phylogenetically grouped into subclade of DFR, designated as *OjDFR1* and was further functionally characterized both *in vitro* and *in vivo*. Three *DFR*-like genes showed different expression patterns among the organs, and *OjDFR1* transcripts were highly correlated with anthocyanin accumulation. Then we performed enzyme assay with recombinant protein and checked their catalytic activities for different substrates. Meanwhile, complementation assay of *OjDFR1* in *Arabidopsis dfr (tt3-1)* mutant displayed that the phenotype of *tt3-1* rosette leaves and seeds was successfully restored. Furthermore, transgenic tobacco plants overexpressing *OjDFR1* revealed that *OjDFR1* could interact with tobacco anthocyanin pathway enzymes *in vivo* to produce darker flower color. Totally, the above results highlight the importance of *OjDFR1* in regulating anthocyanin biosynthesis and bring a better understanding of the *O. japonica* anthocyanin biosynthetic pathway.

MATERIALS & METHODS

Plant materials

O. japonica materials used in this study were grown on the mountain in Shibing, Guizhou Province. For gene isolation and tissue-specific expression analysis, different samples including flowers of four developmental stages (1–4) and six vegetative tissues, *i.e.*, flowers (Fl), roots (Ro), stems (St), leaves (Le), scapes (Sc) and calyxes (Ca) were collected as described in our previous paper (Sun *et al.*, 2019). Arabidopsis mutant (*tt3-1*, AT5G42800, ABRC stock number: CS84) was in Landsberg-0 (Ler) ecotypic background and grown at 22 °C with a 16 h light/8 h dark photoperiod. For anthocyanin measurement and RT-PCR analysis, Arabidopsis seedlings of wild type, mutant and transgenic plants cultivated on 1/2 MS medium containing 3% sucrose were obtained. Tobacco plants (Wild-type and transgenic plants) were grown in the greenhouse, the flowers at full-bloom stage were harvested and then used for later analysis. All the samples above were frozen without delay by liquid nitrogen and kept at –80 °C.

Chemical standards

Dihydroquercetin (DHQ), dihydromyricetin (DHM) and dihydrokaempferol (DHK) were bought from Sigma-Aldrich (St. Louis, MO, USA) and prepared as 10 mg/mL solutions in methanol (chromatographic grade). Cyanidin 3-*O*-glucoside for drawing standard curve was also purchased from Sigma-Aldrich and diluted as one mg/mL solutions in methanol (St. Louis, MO, USA).

Gene cloning and sequence analysis

Flowers of *O. japonica* were selected for RNA extraction by using RNA pure Plant Kit (CW BIO, China) according to the instructions. Subsequently, one µg total RNA was used to synthesize the cDNA through the method described previously (Sun *et al.*, 2019). Specific primers for cloning the full-length gene coding sequence were designed based on the assembled transcriptomic information. After PCR, the products of appropriate length were sub-cloned into the pMD18-T vector (Takara, Japan) and verified by sequencing. The list of primers used for gene cloning is provided in Table S1. Multisequence alignment of deduced protein sequences was analyzed using DNAMAN 6.0. And the Neighbor-Joining phylogenetic tree was constructed with MEGA version 7.0 software using 1,000 bootstrap replicates.

Quantitative real-time PCR analysis

RNA extraction and cDNA synthesis of flowers and other vegetative tissues were performed using the methods above. Primers for real-time amplification were designed by IDT1 and shown in Table S1. Then, qRT-PCR reactions were carried out with the BioRad CFX96 Real-Time PCR System (BIO-RAD, Hercules, CA, USA) and TransStart[®] Green qPCR SuperMix (TRANSGEN, China). Thermal cycling conditions were 95 °C for 60 s, followed by 40 cycles of 95 °C for 5 s and 60 °C for 60 s. The *OjActin* gene and *NtTubA1* gene were chosen as the internal reference for *O. japonica* and tabacum samples respectively. Each experiment sample was conducted in triplicate, and the relative

transcript levels of target genes were analyzed by $2^{-\Delta\Delta Ct}$ method. Meanwhile, melting curve analysis and agarose gel electrophoresis were also employed for confirming the purity of PCR products.

Plasmid construction

In order to obtain His-tagged fusions of OjDFR1, its coding sequence containing the *EcoR* I and *Hind* III restriction enzymes sites was cloned into the pET-32a expression vector. Simultaneously, the complete ORF of *OjDFR1* was also inserted into the binary vector pBI121 that previously digested with *Xba* I and *BamH* I. Then above resulting plasmids were transformed into competent cells and verified by sequencing. Primers used to generate the recombinant are present in [Table S1](#).

Production of recombinant OjDFR1 protein and *in vitro* enzyme assay

The recombinant construct pET-32a-OjDFR1 and empty vector were introduced into *E. coli* strain BL21 (DE3) by the heat shock method. Next day, a single colony containing expression plasmid was inoculated in Luria-Bertani (LB) medium and grown at 37 °C with shaking 200 rpm until OD₆₀₀ reached 0.6. For induction, 0.2 mM of isopropyl-β-d-thiogalactopyranoside (IPTG) was added and the cells were further incubated at 15 °C for 24 h. After that, the cells were harvested centrifugation at 5,000 rpm for 10 min at 4° and resuspended in phosphate-buffered saline (PBS) for sonication, then the debris was removed by centrifugation at 12,000 rpm for 5 min at 4°. Subsequently, the His-tagged OjDFR1 proteins were purified by Ni-NTA pre-packed column (TransGen, China) and its purity was examined by using sodium dodecyl sulfate polyacrylamide gel electrophoresis (SDS-PAGE). Protein concentration was estimated by NanoDrop 1,000 (Thermo scientific, Waltham, MA, USA) Spectrophotometer before *in vitro* enzymatic assays.

Three dihydroflavonols (DHQ, DHM and DHK) were selected as experimental substrates to analyze the substrate specificities of OjDFR1 following the methods depicted previously ([Cheng et al., 2013](#)). Shortly, a 500 μL reaction mixture containing 100 mM TrisHCl buffer (pH 7.0), 20 mM NADPH (50 μL), 10 mg/mL substrate (10 μL) and 35 μg OjDFR1 enzyme extract was conducted at 30 °C for 30 min. And the reaction products were identified by HPLC using a Shimadzu HPLC system with detection at 280 nm. An ACCHROM XUnion C18 column was used with the temperature at 30 °C and 20 μL samples were eluted with 1% H₃PO₄ (solvent A) and methanol (solvent B) according to the program as follows: 0 min, 15% B; 0–20 min, 15–60% B; 20–30 min, 60–15% B at flow rate of one mL/min.

Transformation of OjDFR1 into *A. thaliana* and tobacco

To further characterize the function of *OjDFR1*, the resulting vector pBI121-OjDFR1 and empty expression vector were maintained in *Agrobacterium tumefaciens* strain GV3101 by freeze-thaw method. Next, about 5~6-week old *tt3-1* mutant plants were used for genetic transformation followed the standard floral dipping transformation procedure ([Clough & Bent, 1998](#)). Transformants were screened on 1/2 MS medium with 50 mg L⁻¹ kanamycin to obtain T2 seeds. T2 transgenic seeds as well as wild type and mutant seeds

were then cultured on anthocyanin induction media (1/2 MS medium supplemented with 3% sucrose) to set seedlings for phenotypic investigations and metabolite analysis. Meanwhile, the construct pBI121-OjDFR1 was also transformed into tobacco *via* the method described in previous reports (*Imogen et al., 2006*). The transgenic seedlings of tobacco (T1) were grown in green house, and their flower color was observed after flowering. For detecting the expression levels of *OjDFR1*, Arabidopsis *actin-1* and *NtTubA1* were selected as internal control genes.

Total anthocyanin quantifications by HPLC

The contents of anthocyanin were measured in both Arabidopsis seedlings and tobacco flowers according to previous method reported by predecessors (*Li et al., 2017*). Briefly, 0.1 g samples of seedlings as well as tobacco flowers were mashed into a fine powder and placed into extraction solution (water: methanol: hydrochloric acid = 75:24:1) at 4° C for 12 h. After centrifugation at 12,000 rpm for 5 min at 4° C, the supernatant was collected and filtered by a 0.22 µm nylon membrane filter for the following analysis. Chromatographic detection was performed on a Shimadzu HPLC system with the absorbance at 520 nm based on the procedure described by *Li et al. (2017)*. Total anthocyanin content was quantified by using the external standard curve calibration of cyanidin 3-*O*-glucoside with three biological replicates (*Fanali et al., 2001*).

Statistical analysis

All experiments were repeated at least three times. The data were expressed as mean ± SD (standard deviation). We used a *t*-test to test for significant differences in the anthocyanin contents as well as gene expressions between wild type and transgenic tobaccos (*OjDFR1-4* and *OjDFR1-5*). Difference with $P < 0.05$ was considered statistically significant.

RESULTS

Isolation and phylogenetic analysis of *DFR*-like genes from *Ophiorrhiza japonica*

After *in situ* TBLASTN search of *O. japonica* transcriptomic database, three putative genes which encoded NADP-binding reductases were obtained. Among them, one gene was named as *OjDFR1* (cDNA sequences listed in the [Supplementary Materials](#)), which was most likely to be the *bona fide* *DFR*, because its protein sharing 69.74% identities to petunia *DFR*, and 67.98% identities to *Nicotiana tabacum* *DFR*. On the contrary, other two genes might encode ANR-like and FR proteins on the basis of manual BLASTX search, and thus were tentatively designated as *OjANR1* and *OjFR1*.

Multisequence alignment with AtDFR and NtDFR showed that the three *O. japonica* proteins had the highly conserved NADP-binding motif and the substrate-binding domain at their N terminus ([Fig. 2A](#)). And the amino acid residue (at position 134) which is important for substrate specificity of *DFR* is Asn in *OjDFR1* (encoding 357 amino acid residues with a calculated molecular mass of 39.87 kD) indicating that *OjDFR1* may belong to the Asn-type *DFR* and catalyze the reduction of three dihydroflavonols




Figure 2 Amino acid sequences alignment and phylogenetic analysis of DFR-like proteins in *O. japonica* with proteins from other species. (A) Multiple sequence alignment of *O. japonica* proteins with DFR from other species. The putative NADP-binding site and presumed substrate-binding region were marked by red and green boxes. Black triangles indicated the 134rd and 145rd amino acid residue, which is important for substrate specificity of DFR. (B) Phylogenetic analysis of NADPH-dependent reductase proteins from *O. japonica* and other plant species. Proteins of *O. japonica* were indicated with black rhombus. GenBank accession numbers are as follows: *Agapanthus praecox* ApDFR (BAE78769), *Bromheadia finlaysoniana* BfDFR (AAB62873.1), *Freesia hybrid* FhDFR (KU132393), *Iris hollandica* IhDFR (BAF93856.1), *Lilium hybrida* LhDFR (BAB40789.1), *Oryza sativa* OsDFR (BAA36183.1), *Phoenix dactylifera* PdDFR (XP_008797532.1), *Tulipa gesneriana* TgDFR (BAH98155.1), *Zea mays* ZmDFR1 (NP_001152467.2), *Gentiana triflora* GtDFR (BAA12736.1), *Arabidopsis thaliana* AtDFR (BAD95233.1), *Nicotiana tabacum* NtDFR (NP_001312559.1), *Gerbera jamesonii* GjDFR (KF734593), *Gerbera hybrida* GhDFR (AKN56969.1), *Petunia hybrida* PhDFR (CAA56160), *Vitis vinifera* VvDFR (X75964), *Helianthus annuus* HaDFR (XP_022001438.1), *Callistephus chinensis* CcDFR (CAA91922.1), *Arabidopsis thaliana* AtANR (Q9SEV0.2), *Vitis vinifera* VvLAR1 (AAZ82410), VvLAR2 (AAZ82411), VvANR (BAD89742), *Malus domestica* MdANR (AEL79861.1), *Nicotiana tabacum* NtCCR (A47101), *Saccharum officinarum* SoCCR (AJ231134), *Zea mays* ZmCCR1 (Y13734), ZmCCR2 (Y15069) and *Eucalyptus gunnii* EgCCR (X97433). Full-size DOI: 10.7717/peerj.12323/fig-2

(DHK, DHQ, and DHM) to leucoanthocyanidins. Moreover, phylogenetic analysis between OjFRs and other NADPH-dependent reductases (DFR, LAR, ANR, and CCR) are consistent with the above analyses. As shown in Fig. 2B, OjDFR1 protein was grouped into the eudicots subclade of DFR and exhibited most similar to the DFR from petunia. While, other two genes from *O. japonica* clustered outside the DFR branch, one fell into the subgroup of ANR, and another did not belong to a clear subclade, suggesting they might participate in other NADPH-dependent reduction during flavonoid biosynthesis.

Expression analysis of DFR-like genes in different organs

Before further analysis, all three genes were subjected to analyze their expressions at four flower developmental stages (1–4) and six different organs (Fig. 3A). It was observed that transcripts of three genes all exhibited significant spatial and temporal specificity.

The relative levels of *OjDFR1* transcripts peaked at stage 1 and decreased gradually with




Figure 3 Expression profiles of DFR-like genes in *O. japonica*. (A) Four developmental stages of flowers and different tissues. (B) Relative expression levels of *OjDFR*-like genes at four flower developmental stages. (C) Relative expression levels of *OjDFR*-like genes in different tissues. Results represent means \pm SD from three biological replicates. Asterisks above the bars indicate significant difference between the samples judged by Student's *t*-test (* $P < 0.05$, ** $P < 0.01$).

Full-size DOI: 10.7717/peerj.12323/fig-3

the flower development. For *OjFR1* and *OjANR1* genes, although they were also highly expressed in stage 1, but their transcript levels did not show correlation with the accumulation pattern of anthocyanin (Fig. 3B). Furthermore, expression pattern analyses in various tissues displayed that *OjDFR1* showed higher transcript levels in scapes and calyxes accumulating high levels of anthocyanin than that of *OjFR1* and *OjANR1* (Fig. 3C). These data suggest that *OjDFR1* appears to fulfill crucial roles in the anthocyanin biosynthesis in *O. japonica*.

Heterologous expression in *Escherichia coli* and *in vitro* biochemical characterization

To provide direct evidence that the *OjDFR1* encoded the DFR enzyme, *in vitro* enzyme activity assay was implemented. The cDNA of *OjDFR1* was therefore subcloned into the pET-32a expression vector and soluble recombinant proteins were successfully purified followed by SDS-PAGE identification (Fig. 4A). Next, the purified recombinant proteins were incubated with DHQ, DHM and DHK in the presence of NADPH, and the respective reaction products were characterized by HPLC through comparing to the UV spectra and the relative retention time (Cheng et al., 2013). As shown in Fig. 4B and 4C, HPLC analyses detected at 280 nm revealed that new peak was observed from reactions using DHQ and DHM as substrate, and these two peaks were not observed from




Figure 4 Biochemical assays of recombinant *OjDFR1*. (A) Expression of *OjDFR1* in *E. coli*. (1) Marker (2) Total soluble protein from *E. coli* expressing pET-32a (+) vector (3) Total soluble protein from *E. coli* expressing *OjDFR1* prior to induction by IPTG (4) A total of 24 h after induction (5) Purified *OjDFR1*. Enzymatic reaction contained DHQ (B), DHM (C), and DHK (D) as substrates, NADPH as well as protein extracts from *E. coli* harboring pET-32a and *OjDFR1*. LEUC and LEUD represent leucocyanidin and leucodelphinidin respectively. [Full-size !\[\]\(b345a1c4255362eec3746050dd71ccac_img.jpg\) DOI: 10.7717/peerj.12323/fig-4](https://doi.org/10.7717/peerj.12323/fig-4)

control (protein from *E. coli* carrying the pET-32a empty vector). However, reduction of DHK to the respective leucopelargonidin was not observed (Fig. 4D). In fact, the recombinant *OjFR1* and *OjANR1* were also tested by DHQ, DHM and DHK, but no formation of the respective leucoanthocyanidin was observed (Fig. S1).

Functional analysis of *OjDFR1* in *Arabidopsis tt3-1* mutants

To validate the functional identity of *OjDFR1*, its cDNA was subcloned into the pBI121 expression vector containing the CaMV 35S promoter. *Via* floral dip method, *OjDFR1* was introduced into *Arabidopsis tt3-1* mutant which failed to accumulate brown proanthocyanidins in the seeds and anthocyanins in the cotyledons and hypocotyls. Transgenic plants were selected on MS medium with 50 mg/L kanamycin, and then seeds of wild-type, *tt3-1* as well as T2 transgenic lines were grown for 7 days on medium supplemented with 3% sucrose. Phenotypic investigation showed that transgenic *OjDFR1* plants successfully restored the coloration of their seeds and hypocotyls (Fig. 5A), while the empty vector control was still green (raw data). In order to confirm this phenotype was produced by ectopic expression of *OjDFR1*, RT-PCR was carried out. As expected, *OjDFR1* was highly expressed in transgenic plants, and no amplicons were observed in the control (Fig. 5C).

In addition, 7-day-old seedlings were also used to quantitatively determine the amounts of anthocyanins, and the results indicated that anthocyanins level in transgenic *Arabidopsis* was higher than mutant, and similar to wild type (Fig. 5B). As results present in Fig. 6, anthocyanins were not detected in *tt3-1*, but these absent peaks were fully




Figure 5 Complementation of DFR function in Arabidopsis *tt3-1* mutant. (A) Phenotype of the wild type (Ler), mutant (*tt3-1*) and T2 transgenic lines. (B) Total contents of anthocyanins in Arabidopsis seedlings. (C) Expressional analysis of *OjDFR1* in wild-type, mutant and transgenic lines. Results correspond to means from three biological replicates. Asterisks indicate significant differences between means of wild-type and transgenic plants calculated by Student's *t*-test (** $P < 0.01$, *** $P < 0.001$).

Full-size DOI: 10.7717/peerj.12323/fig-5

complemented in transgenic seedlings expressing *OjDFR1*, though the contents were lower. Totally, our anthocyanin analyses strongly prove that *O. japonica* DFR gene is functional for the biosynthesis of proanthocyanidins and anthocyanins *in vivo*.

***OjDFR1* contributes to darker flower color in transgenic tobacco**

In order to characterize the effects of *OjDFR1* transgene on anthocyanin profiles in flowers, it was overexpressed in tobacco plants. Totally 12 independent transgenic lines that confirmed by gene-specific PCR analysis were obtained. Based on visual observations, transgenic tobacco plants possessed darker pink flowers than wild type (Fig. 7A). Then the presence of *OjDFR1* was examined by RT-PCR (Fig. 7B). Additionally, anthocyanin levels




Figure 6 FHPLC analyses of anthocyanins in *Arabidopsis* seedlings. HPLC chromatograms of the samples from seedlings of wild-type (A), mutant (B) and transgenic lines (C, D).

Full-size DOI: 10.7717/peerj.12323/fig-6




Figure 7 Effect of *OjDFR1* on anthocyanin accumulation in transgenic tobacco flowers. (A) Tobacco flowers of wild-type and transgenic lines. (B) Expression profiles of *OjDFR1* in flowers of transgenic tobacco. (C) Quantitation of anthocyanin accumulation levels in transgenic tobacco flowers with HPLC. Results correspond to means from three biological replicates. Asterisks indicate significant differences between means of wild-type and transgenic plants calculated by Student's *t*-test (* $P < 0.05$).

Full-size DOI: 10.7717/peerj.12323/fig-7

in corollas were also measured through HPLC (Fig. S2). Compared with control, corollas of transgenic tobacco overexpressing *OjDFR1* contained significantly higher amount of anthocyanins on fresh weight basis (Fig. 7C), which implying *OjDFR1* protein might interact with the endogenous enzymes of anthocyanin pathways in tobacco and lead to increased anthocyanin accumulation in transgenic flowers. For further investigate the effect of *OjDFR1* on endogenous tobacco in anthocyanin biosynthetic genes, we performed qPCR analysis. Among these genes (*NtCHS*, *NtCHI*, *NtF3'H*, *NtF3'5'H*, *NtDFR*, *NtANS*, *NtUFGT*, *NtAN1a*, *NtAN1b* and *NtAN2*), the expressions of *NtANS*, *NtUFGT* and *NtAN2* were higher in transgenic tobacco flowers, whereas the transcripts levels of *NtCHS*, *NtCHI*, *NtAN1a* and *NtAN1b* were slightly lower in transgenic flowers than in wild-type (Fig. 8).




Figure 8 Expression analysis of endogenous anthocyanin biosynthetic genes in corollas of transgenic tobacco. (A) Expression profiles of structure genes in corollas of transgenic tobacco. (B) Expression profiles of regulatory genes in corollas of transgenic tobacco. Results represent means \pm SD from three biological replicates. Different letters above the bars indicate significant difference between the samples judged by Student's *t*-test ($P < 0.01$). [Full-size !\[\]\(ba1b80118482ccef74a5d718ca4d7242_img.jpg\) DOI: 10.7717/peerj.12323/fig-8](https://doi.org/10.7717/peerj.12323/fig-8)

DISCUSSION

DFR, one of the members of short chain dehydrogenase/reductase (SDR) superfamily, fulfills important regulatory role in the formation of anthocyanins and controls the carbon flux into different anthocyanin biosynthetic branches which results in various anthocyanin profiles (Li *et al.*, 2017). Although single or multiple *DFR* genes are characterized in many plant species, no detailed identification of *DFR* in *O. japonica* has been performed (Chen *et al.*, 2020). In this study, three NADPH-dependent reductase genes were firstly cloned from *O. japonica*. Among them, OjDFR1 tended to be DFR-like proteins, while other two proteins were likely to be ANR and FR respectively according to BLASTX search. Amino acid alignment showed that all these three proteins had the typical NADPH-binding motif and the substrate-binding domain. Correspondingly, phylogenetic tree of NADP-dependent reductases including DFR, ANR, LAR, CCR revealed that OjDFR1 clustered into the subclade of characterized DFRs, which implying the potential “DFR-like” catalytic activity of OjDFR1. As reported previously, many members of protein family have identical or similar functions *in vitro*, and their different roles in plants are largely due to their specific expression patterns (Vieten *et al.*, 2005), so the expression patterns of above three genes were examined temporally and spatially. During flower development, anthocyanin accumulation in *O. japonica* decreased gradually and calyxes were its dominant tissues for the biosynthesis of anthocyanin (Sun *et al.*, 2019). As results present in Fig. 3, the expressions of OjDFR1 exhibited strong correlations with the accumulation patterns of total anthocyanins in flower development process, but OjANR and OjFR were not. OjDFR1 transcript accumulations were similar to *DFR* in *Ginkgo biloba*, *Chrysanthemum* and *Pyrus communis* which was expressed positively correlated with anthocyanin concentrations (Cheng *et al.*, 2013; Li *et al.*, 2017; Sun *et al.*, 2020). However, in different tissues, relatively higher OjDFR1 expression was observed in Sc than that in Ca, which is not in line with anthocyanin phenotype. These results suggest

that *OjDFR1* is not flower specific and can be involved in the biosynthesis of other flavonoids in *O. japonica* such as proanthocyanin.

Knowledge of DFR biochemical properties is vital for understanding the metabolism of flavonoid, especially its regulation in specific branches. Biochemical studies of DFR have demonstrated that substrate specificity of this enzyme could be determined by the amino acid residue at position 134 (Li et al., 2012). As reported previously, the Asn-type DFRs (contain Asn at position 134) could convert all three dihydroflavonols to the respective leucoanthocyanidin (Forkmann & Ruhнау, 1987). However, to our amazement, *OjDFR1* belonged to Asn-type DFRs, but no leucopelargonidin (LEUP) was observed in the enzyme activity assay (Fig. 4), which is in accord with the fact that pelargonidin glycosides were not detected in *O. japonica*, and this result is similar to the research of DFR in *Agapanthus praecox ssp.orientalis* and *Freesia hybrid* (Mori et al., 2014; Li et al., 2017; Sun et al., 2019). It is possible that the ability of *OjDFR1* catalyzing DHK is lost during the evolution because of the relatively low concentration of DHK in *O. japonica*. Or the residue at position 134 is not an absolute factor to determine the substrate specificity of DFR. Exactly, the DFR from *Grape Hyacinth* which has a point mutation at amino acid residue 134 was still able to utilize all three dihydroflavonol as substrates (Liu et al., 2019). So these results indicate that the residue at position 134 is not an absolute factor to determine the substrate specificity of DFR. Indeed, domain swapping experiments had illustrated that the ability of DFR for catalyzing dihydrokaempferol was influenced by the first 170 amino acids (Johnson et al., 2001). Taken together, it will be necessary and interesting to carry out site-directed mutagenesis analysis for confirming the contribution of a specialized amino acid residue in the substrate-binding region towards the activity and substrate preference of DFRs.

Functionality of *OjDFR1* was further investigated *via* its introduction in Arabidopsis *tt3-1* mutants, and the results indicated that *OjDFR1* could rescue the pigmentation phenotype of *tt3-1* which demonstrated *OjDFR1* as a DFR was functionally active *in vivo* for anthocyanin biosynthesis (Fig. 5A). But results in Fig. 5B revealed that the contents of total anthocyanins in transgenic plants were lower in contrast to wild type, and this can be speculated that the catalytic efficiency of *OjDFR1* might be weaker than the DFR of Arabidopsis. As known, wild-type Arabidopsis accumulate pelargonidin in the seedlings (Sun et al., 2016) which corresponds to peak 3 in our HPLC experiments (Fig. 6). So, apparently, the successful restoration of peak 3 in transgenic Arabidopsis was not consistent to the results of *OjDFR1* biochemical properties assays. On one hand, this contradiction might be ascribed to the very low catalytic efficiency of *OjDFR1* towards DHK which was lost in the course of preparation. On the other hand, the enzymes participated in the biosynthesis of natural product are promiscuous, post-translational modification, coenzyme, the relative concentrations of potential substrates and the internal environment of plants are all the influenced factors to determine their activity *in vivo*. Overall, *OjDFR1* could use DHQ and DHM as substrate in *O. japonica*, DHQ and DHK as substrate in Arabidopsis.

Overexpression was used to evaluate whether or not the *OjDFR1* would perform differently in tobacco cells. As shown in Fig. 7A, the transgenic plants overexpressing

OjDFR1 had darker pink flowers, whereas the control produced pale pink flowers, and this coincided well with accumulation levels of anthocyanin (Fig. 7B). Likewise, the significant phenotype of transgenic tobacco mentioned above was also found in the studies of cranberry and *Populus trichocarpa* which hinted the necessary role of DFR for the formation of flower color (Polashock et al., 2002; Huang et al., 2012). In addition, we observed that although the upper of transgenic corolla were darker pink, the lower part was still white as same to the control. Considering that the position of DFR in anthocyanin pathway, we speculate the downstream enzymes (ANS, UFGT or other enzymes) may also play crucial roles in the formation of transgenic tobacco flower color (Ni et al., 2020). Interestingly, overexpression of *OjDFR1* in tobacco resulted in the up-regulation of *NtANS* and *NtUFGT* as well as the regulatory gene *NtAN2* (Fig. 8A). Previously, it was reported that positive feedback regulation of flavonoid biosynthetic genes via pathway intermediates was existed in *Arabidopsis thaliana* mutant (Pourcel et al., 2013). Therefore, we proposed that the up-regulation of *NtANS* and *NtUFGT* might be due to the positive feedback regulation by anthocyanins accumulation (flavonoid pathway intermediates) resulting from *OjDFR1* overexpression. Meanwhile, it seems that *OjDFR1* overexpression may also affect the regulatory role of *NtAN2*, but whether or not the *OjDFR1* can directly interact with R2R3 MYB proteins is not clear, or that the alteration of *NtAN2* expression was due to feed-back from the increased level of intermediates. Out of expectation, HPLC analysis showed that only Cy-type pigments were produced in transgenic plants (Fig. S2). And the deficiency of Dp-type anthocyanins in host tobacco plants is reported for the lack of DHM (Sun et al., 2016). Inconsistent with the results in transgenic *tt3-1*, Pg-type anthocyanins was not detected in transgenic tobacco, and this further demonstrates that the enzymes involved in natural product biosynthesis such as DFR are extraordinarily complicated. Taken together, overexpression of *OjDFR1* gene in tobacco contributes to significant increase of total anthocyanin which is probably attributed to the up-regulation of *NtANS* and *NtUFGT*.

CONCLUSIONS

Collectively, we carried out a comprehensive analysis of *DFR* gene in *O. japonica*. Our results showed that the identified *OjDFR1* gene was associated with color development in flowers. Meanwhile, the *OjDFR1* enzyme could utilize DHQ as well as DHM as substrates *in vitro*, and could restore the biosynthesis of anthocyanins and proanthocyanidins in *Arabidopsis*. Furthermore, the results obtained from transgenic tobacco demonstrated that *OjDFR1* might be a determinant for the categories of anthocyanins aglycons accumulated in *O. japonica*. Therefore, our findings provide a better understanding of anthocyanin biosynthesis in *O. japonica*.

ADDITIONAL INFORMATION AND DECLARATIONS

Funding

This work was supported by the National Natural Science Foundation of China (Grant No. 31760076), the Joint Fund of the National Natural Science Foundation of

China and the Karst Science Research Center of Guizhou province (U1812401), grants from the department of education of Guizhou Province (KY [2021]059), the Karst Mountain Ecological Security Engineering Research Center (KY [2021]007), and grants from the Guizhou Science and Technology Project ([2017]5726, [2018]1011). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Grant Disclosures

The following grant information was disclosed by the authors:

National Natural Science Foundation of China: 31760076.

National Natural Science Foundation of China and the Karst Science Research Center of Guizhou province: U1812401.

Department of education of Guizhou Province: KY [2021]059.

Karst Mountain Ecological Security Engineering Research Center: KY [2021]007.

Guizhou Science and Technology Project: [2017]5726, [2018]1011.

Competing Interests

The authors declare that they have no competing interests.

Author Contributions

- Wei Sun analyzed the data, authored or reviewed drafts of the paper, and approved the final draft.
- Nana Zhou performed the experiments, analyzed the data, prepared figures and/or tables, and approved the final draft.
- Cai Feng performed the experiments, prepared figures and/or tables, and approved the final draft.
- Shiyu Sun performed the experiments, prepared figures and/or tables, and approved the final draft.
- Ming Tang analyzed the data, authored or reviewed drafts of the paper, and approved the final draft.
- Xiaoxin Tang conceived and designed the experiments, authored or reviewed drafts of the paper, and approved the final draft.
- Zhigang Ju conceived and designed the experiments, authored or reviewed drafts of the paper, and approved the final draft.
- Yin Yi conceived and designed the experiments, authored or reviewed drafts of the paper, and approved the final draft.

Data Availability

The following information was supplied regarding data availability:

The raw measurements are available in the [Supplementary Files](#).

Supplemental Information

Supplemental information for this article can be found online at <http://dx.doi.org/10.7717/peerj.12323#supplemental-information>.

REFERENCES

- Bongue-Bartelsman M, O'Neill SD, Tong Y, Yoder JI. 1994.** Characterization of the gene encoding dihydroflavonol 4-reductase in tomato. *Gene* **138**(1–2):153–157
DOI [10.1016/0378-1119\(94\)90799-4](https://doi.org/10.1016/0378-1119(94)90799-4).
- Chen XF, Liu WL, Huang XY, Fu HH, Wang QX, Wang YY, Cao JG. 2020.** Arg-type dihydroflavonol 4-reductase genes from the fern *Dryopteris erythrosora* play important roles in the biosynthesis of anthocyanins. *PLOS ONE* **15**(5):e0232090
DOI [10.1371/journal.pone.0232090](https://doi.org/10.1371/journal.pone.0232090).
- Chen M, SanMiguel P, Bennetzen JL. 1998.** Sequence organization and conservation in sh2/a1-homologous regions of sorghum and rice. *Genetics* **148**(1):435–443
DOI [10.1093/genetics/148.1.435](https://doi.org/10.1093/genetics/148.1.435).
- Cheng H, Li LL, Cheng SY, Cao FL, Xu F, Yuan HH, Wu CH. 2013.** Molecular cloning and characterization of three genes encoding dihydroflavonol-4-reductase from *Ginkgo biloba* in anthocyanin biosynthetic pathway. *PLOS ONE* **8**(8):e72017 DOI [10.1371/journal.pone.0072017](https://doi.org/10.1371/journal.pone.0072017).
- Clough SJ, Bent AF. 1998.** Floral dip: a simplified method for *Agrobacterium*-mediated transformation of *Arabidopsis thaliana*. *Plant Journal* **16**(6):735–743
DOI [10.1046/j.1365-313x.1998.00343.x](https://doi.org/10.1046/j.1365-313x.1998.00343.x).
- Dharmawansa KVS, David WH, Vasantha R. 2020.** Chemopreventive effect of dietary anthocyanins against gastrointestinal cancers: a review of recent advances and perspectives. *International Journal of Molecular Sciences* **21**(18):6555 DOI [10.3390/ijms21186555](https://doi.org/10.3390/ijms21186555).
- Fanali C, Dugo L, D'Orazio G, Lirangi M, Dacha M, Dugo P, Luigi M. 2001.** Analysis of anthocyanins in commercial fruit juices by using nano-liquid chromatography-electrospray-mass spectrometry and high-performance liquid chromatography with UV-vis detector. *Journal of Separation Science* **34**(2):150–159 DOI [10.1002/jssc.201000665](https://doi.org/10.1002/jssc.201000665).
- Ferreira M, Rius SP, Casati P. 2012.** Flavonoids: biosynthesis, biological functions, and biotechnological applications. *Frontiers in Plant Science* **3**:222 DOI [10.3389/fpls.2012.00222](https://doi.org/10.3389/fpls.2012.00222).
- Feng X, Zhang YT, Wang H, Tian ZD, Xin SY, Zhu PF. 2021.** The dihydroflavonol 4-reductase BoDFR1 drives anthocyanin accumulation in pinkleaved ornamental kale. *Theoretical and Applied Genetics* **134**:159–169.
- Forkmann G, Ruhnau B. 1987.** Distinct substrate specificity of Dihydroflavonol 4-Reductase from flowers of *Petunia hybrida*. *Zeitschrift für Naturforschung C* **42**:1146–1148.
- Helariutta Y, Elomaa P, Kotilainen M, Seppänen P, Teeri TH. 1993.** Cloning of cDNA coding for dihydroflavonol-4-reductase (DFR) and characterization of dfr expression in the corollas of *Gerbera hybrida* var. Regina (Compositae). *Plant Molecular Biology* **22**(2):183–193
DOI [10.1007/BF00014927](https://doi.org/10.1007/BF00014927).
- Holton TA, Cornish EC. 1995.** Genetics and biochemistry of anthocyanin biosynthesis. *Plant Cell* **7**(7):1071–1083 DOI [10.1105/tpc.7.7.1071](https://doi.org/10.1105/tpc.7.7.1071).
- Hoshino A, Johzuka-Hisatomi Y, Iida S. 2001.** Gene duplication and mobile genetic elements in the morning glories. *Gene* **265**:1–10.
- Huang Y, Gou JQ, Jia ZC, Yang L, Sun YM, Xiao XY, Song F, Luo KM. 2012.** Molecular cloning and characterization of two genes encoding Dihydroflavonol-4-Reductase from *Populus trichocarpa*. *PLOS ONE* **7**(2):e30364 DOI [10.1371/journal.pone.0030364](https://doi.org/10.1371/journal.pone.0030364).
- Johnson ET, Ryu S, Yi H, Shin B, Cheong H, Choi G. 2001.** Alteration of a single amino acid changes the substrate specificity of dihydroflavonol 4-reductase. *Plant Journal* **25**(3):325–333
DOI [10.1046/j.1365-313x.2001.00962.x](https://doi.org/10.1046/j.1365-313x.2001.00962.x).

- Johnson ET, Yi H, Shin B, Oh B, Cheong H, Choi G. 1999.** Cymbidium hybrida dihydroflavonol 4-reductase does not efficiently reduce dihydrokaempferol to produce orange pelargonidin-type anthocyanins. *Plant Journal* **19**(1):81–85 DOI [10.1046/j.1365-313x.1999.00502.x](https://doi.org/10.1046/j.1365-313x.1999.00502.x).
- Li L, Ban ZJ, Li XH, Wu MY, Wang AL, Jiang YQ, Jiang YH. 2012.** Differential expression of Anthocyanin biosynthetic genes and transcription factor PcMYB10 in pears (*Pyrus communis* L.). *PLOS ONE* **7**(9):e46070 DOI [10.1371/journal.pone.0046070](https://doi.org/10.1371/journal.pone.0046070).
- Li YQ, Liu XX, Cai XG, Shan XT, Gao RF, Yang S, Han TT, Wang SC, Wang L, Gao X. 2017.** Dihydroflavonol 4-Reductase genes from *Freesia hybrida* play important and partially overlapping roles in the biosynthesis of Flavonoids. *Frontiers in Plant Science* **8**:428 DOI [10.3389/fpls.2017.00428](https://doi.org/10.3389/fpls.2017.00428).
- Liu HL, Lou Q, Ma JR, Su BB, Gao ZZ, Liu YL. 2019.** Cloning and functional characterization of Dihydroflavonol 4-Reductase gene involved in Anthocyanidin biosynthesis of *Grape Hyacinth*. *International Journal of Molecular Sciences* **20**(19):4743 DOI [10.3390/ijms20194743](https://doi.org/10.3390/ijms20194743).
- Meyer P, Heidmann I, Forkmann G, Saedler H. 1987.** A new petunia flower colour generated by transformation of a mutant with a maize gene. *Nature* **330**(6149):677–678 DOI [10.1038/330677a0](https://doi.org/10.1038/330677a0).
- Miyagawa N, Miyahara T, Okamoto M, Hirose Y, Sakaguchi K, Hatano S, Ozeki Y. 2015.** Dihydroflavonol 4-reductase activity is associated with the intensity of flower colors in delphinium. *Plant Biotechnology* **32**:249–255.
- Mori S, Otani M, Kobayashi H, Nakano M. 2014.** Isolation and characterization of the dihydroflavonol 4-reductase gene in the monocotyledonous ornamental *Agapanthus praecox* ssp. *orientalis* (Leighton) Leighton. *Scientia Horticulturae* **166**:1–8.
- Ni J, Ruan RJ, Wang LJ, Jiang ZF, Gu XJ, Chen LS, Xua MJ. 2020.** Functional and correlation analyses of dihydroflavonol-4-reductase genes indicate their roles in regulating anthocyanin changes in *Ginkgo biloba*. *Industrial Crops and Products* **152**:112546.
- Polashock JJ, Griesbach RJ, Sullivan RF, Vorsa N. 2002.** Cloning of a cDNA encoding the cranberry dihydroflavonol-4-reductase (DFR) and expression in transgenic tobacco. *Plant Science* **163**(2):241–251.
- Pourcel L, Irani NG, Koo AJ, Bohorquez-Restrepo A, Howe GA, Grotewold E. 2013.** A chemical complementation approach reveals genes and interactions of flavonoids with other pathways. *Plant Journal* **74**(3):383–397 DOI [10.1111/tbj.12129](https://doi.org/10.1111/tbj.12129).
- Sara C, Pierdomenico P, Silvia G. 2020.** What's behind Purple Tomatoes? Insight into the mechanisms of anthocyanin, synthesis in tomato fruits. *Plant Physiology* **182**(4):1841–1853 DOI [10.1104/pp.19.01530](https://doi.org/10.1104/pp.19.01530).
- Shibata K, Shibata Y, Kasiwagi I. 1919.** Studies on anthocyanins: color variation in anthocyanins. *Journal of the American Chemical Society* **41**:208–220.
- Shirley BW, Hanley S, Goodman HM. 1992.** Effects of ionizing radiation on a plant genome: analysis of two *Arabidopsis transparent testa* mutations. *Plant Cell* **4**(3):333–347 DOI [10.1105/tpc.4.3.333](https://doi.org/10.1105/tpc.4.3.333).
- Sonia DP, Diego AM, Cristina G. 2010.** Flavanols and anthocyanins in cardiovascular health: a review of current evidence. *International Journal of Molecular Sciences* **11**(4):1679–1703 DOI [10.3390/ijms11041679](https://doi.org/10.3390/ijms11041679).
- Sparkes IA, Runions J, Kearns A, Hawes C. 2006.** Rapid, transient expression of fluorescent fusion proteins in tobacco plants and generation of stably transformed plants. *Nature Protocols* **1**(4):2019–2025 DOI [10.1038/nprot.2006.286](https://doi.org/10.1038/nprot.2006.286).

- Sparvoli F, Martin C, Scienza A, Gavazzi G, Tonelli C. 1994.** Cloning and molecular analysis of structural genes involved in flavonoid and stilbene biosynthesis in grape (*Vitis vinifera* L.). *Plant Molecular Biology* **24**(5):743–755 DOI [10.1007/BF00029856](https://doi.org/10.1007/BF00029856).
- Sun HL, Bora P, Da-Hye K, Sangkyu P, Ju-Hee Y, Jae-A J, JeMin L, Jong-Yeol L. 2020.** Cloning and functional characterization of Dihydroflavonol 4-Reductase gene involved in anthocyanin biosynthesis of Chrysanthemum. *International Journal of Molecular Sciences* **21**(21):7960 DOI [10.3390/ijms21217960](https://doi.org/10.3390/ijms21217960).
- Sun W, Liang LJ, Meng XY, Li YQ, Gao FZ, Liu XX, Wang SC, Gao X, Wang L. 2016.** Biochemical and molecular characterization of a flavonoid 3-O-glycosyltransferase responsible for anthocyanins and flavonols biosynthesis in *Freesia hybrida*. *Frontiers in Plant Science* **7**:410 DOI [10.3389/fpls.2016.00410](https://doi.org/10.3389/fpls.2016.00410).
- Sun W, Shen H, Xu H, Tang XX, Tang M, Ju ZG, Yi Y. 2019.** Chalcone isomerase a key enzyme for anthocyanin biosynthesis in *Ophiorrhiza japonica*. *Frontiers in Plant Science* **10**:865 DOI [10.3389/fpls.2019.00865](https://doi.org/10.3389/fpls.2019.00865).
- Tanaka Y, Sasaki N, Ohmiya A. 2008.** Biosynthesis of plant pigments: anthocyanins, betalains and carotenoids. *Plant Journal* **54**(4):733–749 DOI [10.1111/j.1365-313X.2008.03447.x](https://doi.org/10.1111/j.1365-313X.2008.03447.x).
- Vieten A, Vanneste S, Wisniewska J, Benková E, Benjamins R, Beeckman T, Luschnig C, Friml J. 2005.** Functional redundancy of PIN proteins is accompanied by auxin-dependent cross-regulation of PIN expression. *Development* **132**(20):4521–4531 DOI [10.1242/dev.02027](https://doi.org/10.1242/dev.02027).
- Xie DY, Jackson LA, Cooper JD, Ferreira D, Paiva NL. 2004.** Molecular and biochemical analysis of two cDNA clones encoding dihydroflavonol 4-reductase from *Medicago truncatula*. *Plant Physiology* **134**(3):979–994 DOI [10.1104/pp.103.030221](https://doi.org/10.1104/pp.103.030221).
- Xie S, Zhao T, Zhang ZW, Meng JF. 2018.** Reduction of Dihydrokaempferol by *Vitis vinifera* Dihydroflavonol 4-Reductase to produce orange Pelargonidin-Type Anthocyanins. *Journal of Agricultural and Food Chemistry* **66**(13):3524–3532 DOI [10.1021/acs.jafc.7b05766](https://doi.org/10.1021/acs.jafc.7b05766).
- Zhao DQ, Tao J. 2015.** Recent advances on the development and regulation of flower color in ornamental plants. *Frontiers in Plant Science* **6**:261.