

POSTER PRESENTATION

Open Access

# Borrelia turcica in Hyalomma aegyptium ticks in Romania

Z Kalmár\*, G D'Amico, IA Matei, AI Paștiu, DI Mărcuțan, MO Dumitache, AD Mihalca

From The 1st Conference on Neglected Vectors and Vector-Borne Diseases (EurNegVec): with Management Committee and Working Group Meetings of the COST Action TD1303 Cluj-Napoca, Romania. 8-11 April 2014

*Testudo graeca* tortoises are distributed in the south-eastern part (Dobrogea region) of Romania. *T. graeca* is a potential host for the three-host ticks, *Hyalomma aegyptium*. *H. aegyptium* ticks are important from epidemiological point of view as they constitute potential reservoirs for numerous zoonotic bacterial pathogens (*Anaplasma phagocytophilum*, *Ehrlichia canis*, *Coxiella burnetii*). However, *H. aegyptium* was reported to host less studied bacteria, non-Lyme members of genus *Borrelia*. Despite its relatively wide distribution range, the extent of co-distribution of ticks with these bacteria was not investigated in detail. The aim of the present study was to evaluate *H. aegyptium* engorged ticks collected from tortoises in south-eastern Romania for the presence of non-Lyme *Borrelia*. Between 2008 and 2013, 448 *H. aegyptium* ticks were collected from 45 *T. graeca* tortoises located in Dobrogea region in Romania. DNA extraction was performed individually from each tick using a commercial kit. For the total 78 (17.4%) *Borrelia* spp. positive ticks, PCR analysis targeting the intergenic spacer 5S-23S region, *glpQ*, respectively *gyrB* genes, and further sequencing was performed for the further identification. Sequences of *gyrB* and *glpQ* genes showed 99%-100% similarities with reptile-associated *Borrelia turcica*. The most frequently infected stages were males (10.7% of the total males examined or 61.5% from the total infected ticks) followed by females (5.36% of the total females examined or 31% from the total infected ticks) and nymphs (1.34% of the total nymphs examined or 7.7% from the total infected ticks). This is the first report of *Borrelia turcica* in Romania.

This research was supported by grant CNCSIS IDEI PCCE 7/2010 and IDEI PCE 236/2011.

Published: 1 April 2014

doi:10.1186/1756-3305-7-S1-P6

Cite this article as: Kalmár et al.: Borrelia turcica in Hyalomma aegyptium ticks in Romania. Parasites & Vectors 2014 7(Suppl 1):P6.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
[www.biomedcentral.com/submit](http://www.biomedcentral.com/submit)



\* Correspondence: [zsuzsa\\_kalmar@yahoo.com](mailto:zsuzsa_kalmar@yahoo.com)

Department of Parasitology and Parasitic Diseases, University of Agricultural Sciences and Veterinary Medicine Cluj-Napoca, Cluj-Napoca, Romania