

RESEARCH ARTICLE

Identification of function-regulating antibodies targeting the receptor protein tyrosine phosphatase sigma ectodomain

Chia-Lun Wu^{1,2}, Serge Hardy¹, Isabelle Aubry¹, Melissa Landry¹, Allison Haggarty¹, Horacio Uri Saragovi³, Michel L. Tremblay^{1,2*}

1 Rosalind and Morris Goodman Cancer Research Centre, Montréal, Canada, **2** Department of Experimental Medicine, McGill University, Montréal, Canada, **3** Departments of Oncology, Pharmacology & Therapeutics, McGill University, Montréal, Canada

* michel.tremblay@mcgill.ca



OPEN ACCESS

Citation: Wu C-L, Hardy S, Aubry I, Landry M, Haggarty A, Saragovi HU, et al. (2017) Identification of function-regulating antibodies targeting the receptor protein tyrosine phosphatase sigma ectodomain. *PLoS ONE* 12(5): e0178489. <https://doi.org/10.1371/journal.pone.0178489>

Editor: Makoto Sato, Osaka University, JAPAN

Received: March 14, 2017

Accepted: May 13, 2017

Published: May 30, 2017

Copyright: © 2017 Wu et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the manuscript.

Funding: M.L.T. is a Jeanne and Jean-Louis Lévesque chair in Cancer Research. This work was supported by the Canada-Israel Health Research Initiative, jointly funded by the Canadian Institutes of Health Research, the Israel Science Foundation, the International Development Research Centre, and the Azrieli Foundation.

Competing interests: The authors have declared that no competing interests exist.

Abstract

Receptor tyrosine phosphatase sigma (RPTP σ) plays an important role in the regulation of axonal outgrowth and neural regeneration. Recent studies have identified two RPTP σ ligands, chondroitin sulfate proteoglycans (CSPGs) and heparan sulfate proteoglycans (HSPG), which can modulate RPTP σ activity by affecting its dimerization status. Here, we developed a split luciferase assay to monitor RPTP σ dimerization in living cells. Using this system, we demonstrate that heparin, an analog of heparan sulfate, induced the dimerization of RPTP σ , whereas chondroitin sulfate increased RPTP σ activity by inhibiting RPTP σ dimerization. Also, we generated several novel RPTP σ IgG monoclonal antibodies, to identify one that modulates its activity by inducing/stabilizing dimerization in living cells. Lastly, we demonstrate that this antibody promotes neurite outgrowth in SH-SY5Y cells. In summary, we demonstrated that the split luciferase RPTP σ activity assay is a novel high-throughput approach for discovering novel RPTP σ modulators that can promote axonal outgrowth and neural regeneration.

Introduction

Deregulation of protein tyrosine phosphorylation impacts a broad spectrum of human diseases including obesity, vascular diseases, cancers and neural degeneration [1, 2]. Receptor protein tyrosine phosphatase sigma (RPTP σ) is highly expressed throughout neural development and remains important for neural plasticity even in the adult brain [3–7]. However, the mechanisms underlying the transduction of signals mediated by RPTP σ remain unclear. Recently, extracellular matrix proteins CSPGs (chondroitin sulfate proteoglycans) and HSPGs (heparan sulfate proteoglycans) were identified as ligands of RPTP σ , suggesting a potential mechanism of RPTP σ regulation in nerve injury models [4, 6, 7]. These studies have demonstrated that binding of CSPG to RPTP σ receptors expressed on the axon surface induces axonal retraction [4, 6]. In addition to CSPGs and HSPGs, a number of RPTP σ -interacting proteins have been identified. Several synaptic molecules including TrkC, Netrin-G3, and Slitrk1-3 have been

found to interact with the extracellular portion of RPTP σ and are thought to play a role in the regulation of RPTP σ function [8–10].

In contrast to receptor tyrosine kinases, receptor tyrosine phosphatases (RPTPs) are inactive as homodimers. An initial model proposed that a motif of the RPTPs located between the transmembrane segment and the first catalytic domain (D1), named the “wedge domain”, mediates dimer formation and causes RPTP inactivation [11–15]. In this model, the wedge domain of one of the two proteins is proposed to occlude the catalytic site of the D1 domain of the other PTP protein. Several studies support the idea that RPTP σ predominately exists as dimers in vivo [16, 17] suggesting that its ligand(s) regulates receptors dimerization status and thus receptor activity by inducing a radical conformational change of this receptor.

Here, we developed a split firefly luciferase-based sensor system to monitor in real time RPTP σ dimerization/activity in response to various ligands. Using this system, we have validated new anti-RPTP σ antibodies that are capable of modulating axonal outgrowth, and which could be developed as potential therapeutics for the treatment of nerve injury and other neurodegenerative diseases.

Materials and methods

Cell culture, antibodies and reagents

293T/17, COS7, SH-SY5Y cells were routinely maintained in DMEM containing 10% Fetal Bovine Serum and 1% penicillin/streptomycin. The following antibodies were used: Anti-PTPRS clone 1H6 (Abnova), Anti-RPTP σ clone 17G7.2 (MediMabs), Anti-tubulin (Sigma). The following reagents were used in this study: Chondroitin Sulfate (Sigma), Heparin (Sigma), Aggrecan (Sigma) and Chondroitin Sulfate Proteoglycan (CSPG) (Millipore), D-luciferin (ThermoFisher Scientific).

Plasmids and constructs

The murine (NM_011218) and human (BC143287) RPTP σ full-length cDNAs were sub-cloned using PCR-based strategies. The PCR products were digested by EcoRI and SalI restriction enzymes and inserted into pRK5-Nluc, pRK5-Cluc, pRK5-FN and PRK5-CF vectors (gift from David Piwnicka-Worms) [18]. The extracellular domain (AA 1–1370) and the DI-DII domain (AA 1370–1904) of RPTP σ without de wedge domain were made by a similar PCR cloning strategy. In addition, since the RPTP σ antibodies used in this study does not recognize the DIDII region of RPTP σ , we included an HA tag on the DI-DII RPTPs N/C constructs to assess their expressions. All plasmid constructs were verified by sequencing.

Luciferase assay in living cells

293T/17 cells were transfected with Lipofectamine 2000 (Invitrogen) according to the manufacturer’s instructions. 5000 cells were plated in a 96 well plate for 24 hrs followed by transient transfection (0.1 μ g/well) of the various constructs. Following transfection, 1 mM D-luciferin in phenol red free medium was added for 30 min at 37°C. Quantification of the luminescence was carried out using a luminometer (FLUOstar Omega) analysed in real-time for a total of 15 min with data collected at 3 min interval.

Native PAGE

Two micrograms of plasmid DNA was transfected by lipofectamine 2000 in 293T/17 cells. 48 hours post-transfection, cells were treated with purified CSPG then lysed with NP-40 lysis buffer (1% NP-40, 20 mM HEPES, pH 7.4, 150 mM NaCl, 5mM NaF, 1 mM NaPO₄, 10%

Glycerol) containing 1X protease inhibitor (Roche) and 1 mM sodium orthovanadate. Cleared lysates were collected after centrifugation at 12000 rpm for 10 minutes. Protein concentration was determined by BCA (bicinchoninic acid) assay and the lysate supernatants were divided into two portions. One portion was added to 2X SDS-sample buffer (100 mM Tris-Cl (pH 6.8), 4% (w/v) SDS (sodium dodecyl sulfate), 0.2% (w/v) bromophenol blue, 20% (v/v) glycerol, 200 mM β -mercaptoethanol) and the other portion was added to coomassie blue sample buffer (100 mM Tris-Cl, pH6.8, 20% glycerol, 5% Coomassie Brilliant Blue G-250). The protein sample with Coomassie blue sample buffer was subjected to native or SDS PAGE (Polyacrylamide gel electrophoresis) followed by standard western immunoblotting procedures as described below.

Western blot analysis and immunoprecipitation

Proteins were separated by SDS-PAGE and transferred onto polyvinylidene difluoride (PVDF) membrane (Immobilon-P Millipore). The membrane was blocked with 5% BSA in TBST at room temperature then was incubated with primary antibody. For the immunoprecipitation experiment, 500 μ g of cell lysates were incubated with the indicated RPTP σ antibodies conjugated to agarose beads for 4 hr. Immunoprecipitated complex were washed with NP-40 lysis buffer for 3 times followed by western blot analysis.

Membrane protein purification

Six micrograms of RPTP σ expression constructs were transfected into 297T/17 cells in a 10 cm dish using lipofectamine 2000 according to the manufacturer protocol. Ten dishes of transfected cells were rinsed with cold PBS then scraped in 1 ml PBS and collected by centrifugation. Cells were resuspended in membrane wash buffer (0.32 M sucrose, 10 mM Tris-HCl, pH7.5, 5 mM MgCl₂, 50 mM NaCl) and homogenized using a glass Dounce homogenizer using 25 strokes. Different cellular fractions were separated by ultracentrifugation at 20,000 rpm for 10 min at 4°C. Membrane protein fractions were pelleted and washed 3 more times with the same wash buffer and the concentration measured by BCA assay.

In vitro phosphatase assay

50 μ g of the membrane protein fraction overexpressing GFP or RPTP σ N/C was incubated with different amount of Chondroitin sulfate or Heparin at 37°C for 30 minutes followed by the addition of 1 mM p-nitrophenyl phosphate for 30 min. The assay was conducted at 25°C in 96 well plates and the absorbance of p-nitrophenol was monitored at 405 nm every minute using a Varioskan plate reader (Thermo Electron).

Generation of antibodies against human RPTP σ

Specific sequences in the extracellular domain of human RPTP σ were selected to generate peptides. The KLH-linked RPTP σ peptides were injected into 2 months old RPTP σ KO mice in order to induce antibody-dependent immune response. Four weeks after post injection, mice's blood was withdrawn for ELISA analysis. Mice positive to ELISA further injected KLH-linked RPTP σ peptides for the second boost of immune response. Mice reached to 3 months old then were sacrificed to take out blood and spleen for hybridoma generation. Mice were euthanized using isoflurane and carbon dioxide followed by cervical dislocation. All animal procedures were approved by the McGill Animal Care Committee and were conducted according to the Canadian Council of Animal Care ethical guidelines for animal experiments.

ELISA

Epitope peptides from human RPTP σ extracellular region were coated onto 96 well plates overnight. The coated plates were washed with PBS 3 times to remove the excess peptides and 3% BSA solution in PBS was added to the peptide coated plate to block non-specific binding. Supernatant from individual hybridoma medium was added and incubated for 1 hr. Followed by 3 washes, a secondary antibody (1:10000, anti-mouse IgG-HRP) was added, incubated for 1 hr. After 3 washes with PBS, ABTS (2,2'-Azinobis [3-ethylbenzothiazoline-6-sulfonic acid]-diammonium salt) was added and the absorbance read at 410 nm using a Varioskan plate reader (Thermo Electron).

Flow cytometry

Six micrograms pRK5-hRPTP σ was transfected in COS7 cells using lipofectamine 2000. 24 hrs following transfection, cells were detached using 2.5 mM EDTA treatment and blocked by "Fc blocker" (BD) for 30 min on ice. Primary RPTP σ antibody (1H6) (1:100) or hybridomas supernatant was added to label the epitope on the cell surface for one hour on ice. Cells were then washed with cold PBS containing 2% FBS and labeled with anti-mouse FITC (1:100) for one hour on ice. Labeled cells were subjected to FACS analysis. The result was analysed using Flowjo software (Flowjo).

Immunofluorescence

SH-SY5Y human glioma cells were treated with 3% FBS, 10 μ M Retinoic acid for 96hr in order to induce cell differentiation. After 96 hr, cells were washed with PBS and fixed by 4% paraformaldehyde for 5 min on ice. Cells were incubated with 5% BSA for one hour at RT in order to prevent non-specific binding. Primary antibody (IgG, anti-hRPTP σ 1.3H12, anti-hRPTP σ 4.5H5 at 10 μ g/ml each) was applied to cells and incubated for 2 hr at RT. Following PBS washes, cells were incubated with the secondary antibody (1:2000, anti-mouse FITC) for 2 hr at RT. Finally, cells were washed 3 times with PBS and stained with actin dye (Phalloidin) and nucleus dye (DAPI) for 30 min. Samples were mounted and subjected to immunofluorescence analysis.

Neurite growth assay

SH-SY5Y cells were infected with lentivirus carrying GFP. The infection procedures were described in a previous study [19]. GFP positive cells were treated with differentiation medium, DMEM 3% FBS with 10 μ M Retinoic acid, for 96hr in presence or absence of 10 μ g/ml of the indicated antibodies. Cells were subjected to the image analysis using the NeuronJ program. The length of neurite was measured from the cell body to the tip of the neurite.

Results

The RPTP σ split luciferase system

The split luciferase system is particularly well suited for evaluating real-time receptor/ligand interactions in a cellular context [18, 20–22]. Thus, we first developed a novel assay using this luciferase fragment complementation assay to monitor the dimerization status of RPTP σ in live cells in response to various ligands. We generated several different constructs to assess the contribution of different protein segments of RPTP σ . Amino and carboxyl terminal ends of the reporter luciferase gene were independently fused to the carboxyl-terminal end of full-length, truncated and/or tagged forms of RPTP σ as described in Fig 1A. Following validation of their expression in Hek293T (Fig 1B), the basal luciferase activity of these constructs




Fig 1. Description and characterization of RPTP σ constructs for the split luciferase assay in live cells. (A) Schematic of RPTP σ constructs used in this study. Different RPTP σ domains were fused with either NLuc or Cluc fragment independently as indicated. FN: Fibronectin domain; Ig: Immunoglobulin-like domain. (B) Western blot analysis following transfection in 293T/17 cells with various RPTP σ constructs. IB: Immunoblot. (C) 293T/17 cells were transfected with various RPTP σ N/C constructs as indicated and 48 hrs post-transfection, relative luminescence unit (RLU) was determined using a luminometer following the addition of 1mM of D-luciferin. (D) 293T/17 cells were co-transfected with RPTP σ FN/FC and 48 hrs post-transfection, rapamycin was added for 4 hr in order to induce FRB and FKBP association followed by luminescence measurement. Values represent fold changes vs control (DMSO) and the mean of three independent experiments performed in triplicates is given \pm SD. * $p < 0.05$.

<https://doi.org/10.1371/journal.pone.0178489.g001>

indicated that both full length and the extracellular domain of RPTP σ displayed higher levels of luciferase activity, indicating that the extracellular domain of RPTP σ is critical for dimer formation (Fig 1C). We also observed a correlation between the expression levels of the extracellular domain of RPTP σ (exRPTP σ) and its luminescence signal when compared to full length RPTP σ . As the extracellular and transmembrane domains are critical for dimer formation of many RPTPs [11–15], we also clearly show here that removal of these regions (DIDII RPTP σ) affects dimerization (Fig 1C), which also confirms the validity of our approach. Importantly, RPTP σ dimerization is not due to transient overexpression artifact since the DIDII domain of RPTP σ had low luminescence activity. Thus, this result also indicates that RPTP σ dimers formation are driven by the presence of pre-existing extracellular ligands including the previously reported RPTP σ ligands, HSPGs which is known to be expressed on Hek293T cell surface [23].

In order to confirm that RPTP σ dimerization can be induced in this assay, we used the FRB-rapamycin-FKBP dimerization system [18] which induces the dimerization of RPTP σ in

response to rapamycin. Thus, FRB-Nluc and the FKBP-Cluc domains were fused independently with the full length of RPTP σ (Fig 1A and 1B). Although the presence of the FRB and FKBP domains decreased basal luminescence activity, it was still higher than the DIDII domain of RPTP σ ($p = 0.0312$) (Fig 1C). Interestingly, after rapamycin treatment, we detected a 3-fold increase in bioluminescence signal in rapamycin treated cells compared to control cells (Fig 1D), confirming the functionality of RPTP σ dimerization split luciferase assay in live cells.

Modifying RPTP σ dimerization status by its ligands

Since dimerization and monomerization of RPTP σ could be seen in live cells, it opens the possibility that other ligands could modulate the dimerization status of RPTP σ through ligand/receptor interactions. Here we initially tested chondroitin sulfate (CS) and heparin, an analog of heparan sulfate since they are reported ligands of RPTP σ [4, 7, 16]. Moderately increased luciferase signals were seen in Heparin treated RPTP σ N/C expressing cells (Fig 2A). In contrast, addition of CS led to a dose-dependent decrease in luciferase activity (Fig 2B). Thus, these results suggested that these two known ligands interact differently with RPTP σ since CS seemingly acts on RPTP σ to prevent dimerization, whereas Heparin induces dimerization. We further examined the roles of CSPGs in the mediation of RPTP σ monomerization using Aggrecan, a member of the CSPG family protein. Adding this ligand to RPTP σ N/C expressing cells led to a decreased bioluminescence signal (Fig 2C), suggesting that Aggrecan inhibited RPTP σ dimerization in cells. In addition, a native gel showed that dimeric and monomeric forms of RPTP σ both exist in cells and when cells are treated with CSPG, it resulted in a switch from the dimer to the monomer form of RPTP σ (Fig 2D). Taken together our data demonstrates that CSPG and its analog promote RPTP σ monomerization.

Inactivation of the RPTPs CD45 and RPTP α has been shown to be mediated by receptor dimerization [12–15]. However, it is unknown if this mechanism also applies to RPTP σ . Since both CS and heparin have opposite effect on RPTP σ dimerization, we employed an *in vitro* phosphatase assay to test if their binding caused a similar variation in RPTP σ enzymatic activities. To that end, we isolated membrane fractions from cells transfected with full-length RPTP σ proteins followed by a phosphatase assay using the artificial substrate pNPP. Immunoblotting showed that RPTP σ was enriched in the membrane fraction (Fig 3A). In addition, the membrane fraction from RPTP σ -transfected cells displayed a three-fold increase in phosphatase activity compared to cells transfected with a GFP expression vector (Fig 3B), indicating that full-length RPTP σ protein is a functional phosphatase in the membrane fraction. Membrane fractions derived from RPTP σ expressing cells were treated with either heparin or CS and further subjected to phosphatase assay as described above. Although we did not observe significant changes of phosphatase activity in presence of Heparin, CS treatment promoted phosphatase activity (Fig 3D). This strongly suggests that CS acts by inducing RPTP σ monomerization to increase its *in vivo* phosphatase activity.

Antibody against the extracellular domain of RPTP σ promotes axonal growth

In an effort to generate antibodies targeting the extracellular domain of RPTP σ , we generated and screened hybridoma clones to obtain 17 positive clones revealed by ELISA (Fig 4A). Furthermore, COS7 cells transiently overexpressing RPTP σ were analyzed by Fluorescent Activated Cell Sorting (FACS) to identify which of those antibodies produced by these hybridoma cells had the capacity to recognize the native structure of the extracellular domain of RPTP σ and we identified eleven clones that were positive by FACS analysis (Fig 4B).




Fig 2. CSPG analogs prevent RPTP σ dimerization whereas Heparin induces it in living cells. (A-C) 293T/17 cells were co-transfected with RPTP σ N/C and 48 hr post-transfection, cells were treated with the indicated RPTP σ ligands for 5 min followed by luminescence measurement. Values are expressed as percentage vs control (without treatment) and the mean of three independent experiments performed in triplicates is given \pm SD. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$. (D) RPTP σ -transfected 293T/17 cells were treated with the indicated concentration of CSPG for 30 min and protein lysates were separated on a native (*top*) or SDS PAGE (*bottom*) followed by western blot analysis using the indicated antibodies.

<https://doi.org/10.1371/journal.pone.0178489.g002>

Based on ELISA and FACS analysis, 1.3H12 and 4.5H5 antibodies were selected because they displayed the highest sensitivity for RPTP σ . They were purified for further characterization. The two antibodies reacted only with the human form of RPTP σ but not the mouse isoform (Fig 5A). Furthermore, we confirmed that they recognize native protein in immunoprecipitation experiments with strong specificity since both 1.3H12 and 4.5H5 only recognize human RPTP σ but not mouse one (Fig 5B). In addition, immunofluorescence analysis detected the endogenous hRPTP σ expression in SH-SY5Y human glioblastoma cells that when differentiated display neuronal morphology and extensive axonal outgrowth. Since RPTP σ is a cell-surface protein, these antibodies are able to recognize the extracellular




Fig 3. Chondroitin Sulfate increases RPTP σ activity. (A) 293T/17 cells were co-transfected with GFP control or RPTP σ N/C and 48 hrs post-transfection, cells were lysed and membrane fraction were isolated followed by western blot analysis. (B) The membrane fraction was incubated with the phosphatase substrate PNPP and release of PNP was measure using the Varioskan plate reader. Phosphatase activity values are expressed as percentage over GFP control. (C-D) 50 ug of isolated protein membranes fractions were incubated with different amount of Chondroitin Sulfate (CS) or heparin for 30 min at RT followed by the phosphatase activity assay. Values are expressed as percentage vs control (without treatment) and the mean of three independent experiments performed in triplicates is given \pm SD. *, $p < 0.05$.

<https://doi.org/10.1371/journal.pone.0178489.g003>

fragment of RPTP σ in non-permeabilized conditions. RPTP σ exhibited a punctate staining pattern on the axon-like structures of differentiated SH-SY5Y cells stained with 1.3H12 and 4.5H5 antibodies but not IgG control (Fig 5C). This staining pattern is consistent with that of mouse RPTP σ reported in a previous study [9], further confirming that RPTP σ localizes preferentially on the surface of axons.

We tested whether antibodies modulate the dimerization of RPTP σ using the split luciferase assay. Our result shows that the 4.5H5 antibody but not the 1.3H12 antibody significantly increased luminescence, indicating that 4.5H5 was able to significantly induce the dimerization of RPTP σ (Fig 6A). Finally, to determine if these antibodies have the capacity to regulate neurite growth, differentiated SH-SY5Y cells were incubated in the presence of 1.3H12, 4.5H5




Fig 4. Generation of antibodies targeting the extracellular domain of human RPTP σ . (A) ELISA assay using peptides from human RPTP σ extracellular region coated in 96 well plates and incubated with the supernatant of hybridomas. (B) COS7 cells were transfected with human RPTP σ (hRPTP σ) and FACS analysis were performed using the supernatant of 11 positives hybridomas identified by ELISA. IgG is the negative control whereas the commercially available clone 1H6 is the positive control. hRPTP σ positive represent the percentage of transfected cells detected expressing hRPTP σ on the cell surface. All experiments were performed in triplicates for a total of three experiments, which gave similar results. One representative experiment is given \pm SD. * $p < 0.05$, ** $p < 0.01$.

<https://doi.org/10.1371/journal.pone.0178489.g004>




Fig 5. Characterization of purified monoclonal antibodies targeting RPTP σ . COS7 cells were transfected with either human RPTP σ (hRPTP σ) or mouse RPTP σ (HA-mRPTP σ) and 24 hrs post-transfection, western blot (A) and immunoprecipitation (B) analysis was performed using the indicated purified monoclonal antibodies. Antibody 1H6 reacted with both forms was used as positive control. Actin was the loading controls. **C:** Differentiated SH-SY5Y cells were immunostained using the purified 1.3H12 or 4.5H5 monoclonal antibodies for RPTP σ (green), Dapi for the nucleus (blue) and Phalloidin for actin (red). IgG was used as negative control.

<https://doi.org/10.1371/journal.pone.0178489.g005>

and IgG control respectively for 96 hrs. The result shows that SH-SY5Y cells treated with the 4.5H5 antibody bore longer axons than cells treated with the IgG control or 1.3H12 antibody (Fig 6B and 6C). Overall, these data suggest that the 4.5H5 antibody could potentially be used for therapeutic treatment of nerve injury and possibly other neurodegenerative diseases.

Discussion

RPTP σ is one of the three members of the type II sub-family of receptor PTPs [24]. Our group and others have shown over the past decades that RPTP σ is a potential target for the development of novel therapies for spinal cord injury and multiple neurodegenerative diseases [3, 4, 6, 7]. Moreover, we recently reported that RPTP σ inhibition could improve memory [5]. These




Fig 6. RPTP σ 4.5H5 monoclonal antibody induces RPTP σ dimerization and increases neurite outgrowth in SH-SY5Y cells. (A) 297T/17 cells were transfected with hRPTP σ N/C and 48 hrs post-transfection, cells were treated with various amount of the indicated RPTP σ monoclonal antibody followed by luminescence measurement. IgG was used as negative control. Values are expressed as fold changes vs respective IgG control. All experiments were performed in triplicates for a total of three experiments, which gave similar results. One representative experiment is given \pm SD. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$ (B) SH-SY5Y neuroblastoma cells were treated with human RPTP σ antibodies (10 μ g/ml) in the differentiation medium. 96 hr post-treatment cells were subjected to image analysis using the NeuronJ program and neurite length and the distribution of neurite length (C) were determined. A.U.: arbitrary unit. One representative experiment of a total of three experiments is shown. ***, $p < 0.001$.

<https://doi.org/10.1371/journal.pone.0178489.g006>

exciting findings underlie the need for a screening assay that can monitor inhibition of the catalytic activity mediated by RPTP dimerization.

We selected the split luciferase assay, which can be used to monitor protein-protein interactions in intact cells [18, 20–22]. We have demonstrated that this assay can be used to screen for ligands, potentially for small molecule inhibitors, and for anti-RPTP σ antibodies that may regulate RPTP σ dimerization. Surprisingly, validation of this assay also revealed some insight into the structural characteristics of RPTP σ in cells. In live cells, RPTP σ exists predominantly as a dimer. Our results showed that the dimerization of RPTP σ is not solely attributed to the intracellular wedge domain, as previous studies have suggested [13, 15, 25], because the membrane anchored extracellular domain of RPTP σ is still capable of forming a dimer, without the intracellular domain. The DIDII phosphatase domain of RPTP σ without the wedge domain displayed fairly low but noticeable luminescence suggesting that the DIDII domain mainly presents as a monomer. On the other hand, different versions of RPTP σ pairings assayed by the luciferase system suggest that the dimerization of RPTP σ is mainly attributed to the extracellular domain and not to the intracellular domain. It also suggests that the dimerization of RPTP σ is not only regulated by the wedge domain and the DII domain [13, 17], but also by ligands/receptor interactions.

CSPGs and HSPGs are newly identified ligands of RPTP σ [4, 6, 7] and crystallographic analysis suggests that CSPGs prevent RPTP σ dimerization, whereas HSPGs induce RPTP σ clustering [26]. In addition, both CSPGs and HSPGs share the same binding site on the first Ig-like domain of RPTP σ suggesting that they compete for RPTP σ binding, inducing opposite effects on axonal outgrowth [26]. CSPGs inhibit axon outgrowth in RPTP σ WT neurons. This effect was significantly diminished in RPTP σ KO neurons consistent with previous reports [3, 4, 6, 27]. In addition, HSPGs dramatically promoted axon outgrowth in WT but not KO neurons, suggesting that this promoting effect by HSPGs depends on RPTP σ . Both CSPGs and HSPGs have a similar affinity for RPTP σ ; therefore, the CSPGs/HSPGs ratio is the key factor in determining RPTP σ dimerization status.

Using our screening system, we confirmed that CSPG is a ligand of RPTP σ and that CSPGs prevent RPTP σ from dimerizing, resulting in its activation. However, HSPG-mediated RPTP σ dimerization seems moderate in our system. One possible explanation is that RPTP σ exists primarily as a dimer in cells and it is more difficult to induce further dimer formation and inhibition of phosphatase activity.

Our results indicate that the ligand-dependent switch between the monomeric and the dimeric forms of RPTP σ provides an important means of regulating RPTP σ function. There is still much to explore regarding the interaction of CSPGs and HSPGs with RPTP σ from a structural point of view. The crystal structure of the extracellular domain of RPTP σ has been resolved and gives us some structural insights into ligands/receptor interactions [26]. There is still an urgent need to acquire the co-crystal structure of CSPGs/RPTP σ and that of HSPG/RPTP σ to elucidate how CSPGs and HSPGs affect the dimerization status of RPTP σ . Such findings could be applied to pharmaceutical development of small molecule inhibitors or therapeutic antibodies to inhibit RPTP σ activity. In a therapeutic context, inhibitors blocking the interaction between CSPGs and RPTP σ may not be sufficient; a putative RPTP σ -targeted drug may require the capability to induce dimerization of RPTP σ in order to completely inhibit RPTP σ activity. Thus, a monoclonal antibody-based approach, as we proposed here, could be ideal.

The development of small molecule inhibitors has been extremely difficult for researchers studying protein tyrosine phosphatases. Traditional PTP inhibitors mostly target the catalytic domain of PTPs, which is highly conserved among all PTPs; hence they generally suffer from lack of specificity toward its targeted PTPs. Recently, Silver's group has developed a wedge

domain peptide-based RPTP σ inhibitor and showed its effectiveness in a spinal cord injury animal model [28]. Still, the specificity of this inhibitor could be an issue since most receptor tyrosine phosphatases possess a well-conserved wedge domain. An alternative approach is to use specific allosteric inhibitors [29] or, as shown here, antibodies targeted to the extracellular domain of RPTPs.

Antibodies of the IgG class generally mediate receptor dimerization since they are bivalent in nature, facilitating the non-covalent crosslinking of two receptors. However, cumulative evidence indicates that the structures of the receptor ectodomains have to be considered when designing therapeutic antibodies. Indeed, taking lessons from therapeutic antibodies directed against EGFR [30], we recognize that it is similarly challenging to design therapeutic antibodies that not only induce dimerization of RPTP σ , but also simultaneously inactivate it. The 4.5H5 antibody gives us some confidence that this type of high specificity antibody could be found for pharmaceutical applications. Finally, the RPTP σ split luciferase assay would provide a valuable system to screen these types of modulators in a high throughput format.

Collectively, data collected using our RPTP σ split luciferase system not only provide a cell-based platform to examine the genuine properties of RPTP σ in live cells, but also reveals its potential for the screening of ligands and potential therapeutic antibodies. This system could ultimately lead to discovery of novel RPTP σ inhibitors designed for the treatment of patients suffering from spinal cord injury or a broad array of neurodegenerative disorders.

Acknowledgments

We thank David Piwnica-Worms for providing the plasmids for the split luciferase system.

Author Contributions

Conceptualization: CLW HUS MLT.

Funding acquisition: MLT.

Investigation: CLW IA ML AH.

Methodology: CLW HUS MLT.

Supervision: SH MLT.

Validation: CLW IA.

Visualization: CLW SH.

Writing – original draft: CLW SH MLT.

Writing – review & editing: CLW SH IA HUS MLT.

References

1. Tonks NK. Protein tyrosine phosphatases: from genes, to function, to disease. *Nat Rev Mol Cell Biol.* 2006; 7(11):833–46. PMID: [17057753](https://pubmed.ncbi.nlm.nih.gov/17057753/). <https://doi.org/10.1038/nrm2039>
2. Julien SG, Dube N, Hardy S, Tremblay ML. Inside the human cancer tyrosine phosphatome. *Nat Rev Cancer.* 11(1):35–49. PMID: [21179176](https://pubmed.ncbi.nlm.nih.gov/21179176/). <https://doi.org/10.1038/nrc2980>
3. Thompson KM, Uetani N, Manitt C, Elchebly M, Tremblay ML, Kennedy TE. Receptor protein tyrosine phosphatase sigma inhibits axonal regeneration and the rate of axon extension. *Mol Cell Neurosci.* 2003; 23(4):681–92. PMID: [12932447](https://pubmed.ncbi.nlm.nih.gov/12932447/).
4. Shen Y, Tenney AP, Busch SA, Horn KP, Cuascut FX, Liu K, et al. PTPsigma is a receptor for chondroitin sulfate proteoglycan, an inhibitor of neural regeneration. *Science.* 2009; 326(5952):592–6. Epub 2009/10/17. <https://doi.org/10.1126/science.1178310> PMID: [19833921](https://pubmed.ncbi.nlm.nih.gov/19833921/).

5. Horn KE, Xu B, Gobert D, Hamam BN, Thompson KM, Wu CL, et al. Receptor protein tyrosine phosphatase sigma regulates synapse structure, function and plasticity. *J Neurochem*. 2012; 122(1):147–61. PMID: [22519304](https://pubmed.ncbi.nlm.nih.gov/22519304/). <https://doi.org/10.1111/j.1471-4159.2012.07762.x>
6. Fry EJ, Chagnon MJ, Lopez-Vales R, Tremblay ML, David S. Corticospinal tract regeneration after spinal cord injury in receptor protein tyrosine phosphatase sigma deficient mice. *Glia*. 2010; 58(4):423–33. Epub 2009/09/26. <https://doi.org/10.1002/glia.20934> PMID: [19780196](https://pubmed.ncbi.nlm.nih.gov/19780196/).
7. Coles CH, Shen Y, Tenney AP, Siebold C, Sutton GC, Lu W, et al. Proteoglycan-specific molecular switch for RPTPsigma clustering and neuronal extension. *Science*. 2011; 332(6028):484–8. Epub 2011/04/02. <https://doi.org/10.1126/science.1200840> PMID: [21454754](https://pubmed.ncbi.nlm.nih.gov/21454754/).
8. Woo J, Kwon SK, Choi S, Kim S, Lee JR, Dunah AW, et al. Trans-synaptic adhesion between NGL-3 and LAR regulates the formation of excitatory synapses. *Nat Neurosci*. 2009; 12(4):428–37. PMID: [19252495](https://pubmed.ncbi.nlm.nih.gov/19252495/). <https://doi.org/10.1038/nn.2279>
9. Takahashi H, Arstikaitis P, Prasad T, Bartlett TE, Wang YT, Murphy TH, et al. Postsynaptic TrkC and presynaptic PTPsigma function as a bidirectional excitatory synaptic organizing complex. *Neuron*. 69(2):287–303. PMID: [21262467](https://pubmed.ncbi.nlm.nih.gov/21262467/). <https://doi.org/10.1016/j.neuron.2010.12.024>
10. Yim YS, Kwon Y, Nam J, Yoon HI, Lee K, Kim DG, et al. Slitrks control excitatory and inhibitory synapse formation with LAR receptor protein tyrosine phosphatases. *Proc Natl Acad Sci U S A*. 110(10):4057–62. PMID: [23345436](https://pubmed.ncbi.nlm.nih.gov/23345436/). <https://doi.org/10.1073/pnas.1209881110>
11. Desai DM, Sap J, Schlessinger J, Weiss A. Ligand-mediated negative regulation of a chimeric transmembrane receptor tyrosine phosphatase. *Cell*. 1993; 73(3):541–54. PMID: [8490965](https://pubmed.ncbi.nlm.nih.gov/8490965/).
12. Majeti R, Xu Z, Parslow TG, Olson JL, Daikh DI, Killeen N, et al. An inactivating point mutation in the inhibitory wedge of CD45 causes lymphoproliferation and autoimmunity. *Cell*. 2000; 103(7):1059–70. PMID: [11163182](https://pubmed.ncbi.nlm.nih.gov/11163182/).
13. Bilwes AM, den Hertog J, Hunter T, Noel JP. Structural basis for inhibition of receptor protein-tyrosine phosphatase-alpha by dimerization. *Nature*. 1996; 382(6591):555–9. PMID: [8700232](https://pubmed.ncbi.nlm.nih.gov/8700232/).
14. Jiang G, den Hertog J, Su J, Noel J, Sap J, Hunter T. Dimerization inhibits the activity of receptor-like protein-tyrosine phosphatase-alpha. *Nature*. 1999; 401(6753):606–10. PMID: [10524630](https://pubmed.ncbi.nlm.nih.gov/10524630/).
15. Majeti R, Bilwes AM, Noel JP, Hunter T, Weiss A. Dimerization-induced inhibition of receptor protein tyrosine phosphatase function through an inhibitory wedge. *Science*. 1998; 279(5347):88–91. PMID: [9417031](https://pubmed.ncbi.nlm.nih.gov/9417031/).
16. Aricescu AR, McKinnell IW, Halfter W, Stoker AW. Heparan sulfate proteoglycans are ligands for receptor protein tyrosine phosphatase sigma. *Mol Cell Biol*. 2002; 22(6):1881–92. PMID: [11865065](https://pubmed.ncbi.nlm.nih.gov/11865065/). <https://doi.org/10.1128/MCB.22.6.1881-1892.2002>
17. Lee S, Faux C, Nixon J, Alete D, Chilton J, Hawadle M, et al. Dimerization of protein tyrosine phosphatase sigma governs both ligand binding and isoform specificity. *Mol Cell Biol*. 2007; 27(5):1795–808. PMID: [17178832](https://pubmed.ncbi.nlm.nih.gov/17178832/). <https://doi.org/10.1128/MCB.00535-06>
18. Luker KE, Smith MC, Luker GD, Gammon ST, Piwnica-Worms H, Piwnica-Worms D. Kinetics of regulated protein-protein interactions revealed with firefly luciferase complementation imaging in cells and living animals. *Proc Natl Acad Sci U S A*. 2004; 101(33):12288–93. PMID: [15284440](https://pubmed.ncbi.nlm.nih.gov/15284440/). <https://doi.org/10.1073/pnas.0404041101>
19. Hardy S, Wong NN, Muller WJ, Park M, Tremblay ML. Overexpression of the protein tyrosine phosphatase PRL-2 correlates with breast tumor formation and progression. *Cancer Res*. 70(21):8959–67. PMID: [20841483](https://pubmed.ncbi.nlm.nih.gov/20841483/). <https://doi.org/10.1158/0008-5472.CAN-10-2041>
20. Yang KS, Ilagan MX, Piwnica-Worms D, Pike LJ. Luciferase fragment complementation imaging of conformational changes in the epidermal growth factor receptor. *J Biol Chem*. 2009; 284(12):7474–82. Epub 2009/01/28. <https://doi.org/10.1074/jbc.M808041200> PMID: [19171934](https://pubmed.ncbi.nlm.nih.gov/19171934/).
21. Macdonald-Obermann JL, Piwnica-Worms D, Pike LJ. Mechanics of EGF receptor/ErbB2 kinase activation revealed by luciferase fragment complementation imaging. *Proc Natl Acad Sci U S A*. 2012; 109(1):137–42. Epub 2011/12/23. <https://doi.org/10.1073/pnas.1111316109> PMID: [22190492](https://pubmed.ncbi.nlm.nih.gov/22190492/).
22. Macdonald-Obermann JL, Adak S, Landgraf R, Piwnica-Worms D, Pike LJ. Dynamic analysis of the epidermal growth factor (EGF) receptor-ErbB2-ErbB3 protein network by luciferase fragment complementation imaging. *J Biol Chem*. 2013; 288(42):30773–84. Epub 2013/09/10. <https://doi.org/10.1074/jbc.M113.489534> PMID: [24014028](https://pubmed.ncbi.nlm.nih.gov/24014028/).
23. Sarrazin S, Lamanna WC, Esko JD. Heparan sulfate proteoglycans. *Cold Spring Harb Perspect Biol*. 3(7). PMID: [21690215](https://pubmed.ncbi.nlm.nih.gov/21690215/). <https://doi.org/10.1101/cshperspect.a004952>
24. Andersen JN, Mortensen OH, Peters GH, Drake PG, Iversen LF, Olsen OH, et al. Structural and evolutionary relationships among protein tyrosine phosphatase domains. *Mol Cell Biol*. 2001; 21(21):7117–36. PMID: [11585896](https://pubmed.ncbi.nlm.nih.gov/11585896/). <https://doi.org/10.1128/MCB.21.21.7117-7136.2001>

25. Tertoolen LG, Blanchetot C, Jiang G, Overvoorde J, Gadella TW Jr., Hunter T, et al. Dimerization of receptor protein-tyrosine phosphatase alpha in living cells. *BMC Cell Biol.* 2001; 2:8. PMID: [11401727](https://pubmed.ncbi.nlm.nih.gov/11401727/). <https://doi.org/10.1186/1471-2121-2-8>
26. Coles CH, Mitakidis N, Zhang P, Elegheert J, Lu W, Stoker AW, et al. Structural basis for extracellular cis and trans RPTP σ signal competition in synaptogenesis. *Nature communications.* 2014; 5:5209. Epub 2014/11/12. PMID: [25385546](https://pubmed.ncbi.nlm.nih.gov/25385546/).
27. Sapieha PS, Duplan L, Uetani N, Joly S, Tremblay ML, Kennedy TE, et al. Receptor protein tyrosine phosphatase sigma inhibits axon regrowth in the adult injured CNS. *Mol Cell Neurosci.* 2005; 28(4):625–35. PMID: [15797710](https://pubmed.ncbi.nlm.nih.gov/15797710/). <https://doi.org/10.1016/j.mcn.2004.10.011>
28. Lang BT, Cregg JM, DePaul MA, Tran AP, Xu K, Dyck SM, et al. Modulation of the proteoglycan receptor PTP σ promotes recovery after spinal cord injury. *Nature.* 2015; 518(7539):404–8. Epub 2014/12/04. PMID: [25470046](https://pubmed.ncbi.nlm.nih.gov/25470046/).
29. Perron MD, Chowdhury S, Aubry I, Purisima E, Tremblay ML, Saragovi HU. Allosteric noncompetitive small molecule selective inhibitors of CD45 tyrosine phosphatase suppress T-cell receptor signals and inflammation in vivo. *Mol Pharmacol.* 2014; 85(4):553–63. Epub 2014/01/30. <https://doi.org/10.1124/mol.113.089847> PMID: [24473749](https://pubmed.ncbi.nlm.nih.gov/24473749/).
30. Leahy DJ. A molecular view of anti-ErbB monoclonal antibody therapy. *Cancer Cell.* 2008; 13(4):291–3. PMID: [18394550](https://pubmed.ncbi.nlm.nih.gov/18394550/). <https://doi.org/10.1016/j.ccr.2008.03.010>