

Epidemiology, Geographical Distribution, and Outcome Analysis of Patients with Electrical Burns Referred to Shiraz Burn Center, Shiraz, Iran during 2008-2019

Abdolkhalegh Keshavarzi^{1*}, Zahra Rahgozar², Mojtaba Mortazavi³, Ali Dehghani⁴

1. Department of General Surgery, School of Medicine, Shiraz University of Medical Sciences, Shiraz, Iran.
2. Department of Medical Education, Shiraz University of Medical Sciences, Shiraz, Iran.
3. Department of Biotechnology, Institute of Science and High Technology and Environmental Sciences, Graduate University of Advanced Technology, Kerman, Iran.
4. Department of Community Health Nursing, School of Nursing, Jahrom University of Medical Sciences, Jahrom, Iran.

*Corresponding Author:

Abdolkhalegh Keshavarzi

Assistant Professor of General Surgery Ward, Shiraz University of Medical Sciences, Burn and Wound Healing Research Center, Amiralmomenin Burn and Plastic Hospital, Beginning of Sadra Town, Shiraz Iran.

Tel.: +98-71-36223418

Email: iliakeshavarzi@yahoo.com

Received: 2022/01/05

Accepted: 2022/19/07

ABSTRACT

Background: Burn is one of the most significant injuries in industrial and developing societies and is one of the most important traumas leading to hospitalization. The aim of this study was to identify the epidemiology, geographical distribution, and outcome of electric burns in Fars province and to present the distribution map.

Methods: In this descriptive-analytical study, the study population involved all electrical burn victims admitted to Amir al-Momenin and Ghotbeddin Hospitals from 2008 to 2019 in Fars province in the south of Iran. Data were analyzed using SPSS software version 22.

Results: Among a total of 246 patients, the average age was 30.78 ± 11.07 . The highest frequency among educational levels was among under-diploma patients (38.6%), and the majority were employed (87.4%). Also, most of the patients were from urban areas (70.3%). The majority of burn incidences occurred at the workplace (57.7%). Also, among the high voltage patients, 25 patients (30.9%) had an amputation, while among low voltage only 12 patients (16.2%) had an amputation. Non-surgical treatment was applied in 68 (28%) cases, while Escharotomy was performed in 28 (11.4%) patients. There was also a statistically significant association between burn voltage and amputation ($P=0.039$).

Conclusion: Based on our report, the rate of electrical burn injuries in Iran is still high, which underlines the need for stronger efforts in effective prevention, such as better public education and the establishment of strict regulations regarding the distribution and use of electricity.

Keywords: Burn, Electrical Injury, Iran

Please cite this paper as:

Keshavarzi A, Rahgozar Z, Mortazavi M, Dehghani A. Epidemiology, Geographical Distribution, and Outcome Analysis of Patients with Electrical Burns Referred to Shiraz Burn Center, Shiraz, Iran during 2008-2019. *World J Plast Surg.* 2022;11(2):102-109.
doi: 10.52547/wjps.11.2.102

INTRODUCTION

Burn is one of the most common health problems globally, which due to longer hospital stay and complications that occur in the field of the disease and the need for long and multi-stage treatments, impose much economic burden on society¹. A burn is a severe skin injury or other organic tissue that has causes the affected skin cells to die. Burn injuries cause an energy transfer to tissues, associated with various degrees of



tissue destruction depending on the cause of burn². Scald injuries, caused by moist heat, appear more superficial initially. Burn injuries classify according to their depth and size. First-degree burns are injuries involving the uppermost (epidermis) layer of skin because of limited superficial symptoms that resolve without scar formation. Second-degree burns, including superficial partial-thickness (2A) and Deep partial-thickness (2B), require wound care.

Burns has a variety of external sources, including thermal (heat-related), radiation, chemical, electrical, fires, and excessive sun exposure³. In thermal burns, the damage is from the surface to the depth, and in clinical examination, its amount can be estimated, but in contrast, in electric burns, the amount of damage in the depth of the tissue is more than its surface, which may not be seen in the initial examination. Deep, including muscles, may cause myonucleosis and require a fasciotomy⁴. In industrial societies, electrical burns are one of the predominant types of burns in public health issues⁵. Electrical burns are classified as flash burns secondary (electrical current passing through the body) and electric arc (no current passes through the body) or as high voltage (greater than 1000 V), low voltage (less than 1000 V)⁶. The electrical arc is generally considered to be a separate class and often causes superficial burns. On the other hand, electrical burns may lead to extensive and deeper burns and can lead to higher mortality⁷. The electric energy passing through the human body causes very complex reactions in the body's tissues and organs. On the other hand, these reactions affect the organism's histological, anatomical, biochemical and structural composition and, on the other, give rise to a series of electrotraumas.

The length of hospital stays and the cost of surgery and hospitalization in patients with electrical burns are longer than heat burns. Patients with electrical burns are at higher risk for infection, morbidity, and amputation, and the early use of flaps has been shown to reduce complications in these patients⁸. These burns also affect other body organs, including the heart, lungs, eyes, and kidneys⁹. The best treatment is prevention. Many long-term complications, especially in cases of high voltage, are irreversible¹⁰. Burn injuries have major mortality and morbidity, especially in developing countries. Implementation of a prevention strategy is necessary to reduce the mortality and morbidity mortality associated with burn injuries¹¹. Previous studies show that in developed countries burn prevention programs have

been successful¹². Epidemiological studies provide essential information for the development of a well-designed prevention program and significantly contribute to the treatment of patients with electrical burns. A previous study indicate differences in the epidemiology and prevalence reported for electrical burns⁷. In this regard, separate epidemiological studies are required in each region to design prevention programs.

Although some projects have studied the epidemiology of electrical burns in Iran, there is no study on patients with electrical burns in Shiraz in the Fars province. Fars Province is one of the provinces of Iran, located in the southern part of the country. This province is the fourth largest province in Iran with a population of 4,851,274, according to the population of the Statistical Center of Iran in 2016. According to the country divisions of 2021 years, the province was divided into 37 cities.

Thus, considering the high prevalence of burn injury in Iran, we aimed to evaluate the epidemiologic and demographic data related to this traumatic injury. The purpose of this study was to get an epidemiologic point of view and hopefully be able to establish the necessary information on hazardous agents in the settings in which burns occur. Then, the geographical distribution of electric burns in Fars province was followed and its map was presented. Considering the importance of preventing electric burns and considering the high cost and morbidity of these patients, we decided to investigate the effective factors in the outcome of electric burns in patients referred to Shiraz burn centers between 2008 and 2019. The data of this study contribute to the process of documenting the epidemiological features and consequences of burn injuries in Shiraz in the Fars province.

MATERIALS AND METHODS

Study design and setting

The study design in this research was descriptive-analytical. In this study, the census method was utilized.

Patient population and data collection

The study population included all patients who had been referred and admitted to Qutbuddin and Amir Al-Momenin burn centers in Shiraz due to electrical burns and their medical records were available in the archives of these two hospitals. Inclusion criteria

included admission with a diagnosis of electrical burns in Qutbuddin and Amir Al-Momenin hospitals in the 2008–2019 year and the presence of medical records in the two centers. In this study, the patients whose medical records were incomplete were excluded. There was not any age or gender limitation. Data collection was conducted based on the registry of clinical information and by preparing a data collection form. A predesigned checklist was used for data collection. The studied variables in the checklist included age, burn percentage, sex, the geographical location of the residence place, the causative substance of burn, marital status, length of hospital stay, level of education, accident location, and final condition of the patient (mortality rate).

Ethical Consideration

The present study was approved by the Medical Ethics Committee of Shiraz University of Medical Sciences (IR.SUMS.MED.REC.1399.494).

Statistical analysis

After recording all the data in the data collection form, all of them will be entered into SPSS software version 22 (IBM Corp., Armonk, NY, USA) under Windows and then using descriptive and analytical

tests such as the Chi-square test and independent *t*-test and, if necessary nonparametric tests such as Fisher Exact test and Mann-Whitney test data will be analyzed. Data analysis included dispersion indicators of study, frequency distribution tables and graphs, and central tendency variables. Regression will be used to determine the predictive effect of factors on burn outcome. Quantitative data will be displayed as the mean (standard deviation), and qualitative data will be displayed as a number (percentage). In this study, a one-way statistical analysis of variance will be performed. In all analyses a 95% confidence level and two-way *p*-value less than 0.05 will be considered. Also, In order to quantify the distribution of the number of patients in Fars province, data entry into the GIS software environment (ArcGIS 10.2) and classification of different levels were used.

RESULTS

General characteristics

In this descriptive-analytical study, 246 hospitalized electrical burn patients from 2008 to 2019 were studied. Among a total of 246 patients, the average age was 30.78 ± 11.07 . Other demographic characteristics are given in Table 1.

Table 1: Demonstrates the distribution of demographic features in study

Variable	Frequency; N=246
Gender	Male 240 (97.6)
	Female 6 (2.4)
Age	<6 5 (2.4)
	6 – 15 9 (4.4)
	16 – 45 171 (83)
	> 45 21 (10.2)
Educational Level	Illiterate 21 (8.5)
	Under-Diploma 95 (38.6)
	Diploma 66 (26.8)
	Academic 36 (14.6)
	Unknown 28 (11.4)
Occupation	Employed 215 (87.4)
	Freelance 76 (35.3)
	Farmer 10 (4.7)
	Manual Worker 69 (32.1)
	Electrician 21 (9.8)
	Employee of Electricity Department 9 (4.2)
	Unknown 1 (0.5)
Others 16 (7.4)	
Residence	Employee 13 (6)
	Unemployed 28 (11.4)
	Unknown 3 (1.2)
	Urban 173 (70.3)
	Rural 64 (26)
	Suburban Areas 8 (3.3)

Treatment and clinical features

The results of this analysis show that the average total body surface area (TBSA) among our patients was 11% [IQR=15], ranging from 0 to 85%. The median ICU admission, duration for the patients was 8 [IQR=14] days, while the median number of received pack cells for blood transfusion was 3 [IQR=6]. The

most dominant complication was increased CPK in 42 (17.1%) of the patients. The patients' need for the operation was 2 [IQR=4] times, while the median duration of admission was 15 [IQR=22] days. The mortality rate in our study was 3.3%. Non-surgical treatment was applied in 68 (28%) cases (Table 2). Based on our data, among the burn patients, 81

Table 2: Treatment and clinical features of electrical burn patients

Variable	Frequency (%) n=246
ICU Admission	Yes 49 (19.9)
	No 197 (80.1)
Blood transfusion	Yes 80 (32.5)
	No 166 (67.5)
Fresh frozen plasma transfusion	Yes 13 (5.3)
	No 233 (94.7)
Treatment	Non-surgical 68 (28)
	Debridement without graft 6 (2.5)
	Debridement with graft 169 (69.5)
Complications	None 49 (19.9)
	Increased CPK 42 (17.1)
	Fasciotomy 13 (5.3)
Type of surgery	Escharotomy 28 (11.4)
	Fasciotomy + Escharotomy 8 (3.3)
	No 197 (80.1)
Amputation	Yes 49 (19.9)
	Upper limb 34 (13.8)
	Lower limb 7 (2.8)
Outcome	Upper and lower 7 (2.8)
	Live 217 (88.2)
	Deceased 8 (3.3)
	Transfer 21 (8.5)

Table 3: Evaluation of applied treatment based on demographic variables of electric burn patients

Variable	Treatment			P-value
	Non-surgical; n=68	Debridement without graft; n=6	Debridement with graft; n=169	
Age; mean ± Standard deviation	31.7 ± 12.3	32.8 ± 6.2	30.2 ± 10.7	0.575
Gender; n (%)	Male 68 (28.7%)	6 (2.5%)	163 (68.8%)	0.260
	Female 0 (0%)	0 (0%)	6 (100%)	
Education; n (%)	Illiterate 2 (9.5%)	0 (0%)	19 (90.5%)	0.384
	Under diploma 31 (32.6%)	2 (2.1%)	62 (65.3%)	
	Diploma 20 (30.3%)	3 (4.5%)	43 (65.2%)	
	Academic 7 (20.6%)	1 (2.9%)	26 (76.5%)	
	Unknown 8 (29.6%)	0 (0%)	19 (70.4%)	
Occupation; n (%)	Employed 62 (29.1%)	6 (2.8%)	145 (68.1%)	0.537
	Unemployed 5 (17.9%)	0 (0%)	23 (82.1%)	
	Unknown 1 (50%)	0 (0%)	1 (50%)	
Residence; n (%)	Urban 54 (31.6%)	5 (2.9%)	112 (65.5%)	0.306
	Rural 13 (20.3%)	1 (1.6%)	50 (78.1%)	
	Suburban 1 (12.5%)	0 (0%)	7 (87.5%)	

(52.3%) were categorized as high voltage, and 74 (47.7%) were due to low voltage. Also, among the high voltage patients, 25 (30.9%) had an amputation, while among low voltage only 12 (16.2%) had an amputation.

The patient's features were evaluated based on the treatment method, the data of which are shown in Table 3. As demonstrated, none of the variables had a significant correlation with the treatment method of the patients.

The age of the patients was evaluated based on their outcome, and based on statistical analysis, there was no significant difference between the two groups ($P=0.625$) (Table 4).

In this study, the voltage was considered a dependent variable, and the rest of the variables were entered as independent variables in the stepwise regression model. The results showed that the degree, duration of admission, ICU, and body part burn variables entered the regression model and explained 31% of the data

changes. Regression analysis of variance showed that regression was significant at the level of one percent.

Dispersion studies and maps

One of the targets of this study is to identify the geographical distribution of electric burns in Fars province and provide a map. In order to quantitatively investigate the distribution of the number of patients in Fars province, data entry into the GIS software environment was used. The results showed that the distribution of patients in the province is different, so the highest number of people with burns was observed in Shiraz with 69 cases. Moreover, Farashband, Pasargad, Arsanjan, Kharameh, and Sarvestan cities had the lowest reports (Figure 1)

However, based on the population ratio, the highest frequency of people with burns was related to Mamasani compared to the population, and Shiraz was an intermediate based on the population ratio (Figure 2).




Figure 1: Distribution map of the number of people with electric burns in Fars Province

greatly depending on the extent of the burns caused by the electric arch. Electrical burns are among the most serious and costly diseases, requiring a lengthy hospital stay and multiple surgical procedures. In our study, the median hospital stay for surgically treated patients was 15 days.

Treatment aims to achieve skin coverage to avoid infection and allow for early mobilization. Adults are more likely to be injured by electricity due to their work, while children are more likely to be injured unintentionally. Electrical injury has a wide range of severity, ranging from mild tissue damage to extreme multi-organ involvement, with both delayed and occult complications and death. Decompression is needed if signs and symptoms of compartment syndrome are present. Upper limb and trunk escharotomy and fasciotomies were performed to 36 relieve compartment pressure in the upper limb and improve ventilation. Amputations are required when the nerves, all blood vessels, tendons, and skin are damaged. In our study, Non-surgical treatment was applied in 68 (28%) cases, while Escharotomy was performed in 28 (11.4%) of the patients. Current that passes through the heart or thorax can cause direct myocardial and cardiac arrhythmias damage, while current that passes through the brain can cause respiratory seizures, arrest, brain injury, and paralysis. Cataracts can be caused by current passing too close to the eyes¹⁵. Our study's mortality rate was 3.3%, and the ICU admission rate in our study was 19.9%. It has been suggested that burn patients admitted late to the hospital have a higher mortality rate^{19,20}. However, a study by Nursal et al.¹⁸ discovered that delayed admission had no statistically significant relationship with the length of stay in the hospital, the cost of treatment, the number of complications, or mortality. These variations may be attributed to the quality of the initial resuscitation provided by other local hospitals. Our study group's mortality rate was around 3.3%, which is lower than in industrialized countries^{21, 22}. However, the other study in the American Burn Association's patient registry shows a mean burn size of 12% of TBSA, which is higher than our figure of 11 percent of TBSA²². The patients' age was evaluated based on their outcome, and based on statistical analysis; there was no significant difference between the two groups. However, other studies have shown that survivors' mean age was lower than no survivors¹⁸. Also, specific burns, such as those in the hands, face, neck, and perineum, have been linked to a worse prognosis¹⁸.

Spatial analysis of the frequency of electric burns showed that the Sarchahan and Mamasani districts, which are far from the center of the province and are some of the suburban cities of the province, have the highest percentage of burns compared to the population. This percentage is somewhat lower in central cities and therefore, educational programs are needed to raise awareness in the suburban cities of the province.

One of the limitations of the current study is the small number of patients with electrical burns. Also, the inclusion of the injured body part (hands, feet, face, etc.) and degree of burn in the information record form could better mention the details for a clearer determination of the factors affecting the burn percentage, length of hospital stay, and etc.

CONCLUSION

Awareness of the functional and aesthetic disabling effects of electrical burns, despite careful treatment, emphasizes the critical importance of accident prevention. To minimize tissue damage and avoid later complications, it is also necessary to perfect tissue-saving surgical procedures threatened by secondary necrosis. Electrical accidents can be prevented if proper safety precautions are taken. Work requiring interaction with electricity should be carefully prepared to ensure that anyone responsible for the different tasks is qualified to do so. At all times and all places, safety should be promoted from the home to the workplace. To prevent these accidents, education and adherence to safety measures and common sense and consideration of the possible dangers of electricity are still needed.

ACKNOWLEDGEMENTS

The authors would like to express their thanks to all colleagues and officials of Shiraz Burn Centers (Amir Al-Momenin and Ghotbeddin Shirazi burn hospitals) as well as the Vice Chancellor for Research of Shiraz University of Medical Sciences for their sincere cooperation. This research was supported by the Research Department at Shiraz University of Medical Sciences, Shiraz, Iran.

CONFLICT OF INTEREST

The authors declare that they have no known competing financial interests or personal relationships

that could have appeared to influence the work reported in this paper.

REFERENCES

1. Ansari-Lari M, Askarian M. Epidemiology of burns presenting to an emergency department in Shiraz, South Iran. *Burns* 2003;**29**(6):579-81. doi: 10.1016/s0305-4179(03)00066-4.
2. Jeschke MG, van Baar ME, Choudhry MA, Chung KK, Gibran NS, Logsetty S. Burn injury. *Nat Rev Dis Primers* 2020;**6**(1):1-25. doi: 10.1038/s41572-020-0145-5.
3. Rosdahl CB, Kowalski MT, editors. Textbook of basic nursing. Lippincott Williams & Wilkins; 2008.
4. Pannucci CJ, Osborne NH, Jaber RM, Cederna PS, Wahl WL. Early fasciotomy in electrically injured patients as a marker for injury severity and deep venous thrombosis risk: an analysis of the National Burn Repository. *J Burn Care Res* 2010;**31**(6):882-7. doi: 10.1097/BCR.0b013e3181f93597.
5. Shih JG, Shahrokhi S, Jeschke MG. Review of adult electrical burn injury outcomes worldwide: an analysis of low-voltage vs high-voltage electrical injury. *J Burn Care Res* 2017;**38**(1):e293-8. doi: 10.1097/BCR.0000000000000373.
6. Arnoldo BD, Purdue GF, Kowalske K, Helm PA, Burris A, Hunt JL. Electrical injuries: a 20-year review. *J Burn Care Rehabil* 2004;**25**(6):479-84. doi: 10.1097/01.bcr.0000144536.22284.5c.
7. Ding H, Huang M, Li D, Lin Y, Qian W. Epidemiology of electrical burns: a 10-year retrospective analysis of 376 cases at a burn centre in South China. *J Int Med Res* 2020;**48**(3). doi: 10.1177/0300060519891325.
8. Karimi H, Akhoondinasab MR, Kazem-Zadeh J, Dayani AR. Comparison of the results of early flap coverage with late flap coverage in high-voltage electrical injury. *J Burn Care Res* 2017;**38**(2):e568-73. doi: 10.1097/BCR.0000000000000422.
9. Piotrowski A, Fillet AM, Perez P, Walkowiak P, Simon D, Corniere MJ, Cabanes PA, Lambrozo J. Outcome of occupational electrical injuries among French electric company workers: a retrospective report of 311 cases, 1996-2005. *Burns* 2014;**40**(3):480-8. doi: 10.1016/j.burns.2013.08.008.
10. Edlich R, Farinholt HM, Winters KL, Britt LD, Long III WB. Modern concepts of treatment and prevention of electrical burns. *J Long Term Eff Med Implants* 2005;**15**(5). 1 - 12. doi: 10.1615/jlongtermeffmedimplants.v15.i5.50.
11. Stylianou N, Buchan I, Dunn KW. A review of the international Burn Injury Database (iBID) for England and Wales: descriptive analysis of burn injuries 2003-2011. *BMJ open* 2015;**5**(2):e006184. doi: 10.1136/bmjopen-2014-006184.
12. Alipour J, Mehdipour Y, Karimi A. Epidemiology and outcome analysis of 3030 burn patients with an ICD-10 approach. *Ann Burns Fire Disasters* 2020;**33**(1):3. PMID: 32523489
13. Groohi B, Rossignol AM, Barrero SP, Alaghehbandan R. Suicidal behavior by burns among adolescents in Kurdistan, Iran: a social tragedy. *Crisis* 2006;**27**(1):16. DOI: 10.1027/0227-5910.27.1.16
14. Mohammadi AA, Keshavarzi A, Erfani A, Modarresi MS, Shahriarirad R, Ranjbar K. Evaluation of epilepsy and burn patterns in a tertiary hospital in southwestern Iran. *Epilepsy Behav* 2020;**111**:107213. doi: 10.1016/j.yebeh.2020.107213.
15. Zikaj G, Xhepa G, Belba G, Kola N, Isaraj S. Electrical burns and their treatment in a tertiary hospital in albania. *Open Access Maced J Med Sci* 2018;**6**(5):835. doi: 10.3889/oamjms.2018.206.
16. Buja Z, Arifi H, Hoxha E. Electrical burn injuries. An eight-year review. *Ann Burns Fire Disasters* 2010;**23**(1):4. PMID: 21991188
17. Patil SB, Khare NA, Jaiswal S, Jain A, Chitranshi A, Math M. Changing patterns in electrical burn injuries in a developing country: should prevention programs focus on the rural population?. *J Burn Care Res* 2010;**31**(6):931-4. doi: 10.1097/BCR.0b013e3181f93924.
18. Nursal TZ, Yildirim S, Tarim A, Caliskan K, Ezer A, Noyan T. Burns in southern Turkey: electrical burns remain a major problem. *J Burn Care Rehabil* 2003;**24**(5):309-14. doi: 10.1097/01.BCR.0000085876.28504.EE.
19. Haberal M, Uçar U, Bilgin N. Epidemiological survey of burns treated in Ankara, Turkey and desirable burn-prevention strategies. *Burns* 1995;**21**(8):601-6. doi: 10.1016/0305-4179(95)00044-c.
20. El Danaf A. Burn variables influencing survival: a study of 144 patients. *Burns* 1995;**21**(7):517-20. doi: 10.1016/0305-4179(95)00035-a.
21. JC, L.J.P. and p.o.b.m.N.Y.C. Livingstone, Burns and scalds: aetiology and prevention. 1996: p. 3-25.
22. Saffle JR, Davis B, Williams P, American Burn Association Registry Participant Group. Recent outcomes in the treatment of burn injury in the United States: a report from the American Burn Association Patient Registry. *J Burn Care Rehabil* 1995;**16**(suppl_3_pt_1):219-32. doi: 10.1097/00004630-199505000-00002.