

ORAL PRESENTATION

Open Access

Decorrelation of low-frequency neural activity by inhibitory feedback

Tom Tetzlaff^{1*}, Moritz Helias², Gaute T Einevoll¹, Markus Diesmann^{2,3}

From Nineteenth Annual Computational Neuroscience Meeting: CNS*2010
San Antonio, TX, USA. 24-30 July 2010

To correctly judge the functional role of cooperative neural activity it is essential to understand how neural correlations are determined by the structure and dynamics of neural networks. Shared presynaptic input is one of the major sources of correlated synaptic activity in such systems. In the asynchronous state of recurrent neural network models, however, spike correlations are considerably smaller than what one would expect based on the amount of shared presynaptic sources [1,2]. A similar lack of correlations in the spiking activity of neighbouring cortical neurons has been observed experimentally [3]. Recently, it has been pointed out that shared-input correlations can be actively suppressed by the dynamics of recurrent networks [4]. Here, we show that both in networks with purely inhibitory coupling (Fig. 1A) and in those with mixed excitatory-inhibitory coupling (Fig. 1B) this active decorrelation affects


mainly the activity at low frequencies (<20 Hz). High-frequency activity, in contrast, is rather unaffected. Simulations rule out that this phenomenon is the result of refractoriness. By means of a simple linear population-rate model we demonstrate that the effect is essentially explained by inhibitory feedback.

Acknowledgements

Supported by the Research Council of Norway (eVITA, Notur), the Helmholtz Alliance on Systems Biology, the Next-Generation Supercomputer Project of MEXT, EU Grant 15879 (FACETS), DIP F1.2, and BMBF Grant 01GQ0420 to BCCN Freiburg. All network simulations were carried out using NEST (<http://www.nest-initiative.org>).

Author details

¹Department of Mathematical Sciences and Technology, Norwegian University of Life Sciences, Ås, Norway. ²RIKEN Brain Science Institute, Wako City, Japan. ³Brain and Neural Systems Team, RIKEN Computational Science Research Program, Wako City, Japan.



* Correspondence: tom.tetzlaff@umb.no

¹Department of Mathematical Sciences and Technology, Norwegian University of Life Sciences, Ås, Norway

Published: 20 July 2010

References

1. Tetzlaff T, Rotter S, Stark E, Abeles M, Aertsen A, Diesmann M: **Dependence of neuronal correlations on filter characteristics and marginal spike-train statistics.** *Neural Comput* 2008, **20**(9):2133-2184.
2. Kriener B, Tetzlaff T, Aertsen A, Diesmann M, Rotter S: **Correlations and population dynamics in cortical networks.** *Neural Comput* 2008, **20**(9):2185-2226.
3. Ecker AS, Berens P, Keliris GA, Bethge M, Logothetis NK, Tolias AS: **Decorrelated Neuronal Firing in Cortical Microcircuits.** *Science* 2010, **327**(5965):584-587.
4. Renart A, de la Rocha J, Bartho P, Hollender L, Parga N, Reyes A, Harris KD: **The Asynchronous State in Cortical Circuits.** *Science* 2010, **327**(5965):587-590.

doi:10.1186/1471-2202-11-S1-O11

Cite this article as: Tetzlaff *et al.*: Decorrelation of low-frequency neural activity by inhibitory feedback. *BMC Neuroscience* 2010 **11**(Suppl 1):O11.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

