

CORRECTION

Correction: Conjunctivitis, the key clinical characteristic of adult rubella in Japan during two large outbreaks, 2012–2013 and 2018–2019

Hidetoshi Nomoto, Masahiro Ishikane, Takato Nakamoto, Masayuki Ohta, Shinichiro Morioka, Kei Yamamoto, Satoshi Kutsuna, Shunsuke Tezuka, Junwa Kunimatsu, Norio Ohmagari

The following information is missing from the Funding statement: This work is supported by research grants from the Emerging/Re-emerging Infectious Diseases Project of Japan, the Japan Agency for Medical Research and Development, AMED (20fk0108095h0002) awarded to SK.

Reference

1. Nomoto H, Ishikane M, Nakamoto T, Ohta M, Morioka S, Yamamoto K, et al. (2020) Conjunctivitis, the key clinical characteristic of adult rubella in Japan during two large outbreaks, 2012–2013 and 2018–2019. PLoS ONE 15(4): e0231966. <https://doi.org/10.1371/journal.pone.0231966> PMID: 32330153



OPEN ACCESS

Citation: Nomoto H, Ishikane M, Nakamoto T, Ohta M, Morioka S, Yamamoto K, et al. (2022) Correction: Conjunctivitis, the key clinical characteristic of adult rubella in Japan during two large outbreaks, 2012–2013 and 2018–2019. PLoS ONE 17(8): e0272880. <https://doi.org/10.1371/journal.pone.0272880>

Published: August 4, 2022

Copyright: © 2022 Nomoto et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.