



Taxonomic Paper

# Incidence of pests and viral disease on pepino (*Solanum muricatum* Ait.) in Kanagawa Prefecture, Japan

Ok-Kyung Kim<sup>‡</sup>, Tadashi Ishikawa<sup>§</sup>, Yoshihiro Yamada<sup>§</sup>, Takuma Sato<sup>‡</sup>, Hirosuke Shinohara<sup>‡</sup>, Ken Takahata<sup>|</sup>

<sup>‡</sup> Laboratory of Plant Pathology, Faculty of Agriculture, Tokyo University of Agriculture, Atsugi-shi, Kanagawa, Japan

<sup>§</sup> Laboratory of Entomology, Faculty of Agriculture, Tokyo University of Agriculture, Atsugi-shi, Kanagawa, Japan

<sup>|</sup> Laboratory of Vegetables, Faculty of Agriculture, Tokyo University of Agriculture, Atsugi-shi, Kanagawa, Japan

Corresponding author: Tadashi Ishikawa ([chuishikawa@gmail.com](mailto:chuishikawa@gmail.com))

Academic editor: Jenő Kontschán

Received: 05 Jul 2017 | Accepted: 17 Aug 2017 | Published: 22 Aug 2017

Citation: Kim O, Ishikawa T, Yamada Y, Sato T, Shinohara H, Takahata K (2017) Incidence of pests and viral disease on pepino (*Solanum muricatum* Ait.) in Kanagawa Prefecture, Japan. Biodiversity Data Journal 5: e14879. <https://doi.org/10.3897/BDJ.5.e14879>

## Abstract

## Background

The solanaceous fruit crop pepino (*Solanum muricatum* Ait.), originating in the Andes, is grown commercially in South American countries and New Zealand. In these areas, pests and diseases of pepino have been identified well; however, to date, these have seldom been investigated in detail in Japan. Herein, we attempt to reconstruct an agricultural production system for commercial pepino crops in Japan, and evaluate the incidence of pests and viral diseases on pepino. The findings of this study will facilitate in developing a better crop system for the commercial cultivation of healthy pepino fruits.

## New information

A total of 11 species, comprising nine insects and two mites, were recognized as pests of pepino plants in our experimental fields in Kanagawa Prefecture, central Honshu, Japan. Of these pest species, the two-spotted spider mite *Tetranychus urticae* Koch, 1836 and the cotton aphid *Aphis gossypii* Glover, 1877, were remarkably abundant than the other pest species. Eventually, 13 species, including two previously recorded, are currently recognized as the pests of pepino in Japan. With regard to viruses, we tested two species *Alfalfa mosaic virus* (AMV) and *Cucumber mosaic virus* (CMV), as well as three genera *Carlavirus*, *Potexvirus*, and *Potyvirus*. No virus was detected in symptomatic pepino leaves collected in our experimental fields. This is a first report on the identification of pests on pepino plants in Kanagawa Prefecture, Japan and elucidates the relationship between currently occurring pests of pepino plants and potential viral pathogens that they can transmit.

## Keywords

insects, mites, virus vector, virus, sweet cucumber

## Introduction

*Pepino*, the Spanish name for sweet cucumber, (*Solanum muricatum* Ait.), is a solanaceous plant cultivated as a fruit crop. It originated in the Andes, became popular in several countries and regions of South America (Heiser 1964), and then it was introduced to Central America and New Zealand. In Japan, cultivation of pepino began in 1984 based on pepino fruits imported from New Zealand in 1983 (Sakata 2011). Since then, pepino caused a major boom and its cultivation rapidly spread throughout Japan within a few years. However, the production gradually declined prior to 1990 due to the low soluble solids content (Brix) in the Japanese pepino fruits (less than 8°Brix) (Sakata 2011). Currently, farmers in Japan do not grow pepino, except for people with gardening as a hobby, who cultivate pepino.

In 2016, our research team began a project for regional development “Launching of Nodai-branded *Pepino Crop*” conducted by Faculty of Agriculture, Tokyo University of Agriculture (TUA; Nodai is a Japanese abbreviated name of the university). The main purpose of this project was to produce high quality and flavorsome pepino fruits with sufficient soluble solids content. As a recent achievement of this project, Takahata (2017) succeeded in increasing the soluble solids content of pepino fruits by using a washer ring (metal washer) at the bottom of the stem. This technique strongly contributes to improving the quality and flavor of pepino fruits.

To date, at least 24 species of insect and mite pests on pepino (Larraín 2002) and one virus infected to pepino (Jones et al. 1980) have been recorded in the native range of pepino, the Andes. In contrast, in regions where pepino was introduced other than Japan, such as New Zealand, China and Turkey, there are only a few records on the pests (two species: Galbreath and Clearwater 1983, Akyazi 2012) and viruses (two species: Thomas et al. 1980, Abouelnasr et al. 2014) in the academic literature. In Japan, little has been known in detail about the pests and viruses of pepino, except that inadequately identified pests such as spider mites and aphids damage to pepino.

It is important to establish solid pest control in its commercial cultivation to produce high quality and stable pepinos. Unfortunately, however, no pesticides applicable to pepino plants have been registered in Japan; this could be attributed to the few detailed studies on pests and diseases of pepino. Therefore, our research team has tried to comprehensively elucidate the pests and viral diseases of pepino in this project in order to contribute to the accumulation of basic knowledge toward the establishment of its pest control. This paper documents the results of our field survey on pests and diseases of pepino in Kanagawa Prefecture, central Japan.

## Materials and methods

### Study sites

This study was conducted at the Atsugi Campus (35.432N 139.346E; at altitudes between 25 and 62 meters above sea level) of Tokyo University of Agriculture (TUA), Atsugi City, Kanagawa Prefecture, Japan, which is surrounded by residential quarters and a woody and grassy park (Fig. 1). The total site area of the campus is approximately 17.3 ha, within which several greenhouses and open fields for experimental use are present. The study site is located in a warm-temperate climate zone and has an annual mean temperature of 15.3 °C and annual mean precipitation of 1,729.9 mm (Japan Meteorological Agency 2017). In the campus, three survey plots were set (Fig. 1); one of these was an open field (approximately 70 m<sup>2</sup>), where 40 pepino plants were cultivated (Plot A) (Fig. 2); another was a greenhouse (approximately 53 m<sup>2</sup>), in which 60 pot pepino plants were grown (Plot B) (Fig. 3); and the other was also a greenhouse (approximately 90 m<sup>2</sup>), in which approximately 400 pot pepino plants were grown (Plot C) (Fig. 4). In these plots, acaricides were applied approximately every two weeks in order to prevent pepino plants from withering due to mites, except for 20 pepino plants in Plot A; when the density of mites became high, appropriate chemicals were sprayed. All the plots were located within a radius of 250 m.




Figure 1. [doi](#)

Aerial photograph of the Atsugi Campus of Tokyo University of Agriculture (TUA) and surrounding residential quarters, with locations of the three survey plots (taken in 2007 by the Geospatial Information Authority of Japan).



Figure 2. [doi](#)

An open field, one of the survey plots in the Atsugi Campus of Tokyo University of Agriculture (TUA), indicated as Plot A in Fig. 1



Figure 3. [doi](#)

A greenhouse, one of the survey plots in the Atsugi Campus of Tokyo University of Agriculture (TUA), indicated as Plot B in Fig. 1



Figure 4. [doi](#)

A greenhouse, one of the survey plots in the Atsugi Campus of Tokyo University of Agriculture (TUA), indicated as Plot C in Fig. 1.

## Sampling methods for insects and mites

All specimens were collected by beating the leaves and branches of pepino plants after observation in field. A total of 34 collections were performed in the three plots from August 30, 2016 to January 21, 2017, for a maximum of 3 h/day in the daytime. The collected insects were killed immediately after capture, using ethyl acetate; aphids, lepidopteran larvae, and mites were fixed in plastic bottles filled with 70–80% ethanol. All specimens were prepared as dry mounted, slide mounted, or ethanol preserved for morphological examination.

## Identification methods for insects and mites

Identification of insect and mite specimens was performed using stereoscopic microscopes (Olympus SZ60 and Olympus SZX16, Tokyo, Japan) and optical microscopes (Olympus BH-2 and Olympus BX41, Tokyo, Japan) by TI and YY according to the following literature: Ehara and Gotoh (2009), Furukawa (2005), Harada and Takizawa (2012), Iwasaki et al. (2000), Kawai (1980), Masumoto and Okajima (2013), Matsumoto (2008), Moritsu (1983), Tanaka and Uesato (2012), Umeya and Okada (2003), Yasunaga et al. (2001), Yasunaga et al. (2015), along with the original descriptions and/or redescriptions of corresponding species if necessary. Collected specimens were regarded as pests in case these were directly damaging insects or mites on pepino plants, were known as pests of pepino in the native range and introduced regions of pepino other than Japan, or were known as pests of major solanaceous crops such as tomato, eggplant, green pepper, and potato, in Japan, with a reference to The Japanese Society of Applied Entomology and Zoology (2006). All examined specimens are preserved in the Insect Collection (IC) at the Laboratory of Entomology, TUA (LETUA).

## Observation of virus-like diseases and virus detection

We surveyed whether pepino plants showed symptoms of virus infection such as mosaic, mottle, necrosis, or chlorosis. The symptomatic leaves were collected and used for virus detection as follows: Total RNA was extracted from the samples using Trizol reagent (Invitrogen Corp., Carlsbad, CA) according to the manufacturer's instructions. Total RNA was used as a template for first-strand cDNA synthesis by ReverTra Ace - $\alpha$ -<sup>®</sup> kit (TOYOBO Co., Ltd., Osaka, Japan) followed by DNA amplification using TaKaRa Ex Taq<sup>TM</sup> PCR buffer (Takara Bio Inc., Otsu, Japan) with genus-specific or species-specific primers (Table 1). *Alfalfa mosaic virus* (AMV, genus *Alfamovirus*) and *Cucumber mosaic virus* (CMV, genus *Cucumovirus*) had been reported from pepino plants grown in Kanagawa Prefecture, Japan, additionally, *Pepino mosaic virus* (PeMV, genus *Potexvirus*) and two carlavirus (Potato virus H (PVH) named tentatively and Potato virus S (PVS) reclassified) had been detected from symptomatic and asymptomatic pepino plants in abroad, respectively. Even though any potyvirus had not been reported from pepino plants so far, we tried here detecting whether it was occurred or not from our samples. Reverse transcription-polymerase chain reaction (RT-PCR) products were analyzed by electrophoresis in a 2% agarose gel.

**Table 1.**

Primers used for virus detection in this study.

Target virus	Primer	Strand	Sequence (5' to 3')	Expected amplicon size (bp)	Reference
<i>Alfalfa mosaic virus</i> (AMV)	AMV-F2	+	ATCATGAGTTCTTCACAAAAGAA	670	Xu and Nie (2006)
	AMV-R2	-	TCAATGACGATCAAGATCGTC		
<i>Cucumber mosaic virus</i> (CMV)	CPTALL-5	+	YASYTTDRGGTTCAATTCC	950	Choi et al. (1999)
	CPTALL-3	-	GACTGACCATTAGCCG		
Genus <i>Carlavirus</i>	Carla-uni	+	GGAGTAACCGAGGTGATACC	120	Badge et al. (1996) modified
	oligo dT	-	T <sub>18</sub>		
Genus <i>Potexvirus</i>	Potex 5	+	CAYCARCARGCMAARGAYGA	600	van der Vlugt and Berendsen (2002)
	Potex 2RC	-	AGCATRGNCNSRCTCYTG		
Genus <i>Potyvirus</i>	ClFor	+	GGIVVIGTIGGIWSIGGIAARTCIAC	700	Ha et al. (2008)
	ClRev	-	ACICCRTTYTCDATDATRTTIGTIGC		

## Data resources

In this study, a total of 498 specimens of insects and mites were collected from pepino plants on the three studied plots. Of these specimens, 459 individuals belonging to 11 species were recognized as pests of pepino. They consisted of nine insect species belonging to eight families of five orders and two mite species in two families of one order (Table 2). The remaining specimens (39 individuals) were identified as predatory mites attacking other mites (Acari: Phytoseiidae), parasitic wasps of certain insects (Hymenoptera: Braconidae), and incidental visitors to pepino plants, including grasshoppers (Orthoptera: Tettigoniidae), seed bugs (Hemiptera: Lygaeoidea), and butterflies (Lepidoptera: Papilionidae).

**Table 2.**

List of insect and mite pests found on pepino plants in the Atsugi Campus of Tokyo University of Agriculture (TUA), Kanagawa, Japan.

Class	Order	Family	Species	Suvery plots
Insecta	Thysanoptera	Thripidae	<i>Frankliniella intonsa</i> (Trybom, 1895)	A C
Insecta	Hemiptera	Aleyrodidae	<i>Bemisia tabaci</i> (Gennadius, 1889)	B C

Insecta	Hemiptera	Aphididae	<i>Aphis gossypii</i> Glover, 1877	A C
Insecta	Hemiptera	Pseudococcidae	<i>Phenacoccus solani</i> Ferris, 1918	C
Insecta	Hemiptera	Miridae	<i>Campylomma livida</i> Reuter, 1885	A
Insecta	Coleoptera	Chrysomelidae	<i>Epitrix hirtipennis</i> (Melsheimer, 1847)	A C
Insecta	Diptera	Agromyzidae	<i>Liriomyza sativae</i> Blanchard, 1938	C
Insecta	Lepidoptera	Noctuidae	<i>Spodoptera litura</i> (Fabricius, 1775)	A B
Insecta	Lepidoptera	Noctuidae	<i>Trichoplusia ni</i> (Hübner, 1803)	B
Arachnida	Acari	Tarsonemidae	<i>Polyphagotarsonemus latus</i> (Banks, 1904)	C
Arachnida	Acari	Tetranychidae	<i>Tetranychus urticae</i> Koch, 1836	A C

Of these 11 pest species, six were detected from Plot A, three from Plot B, and eight from Plot C (Table 2). The plant bug species *Campylomma livida* Reuter, 1885 was found only in the open field (Plot A). Five species, the cotton whitefly *Bemisia tabaci* (Gennadius, 1889), the mealy bug *Phenacoccus solani* Ferris, 1918, the tomato leaf miner *Liriomyza sativae* Blanchard, 1938, the plusiine noctuid caterpillar *Trichoplusia ni* (Hübner, 1803), and the broad mite *Polyphagotarsonemus latus* (Banks, 1904), were found only in greenhouses (Plots B and C); the former two are well known as glasshouse pests in Japan. No pest species were common in all the three plots. On an empirical basis, through our survey in the plots, two pest species, the two-spotted spider mite *Tetranychus urticae* Koch, 1836 and the cotton aphid *Aphis gossypii* Glover, 1877, were much more abundant on pepino plants than the other pest species recognized, with several hundreds of these two species on each pepino plant.



Figure 5. doi

Virus-like symptoms on pepino plants: necrotic spots on pepino leaves in a greenhouse (Fig. 1, Plot C).

Regarding virus-like diseases in our research fields, pepino leaves showing necrosis were rarely found in the greenhouse (Fig. 5). However, upper leaves showing mottle symptoms with deformation were remarkably observed only in the acaricide-untreated pepinos in the open field (Plot A) during September to October (Fig. 6). Those symptomatic leaves were tested for the detection of *Alfalfa mosaic virus* (AMV) and *Cucumber mosaic virus* (CMV), as well as the genera *Carlavirus*, *Potexvirus*, and *Potyvirus*. All tested pepino leaves showed no infection of the above-mentioned viruses (data not shown).



Figure 6. [doi](#)

Virus-like symptoms on pepino plants: mottle and deformation on young pepino leaves at an open field (Fig. 1, Plot A).

## Checklist of insect and mite pests of pepino in Kanagawa, Japan

### Class Insecta Linnaeus, 1758

### Order Thysanoptera Haliday, 1836

#### Family Thripidae Stevens, 1829

##### *Frankliniella intonsa* (Trybom, 1895)

#### Materials

- a. namePublishedIn: 1895; kingdom: Animalia; phylum: Arthropoda; class: Insecta; order: Thysanoptera; family: Thripidae; genus: *Frankliniella*; specificEpithet: *intonsa*; scientificNameAuthorship: Trybom; country: Japan; stateProvince: Kanagawa; municipality: Atsugi-shi; locality: Atsugi Campus, Tokyo University of Agriculture, Funako; minimumElevationInMeters: 49; maximumElevationInMeters: 49; decimalLatitude:

- 35.431707; decimalLongitude: 139.345165; geodeticDatum: WGS84; samplingProtocol: beating of leaves and branches (including visual searches); eventDate: 2016-09-29; individualCount: 1; lifeStage: adult; recordedBy: Y. Yamada; otherCatalogNumbers: 2017-00005; identifiedBy: Y. Yamada; datelidentified: 2017; institutionCode: LETUA; collectionCode: IC
- b. namePublishedIn: 1895; kingdom: Animalia; phylum: Arthropoda; class: Insecta; order: Thysanoptera; family: Thripidae; genus: *Frankliniella*; specificEpithet: *intonsa*; scientificNameAuthorship: Trybom; country: Japan; stateProvince: Kanagawa; municipality: Atsugi-shi; locality: Atsugi Campus, Tokyo University of Agriculture, Funako; minimumElevationInMeters: 42; maximumElevationInMeters: 42; decimalLatitude: 35.428874; decimalLongitude: 139.34929; geodeticDatum: WGS84; samplingProtocol: beating of leaves and branches (including visual searches); eventDate: 2017-01-07; individualCount: 3; lifeStage: adult; recordedBy: Y. Yamada; otherCatalogNumbers: 2017-00006 | 2017-00007 | 2017-00008; identifiedBy: Y. Yamada; datelidentified: 2017; institutionCode: LETUA; collectionCode: IC
- c. namePublishedIn: 1895; kingdom: Animalia; phylum: Arthropoda; class: Insecta; order: Thysanoptera; family: Thripidae; genus: *Frankliniella*; specificEpithet: *intonsa*; scientificNameAuthorship: Trybom; country: Japan; stateProvince: Kanagawa; municipality: Atsugi-shi; locality: Atsugi Campus, Tokyo University of Agriculture, Funako; minimumElevationInMeters: 42; maximumElevationInMeters: 42; decimalLatitude: 35.428874; decimalLongitude: 139.34929; geodeticDatum: WGS84; samplingProtocol: beating of leaves and branches (including visual searches); eventDate: 2017-01-09; individualCount: 1; lifeStage: adult; recordedBy: Y. Yamada; otherCatalogNumbers: 2017-00009; identifiedBy: Y. Yamada; datelidentified: 2017; institutionCode: LETUA; collectionCode: IC

## Order Hemiptera Linnaeus, 1758

### Family Aleyrodidae Westwood, 1840

#### *Bemisia tabaci* (Gennadius, 1889)

##### Materials

- a. namePublishedIn: 1889; kingdom: Animalia; phylum: Arthropoda; class: Insecta; order: Hemiptera; family: Aleyrodidae; genus: *Bemisia*; specificEpithet: *tabaci*; scientificNameAuthorship: Gennadius; country: Japan; stateProvince: Kanagawa; municipality: Atsugi-shi; locality: Atsugi Campus, Tokyo University of Agriculture, Funako; minimumElevationInMeters: 27; maximumElevationInMeters: 27; decimalLatitude: 35.43043; decimalLongitude: 139.349516; geodeticDatum: WGS84; samplingProtocol: beating of leaves and branches (including visual searches); eventDate: 2016-12-18; individualCount: 1; lifeStage: adult; recordedBy: K.Takahata; otherCatalogNumbers: 2017-00010; identifiedBy: Y. Yamada; datelidentified: 2017; institutionCode: LETUA; collectionCode: IC
- b. namePublishedIn: 1889; kingdom: Animalia; phylum: Arthropoda; class: Insecta; order: Hemiptera; family: Aleyrodidae; genus: *Bemisia*; specificEpithet: *tabaci*; scientificNameAuthorship: Gennadius; country: Japan; stateProvince: Kanagawa; municipality: Atsugi-shi; locality: Atsugi Campus, Tokyo University of Agriculture, Funako; minimumElevationInMeters: 42; maximumElevationInMeters: 42; decimalLatitude:

- 35.428874; decimalLongitude: 139.34929; geodeticDatum: WGS84; samplingProtocol: beating of leaves and branches (including visual searches); eventDate: 2016-12-19; individualCount: 3; lifeStage: adult; recordedBy: Y. Yamada; otherCatalogNumbers: 2017-00011 | 2017-00012 | 2017-00013; identifiedBy: Y. Yamada; datelidentified: 2017; institutionCode: LETUA; collectionCode: IC
- c. namePublishedIn: 1889; kingdom: Animalia; phylum: Arthropoda; class: Insecta; order: Hemiptera; family: Aleyrodidae; genus: *Bemisia*; specificEpithet: *tabaci*; scientificNameAuthorship: Gennadius; country: Japan; stateProvince: Kanagawa; municipality: Atsugi-shi; locality: Atsugi Campus, Tokyo University of Agriculture, Funako; minimumElevationInMeters: 42; maximumElevationInMeters: 42; decimalLatitude: 35.428874; decimalLongitude: 139.34929; geodeticDatum: WGS84; samplingProtocol: beating of leaves and branches (including visual searches); eventDate: 2017-01-09; individualCount: 2; lifeStage: adult; recordedBy: Y. Yamada; otherCatalogNumbers: 2017-00014 | 2017-00015; identifiedBy: Y. Yamada; datelidentified: 2017; institutionCode: LETUA; collectionCode: IC

## Family Aphididae Latreille, 1802

### *Aphis gossypii* Glover, 1877

#### Materials

- a. namePublishedIn: 1877; kingdom: Animalia; phylum: Arthropoda; class: Insecta; order: Hemiptera; family: Aphididae; genus: *Aphis*; specificEpithet: *gossypii*; scientificNameAuthorship: Glover; country: Japan; stateProvince: Kanagawa; municipality: Atsugi-shi; locality: Atsugi Campus, Tokyo University of Agriculture, Funako; minimumElevationInMeters: 49; maximumElevationInMeters: 49; decimalLatitude: 35.431707; decimalLongitude: 139.345165; geodeticDatum: WGS84; samplingProtocol: beating of leaves and branches (including visual searches); eventDate: 2016-08-30; individualCount: 3; lifeStage: adult; recordedBy: T. Ishikawa; otherCatalogNumbers: 2017-00016 | 2017-00017 | 2017-00018; identifiedBy: Y. Yamada; datelidentified: 2017; institutionCode: LETUA; collectionCode: IC
- b. namePublishedIn: 1877; kingdom: Animalia; phylum: Arthropoda; class: Insecta; order: Hemiptera; family: Aphididae; genus: *Aphis*; specificEpithet: *gossypii*; scientificNameAuthorship: Glover; country: Japan; stateProvince: Kanagawa; municipality: Atsugi-shi; locality: Atsugi Campus, Tokyo University of Agriculture, Funako; minimumElevationInMeters: 49; maximumElevationInMeters: 49; decimalLatitude: 35.431707; decimalLongitude: 139.345165; geodeticDatum: WGS84; samplingProtocol: beating of leaves and branches (including visual searches); eventDate: 2016-09-28; individualCount: 3; lifeStage: adult; recordedBy: Y. Yamada; otherCatalogNumbers: 2017-00019 | 2017-00020 | 2017-00021; identifiedBy: Y. Yamada; datelidentified: 2017; institutionCode: LETUA; collectionCode: IC
- c. namePublishedIn: 1877; kingdom: Animalia; phylum: Arthropoda; class: Insecta; order: Hemiptera; family: Aphididae; genus: *Aphis*; specificEpithet: *gossypii*; scientificNameAuthorship: Glover; country: Japan; stateProvince: Kanagawa; municipality: Atsugi-shi; locality: Atsugi Campus, Tokyo University of Agriculture, Funako; minimumElevationInMeters: 49; maximumElevationInMeters: 49; decimalLatitude: 35.431707; decimalLongitude: 139.345165; geodeticDatum: WGS84; samplingProtocol: beating of leaves and branches (including visual searches); eventDate: 2016-10-06; individualCount: 1; lifeStage: adult; recordedBy: Y. Yamada; otherCatalogNumbers:

- 2017-00022; identifiedBy: Y. Yamada; dateIdentified: 2017; institutionCode: LETUA; collectionCode: IC
- d. namePublishedIn: 1877; kingdom: Animalia; phylum: Arthropoda; class: Insecta; order: Hemiptera; family: Aphididae; genus: *Aphis*; specificEpithet: *gossypii*; scientificNameAuthorship: Glover; country: Japan; stateProvince: Kanagawa; municipality: Atsugi-shi; locality: Atsugi Campus, Tokyo University of Agriculture, Funako; minimumElevationInMeters: 49; maximumElevationInMeters: 49; decimalLatitude: 35.431707; decimalLongitude: 139.345165; geodeticDatum: WGS84; samplingProtocol: beating of leaves and branches (including visual searches); eventDate: 2016-11-08; individualCount: 1; lifeStage: adult; recordedBy: Y. Yamada; otherCatalogNumbers: 2017-00023; identifiedBy: Y. Yamada; dateIdentified: 2017; institutionCode: LETUA; collectionCode: IC
- e. namePublishedIn: 1877; kingdom: Animalia; phylum: Arthropoda; class: Insecta; order: Hemiptera; family: Aphididae; genus: *Aphis*; specificEpithet: *gossypii*; scientificNameAuthorship: Glover; country: Japan; stateProvince: Kanagawa; municipality: Atsugi-shi; locality: Atsugi Campus, Tokyo University of Agriculture, Funako; minimumElevationInMeters: 49; maximumElevationInMeters: 49; decimalLatitude: 35.431707; decimalLongitude: 139.345165; geodeticDatum: WGS84; samplingProtocol: beating of leaves and branches (including visual searches); eventDate: 2016-11-10; individualCount: 1; lifeStage: adult; recordedBy: Y. Yamada; otherCatalogNumbers: 2017-00024; identifiedBy: Y. Yamada; dateIdentified: 2017; institutionCode: LETUA; collectionCode: IC
- f. namePublishedIn: 1877; kingdom: Animalia; phylum: Arthropoda; class: Insecta; order: Hemiptera; family: Aphididae; genus: *Aphis*; specificEpithet: *gossypii*; scientificNameAuthorship: Glover; country: Japan; stateProvince: Kanagawa; municipality: Atsugi-shi; locality: Atsugi Campus, Tokyo University of Agriculture, Funako; minimumElevationInMeters: 49; maximumElevationInMeters: 49; decimalLatitude: 35.431707; decimalLongitude: 139.345165; geodeticDatum: WGS84; samplingProtocol: beating of leaves and branches (including visual searches); eventDate: 2016-11-17; individualCount: 1; lifeStage: adult; recordedBy: Y. Yamada; otherCatalogNumbers: 2017-00025; identifiedBy: Y. Yamada; dateIdentified: 2017; institutionCode: LETUA; collectionCode: IC
- g. namePublishedIn: 1877; kingdom: Animalia; phylum: Arthropoda; class: Insecta; order: Hemiptera; family: Aphididae; genus: *Aphis*; specificEpithet: *gossypii*; scientificNameAuthorship: Glover; country: Japan; stateProvince: Kanagawa; municipality: Atsugi-shi; locality: Atsugi Campus, Tokyo University of Agriculture, Funako; minimumElevationInMeters: 49; maximumElevationInMeters: 49; decimalLatitude: 35.431707; decimalLongitude: 139.345165; geodeticDatum: WGS84; samplingProtocol: beating of leaves and branches (including visual searches); eventDate: 2016-11-30; individualCount: 4; lifeStage: adult; recordedBy: Y. Yamada; otherCatalogNumbers: 2017-00026 | 2017-00027 | 2017-00028 | 2017-00029; identifiedBy: Y. Yamada; dateIdentified: 2017; institutionCode: LETUA; collectionCode: IC
- h. namePublishedIn: 1877; kingdom: Animalia; phylum: Arthropoda; class: Insecta; order: Hemiptera; family: Aphididae; genus: *Aphis*; specificEpithet: *gossypii*; scientificNameAuthorship: Glover; country: Japan; stateProvince: Kanagawa; municipality: Atsugi-shi; locality: Atsugi Campus, Tokyo University of Agriculture, Funako; minimumElevationInMeters: 42; maximumElevationInMeters: 42; decimalLatitude: 35.428874; decimalLongitude: 139.34929; geodeticDatum: WGS84; samplingProtocol: beating of leaves and branches (including visual searches); eventDate: 2016-12-26; individualCount: 17; lifeStage: adult; recordedBy: Y. Yamada; otherCatalogNumbers:

- 2017-00030 | 2017-00031 | 2017-00032 | 2017-00033 | 2017-00034 | 2017-00035 |  
2017-00036 | 2017-00037 | 2017-00038 | 2017-00039 | 2017-00040 | 2017-00041 |  
2017-00042 | 2017-00043 | 2017-00044 | 2017-00047 | 2017-00048; identifiedBy: Y.  
Yamada; datelidentified: 2017; institutionCode: LETUA; collectionCode: IC
- i. namePublishedIn: 1877; kingdom: Animalia; phylum: Arthropoda; class: Insecta; order:  
Hemiptera; family: Aphididae; genus: *Aphis*; specificEpithet: *gossypii*;  
scientificNameAuthorship: Glover; country: Japan; stateProvince: Kanagawa; municipality:  
Atsugi-shi; locality: Atsugi Campus, Tokyo University of Agriculture, Funako;  
minimumElevationInMeters: 42; maximumElevationInMeters: 42; decimalLatitude:  
35.428874; decimalLongitude: 139.34929; geodeticDatum: WGS84; samplingProtocol:  
beating of leaves and branches (including visual searches); eventDate: 2016-12-26;  
individualCount: 2; lifeStage: nymph; recordedBy: Y. Yamada; otherCatalogNumbers:  
2017-00045 | 2017-00046; identifiedBy: Y. Yamada; datelidentified: 2017; institutionCode:  
LETUA; collectionCode: IC
- j. namePublishedIn: 1877; kingdom: Animalia; phylum: Arthropoda; class: Insecta; order:  
Hemiptera; family: Aphididae; genus: *Aphis*; specificEpithet: *gossypii*;  
scientificNameAuthorship: Glover; country: Japan; stateProvince: Kanagawa; municipality:  
Atsugi-shi; locality: Atsugi Campus, Tokyo University of Agriculture, Funako;  
minimumElevationInMeters: 42; maximumElevationInMeters: 42; decimalLatitude:  
35.428874; decimalLongitude: 139.34929; geodeticDatum: WGS84; samplingProtocol:  
beating of leaves and branches (including visual searches); eventDate: 2016-12-30;  
individualCount: 30; lifeStage: adult; recordedBy: Y. Yamada; otherCatalogNumbers:  
2017-00049 | 2017-00050 | 2017-00051 | 2017-00052 | 2017-00053 | 2017-00054 |  
2017-00055 | 2017-00056 | 2017-00057 | 2017-00058 | 2017-00059 | 2017-00060 |  
2017-00061 | 2017-00062 | 2017-00063 | 2017-00064 | 2017-00065 | 2017-00066 |  
2017-00067 | 2017-00068 | 2017-00069 | 2017-00070 | 2017-00071 | 2017-00072 |  
2017-00073 | 2017-00074 | 2017-00075 | 2017-00078 | 2017-00079 | 2017-00080;  
identifiedBy: Y. Yamada; datelidentified: 2017; institutionCode: LETUA; collectionCode: IC
- k. namePublishedIn: 1877; kingdom: Animalia; phylum: Arthropoda; class: Insecta; order:  
Hemiptera; family: Aphididae; genus: *Aphis*; specificEpithet: *gossypii*;  
scientificNameAuthorship: Glover; country: Japan; stateProvince: Kanagawa; municipality:  
Atsugi-shi; locality: Atsugi Campus, Tokyo University of Agriculture, Funako;  
minimumElevationInMeters: 42; maximumElevationInMeters: 42; decimalLatitude:  
35.428874; decimalLongitude: 139.34929; geodeticDatum: WGS84; samplingProtocol:  
beating of leaves and branches (including visual searches); eventDate: 2016-12-30;  
individualCount: 2; lifeStage: nymph; recordedBy: Y. Yamada; otherCatalogNumbers:  
2017-00076 | 2017-00077; identifiedBy: Y. Yamada; datelidentified: 2017; institutionCode:  
LETUA; collectionCode: IC
- l. namePublishedIn: 1877; kingdom: Animalia; phylum: Arthropoda; class: Insecta; order:  
Hemiptera; family: Aphididae; genus: *Aphis*; specificEpithet: *gossypii*;  
scientificNameAuthorship: Glover; country: Japan; stateProvince: Kanagawa; municipality:  
Atsugi-shi; locality: Atsugi Campus, Tokyo University of Agriculture, Funako;  
minimumElevationInMeters: 42; maximumElevationInMeters: 42; decimalLatitude:  
35.428874; decimalLongitude: 139.34929; geodeticDatum: WGS84; samplingProtocol:  
beating of leaves and branches (including visual searches); eventDate: 2017-01-03;  
individualCount: 48; lifeStage: adult; recordedBy: Y. Yamada; otherCatalogNumbers:  
2017-00081 | 2017-00082 | 2017-00083 | 2017-00084 | 2017-00085 | 2017-00086 |  
2017-00087 | 2017-00088 | 2017-00089 | 2017-00090 | 2017-00091 | 2017-00092 |  
2017-00093 | 2017-00094 | 2017-00095 | 2017-00096 | 2017-00097 | 2017-00098 |  
2017-00099 | 2017-00100 | 2017-00101 | 2017-00102 | 2017-00103 | 2017-00104 |

- 2017-00105 | 2017-00106 | 2017-00107 | 2017-00108 | 2017-00109 | 2017-00110 |  
2017-00111 | 2017-00112 | 2017-00113 | 2017-00114 | 2017-00115 | 2017-00116 |  
2017-00117 | 2017-00118 | 2017-00119 | 2017-00120 | 2017-00121 | 2017-00122 |  
2017-00131 | 2017-00132 | 2017-00133 | 2017-00134 | 2017-00135 | 2017-00136;  
identifiedBy: Y. Yamada; datelidentified: 2017; institutionCode: LETUA; collectionCode: IC  
m. namePublishedIn: 1877; kingdom: Animalia; phylum: Arthropoda; class: Insecta; order:  
Hemiptera; family: Aphididae; genus: *Aphis*; specificEpithet: *gossypii*;  
scientificNameAuthorship: Glover; country: Japan; stateProvince: Kanagawa; municipality:  
Atsugi-shi; locality: Atsugi Campus, Tokyo University of Agriculture, Funako;  
minimumElevationInMeters: 42; maximumElevationInMeters: 42; decimalLatitude:  
35.428874; decimalLongitude: 139.34929; geodeticDatum: WGS84; samplingProtocol:  
beating of leaves and branches (including visual searches); eventDate: 2017-01-03;  
individualCount: 8; lifeStage: nymph; recordedBy: Y. Yamada; otherCatalogNumbers:  
2017-00123 | 2017-00124 | 2017-00125 | 2017-00126 | 2017-00127 | 2017-00128 |  
2017-00129 | 2017-00130; identifiedBy: Y. Yamada; datelidentified: 2017; institutionCode:  
LETUA; collectionCode: IC  
n. namePublishedIn: 1877; kingdom: Animalia; phylum: Arthropoda; class: Insecta; order:  
Hemiptera; family: Aphididae; genus: *Aphis*; specificEpithet: *gossypii*;  
scientificNameAuthorship: Glover; country: Japan; stateProvince: Kanagawa; municipality:  
Atsugi-shi; locality: Atsugi Campus, Tokyo University of Agriculture, Funako;  
minimumElevationInMeters: 42; maximumElevationInMeters: 42; decimalLatitude:  
35.428874; decimalLongitude: 139.34929; geodeticDatum: WGS84; samplingProtocol:  
beating of leaves and branches (including visual searches); eventDate: 2017-01-05;  
individualCount: 36; lifeStage: adult; recordedBy: Y. Yamada; otherCatalogNumbers:  
2017-00137 | 2017-00138 | 2017-00139 | 2017-00140 | 2017-00141 | 2017-00142 |  
2017-00143 | 2017-00144 | 2017-00145 | 2017-00146 | 2017-00147 | 2017-00148 |  
2017-00149 | 2017-00150 | 2017-00151 | 2017-00152 | 2017-00153 | 2017-00154 |  
2017-00155 | 2017-00156 | 2017-00157 | 2017-00158 | 2017-00159 | 2017-00160 |  
2017-00161 | 2017-00162 | 2017-00163 | 2017-00164 | 2017-00165 | 2017-00166 |  
2017-00167 | 2017-00168 | 2017-00169 | 2017-00170 | 2017-00171 | 2017-00172;  
identifiedBy: Y. Yamada; datelidentified: 2017; institutionCode: LETUA; collectionCode: IC  
o. namePublishedIn: 1877; kingdom: Animalia; phylum: Arthropoda; class: Insecta; order:  
Hemiptera; family: Aphididae; genus: *Aphis*; specificEpithet: *gossypii*;  
scientificNameAuthorship: Glover; country: Japan; stateProvince: Kanagawa; municipality:  
Atsugi-shi; locality: Atsugi Campus, Tokyo University of Agriculture, Funako;  
minimumElevationInMeters: 42; maximumElevationInMeters: 42; decimalLatitude:  
35.428874; decimalLongitude: 139.34929; geodeticDatum: WGS84; samplingProtocol:  
beating of leaves and branches (including visual searches); eventDate: 2017-01-07;  
individualCount: 25; lifeStage: adult; recordedBy: Y. Yamada; otherCatalogNumbers:  
2017-00173 | 2017-00174 | 2017-00175 | 2017-00176 | 2017-00177 | 2017-00178 |  
2017-00179 | 2017-00180 | 2017-00181 | 2017-00182 | 2017-00183 | 2017-00184 |  
2017-00185 | 2017-00186 | 2017-00187 | 2017-00188 | 2017-00189 | 2017-00190 |  
2017-00191 | 2017-00192 | 2017-00193 | 2017-00194 | 2017-00195 | 2017-00196 |  
2017-00197; identifiedBy: Y. Yamada; datelidentified: 2017; institutionCode: LETUA;  
collectionCode: IC  
p. namePublishedIn: 1877; kingdom: Animalia; phylum: Arthropoda; class: Insecta; order:  
Hemiptera; family: Aphididae; genus: *Aphis*; specificEpithet: *gossypii*;  
scientificNameAuthorship: Glover; country: Japan; stateProvince: Kanagawa; municipality:  
Atsugi-shi; locality: Atsugi Campus, Tokyo University of Agriculture, Funako;  
minimumElevationInMeters: 42; maximumElevationInMeters: 42; decimalLatitude:

- 35.428874; decimalLongitude: 139.34929; geodeticDatum: WGS84; samplingProtocol: beating of leaves and branches (including visual searches); eventDate: 2017-01-09; individualCount: 4; lifeStage: adult; recordedBy: Y. Yamada; otherCatalogNumbers: 2017-00198 | 2017-00199 | 2017-00200 | 2017-00201; identifiedBy: Y. Yamada; dateIdentified: 2017; institutionCode: LETUA; collectionCode: IC
- q. namePublishedIn: 1877; kingdom: Animalia; phylum: Arthropoda; class: Insecta; order: Hemiptera; family: Aphididae; genus: *Aphis*; specificEpithet: *gossypii*; scientificNameAuthorship: Glover; country: Japan; stateProvince: Kanagawa; municipality: Atsugi-shi; locality: Atsugi Campus, Tokyo University of Agriculture, Funako; minimumElevationInMeters: 42; maximumElevationInMeters: 42; decimalLatitude: 35.428874; decimalLongitude: 139.34929; geodeticDatum: WGS84; samplingProtocol: beating of leaves and branches (including visual searches); eventDate: 2017-01-12; individualCount: 43; lifeStage: adult; recordedBy: Y. Yamada; otherCatalogNumbers: 2017-00202 | 2017-00203 | 2017-00204 | 2017-00205 | 2017-00206 | 2017-00207 | 2017-00208 | 2017-00209 | 2017-00210 | 2017-00211 | 2017-00212 | 2017-00213 | 2017-00214 | 2017-00215 | 2017-00216 | 2017-00217 | 2017-00218 | 2017-00219 | 2017-00220 | 2017-00221 | 2017-00222 | 2017-00223 | 2017-00224 | 2017-00225 | 2017-00226 | 2017-00227 | 2017-00228 | 2017-00229 | 2017-00230 | 2017-00231 | 2017-00232 | 2017-00233 | 2017-00234 | 2017-00235 | 2017-00236 | 2017-00237 | 2017-00238 | 2017-00239 | 2017-00240 | 2017-00241 | 2017-00242 | 2017-00243 | 2017-00244; identifiedBy: Y. Yamada; dateIdentified: 2017; institutionCode: LETUA; collectionCode: IC
- r. namePublishedIn: 1877; kingdom: Animalia; phylum: Arthropoda; class: Insecta; order: Hemiptera; family: Aphididae; genus: *Aphis*; specificEpithet: *gossypii*; scientificNameAuthorship: Glover; country: Japan; stateProvince: Kanagawa; municipality: Atsugi-shi; locality: Atsugi Campus, Tokyo University of Agriculture, Funako; minimumElevationInMeters: 42; maximumElevationInMeters: 42; decimalLatitude: 35.428874; decimalLongitude: 139.34929; geodeticDatum: WGS84; samplingProtocol: beating of leaves and branches (including visual searches); eventDate: 2017-01-16; individualCount: 22; lifeStage: adult; recordedBy: Y. Yamada; otherCatalogNumbers: 2017-00245 | 2017-00246 | 2017-00247 | 2017-00248 | 2017-00249 | 2017-00250 | 2017-00251 | 2017-00252 | 2017-00253 | 2017-00254 | 2017-00255 | 2017-00256 | 2017-00257 | 2017-00258 | 2017-00259 | 2017-00260 | 2017-00261 | 2017-00262 | 2017-00263 | 2017-00264 | 2017-00265 | 2017-00266; identifiedBy: Y. Yamada; dateIdentified: 2017; institutionCode: LETUA; collectionCode: IC
- s. namePublishedIn: 1877; kingdom: Animalia; phylum: Arthropoda; class: Insecta; order: Hemiptera; family: Aphididae; genus: *Aphis*; specificEpithet: *gossypii*; scientificNameAuthorship: Glover; country: Japan; stateProvince: Kanagawa; municipality: Atsugi-shi; locality: Atsugi Campus, Tokyo University of Agriculture, Funako; minimumElevationInMeters: 42; maximumElevationInMeters: 42; decimalLatitude: 35.428874; decimalLongitude: 139.34929; geodeticDatum: WGS84; samplingProtocol: beating of leaves and branches (including visual searches); eventDate: 2017-01-21; individualCount: 3; lifeStage: adult; recordedBy: Y. Yamada; otherCatalogNumbers: 2017-00267 | 2017-00268 | 2017-00269; identifiedBy: Y. Yamada; dateIdentified: 2017; institutionCode: LETUA; collectionCode: IC

## Family Pseudococcidae Heymons, 1915

### *Phenacoccus solani* Ferris, 1918

#### Materials

- a. namePublishedIn: 1918; kingdom: Animalia; phylum: Arthropoda; class: Insecta; order: Hemiptera; family: Pseudococcidae; genus: *Phenacoccus*; specificEpithet: *solani*; scientificNameAuthorship: Ferris; country: Japan; stateProvince: Kanagawa; municipality: Atsugi-shi; locality: Atsugi Campus, Tokyo University of Agriculture, Funako; minimumElevationInMeters: 42; maximumElevationInMeters: 42; decimalLatitude: 35.428874; decimalLongitude: 139.34929; geodeticDatum: WGS84; samplingProtocol: beating of leaves and branches (including visual searches); eventDate: 2016-12-30; individualCount: 1; lifeStage: nymph; recordedBy: Y. Yamada; otherCatalogNumbers: 2017-00270; identifiedBy: Y. Yamada; dateIdentified: 2017; institutionCode: LETUA; collectionCode: IC
- b. namePublishedIn: 1918; kingdom: Animalia; phylum: Arthropoda; class: Insecta; order: Hemiptera; family: Pseudococcidae; genus: *Phenacoccus*; specificEpithet: *solani*; scientificNameAuthorship: Ferris; country: Japan; stateProvince: Kanagawa; municipality: Atsugi-shi; locality: Atsugi Campus, Tokyo University of Agriculture, Funako; minimumElevationInMeters: 42; maximumElevationInMeters: 42; decimalLatitude: 35.428874; decimalLongitude: 139.34929; geodeticDatum: WGS84; samplingProtocol: beating of leaves and branches (including visual searches); eventDate: 2017-01-15; individualCount: 2; lifeStage: adult; recordedBy: Y. Yamada; otherCatalogNumbers: 2017-00271 | 2017-00272; identifiedBy: Y. Yamada; dateIdentified: 2017; institutionCode: LETUA; collectionCode: IC

## Family Miridae Hahn, 1833

### *Campylomma livida* Reuter, 1885

#### Materials

- a. namePublishedIn: 1885; kingdom: Animalia; phylum: Arthropoda; class: Insecta; order: Hemiptera; family: Miridae; genus: *Campylomma*; specificEpithet: *livida*; scientificNameAuthorship: Reuter; country: Japan; stateProvince: Kanagawa; municipality: Atsugi-shi; locality: Atsugi Campus, Tokyo University of Agriculture, Funako; minimumElevationInMeters: 49; maximumElevationInMeters: 49; decimalLatitude: 35.431707; decimalLongitude: 139.345165; geodeticDatum: WGS84; samplingProtocol: beating of leaves and branches (including visual searches); eventDate: 2016-11-10; individualCount: 1; lifeStage: nymph; recordedBy: Y. Yamada; otherCatalogNumbers: 2017-00273; identifiedBy: Y. Yamada; dateIdentified: 2017; institutionCode: LETUA; collectionCode: IC
- b. namePublishedIn: 1885; kingdom: Animalia; phylum: Arthropoda; class: Insecta; order: Hemiptera; family: Miridae; genus: *Campylomma*; specificEpithet: *livida*; scientificNameAuthorship: Reuter; country: Japan; stateProvince: Kanagawa; municipality: Atsugi-shi; locality: Atsugi Campus, Tokyo University of Agriculture, Funako; minimumElevationInMeters: 49; maximumElevationInMeters: 49; decimalLatitude: 35.431707; decimalLongitude: 139.345165; geodeticDatum: WGS84; samplingProtocol:

- beating of leaves and branches (including visual searches); eventDate: 2016-11-10; individualCount: 1; lifeStage: adult; recordedBy: Y. Yamada; otherCatalogNumbers: 2017-00274; identifiedBy: Y. Yamada; datelidentified: 2017; institutionCode: LETUA; collectionCode: IC
- c. namePublishedIn: 1885; kingdom: Animalia; phylum: Arthropoda; class: Insecta; order: Hemiptera; family: Miridae; genus: *Campylomma*; specificEpithet: *lividia*; scientificNameAuthorship: Reuter; country: Japan; stateProvince: Kanagawa; municipality: Atsugi-shi; locality: Atsugi Campus, Tokyo University of Agriculture, Funako; minimumElevationInMeters: 49; maximumElevationInMeters: 49; decimalLatitude: 35.431707; decimalLongitude: 139.345165; geodeticDatum: WGS84; samplingProtocol: beating of leaves and branches (including visual searches); eventDate: 2016-11-15; individualCount: 1; lifeStage: adult; recordedBy: Y. Yamada; otherCatalogNumbers: 2017-00275; identifiedBy: Y. Yamada; datelidentified: 2017; institutionCode: LETUA; collectionCode: IC
- d. namePublishedIn: 1885; kingdom: Animalia; phylum: Arthropoda; class: Insecta; order: Hemiptera; family: Miridae; genus: *Campylomma*; specificEpithet: *lividia*; scientificNameAuthorship: Reuter; country: Japan; stateProvince: Kanagawa; municipality: Atsugi-shi; locality: Atsugi Campus, Tokyo University of Agriculture, Funako; minimumElevationInMeters: 49; maximumElevationInMeters: 49; decimalLatitude: 35.431707; decimalLongitude: 139.345165; geodeticDatum: WGS84; samplingProtocol: beating of leaves and branches (including visual searches); eventDate: 2016-11-17; individualCount: 1; lifeStage: adult; recordedBy: Y. Yamada; otherCatalogNumbers: 2017-00276; identifiedBy: Y. Yamada; datelidentified: 2017; institutionCode: LETUA; collectionCode: IC
- e. namePublishedIn: 1885; kingdom: Animalia; phylum: Arthropoda; class: Insecta; order: Hemiptera; family: Miridae; genus: *Campylomma*; specificEpithet: *lividia*; scientificNameAuthorship: Reuter; country: Japan; stateProvince: Kanagawa; municipality: Atsugi-shi; locality: Atsugi Campus, Tokyo University of Agriculture, Funako; minimumElevationInMeters: 49; maximumElevationInMeters: 49; decimalLatitude: 35.431707; decimalLongitude: 139.345165; geodeticDatum: WGS84; samplingProtocol: beating of leaves and branches (including visual searches); eventDate: 2016-11-17; individualCount: 1; lifeStage: nymph; recordedBy: Y. Yamada; otherCatalogNumbers: 2017-00277; identifiedBy: Y. Yamada; datelidentified: 2017; institutionCode: LETUA; collectionCode: IC
- f. namePublishedIn: 1885; kingdom: Animalia; phylum: Arthropoda; class: Insecta; order: Hemiptera; family: Miridae; genus: *Campylomma*; specificEpithet: *lividia*; scientificNameAuthorship: Reuter; country: Japan; stateProvince: Kanagawa; municipality: Atsugi-shi; locality: Atsugi Campus, Tokyo University of Agriculture, Funako; minimumElevationInMeters: 49; maximumElevationInMeters: 49; decimalLatitude: 35.431707; decimalLongitude: 139.345165; geodeticDatum: WGS84; samplingProtocol: beating of leaves and branches (including visual searches); eventDate: 2016-11-23; individualCount: 1; lifeStage: nymph; recordedBy: Y. Yamada; otherCatalogNumbers: 2017-00278; identifiedBy: Y. Yamada; datelidentified: 2017; institutionCode: LETUA; collectionCode: IC
- g. namePublishedIn: 1885; kingdom: Animalia; phylum: Arthropoda; class: Insecta; order: Hemiptera; family: Miridae; genus: *Campylomma*; specificEpithet: *lividia*; scientificNameAuthorship: Reuter; country: Japan; stateProvince: Kanagawa; municipality: Atsugi-shi; locality: Atsugi Campus, Tokyo University of Agriculture, Funako; minimumElevationInMeters: 49; maximumElevationInMeters: 49; decimalLatitude: 35.431707; decimalLongitude: 139.345165; geodeticDatum: WGS84; samplingProtocol:

- beating of leaves and branches (including visual searches); eventDate: 2016-11-23; individualCount: 1; lifeStage: adult; recordedBy: Y. Yamada; otherCatalogNumbers: 2017-00279; identifiedBy: Y. Yamada; dateIdentified: 2017; institutionCode: LETUA; collectionCode: IC
- h. namePublishedIn: 1885; kingdom: Animalia; phylum: Arthropoda; class: Insecta; order: Hemiptera; family: Miridae; genus: *Campylomma*; specificEpithet: *lividia*; scientificNameAuthorship: Reuter; country: Japan; stateProvince: Kanagawa; municipality: Atsugi-shi; locality: Atsugi Campus, Tokyo University of Agriculture, Funako; minimumElevationInMeters: 49; maximumElevationInMeters: 49; decimalLatitude: 35.431707; decimalLongitude: 139.345165; geodeticDatum: WGS84; samplingProtocol: beating of leaves and branches (including visual searches); eventDate: 2016-11-25; individualCount: 1; lifeStage: adult; recordedBy: Y. Yamada; otherCatalogNumbers: 2017-00280; identifiedBy: Y. Yamada; dateIdentified: 2017; institutionCode: LETUA; collectionCode: IC
  - i. namePublishedIn: 1885; kingdom: Animalia; phylum: Arthropoda; class: Insecta; order: Hemiptera; family: Miridae; genus: *Campylomma*; specificEpithet: *lividia*; scientificNameAuthorship: Reuter; country: Japan; stateProvince: Kanagawa; municipality: Atsugi-shi; locality: Atsugi Campus, Tokyo University of Agriculture, Funako; minimumElevationInMeters: 49; maximumElevationInMeters: 49; decimalLatitude: 35.431707; decimalLongitude: 139.345165; geodeticDatum: WGS84; samplingProtocol: beating of leaves and branches (including visual searches); eventDate: 2016-11-30; individualCount: 1; lifeStage: adult; recordedBy: Y. Yamada; otherCatalogNumbers: 2017-00281; identifiedBy: Y. Yamada; dateIdentified: 2017; institutionCode: LETUA; collectionCode: IC
  - j. namePublishedIn: 1885; kingdom: Animalia; phylum: Arthropoda; class: Insecta; order: Hemiptera; family: Miridae; genus: *Campylomma*; specificEpithet: *lividia*; scientificNameAuthorship: Reuter; country: Japan; stateProvince: Kanagawa; municipality: Atsugi-shi; locality: Atsugi Campus, Tokyo University of Agriculture, Funako; minimumElevationInMeters: 49; maximumElevationInMeters: 49; decimalLatitude: 35.431707; decimalLongitude: 139.345165; geodeticDatum: WGS84; samplingProtocol: beating of leaves and branches (including visual searches); eventDate: 2016-11-30; individualCount: 2; lifeStage: nymph; recordedBy: Y. Yamada; otherCatalogNumbers: 2017-00282 | 2017-00283; identifiedBy: Y. Yamada; dateIdentified: 2017; institutionCode: LETUA; collectionCode: IC

## Order Coleoptera Linnaeus, 1758

### Family Chrysomelidae Latreille, 1802

#### *Epitrix hirtipennis* (Melsheimer, 1847)

##### Materials

- a. namePublishedIn: 1847; kingdom: Animalia; phylum: Arthropoda; class: Insecta; order: Coleoptera; family: Chrysomelidae; genus: *Epitrix*; specificEpithet: *hirtipennis*; scientificNameAuthorship: Melsheimer; country: Japan; stateProvince: Kanagawa; municipality: Atsugi-shi; locality: Atsugi Campus, Tokyo University of Agriculture, Funako; minimumElevationInMeters: 49; maximumElevationInMeters: 49; decimalLatitude: 35.431707; decimalLongitude: 139.345165; geodeticDatum: WGS84; samplingProtocol:

- beating of leaves and branches (including visual searches); eventDate: 2016-08-30; individualCount: 12; lifeStage: adult; recordedBy: T. Ishikawa; otherCatalogNumbers: 2017-00284 | 2017-00285 | 2017-00286 | 2017-00287 | 2017-00288 | 2017-00289 | 2017-00290 | 2017-00291 | 2017-00292 | 2017-00293 | 2017-00294 | 2017-00295; identifiedBy: Y. Yamada; datelidentified: 2017; institutionCode: LETUA; collectionCode: IC
- b. namePublishedIn: 1847; kingdom: Animalia; phylum: Arthropoda; class: Insecta; order: Coleoptera; family: Chrysomelidae; genus: *Epitrix*; specificEpithet: *hirtipennis*; scientificNameAuthorship: Melsheimer; country: Japan; stateProvince: Kanagawa; municipality: Atsugi-shi; locality: Atsugi Campus, Tokyo University of Agriculture, Funako; minimumElevationInMeters: 49; maximumElevationInMeters: 49; decimalLatitude: 35.431707; decimalLongitude: 139.345165; geodeticDatum: WGS84; samplingProtocol: beating of leaves and branches (including visual searches); eventDate: 2016-09-28; individualCount: 4; lifeStage: adult; recordedBy: Y. Yamada; otherCatalogNumbers: 2017-00296 | 2017-00297 | 2017-00298 | 2017-00299; identifiedBy: Y. Yamada; datelidentified: 2017; institutionCode: LETUA; collectionCode: IC
- c. namePublishedIn: 1847; kingdom: Animalia; phylum: Arthropoda; class: Insecta; order: Coleoptera; family: Chrysomelidae; genus: *Epitrix*; specificEpithet: *hirtipennis*; scientificNameAuthorship: Melsheimer; country: Japan; stateProvince: Kanagawa; municipality: Atsugi-shi; locality: Atsugi Campus, Tokyo University of Agriculture, Funako; minimumElevationInMeters: 49; maximumElevationInMeters: 49; decimalLatitude: 35.431707; decimalLongitude: 139.345165; geodeticDatum: WGS84; samplingProtocol: beating of leaves and branches (including visual searches); eventDate: 2016-09-29; individualCount: 5; lifeStage: adult; recordedBy: Y. Yamada; otherCatalogNumbers: 2017-00300 | 2017-00301 | 2017-00302 | 2017-00303 | 2017-00304; identifiedBy: Y. Yamada; datelidentified: 2017; institutionCode: LETUA; collectionCode: IC
- d. namePublishedIn: 1847; kingdom: Animalia; phylum: Arthropoda; class: Insecta; order: Coleoptera; family: Chrysomelidae; genus: *Epitrix*; specificEpithet: *hirtipennis*; scientificNameAuthorship: Melsheimer; country: Japan; stateProvince: Kanagawa; municipality: Atsugi-shi; locality: Atsugi Campus, Tokyo University of Agriculture, Funako; minimumElevationInMeters: 49; maximumElevationInMeters: 49; decimalLatitude: 35.431707; decimalLongitude: 139.345165; geodeticDatum: WGS84; samplingProtocol: beating of leaves and branches (including visual searches); eventDate: 2016-09-30; individualCount: 8; lifeStage: adult; recordedBy: Y. Yamada; otherCatalogNumbers: 2017-00305 | 2017-00306 | 2017-00307 | 2017-00308 | 2017-00309 | 2017-00310 | 2017-00311 | 2017-00312; identifiedBy: Y. Yamada; datelidentified: 2017; institutionCode: LETUA; collectionCode: IC
- e. namePublishedIn: 1847; kingdom: Animalia; phylum: Arthropoda; class: Insecta; order: Coleoptera; family: Chrysomelidae; genus: *Epitrix*; specificEpithet: *hirtipennis*; scientificNameAuthorship: Melsheimer; country: Japan; stateProvince: Kanagawa; municipality: Atsugi-shi; locality: Atsugi Campus, Tokyo University of Agriculture, Funako; minimumElevationInMeters: 49; maximumElevationInMeters: 49; decimalLatitude: 35.431707; decimalLongitude: 139.345165; geodeticDatum: WGS84; samplingProtocol: beating of leaves and branches (including visual searches); eventDate: 2016-10-01; individualCount: 7; lifeStage: adult; recordedBy: Y. Yamada; otherCatalogNumbers: 2017-00313 | 2017-00314 | 2017-00315 | 2017-00316 | 2017-00317 | 2017-00318 | 2017-00319; identifiedBy: Y. Yamada; datelidentified: 2017; institutionCode: LETUA; collectionCode: IC
- f. namePublishedIn: 1847; kingdom: Animalia; phylum: Arthropoda; class: Insecta; order: Coleoptera; family: Chrysomelidae; genus: *Epitrix*; specificEpithet: *hirtipennis*; scientificNameAuthorship: Melsheimer; country: Japan; stateProvince: Kanagawa;

- municipality: Atsugi-shi; locality: Atsugi Campus, Tokyo University of Agriculture, Funako; minimumElevationInMeters: 49; maximumElevationInMeters: 49; decimalLatitude: 35.431707; decimalLongitude: 139.345165; geodeticDatum: WGS84; samplingProtocol: beating of leaves and branches (including visual searches); eventDate: 2016-10-04; individualCount: 4; lifeStage: adult; recordedBy: Y. Yamada; otherCatalogNumbers: 2017-00320 | 2017-00321 | 2017-00322 | 2017-00323; identifiedBy: Y. Yamada; datelIdentified: 2017; institutionCode: LETUA; collectionCode: IC
- g. namePublishedIn: 1847; kingdom: Animalia; phylum: Arthropoda; class: Insecta; order: Coleoptera; family: Chrysomelidae; genus: *Epitrix*; specificEpithet: *hirtipennis*; scientificNameAuthorship: Melsheimer; country: Japan; stateProvince: Kanagawa; municipality: Atsugi-shi; locality: Atsugi Campus, Tokyo University of Agriculture, Funako; minimumElevationInMeters: 49; maximumElevationInMeters: 49; decimalLatitude: 35.431707; decimalLongitude: 139.345165; geodeticDatum: WGS84; samplingProtocol: beating of leaves and branches (including visual searches); eventDate: 2016-10-14; individualCount: 4; lifeStage: adult; recordedBy: Y. Yamada; otherCatalogNumbers: 2017-00324 | 2017-00325 | 2017-00326 | 2017-00327; identifiedBy: Y. Yamada; datelIdentified: 2017; institutionCode: LETUA; collectionCode: IC
- h. namePublishedIn: 1847; kingdom: Animalia; phylum: Arthropoda; class: Insecta; order: Coleoptera; family: Chrysomelidae; genus: *Epitrix*; specificEpithet: *hirtipennis*; scientificNameAuthorship: Melsheimer; country: Japan; stateProvince: Kanagawa; municipality: Atsugi-shi; locality: Atsugi Campus, Tokyo University of Agriculture, Funako; minimumElevationInMeters: 49; maximumElevationInMeters: 49; decimalLatitude: 35.431707; decimalLongitude: 139.345165; geodeticDatum: WGS84; samplingProtocol: beating of leaves and branches (including visual searches); eventDate: 2016-10-15; individualCount: 4; lifeStage: adult; recordedBy: Y. Yamada; otherCatalogNumbers: 2017-00328 | 2017-00329 | 2017-00330 | 2017-00331; identifiedBy: Y. Yamada; datelIdentified: 2017; institutionCode: LETUA; collectionCode: IC
- i. namePublishedIn: 1847; kingdom: Animalia; phylum: Arthropoda; class: Insecta; order: Coleoptera; family: Chrysomelidae; genus: *Epitrix*; specificEpithet: *hirtipennis*; scientificNameAuthorship: Melsheimer; country: Japan; stateProvince: Kanagawa; municipality: Atsugi-shi; locality: Atsugi Campus, Tokyo University of Agriculture, Funako; minimumElevationInMeters: 49; maximumElevationInMeters: 49; decimalLatitude: 35.431707; decimalLongitude: 139.345165; geodeticDatum: WGS84; samplingProtocol: beating of leaves and branches (including visual searches); eventDate: 2016-10-18; individualCount: 8; lifeStage: adult; recordedBy: Y. Yamada; otherCatalogNumbers: 2017-00332 | 2017-00333 | 2017-00334 | 2017-00335 | 2017-00336 | 2017-00337 | 2017-00338 | 2017-00339; identifiedBy: Y. Yamada; datelIdentified: 2017; institutionCode: LETUA; collectionCode: IC
- j. namePublishedIn: 1847; kingdom: Animalia; phylum: Arthropoda; class: Insecta; order: Coleoptera; family: Chrysomelidae; genus: *Epitrix*; specificEpithet: *hirtipennis*; scientificNameAuthorship: Melsheimer; country: Japan; stateProvince: Kanagawa; municipality: Atsugi-shi; locality: Atsugi Campus, Tokyo University of Agriculture, Funako; minimumElevationInMeters: 49; maximumElevationInMeters: 49; decimalLatitude: 35.431707; decimalLongitude: 139.345165; geodeticDatum: WGS84; samplingProtocol: beating of leaves and branches (including visual searches); eventDate: 2016-10-20; individualCount: 1; lifeStage: adult; recordedBy: Y. Yamada; otherCatalogNumbers: 2017-00340; identifiedBy: Y. Yamada; datelIdentified: 2017; institutionCode: LETUA; collectionCode: IC
- k. namePublishedIn: 1847; kingdom: Animalia; phylum: Arthropoda; class: Insecta; order: Coleoptera; family: Chrysomelidae; genus: *Epitrix*; specificEpithet: *hirtipennis*;

- scientificNameAuthorship: Melsheimer; country: Japan; stateProvince: Kanagawa; municipality: Atsugi-shi; locality: Atsugi Campus, Tokyo University of Agriculture, Funako; minimumElevationInMeters: 49; maximumElevationInMeters: 49; decimalLatitude: 35.431707; decimalLongitude: 139.345165; geodeticDatum: WGS84; samplingProtocol: beating of leaves and branches (including visual searches); eventDate: 2016-10-24; individualCount: 1; lifeStage: adult; recordedBy: Y. Yamada; otherCatalogNumbers: 2017-00341; identifiedBy: Y. Yamada; datelIdentified: 2017; institutionCode: LETUA; collectionCode: IC
- l. namePublishedIn: 1847; kingdom: Animalia; phylum: Arthropoda; class: Insecta; order: Coleoptera; family: Chrysomelidae; genus: *Epitrix*; specificEpithet: *hirtipennis*; scientificNameAuthorship: Melsheimer; country: Japan; stateProvince: Kanagawa; municipality: Atsugi-shi; locality: Atsugi Campus, Tokyo University of Agriculture, Funako; minimumElevationInMeters: 49; maximumElevationInMeters: 49; decimalLatitude: 35.431707; decimalLongitude: 139.345165; geodeticDatum: WGS84; samplingProtocol: beating of leaves and branches (including visual searches); eventDate: 2016-11-03; individualCount: 4; lifeStage: adult; recordedBy: Y. Yamada; otherCatalogNumbers: 2017-00342 | 2017-00343 | 2017-00344 | 2017-00345; identifiedBy: Y. Yamada; datelIdentified: 2017; institutionCode: LETUA; collectionCode: IC
- m. namePublishedIn: 1847; kingdom: Animalia; phylum: Arthropoda; class: Insecta; order: Coleoptera; family: Chrysomelidae; genus: *Epitrix*; specificEpithet: *hirtipennis*; scientificNameAuthorship: Melsheimer; country: Japan; stateProvince: Kanagawa; municipality: Atsugi-shi; locality: Atsugi Campus, Tokyo University of Agriculture, Funako; minimumElevationInMeters: 49; maximumElevationInMeters: 49; decimalLatitude: 35.431707; decimalLongitude: 139.345165; geodeticDatum: WGS84; samplingProtocol: beating of leaves and branches (including visual searches); eventDate: 2016-11-08; individualCount: 2; lifeStage: adult; recordedBy: Y. Yamada; otherCatalogNumbers: 2017-00346 | 2017-00347; identifiedBy: Y. Yamada; datelIdentified: 2017; institutionCode: LETUA; collectionCode: IC
- n. namePublishedIn: 1847; kingdom: Animalia; phylum: Arthropoda; class: Insecta; order: Coleoptera; family: Chrysomelidae; genus: *Epitrix*; specificEpithet: *hirtipennis*; scientificNameAuthorship: Melsheimer; country: Japan; stateProvince: Kanagawa; municipality: Atsugi-shi; locality: Atsugi Campus, Tokyo University of Agriculture, Funako; minimumElevationInMeters: 49; maximumElevationInMeters: 49; decimalLatitude: 35.431707; decimalLongitude: 139.345165; geodeticDatum: WGS84; samplingProtocol: beating of leaves and branches (including visual searches); eventDate: 2016-11-10; individualCount: 1; lifeStage: adult; recordedBy: Y. Yamada; otherCatalogNumbers: 2017-00348; identifiedBy: Y. Yamada; datelIdentified: 2017; institutionCode: LETUA; collectionCode: IC
- o. namePublishedIn: 1847; kingdom: Animalia; phylum: Arthropoda; class: Insecta; order: Coleoptera; family: Chrysomelidae; genus: *Epitrix*; specificEpithet: *hirtipennis*; scientificNameAuthorship: Melsheimer; country: Japan; stateProvince: Kanagawa; municipality: Atsugi-shi; locality: Atsugi Campus, Tokyo University of Agriculture, Funako; minimumElevationInMeters: 49; maximumElevationInMeters: 49; decimalLatitude: 35.431707; decimalLongitude: 139.345165; geodeticDatum: WGS84; samplingProtocol: beating of leaves and branches (including visual searches); eventDate: 2016-11-15; individualCount: 1; lifeStage: adult; recordedBy: Y. Yamada; otherCatalogNumbers: 2017-00349; identifiedBy: Y. Yamada; datelIdentified: 2017; institutionCode: LETUA; collectionCode: IC
- p. namePublishedIn: 1847; kingdom: Animalia; phylum: Arthropoda; class: Insecta; order: Coleoptera; family: Chrysomelidae; genus: *Epitrix*; specificEpithet: *hirtipennis*;

- scientificNameAuthorship: Melsheimer; country: Japan; stateProvince: Kanagawa; municipality: Atsugi-shi; locality: Atsugi Campus, Tokyo University of Agriculture, Funako; minimumElevationInMeters: 49; maximumElevationInMeters: 49; decimalLatitude: 35.431707; decimalLongitude: 139.345165; geodeticDatum: WGS84; samplingProtocol: beating of leaves and branches (including visual searches); eventDate: 2016-11-17; individualCount: 1; lifeStage: adult; recordedBy: Y. Yamada; otherCatalogNumbers: 2017-00350; identifiedBy: Y. Yamada; datelIdentified: 2017; institutionCode: LETUA; collectionCode: IC
- q. namePublishedIn: 1847; kingdom: Animalia; phylum: Arthropoda; class: Insecta; order: Coleoptera; family: Chrysomelidae; genus: *Epitrix*; specificEpithet: *hirtipennis*; scientificNameAuthorship: Melsheimer; country: Japan; stateProvince: Kanagawa; municipality: Atsugi-shi; locality: Atsugi Campus, Tokyo University of Agriculture, Funako; minimumElevationInMeters: 49; maximumElevationInMeters: 49; decimalLatitude: 35.431707; decimalLongitude: 139.345165; geodeticDatum: WGS84; samplingProtocol: beating of leaves and branches (including visual searches); eventDate: 2016-11-25; individualCount: 2; lifeStage: adult; recordedBy: Y. Yamada; otherCatalogNumbers: 2017-00351 | 2017-00352; identifiedBy: Y. Yamada; datelIdentified: 2017; institutionCode: LETUA; collectionCode: IC

**Notes:** Known as a recent alien species to Japan (Harada and Takizawa 2012)

## Order Lepidoptera Linnaeus, 1758

### Family Noctuidae Latreille, 1809

#### *Spodoptera litura* (Fabricius, 1775)

##### Materials

- a. namePublishedIn: 1775; kingdom: Animalia; phylum: Arthropoda; class: Insecta; order: Lepidoptera; family: Noctuidae; genus: *Spodoptera*; specificEpithet: *litura*; scientificNameAuthorship: Fabricius; country: Japan; stateProvince: Kanagawa; municipality: Atsugi-shi; locality: Atsugi Campus, Tokyo University of Agriculture, Funako; minimumElevationInMeters: 49; maximumElevationInMeters: 49; decimalLatitude: 35.431707; decimalLongitude: 139.345165; geodeticDatum: WGS84; samplingProtocol: beating of leaves and branches (including visual searches); eventDate: 2016-10-24; individualCount: 1; lifeStage: larva; recordedBy: Y. Yamada; otherCatalogNumbers: 2017-00353; identifiedBy: T. Ishikawa; datelIdentified: 2017; institutionCode: LETUA; collectionCode: IC
- b. namePublishedIn: 1775; kingdom: Animalia; phylum: Arthropoda; class: Insecta; order: Lepidoptera; family: Noctuidae; genus: *Spodoptera*; specificEpithet: *litura*; scientificNameAuthorship: Fabricius; country: Japan; stateProvince: Kanagawa; municipality: Atsugi-shi; locality: Atsugi Campus, Tokyo University of Agriculture, Funako; minimumElevationInMeters: 27; maximumElevationInMeters: 27; decimalLatitude: 35.43043; decimalLongitude: 139.349516; geodeticDatum: WGS84; samplingProtocol: beating of leaves and branches (including visual searches); eventDate: 2016-11-18; individualCount: 1; lifeStage: larva; recordedBy: Y. Yamada; otherCatalogNumbers: 2017-00354; identifiedBy: T. Ishikawa; datelIdentified: 2017; institutionCode: LETUA; collectionCode: IC

## ***Trichoplusia ni* (Hübner, 1803)**

### **Materials**

- a. namePublishedIn: 1803; kingdom: Animalia; phylum: Arthropoda; class: Insecta; order: Lepidoptera; family: Noctuidae; genus: *Trichoplusia*; specificEpithet: ni; scientificNameAuthorship: Hübner; country: Japan; stateProvince: Kanagawa; municipality: Atsugi-shi; locality: Atsugi Campus, Tokyo University of Agriculture, Funako; minimumElevationInMeters: 27; maximumElevationInMeters: 27; decimalLatitude: 35.43043; decimalLongitude: 139.349516; geodeticDatum: WGS84; samplingProtocol: beating of leaves and branches (including visual searches); eventDate: 2016-11-08; individualCount: 1; lifeStage: larva; recordedBy: K. Niwa; otherCatalogNumbers: 2017-00355; identifiedBy: T. Ishikawa; datelidentified: 2017; institutionCode: LETUA; collectionCode: IC
- b. namePublishedIn: 1803; kingdom: Animalia; phylum: Arthropoda; class: Insecta; order: Lepidoptera; family: Noctuidae; genus: *Trichoplusia*; specificEpithet: ni; scientificNameAuthorship: Hübner; country: Japan; stateProvince: Kanagawa; municipality: Atsugi-shi; locality: Atsugi Campus, Tokyo University of Agriculture, Funako; minimumElevationInMeters: 27; maximumElevationInMeters: 27; decimalLatitude: 35.43043; decimalLongitude: 139.349516; geodeticDatum: WGS84; samplingProtocol: beating of leaves and branches (including visual searches); eventDate: 2016-11-18; individualCount: 8; lifeStage: larva; recordedBy: K. Niwa; otherCatalogNumbers: 2017-00356 | 2017-00357 | 2017-00358 | 2017-00359 | 2017-00360 | 2017-00361 | 2017-00362 | 2017-00363; identifiedBy: T. Ishikawa; datelidentified: 2017; institutionCode: LETUA; collectionCode: IC

## **Order Diptera Linnaeus, 1758**

### **Family Agromyzidae Fallén, 1823**

#### ***Liriomyza sativae* Blanchard, 1938**

### **Materials**

- a. namePublishedIn: 1938; kingdom: Animalia; phylum: Arthropoda; class: Insecta; order: Diptera; family: Agromyzidae; genus: *Liriomyza*; specificEpithet: *sativae*; scientificNameAuthorship: Blanchard; country: Japan; stateProvince: Kanagawa; municipality: Atsugi-shi; locality: Atsugi Campus, Tokyo University of Agriculture, Funako; minimumElevationInMeters: 42; maximumElevationInMeters: 42; decimalLatitude: 35.428874; decimalLongitude: 139.34929; geodeticDatum: WGS84; samplingProtocol: beating of leaves and branches (including visual searches); eventDate: 2016-11-17; individualCount: 4; lifeStage: adult; recordedBy: O-k. Kim; otherCatalogNumbers: 2017-00364 | 2017-00365 | 2017-00366 | 2017-00367; identifiedBy: Y. Yamada; datelidentified: 2017; institutionCode: LETUA; collectionCode: IC
- b. namePublishedIn: 1938; kingdom: Animalia; phylum: Arthropoda; class: Insecta; order: Diptera; family: Agromyzidae; genus: *Liriomyza*; specificEpithet: *sativae*; scientificNameAuthorship: Blanchard; country: Japan; stateProvince: Kanagawa; municipality: Atsugi-shi; locality: Atsugi Campus, Tokyo University of Agriculture, Funako; minimumElevationInMeters: 42; maximumElevationInMeters: 42; decimalLatitude:

- 35.428874; decimalLongitude: 139.34929; geodeticDatum: WGS84; samplingProtocol: beating of leaves and branches (including visual searches); eventDate: 2016-12-15; individualCount: 5; lifeStage: adult; recordedBy: Y. Yamada; otherCatalogNumbers: 2017-00368 | 2017-00369 | 2017-00370 | 2017-00371 | 2017-00372; identifiedBy: Y. Yamada; datelidentified: 2017; institutionCode: LETUA; collectionCode: IC
- c. namePublishedIn: 1938; kingdom: Animalia; phylum: Arthropoda; class: Insecta; order: Diptera; family: Agromyzidae; genus: *Liriomyza*; specificEpithet: *sativae*; scientificNameAuthorship: Blanchard; country: Japan; stateProvince: Kanagawa; municipality: Atsugi-shi; locality: Atsugi Campus, Tokyo University of Agriculture, Funako; minimumElevationInMeters: 42; maximumElevationInMeters: 42; decimalLatitude: 35.428874; decimalLongitude: 139.34929; geodeticDatum: WGS84; samplingProtocol: beating of leaves and branches (including visual searches); eventDate: 2016-12-19; individualCount: 1; lifeStage: adult; recordedBy: Y. Yamada; otherCatalogNumbers: 2017-00373; identifiedBy: Y. Yamada; datelidentified: 2017; institutionCode: LETUA; collectionCode: IC
- d. namePublishedIn: 1938; kingdom: Animalia; phylum: Arthropoda; class: Insecta; order: Diptera; family: Agromyzidae; genus: *Liriomyza*; specificEpithet: *sativae*; scientificNameAuthorship: Blanchard; country: Japan; stateProvince: Kanagawa; municipality: Atsugi-shi; locality: Atsugi Campus, Tokyo University of Agriculture, Funako; minimumElevationInMeters: 42; maximumElevationInMeters: 42; decimalLatitude: 35.428874; decimalLongitude: 139.34929; geodeticDatum: WGS84; samplingProtocol: beating of leaves and branches (including visual searches); eventDate: 2017-01-07; individualCount: 13; lifeStage: adult; recordedBy: Y. Yamada; otherCatalogNumbers: 2017-00374 | 2017-00375 | 2017-00376 | 2017-00377 | 2017-00378 | 2017-00379 | 2017-00380 | 2017-00381 | 2017-00382 | 2017-00383 | 2017-00384 | 2017-00385 | 2017-00386; identifiedBy: Y. Yamada; datelidentified: 2017; institutionCode: LETUA; collectionCode: IC

## Class Arachnida Lamarck, 1801

### Order Acari Leach, 1817

#### Family Tarsonemidae Kramer, 1877

##### *Polyphagotarsonemus latus* (Banks, 1904)

#### Material

- a. namePublishedIn: 1904; kingdom: Animalia; phylum: Arthropoda; class: Arachnida; order: Acari; family: Tarsonemidae; genus: *Polyphagotarsonemus*; specificEpithet: *latus*; scientificNameAuthorship: Banks; country: Japan; stateProvince: Kanagawa; municipality: Atsugi-shi; locality: Atsugi Campus, Tokyo University of Agriculture, Funako; minimumElevationInMeters: 42; maximumElevationInMeters: 42; decimalLatitude: 35.428874; decimalLongitude: 139.34929; geodeticDatum: WGS84; samplingProtocol: beating of leaves and branches (including visual searches); eventDate: 2017-01-07; individualCount: 1; lifeStage: adult; recordedBy: Y. Yamada; otherCatalogNumbers: 2017-00387; identifiedBy: Y. Yamada; datelidentified: 2017; institutionCode: LETUA; collectionCode: IC

## Family Tetranychidae Donnadiieu, 1875

### *Tetranychus urticae* Koch, 1836

#### Materials

- a. namePublishedIn: 1836; kingdom: Animalia; phylum: Arthropoda; class: Arachnida; order: Acari; family: Tetranychidae; genus: *Tetranychus*; specificEpithet: *urticae*; scientificNameAuthorship: Koch; country: Japan; stateProvince: Kanagawa; municipality: Atsugi-shi; locality: Atsugi Campus, Tokyo University of Agriculture, Funako; minimumElevationInMeters: 49; maximumElevationInMeters: 49; decimalLatitude: 35.431707; decimalLongitude: 139.345165; geodeticDatum: WGS84; samplingProtocol: beating of leaves and branches (including visual searches); eventDate: 2016-08-30; individualCount: 8; lifeStage: adult; recordedBy: T. Ishikawa; otherCatalogNumbers: 2017-00388 | 2017-00389 | 2017-00390 | 2017-00391 | 2017-00392 | 2017-00393 | 2017-00394 | 2017-00395; identifiedBy: Y. Yamada; datelidentified: 2017; institutionCode: LETUA; collectionCode: IC
- b. namePublishedIn: 1836; kingdom: Animalia; phylum: Arthropoda; class: Arachnida; order: Acari; family: Tetranychidae; genus: *Tetranychus*; specificEpithet: *urticae*; scientificNameAuthorship: Koch; country: Japan; stateProvince: Kanagawa; municipality: Atsugi-shi; locality: Atsugi Campus, Tokyo University of Agriculture, Funako; minimumElevationInMeters: 42; maximumElevationInMeters: 42; decimalLatitude: 35.428874; decimalLongitude: 139.34929; geodeticDatum: WGS84; samplingProtocol: beating of leaves and branches (including visual searches); eventDate: 2017-01-07; individualCount: 13; lifeStage: adult; recordedBy: Y. Yamada; otherCatalogNumbers: 2017-00396 | 2017-00397 | 2017-00398 | 2017-00399 | 2017-00400 | 2017-00401 | 2017-00402 | 2017-00403 | 2017-00404 | 2017-00405 | 2017-00406 | 2017-00407 | 2017-00408; identifiedBy: Y. Yamada; datelidentified: 2017; institutionCode: LETUA; collectionCode: IC
- c. namePublishedIn: 1836; kingdom: Animalia; phylum: Arthropoda; class: Arachnida; order: Acari; family: Tetranychidae; genus: *Tetranychus*; specificEpithet: *urticae*; scientificNameAuthorship: Koch; country: Japan; stateProvince: Kanagawa; municipality: Atsugi-shi; locality: Atsugi Campus, Tokyo University of Agriculture, Funako; minimumElevationInMeters: 42; maximumElevationInMeters: 42; decimalLatitude: 35.428874; decimalLongitude: 139.34929; geodeticDatum: WGS84; samplingProtocol: beating of leaves and branches (including visual searches); eventDate: 2017-01-09; individualCount: 18; lifeStage: adult; recordedBy: Y. Yamada; otherCatalogNumbers: 2017-00409 | 2017-00410 | 2017-00411 | 2017-00412 | 2017-00413 | 2017-00414 | 2017-00415 | 2017-00416 | 2017-00417 | 2017-00418 | 2017-00419 | 2017-00420 | 2017-00421 | 2017-00422 | 2017-00423 | 2017-00424 | 2017-00425 | 2017-00426; identifiedBy: Y. Yamada; datelidentified: 2017; institutionCode: LETUA; collectionCode: IC
- d. namePublishedIn: 1836; kingdom: Animalia; phylum: Arthropoda; class: Arachnida; order: Acari; family: Tetranychidae; genus: *Tetranychus*; specificEpithet: *urticae*; scientificNameAuthorship: Koch; country: Japan; stateProvince: Kanagawa; municipality: Atsugi-shi; locality: Atsugi Campus, Tokyo University of Agriculture, Funako; minimumElevationInMeters: 42; maximumElevationInMeters: 42; decimalLatitude: 35.428874; decimalLongitude: 139.34929; geodeticDatum: WGS84; samplingProtocol: beating of leaves and branches (including visual searches); eventDate: 2017-01-09; individualCount: 2; lifeStage: larva; recordedBy: Y. Yamada; otherCatalogNumbers:

- 2017-00427 | 2017-00428; identifiedBy: Y. Yamada; dateIdentified: 2017; institutionCode: LETUA; collectionCode: IC
- e. namePublishedIn: 1836; kingdom: Animalia; phylum: Arthropoda; class: Arachnida; order: Acari; family: Tetranychidae; genus: *Tetranychus*; specificEpithet: *urticae*; scientificNameAuthorship: Koch; country: Japan; stateProvince: Kanagawa; municipality: Atsugi-shi; locality: Atsugi Campus, Tokyo University of Agriculture, Funako; minimumElevationInMeters: 42; maximumElevationInMeters: 42; decimalLatitude: 35.428874; decimalLongitude: 139.34929; geodeticDatum: WGS84; samplingProtocol: beating of leaves and branches (including visual searches); eventDate: 2017-01-12; individualCount: 15; lifeStage: adult; recordedBy: Y. Yamada; otherCatalogNumbers: 2017-00429 | 2017-00430 | 2017-00431 | 2017-00432 | 2017-00433 | 2017-00434 | 2017-00435 | 2017-00436 | 2017-00437 | 2017-00438 | 2017-00439 | 2017-00440 | 2017-00441 | 2017-00442 | 2017-00443; identifiedBy: Y. Yamada; dateIdentified: 2017; institutionCode: LETUA; collectionCode: IC
- f. namePublishedIn: 1836; kingdom: Animalia; phylum: Arthropoda; class: Arachnida; order: Acari; family: Tetranychidae; genus: *Tetranychus*; specificEpithet: *urticae*; scientificNameAuthorship: Koch; country: Japan; stateProvince: Kanagawa; municipality: Atsugi-shi; locality: Atsugi Campus, Tokyo University of Agriculture, Funako; minimumElevationInMeters: 42; maximumElevationInMeters: 42; decimalLatitude: 35.428874; decimalLongitude: 139.34929; geodeticDatum: WGS84; samplingProtocol: beating of leaves and branches (including visual searches); eventDate: 2017-01-16; individualCount: 8; lifeStage: adult; recordedBy: Y. Yamada; otherCatalogNumbers: 2017-00444 | 2017-00445 | 2017-00446 | 2017-00447 | 2017-00448 | 2017-00449 | 2017-00450 | 2017-00451; identifiedBy: Y. Yamada; dateIdentified: 2017; institutionCode: LETUA; collectionCode: IC
- g. namePublishedIn: 1836; kingdom: Animalia; phylum: Arthropoda; class: Arachnida; order: Acari; family: Tetranychidae; genus: *Tetranychus*; specificEpithet: *urticae*; scientificNameAuthorship: Koch; country: Japan; stateProvince: Kanagawa; municipality: Atsugi-shi; locality: Atsugi Campus, Tokyo University of Agriculture, Funako; minimumElevationInMeters: 42; maximumElevationInMeters: 42; decimalLatitude: 35.428874; decimalLongitude: 139.34929; geodeticDatum: WGS84; samplingProtocol: beating of leaves and branches (including visual searches); eventDate: 2017-01-21; individualCount: 12; lifeStage: adult; recordedBy: Y. Yamada; otherCatalogNumbers: 2017-00452 | 2017-00453 | 2017-00454 | 2017-00455 | 2017-00456 | 2017-00457 | 2017-00458 | 2017-00459 | 2017-00460 | 2017-00461 | 2017-00462 | 2017-00463; identifiedBy: Y. Yamada; dateIdentified: 2017; institutionCode: LETUA; collectionCode: IC

## Discussion

Prior to the present study, nine groups of insects and mites have been known as pests of pepino plants in Japan (Table 3), including those that are inadequately identified, such as *Helicoverpa* spp. at the genus level, aphids at the family level (Aphididae), and mites at the order level (Acari). Among these, three groups were identified at the species level, comprising the greenhouse whitefly *Trialeurodes vaporariorum* (Westwood, 1856), the potato tuberworm *Phthorimaea operculella* (Zeller, 1873), and the two-spotted spider mite *Tetranychus urticae* Koch, 1836. In the present study, our survey in the campus revealed the presence of 11 species of insect and mite pests on pepino as mentioned above. Only *T. urticae* is common in the previous records as well as the results of the present study; in

addition, *T. vaporariorum* and *P. operculella* were not found in our survey. Currently, 13 species of insects and mites are recognized as pests of pepino plants in Japan; all of these are also the pests of major solanaceous crops such as tomato, eggplant, green pepper, and potato (The Japanese Society of Applied Entomology and Zoology 2006). It is likely that *T. vaporariorum* and *P. operculella* may be detected on pepino from the campus, because the two species are widely distributed in Kanagawa Prefecture (Matsumoto 2004, Nakajima and Yamamoto 2004) where our study was carried out.

Table 3.

Insect and mite pests of pepino plants previously recorded in Japan.

Class	Order	Family	Group (of species and its allies)	Species	References
Insecta	Hemiptera	Aleyrodidae	-	<i>Trialeurodes vaporariorum</i> (Westwood, 1856)	Furusato (1984), Takahashi (1985), Takagi (1985), Kita (1986), Odagiri et al. (1986), Ozawa (1986)
Insecta	Hemiptera	Apididae	aphids	-	Takagi (1985), Takahashi (1985), Kita (1986), Odagiri et al. (1986), Ozawa (1986), Takahashi (1986)
Insecta	Lepidoptera	Noctuidae	<i>Helicoverpa</i> spp.	-	Odagiri et al. (1986)
Insecta	Lepidoptera	Noctuidae	<i>Spodoptera</i> spp.	-	Odagiri et al. (1986)
Insecta	Lepidoptera	Gelechiidae	-	<i>Phthorimaea operculella</i> (Zeller, 1873)	Ozawa (1986)
Insecta	Lepidoptera	unspecified	green caterpillars	-	Furusato (1986)
Arachnida	Acari	unspecified	mites	-	Furusato (1984), Kita (1986), Odagiri et al. (1986)
Arachnida	Acari	Tetranychidae	spider mites	-	Takahashi (1985), Takagi (1985), Ozawa (1986), Takahashi (1986)
Arachnida	Acari	Tetranychidae	-	<i>Tetranychus urticae</i> Koch, 1836	Ozawa (1986)

Currently, 25 species of insects and mites have been reported as pests of pepino plants worldwide (Table 4) (Galbreath and Clearwater 1983, Larraín 2002, Akyazi 2012). Among these pests, 16 species are distributed in Japan, but only three species, *Trichoplusia ni*, *Polyphagotarsonemus latus*, and *Tetranychus urticae*, were evaluated as pests of pepino in the present study. This indicates that the remaining 13 species are very likely to be

potential pests on pepino in Japan. Therefore, at least 26 insect and mite species, including the 13 currently known and the 13 potential ones in Japan, will be recognized as pests of pepino in the near future if the cultivation of pepinos spreads throughout the Japanese Archipelago.

Table 4.

Insect and mite pests of pepino plants previously recorded worldwide (excluding Japan).

Class	Order	Family	Species	Country recorded as a pest	References	Notes
Insecta	Orthoptera	Acrididae	<i>Schistocerca cancellata</i> (Serville, 1838)	Chili	Larraín (2002)	
Insecta	Thysanoptera	Thripidae	<i>Frankliniella occidentalis</i> (Pergande, 1895)	Chili	Larraín (2002)	Distributed in Japan
Insecta	Thysanoptera	Thripidae	<i>Thrips tabaci</i> Lindeman, 1889	Chili	Larraín (2002)	Distributed in Japan
Insecta	Hemiptera	Aphididae	<i>Aulacorthum solani</i> (Kaltenbach, 1843)	Chili	Larraín (2002)	Distributed in Japan
Insecta	Hemiptera	Aphididae	<i>Macrosiphum euphorbiae</i> (Thomas, 1878)	Chili	Larraín (2002)	Distributed in Japan
Insecta	Hemiptera	Aphididae	<i>Myzus persicae</i> (Sulzer, 1776)	Chili	Larraín (2002)	Distributed in Japan
Insecta	Hemiptera	Pseudococcidae	<i>Phenacoccus solenopsis</i> Tinsley, 1898	Chili	Larraín (2002)	Distributed in Japan
Insecta	Hemiptera	Pseudococcidae	<i>Pseudococcus viburni</i> (Signoret, 1875)	Chili	Larraín (2002)	Distributed in Japan
Insecta	Hemiptera	Psyllidae	<i>Russelliana solanicola</i> Tuthill, 1959	Chili	Larraín (2002)	
Insecta	Hemiptera	Trioziidae	<i>Trioza chenopodii</i> Reuter, 1876	Chili	Larraín (2002)	Distributed in Japan
Insecta	Hemiptera	Cicadellidae	<i>Xerophloea viridis</i> (Fabricius, 1794)	Chili	Larraín (2002)	
Insecta	Hemiptera	Cicadellidae	<i>Paratanus exitiosus</i> (Beamer, 1943)	Chili	Larraín (2002)	
Insecta	Diptera	Agromyzidae	<i>Liriomyza huidobrensis</i> (Blanchard, 1926)	Chili	Larraín (2002)	Distributed in Japan

Insecta	Diptera	Tephritidae	<i>Rhagoletis nova</i> (Schiner, 1868)	Chili	Larraín (2002)	
Insecta	Lepidoptera	Noctuidae	<i>Agrotis bilitura</i> Guenée, 1852	Chili	Larraín (2002)	
Insecta	Lepidoptera	Noctuidae	<i>Copitarsia turbata</i> (Herrich-Schaeffer, 1855)	Chili	Larraín (2002)	
Insecta	Lepidoptera	Noctuidae	<i>Trichoplusia ni</i> (Hübner, [1803])	Chili	Larraín (2002)	Recognized as a pest in Japan by this study
Insecta	Lepidoptera	Sphingidae	<i>Manduca sexta</i> (Linnaeus, 1763)	Chili	Larraín (2002)	Distributed in Japan
Insecta	Lepidoptera	Gelechiidae	<i>Phthorimaea operculella</i> (Zeller, 1873)	Chili	Larraín (2002)	Distributed in Japan
Insecta	Lepidoptera	Gelechiidae	<i>Symmetrischema tangolias</i> (Gyen, 1913)	Chili	Larraín (2002)	
Insecta	Lepidoptera	Gelechiidae	<i>Tuta absoluta</i> (Meyrick, 1917)	Chili	Larraín (2002)	Distributed in Japan
Insecta	Lepidoptera	Crambidae	<i>Sceliodes cordalis</i> (Doubleday, 1843)	New Zealand	Galbreath and Clearwater (1983)	
Arachnida	Acari	Eriophyidae	<i>Aculops lycopersici</i> (Tryon, 1917)	Chili, Turkey	Larraín (2002), Akyazi (2012)	Distributed in Japan
Arachnida	Acari	Tarsonemidae	<i>Polyphagotarsonemus latus</i> (Banks, 1904)	Chili	Larraín (2002)	Recognized as a pest in Japan by this study
Arachnida	Acari	Tetranychidae	<i>Tetranychus urticae</i> Koch, 1836	Chili	Larraín (2002)	Recognized as a pest in Japan by this study

The presence of pests can directly affect agricultural production and it may contribute to the transmission of plant viruses followed by economic losses. *Bemisia tabaci*, well-known as one of the most important whiteflies in terms of virus transmission, is widely distributed in the world, and is a vector of viruses of the genera *Begomovirus*, *Carlavirus*, *Crinivirus*, *Ipomovirus*, and *Torradovirus* (Navas-Castillo et al. 2011). Of these transmissible viruses, *Tomato yellow leaf curl virus* (TYLCV; *Begomovirus*) is the most devastating causal virus on tomato crops in many tropical, subtropical and temperate regions worldwide (Moriones and Navas-Castillo 2000). In Japan, TYLCV spread along with whitefly and indicated its

dispersion throughout 38 prefectures by 2014 since the occurrence of TYLCV on tomato was first reported in Japan (Kato et al. 1998, Matsuura and Hoshino 2008, Ohnishi et al. 2016). However, there is no report of TYLCV incidence on pepino plants to date. *Pepino* can be regarded as a potential host for TYLCV through host adaptation or mutant as long as TYLCV-acquired vectors are present. Therefore, continuous monitoring of the distribution of both TYLCV and its vector is required. Some other species of insects and mites found on pepino plants in our research fields are also regarded as virus vectors (Table 5). Although some transmitted viruses cannot infect pepino plants, they may provide a habitat for virus vectors, which may impact crop ecosystems as well as virus-vector systems.

Table 5.

Insects or mites of pepino plants involved in virus transmission

Insect vector	Transmissible genus or species of virus	Reference
<i>Frankliniella intonsa</i>	<i>Groundnut ringspot virus</i>	Riley et al. (2011)
	<i>Impatiens necrotic spot virus</i>	
	<i>Tomato chlorotic spot virus</i>	
	<i>Tomato spotted wilt virus</i>	
<i>Bemisia tabaci</i>	Genus <i>Begomovirus</i>	Navas-Castillo et al. (2011)
	Genus <i>Carlavirus</i>	
	Genus <i>Crinivirus</i>	
	Genus <i>Ipomovirus</i>	
	Genus <i>Torradovirus</i>	
<i>Aphis gossypii</i>	<i>Cucumber mosaic virus</i> (CMV)	Pinto et al. (2008)
	<i>Papaya ringspot virus</i> (PRSV)	
	<i>Tobacco ringspot virus</i> (TRSV)	
	<i>Zucchini yellow mosaic virus</i> (ZYMV)	
<i>Epitrix hirtipennis</i>	<i>Tobacco ringspot virus</i> (TRSV)	Stace-Smith (1985)
<i>Liriomyza sativae</i>	<i>Celery mosaic virus</i> (CeMV)	Zitter and Tsai (1977)
	<i>Watermelon mosaic virus</i> strain 1*1	
	<i>Watermelon mosaic virus</i> strain 2	
<i>Tetranychus urticae</i>	<i>Tobacco mosaic virus</i> (TMV)*2	Orlob (1968)
	<i>Potato virus X</i> (PVX)*2	

We found virus-like symptoms, showing mottle and deformation on pepino leaves. However, none of the tested viruses were detected in any plant. Our internet-based image searching results showed that the symptoms were similar to those on pepper or chili plants by *Polyphagotarsonemus latus*. Grinberg et al. (2005) described that young leaves were usually affected by *P. latus* and consequently showed distortion and leaf-curl downwards. This suggests that our symptomatic leaves might be caused by *P. latus*. Further research is required to reveal any effect of *P. latus* on pepino.

No related virus was detected in the present study, whereas two virus species had been detected from pepino plants in Japan: *Alfalfa mosaic virus* (AMV) and *Cucumber mosaic virus* (CMV) inducing chlorotic ring spots and mosaic symptoms on pepino plants, respectively (Honda et al. 1986). Moreover, *Pepino mosaic virus* (PepMV) was first isolated from pepinos showing yellow mosaic symptoms in Peru in 1974 (Jones et al. 1980). Currently, PepMV has become a major pathogen of tomato plants worldwide and is one of the quarantine pathogens strictly prohibited from entry into Japan. *Pepino* latent virus (PePLV; later reclassified as *Potato virus S* (PVS)) was detected from pepino cuttings in New Zealand, even though pepino plants with no symptoms had been imported from Chile (Thomas et al. 1980). With the increase in pepino production in China, *Potato virus H* (PVH) infected pepino, with no obvious symptoms (Abouelnasr et al. 2014) (Table 6). Pests and pathogens in many crops cause tremendous losses both in terms of quantity and quality. Monitoring, detection, and identification of pests and pathogens prior to the introduction of a new crop or during production are important to efficiently regulate their future damage. Our findings will aid in understanding the incidence of pests and viral diseases on pepino plants and developing better crop production systems to combat pests and diseases.

Table 6.

Viruses infecting pepino plants naturally around the world.

Family	Genus	Species	Acronym	Symptoms on pepino	First reported country	Reference
<i>Alphaflexiviridae</i>	<i>Potexvirus</i>	<i>Pepino mosaic virus</i>	PepMV	yellow mosaic in young leaves	Peru	Jones et al. (1980)
<i>Betaflexiviridae</i>	<i>Carlavirus</i>	<i>Potato virus H</i>	PVH	symptomless	China	Abouelnasr et al. (2014)
		<i>Pepino latent virus*3</i>	PePLV	symptomless	New Zealand	Thomas et al. (1980)
<i>Bromoviridae</i>	<i>Alfamovirus</i>	<i>Alfalfa mosaic virus</i>	AMV	chlorotic ring spot	Japan	Honda et al. (1986)
	<i>Cucumovirus</i>	<i>Cucumber mosaic virus</i>	CMV	mosaic	Japan	Honda et al. (1986)

## Acknowledgements

We are highly grateful to all the members of our project for kindly offering materials, supporting our field investigations and/or providing valuable comments on the manuscript. We also thank the members of the Laboratory of Vegetables and the Laboratory of Plant Pathology for their committed cultivation and management of pepino plants in our experimental fields. We are also indebted to Jenő Kontschán (Hungarian Academy of Sciences, Hungary) and Alina Avanesyan (Grand View University, USA) for their critical reading of the manuscript and for giving valuable comments. This study was financially supported by the Strategic Research Project from Tokyo University of Agriculture, Tokyo, Japan. We would like to thank Editage ([www.editage.jp](http://www.editage.jp)) for English language editing.

## References

- Abouelnasr H, Li Y, Zhang Z, Liu J, Li S, Li D, Yu J, McBeath JH, Han C (2014) First report of Potato virus H on *Solanum muricatum* in China. *Plant Disease* 98: 1016. <https://doi.org/10.1094/PDIS-01-14-0024-PDN>
- Akyazi R (2012) First report of *Aculops lycopersici* (Tryon, 1917) (Acarı: Eriophyidae) on Pepino in Turkey. *Journal of Entomological and Acarological Research* 2012 44: e20. <https://doi.org/10.4081/jear.2012.e20>
- Badge J, Brunt A, Carson R, Dagless E, Karamagioli M, Phillips S, Seal S, Turner R, Foster GD (1996) A carlavirus-specific PCR primer and partial nucleotide sequence provides further evidence for the recognition of cowpea mild mottle virus as a whitefly-transmitted carlavirus. *European Journal of Plant Pathology* 102: 305-310. <https://doi.org/10.1007/BF01877970>
- Choi SK, Choi JK, Park WM, Ryu KH (1999) RT-PCR detection and identification of three species of cucumoviruses with a genus-specific single pair of primers. *Journal of Virological Methods* 83: 67-73. [https://doi.org/10.1016/S0166-0934\(99\)00106-8](https://doi.org/10.1016/S0166-0934(99)00106-8)
- Dolby CA, Jones RAC (1988) The relationship between the Andean strain of potato virus S and pepino latent virus. *Annals of Applied Biology* 112: 231-234. <https://doi.org/10.1111/j.1744-7348.1988.tb02058.x>
- Ehara S, Gotoh T (Eds) (2009) Colored Guide to the Plant Mites of Japan. Zenkoku Noson Kyoku Kyokai, Tokyo, vi+349 pp. [In Japanese]. [ISBN 4-05-402370-3]
- Furukawa M (2005) Noctuidae. In: Ishiwata S, Hanada S, Hayashi F, Yamazaki T, Uemura Y, Aoki N, Hayashi M, Nozaki T, Fukuda H, Kishida Y, Hayashi N, Shinohara A, Shinonaga T (Eds) *Insect Larvae of Japan*. Gakken, Tokyo, 217-225 pp. [In Japanese]. [ISBN 4-05-402370-3].
- Furusato K (1984) Atarashii-Yasai "Pepino" to sono-Tsukuri-kata [A new vegetable "pepino" and its cultivation]. *Nogyo Gijutsu Kenkyu* 38 (7): 36-37. [In Japanese].
- Furusato K (1986) Peru ni-okeru Pepino no Saibai-jōkyō [The situation in cultivation of pepino in Peru]. *Nogyo Gijutsu Kenkyu* 40 (6): 46-47. [In Japanese].
- Galbreath RA, Clearwater JR (1983) Pheromone monitoring of *Sceliodes cordalis*, a pest of pepino. In: Hartley M (Ed.) *Proceedings of 36th New Zealand Weed and Pest Control Conference*. Hastings, 9-11 August 1983. The New Zealand Weed and Pest Control Society Inc., Palmerston North, New Zealand, 128-130 pp.

- Grinberg M, Perl-Treves R, Palevsky E, Shomer I, Soroker V (2005) Interaction between cucumber plants and the broad mite, *Polyphagotarsonemus latus* from damage to defense gene expression. *Entomologia Experimentalis et Applicata* 115: 135-144. <https://doi.org/10.1111/j.1570-7458.2005.00275.x>
- Ha C, Coombs S, Revill PA, Harding RM, Vu M, Dale JL (2008) Design and application of two novel degenerate primer pairs for the detection and complete genomic characterization of potyviruses. *Archives of Virology* 153: 25-36. <https://doi.org/10.1007/s00705-007-1053-7>
- Harada H, Takizawa H (2012) Occurrence of *Epitrix hirtipennis* (Melsheimer) (Coleoptera: Chrysomelidae), an alien insect pest, in Japan. *Japanese Journal of Applied Entomology and Zoology* 56: 117-120. [In Japanese with English summary]. <https://doi.org/10.1303/jjaez.2012.117>
- Heiser CB (1964) Origin and variability of the pepino (*Solanum muricatum*): A preliminary report. *Baileya* 12: 151-158.
- Honda Y, Hanada K, Ushiyama K, Zenbayashi R, Tochihara H (1986) Alfalfa mosaic virus and cucumber mosaic virus isolated from pepino (*Solanum muricatum*). *Annals of the Phytopathological Society of Japan* 52: 870-873. [In Japanese with English abstract]. <https://doi.org/10.3186/ijphytopath.52.870>
- Iwasaki A, Kasugai K, Iwaizumi R, Sasakawa M (2000) A newly recorded pest, *Liriomyza sativae* Blanchard in Japan. *Plant Protection* 54 (4): 142-147. [In Japanese].
- Japan Meteorological Agency (2017) <http://www.jma.go.jp/jma/index.html>. Accessed on: 2017-3-29.
- Jones RAC, Koenig R, Lesemann D (1980) Pepino mosaic virus, a new potexvirus from pepino (*Solanum muricatum*). *Annals of Applied Biology* 94: 61-68. <https://doi.org/10.1111/j.1744-7348.1980.tb03896.x>
- Kato K, Onuki M, Fuji S, Hanada K (1998) The first occurrence of tomato yellow leaf curl virus in tomato (*Lycopersicon esculentum* Mill.) in Japan. *Annals of the Phytopathological Society of Japan* 64: 552-559. [In Japanese]. <https://doi.org/10.3186/ijphytopath.64.552>
- Kawai S (1980) Scale Insects of Japan in Colors. Zenkoku Noson Kyoiku Kyokai, Tokyo, viii+455 pp. [In Japanese]. [ISBN 4-88137-011-1]
- Kita N (1986) Pepino no Saibai-gijutsu [Cultivation technique of pepino]. *Yasai Engei Gijutsu* 13 (9): 6-10. [In Japanese].
- Larraín SP (2002) Incidencia de insectos y ácaros plagas en pepino dulce (*Solanum muricatum* Ait.) culticado en la IV Región. *Agricultura Técnica* 61: 15-26. [In Spanish]. <https://doi.org/10.4067/S036528072002000100002>
- Masumoto M, Okajima S (2013) Review of the genus *Thrips* and related genera (Thysanoptera, Thripidae) from Japan. *Zootaxa* 3678: 1-65. <https://doi.org/10.11646/zootaxa.3678.1.1>
- Matsumoto K (2004) Hemiptera (Sternorrhyncha: Psylloidea & Aleyrodoidea). In: Kanagawa Konchū Danwakai Insect Fauna of Kanagawa, I (General Remarks, Protura, Diplura, Collembola, Archaeognatha, Thysanura, Ephemeroptera, Odonata, Plecoptera, Grylloblattodea, Orthoptera, Phasmida, Dermaptera, Mantodea, Blattaria, Isoptera, Psocoptera, Mallophaga, Anoplura, Hemiptera). Kanagawa Konchū Danwakai, Odawara, 273-282 pp. [In Japanese]. [ISBN 4-9902315-1-1].
- Matsumoto Y (2008) A Guide Illustrated Book of Aphids. Zenkoku Noson Kyoiku Kyokai, Tokyo, (10)+239 pp. [In Japanese]. [ISBN 978-4-88137-134-3]

- Matsuura S, Hoshino S (2008) Comparative spatial dispersal of *Tomato yellow leaf curl virus* vectored by B and Q biotypes of *Bemisia tabaci* in tomato glasshouses. *Phytoparasitica* 36: 42-51. <https://doi.org/10.1007/BF02980747>
- Moriones E, Navas-Castillo J (2000) Tomato yellow leaf curl virus, an emerging virus complex causing epidemics worldwide. *Virus Research* 71: 123-134. [https://doi.org/10.1016/S0168-1702\(00\)00193-3](https://doi.org/10.1016/S0168-1702(00)00193-3)
- Moritsu M (1983) Aphids of Japan in Colors. Zenkoku Noson Kyoiku Kyokai, Tokyo, 545 pp. [In Japanese]. [ISBN 4-88137-017-0]
- Nakajima H, Yamamoto M (2004) Lepidoptera (excl. Hesperioidea & Papilioidea). In: Kanagawa Konchū Danwakai Insect Fauna of Kanagawa, III (Mecoptera, Siphonaptera, Diptera, Lepidoptera, Trichoptera, Hymenoptera). Kanagawa Konchū Danwakai, Odawara, 907-1158 pp. [In Japanese]. [ISBN 4-9902315-1-1].
- Navas-Castillo J, Fiallo-Olivé E, Sánchez-Campos S (2011) Emerging virus diseases transmitted by whiteflies. *Annual Review of Phytopathology* 49: 219-248. <https://doi.org/10.1146/annurev-phyto-072910-095235>
- Odagiri T, Nishiyama T, Inaba K, Fukuoka N, Saida Y (1986) Cultivation management and plant physiology and ecology. 12. Establishment of cultivation system for pepino. *Sosai Shiken Seisekisho* 1986: 120-122. [In Japanese].
- Ohnishi J, Yamaguchi H, Saito A (2016) Analysis of the Mild strain of tomato yellow leaf curl virus, which overcomes *Ty-2* gene-mediated resistance in tomato line H24. *Archives of Virology* 161: 2207-2217. <https://doi.org/10.1007/s00705-016-2898-4>
- Orlob GB (1968) Relationship between *Tetranychus urticae* Koch and some plant viruses. *Virology* 35: 121-133. [https://doi.org/10.1016/0042-6822\(68\)90312-7](https://doi.org/10.1016/0042-6822(68)90312-7)
- Ozawa A (1986) Pepino no Seitaiteki-tokusei to Saibaijō no Ryūi-ten [Ecological properties and points of attention for cultivation in pepino]. *Shisetsu Engei* 28 (1): 26-29. [In Japanese].
- Pinto ZV, Rezende JAM, Yuki VA, Piedade SMS (2008) Ability of *Aphis gossypii* and *Myzus persicae* to transmit Cucumber mosaic virusin single and mixed infection with two potyviruses to zucchini squash. *Summa Phytopathologica Summa Phytopathologica* 34: 183-185. <https://doi.org/10.1590/S0100-54052008000200016>
- Purcifull D, Hiebert E, Edwardson J (1984) Description of Plant Viruses. No. 293. Watermelon mosaic virus 2. <http://www.dpvweb.net/dpv/showdpv.php?dpvno=293>. Accessed on: 2017-6-29.
- Riley DG, Joseph SV, Srinivasan R, Diffie S (2011) Thrips vectors of tospoviruses. *Journal of Integrated Pest Management* 1 (2): 1-10. <https://doi.org/10.1603/IPM10020>
- Sakata Y (2011) Pepino. In: Nosangyoson-bunkakyokai Encyclopedia of growing vegetable 20. 2nd edition. Nosangyoson-bunkakyokai, Tokyo, 305-309 pp. [In Japanese].
- Stace-Smith R (1985) Description of Plant Viruses. No. 309. Tobacco ringspot virus . <http://www.dpvweb.net/dpv/showdpv.php?dpvno=309>. Accessed on: 2017-6-29.
- Takagi K (1985) Kasai Pepino [Pepino, a fruit vegetable]. *Garden Life* 3: 67-70. [In Japanese].
- Takahashi M (1985) Pepino. In: Kiyota I (Ed.) *Yûbô-shin-yasai no Tsukuri-kata* [Cultivation methods for new prospective vegetables]. Rural Culture Association Japan, Tokyo, 112-120 pp. [In Japanese]. [ISBN 4-540-85003-2].

- Takahashi M (1986) Shin-dônyû-kasai Pepino no Tokusei to Riyô-hô [Property and usage in pepino, a newly introduced vegetable]. *Kongetsu no Noyaku* 30 (1): 42-46. [In Japanese].
- Takahata K (2017) Effects of stem constriction using steel washer rings on the soluble solids content of pepino (*Solanum muricatum* Ait.) fruit. *The Horticulture Journal Preview* <https://doi.org/10.2503/hortj.OKD-034>
- Tanaka H, Uesato T (2012) New records of some potential pest mealybugs (Hemiptera: Coccoidea: Pseudococcidae) in Japan. *Applied Entomology and Zoology* 47: 413-419. <https://doi.org/10.1007/s13355-012-0134-6>
- The Japanese Society of Applied Entomology and Zoology (2006) Major Insect and Other Pests of Economic Plants in Japan. Revised Edition. The Japanese Society of Applied Entomology and Zoology, Tokyo, v+387 pp.
- Thomas W, Mohamed NA, Fry ME (1980) Properties of a carlavirus causing a latent infection of pepino (*Solanum muricatum*). *Annals of Applied Biology* 95: 191-196. <https://doi.org/10.1111/j.1744-7348.1980.tb04738.x>
- Umeya K, Okada T (Eds) (2003) Agricultural Insect Pests in Japan. Zenkoku Noson Kyoiku Kyokai, Tokyo, 1203 pp. [In Japanese]. [ISBN 4-88137-103-7]
- van der Vlugt RAA, Berendsen M (2002) Development of a general potexvirus detection method. *European Journal of Plant Pathology* 108: 367-371. <https://doi.org/10.1023/A:1015644409484>
- Xu H, Nie J (2006) Identification, characterization, and molecular detection of Alfalfa mosaic virus in Potato. *Phytopathology* 96: 1237-1242. <https://doi.org/10.1094/PHYTO-96-1237>
- Yasunaga T, Schuh RT, Duwal RK (2015) Taxonomic review of the plant bug genus *Campylomma* Reuter from Japan (Heteroptera: Miridae: Phylinae: Nasocorini), with descriptions of two new species. *Tijdschrift voor Entomologie* 158: 49-69. <https://doi.org/10.1163/22119434-00002046>
- Yasunaga T, Takai M, Kawasawa T (Eds) (2001) A Field Guide to Japanese Bugs – Terrestrial Heteropterans. Vol. 2. Zenkoku Noson Kyoiku Kyokai, Tokyo, 350 pp. [In Japanese]. [ISBN 4-88137-089-8]
- Zitter TA, Tsai JH (1977) Transmission of three potyviruses by the leaf miner, *Liriomyza sativae* (Diptera: Agromyzidae). *Plant Disease Reporter* 61: 1025-1029.

## Endnotes

\*<sup>1</sup> Reclassified as PRSV (Purcifull et al. 1984)

\*<sup>2</sup> Experimentally transmissible

\*<sup>3</sup> Reclassified as an isolate of *Potato virus S* (PVS) based on biological and serological characteristics (Dolby and Jones 1988).