

ICD-1/BTF3 antagonizes SKN-1-mediated endoderm specification in *Caenorhabditis elegans*

Chee Kiang Ewe¹, Yamila N Torres Cleuren^{1,2}, Geneva Alok¹ and Joel H Rothman^{1§}

¹Department of MCD Biology and Neuroscience Research Institute, University of California Santa Barbara, CA, USA


²School of Biological Sciences, University of Auckland, Auckland, New Zealand

[§]To whom correspondence should be addressed: joel.rothman@lifesci.ucsb.edu

A

MDSKAI AERIKKLQAQQEHVRIGGGKGT**PRRKKKV**IHKTA AADDKKLQSNLKKLSVTNIPGIEEVNMIKDDG
TVIHFNNPKVQTSVPANTFSVTGSADNKQITEMLPGILNQLGPESLTHLKKLANNVTKLGPDKGGEDEDVP
ELVGDFDAASKNETKADEQ

B



C




Figure 1: A) Amino acid sequence of ICD-1. The putative nuclear localization signal is highlighted in red. B) The effects of *icd-1* RNAi on N2, *skn-1(zu67)*, *mom-2(or42)*, and *mom-4(or39)* absence-of-endoderm phenotype. At least three replicates were performed per experiment with >200 embryos scored per experiment. Student t-test (NS p-value > 0.05, ** p-value ≤ 0.01, *** p-value ≤ 0.001). C) Hypothesized model of ICD-1 function in endoderm specification, positing that it antagonizes the SKN-1 input upstream of END-1/3.

Description

The entire *C. elegans* intestine is derived from a single endodermal progenitor cell (E), the posterior daughter arising from the asymmetric division of the EMS blastomere. During early embryonic development, maternally provided SKN-1/Nrf2 activates the mesendoderm gene regulatory network (GRN) in both E and its sister, MS. A triply redundant Wnt/MAPK/Src signaling system from the neighboring P₂ blastomere polarizes EMS, resulting in activation of E fate on the side contacting it. In MS, and in an un signaled E cell, POP-1/Tcf represses expression of the redundant endoderm specifying factors, the END-1 and -3 GATA-type transcription factors. In a normal E cell, Wnt (initiated by the MOM-2/Wnt ligand) and MAPK signaling (through the MOM-4 MAPKKK) converge on POP-1 to convert it from a repressor to an activator of the *end* genes which, in collaboration with SKN-1, activates E cell fate (Thorpe *et al.* 1997; Maduro and Rothman 2002; McGhee 2007; Maduro 2017).

Basal transcription factor 3 (BTF3) facilitates transfer of nascent polypeptide chains into mitochondria and regulates transcription in plants and animals (Jamil *et al.* 2015). We previously found that the *C. elegans* BTF-3 orthologue, ICD-1, is required to prevent apoptosis: eliminating *icd-1* leads to increased cell death in embryos and larvae (Bloss *et al.* 2003). However, consistent with its potential function as a transcription factor, ICD-1 contains a putative nuclear localization signal in the N-terminus (Fig.1A) (Lange *et al.* 2007). Here, we report that ICD-1 performs a function in endoderm

specification. We found that *icd-1* RNAi does not affect endoderm specification in a wild-type N2 background or the gut-less phenotype of *skn-1(-)* embryos. However, knockdown of *icd-1* strongly suppresses the absence of gut in *mom-2/Wnt(-)* embryos (*mom-2(or42)*: 26.0% ± s.d. 4.5% with gut vs. *mom-2(or42); icd-1(RNAi)*: 75.9% ± s.d. 2.6%). Similarly, depleting ICD-1 rescues the gut-less phenotype of *mom-4/Tak1(-)* embryos (*mom-4(or39)*: 67.9% ± s.d. 6.3% with gut vs. *mom-4(or39); icd-1(RNAi)*: 95.5% ± s.d. 1.2%) (Fig. 1B). Our findings suggest a model in which ICD-1 antagonizes the SKN-1 input, perhaps by preventing SKN-1 from binding to *end-1/3* promoters. This possibility is also consistent with the finding that competition between ICD-1 and SKN-1 is seen in the context of the unfolded protein response (UPR). SKN-1 binds to and activates *hsp-4*, which codes for an endoplasmic reticulum chaperone BiP (Glover-Cutter *et al.* 2013). Depleting ICD-1 results in upregulation of *hsp-4* and activation of the UPR (Arsenovic *et al.* 2012), suggesting that ICD-1 and SKN-1 perform opposing functions in other contexts. In this hypothesized model, ICD-1 may act to fine-tune developmental signals, thereby ensuring proper specification and differentiation of endoderm (Fig. 1C).

Reagents

Strains

JJ185 *dpy-13(e184) skn-1(zu67) IV; mDp1 (IV;f)*

JR3936 *dpy-13(e184) skn-1(zu67) IV/nT1 [qIs51] (IV;V)*

EU384 *dpy-11(e1180) mom-2(or42) V/nT1 [let-(m435)] (IV;V)*.

EU414 *unc-13(e1091) mom-4(or39)/hT2 I; +/hT2 [bli-4(e937) let-(h661)] III*.

RNAi and quantification of endoderm specification

E. coli HT115 expressing *icd-1* dsRNA was obtained from the Ahringer RNAi library (Kamath *et al.* 2003). RNAi experiments for embryonic endoderm specification were performed as described (Torres Cleuren *et al.* 2019). In brief, bacteria were grown at 37°C in LB containing 50 µg/ml ampicillin. The overnight culture was then diluted 1:10. After 4 hours of incubation at 37°C, 1 mM of IPTG was added and 60 µl was seeded onto 35 mm agar plates containing 1 mM IPTG and 25 µg/ml carbenicillin. Seeded plates were allowed to dry overnight before use. 20-30 L4 or young adults were placed on the seeded RNAi plate. 24 hours later, they were transferred to a fresh RNAi plate and allowed to lay eggs for four hours. The adults were then removed, leaving the embryos to develop for an extra 5-7 hours. Embryos expressing birefringent gut granules were quantified and imaged on an agar pad using a Nikon Ti-E inverted microscope under dark field with polarized light (Clokey and Jacobson 1986; Hermann *et al.* 2005). All experiments were performed at 20°C.

Acknowledgments: Some strains used in this study were provided by the Caenorhabditis Genetics Center, which is funded by NIH Office of Research Infrastructure Programs (P40 OD010440). The authors thank members of Rothman lab, especially Pradeep M. Joshi, for helpful advice and feedback.

References

- Arsenovic P. T., A. T. Maldonado, V. D. Colletuori, and T. A. Bloss, 2012 Depletion of the *C. elegans* NAC engages the unfolded protein response, resulting in increased chaperone expression and apoptosis. *PLoS One* 7: e44038. DOI: 10.1371/journal.pone.0044038
- Bloss T. A., E. S. Witze, and J. H. Rothman, 2003 Suppression of CED-3-independent apoptosis by mitochondrial βNAC in *Caenorhabditis elegans*. *Nature* 424: 1066–1071. DOI: 10.1038/nature01920
- Clokey G. V, and L. A. Jacobson, 1986 The autofluorescent “lipofuscin granules” in the intestinal cells of *Caenorhabditis elegans* are secondary lysosomes. *Mech. Ageing Dev.* 35: 79–94. DOI: 10.1016/0047-6374(86)90068-0 | PMID: 3736133.
- Glover-Cutter K. M., S. Lin, and T. K. Blackwell, 2013 Integration of the Unfolded Protein and Oxidative Stress Responses through SKN-1/Nrf. (D. A. Garsin, Ed.). *PLoS Genet.* 9: e1003701. DOI: 10.1371/journal.pgen.1003701
- Hermann G. J., L. K. Schroeder, C. A. Hieb, A. M. Kershner, B. M. Rabbitts, et al., 2005 Genetic Analysis of Lysosomal Trafficking in *Caenorhabditis elegans*. *Mol. Biol. Cell* 16: 3273–3288. DOI: 10.1091/mbc.E05
- Jamil M., W. Wang, M. Xu, and J. Tu, 2015 Exploring the roles of basal transcription factor 3 in eukaryotic growth and development. *Biotechnol. Genet. Eng. Rev.* 31: 21–45. DOI: 10.1080/02648725.2015.1080064
- Kamath R. S., A. G. Fraser, Y. Dong, G. Poulin, R. Durbin, et al., 2003 Systematic functional analysis of the *Caenorhabditis elegans* genome using RNAi. *Nature* 421: 231–237. DOI: 10.1038/nature01278
- Lange A., R. E. Mills, C. J. Lange, M. Stewart, S. E. Devine, et al., 2007 Classical nuclear localization signals: definition, function, and interaction with importin alpha. *J. Biol. Chem.* 282: 5101–5. DOI: 10.1074/jbc.R600026200
- Maduro M. F., and J. H. Rothman, 2002 Making Worm Guts: The Gene Regulatory Network of the *Caenorhabditis elegans* Endoderm. *Dev. Biol.* 246: 68–85. DOI: 10.1006/DBIO.2002.0655

Maduro M. F., 2017 Gut development in *C. elegans*. *Semin. Cell Dev. Biol.* 66: 3–11. DOI: 10.1016/j.semcdb.2017.01.001

McGhee J., 2007 The *C. elegans* intestine. *WormBook* 1–36. <https://doi.org/10.1895/wormbook.1.133.1> DOI: 10.1895/wormbook.1.133.1

Thorpe C. J., A. Schlesinger, J. C. Carter, and B. Bowerman, 1997 Wnt signaling polarizes an early *C. elegans* blastomere to distinguish endoderm from mesoderm. *Cell* 90: 695–705. DOI: 10.1016/s0092-8674(00)80530-9 | PMID: 9288749.

Torres Cleuren Y. N., C. K. Ewe, K. C. Chipman, E. R. Mears, C. G. Wood, et al., 2019 Extensive intraspecies cryptic variation in an ancient embryonic gene regulatory network. *Elife* 8. DOI: 10.7554/eLife.48220

Funding: This research was supported by NIH grants # HD081266 and HD082347 to J.H.R.

Author Contributions: Chee Kiang Ewe: Conceptualization, Investigation, Writing - original draft, Writing - review and editing. Yamila N Torres Cleuren: Conceptualization, Funding acquisition, Investigation, Writing - review and editing. Geneva Alok: Investigation, Resources, Writing - review and editing. Joel H Rothman: Conceptualization, Funding acquisition, Writing - review and editing.

Reviewed By: Anonymous

History: Received September 3, 2019 Accepted September 26, 2019 Published October 4, 2019

Copyright: © 2019 by the authors. This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International (CC BY 4.0) License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Citation: Ewe, CK; Torres Cleuren, YN; Alok, G; Rothman, JH (2019). ICD-1/BTF3 antagonizes SKN-1-mediated endoderm specification in *Caenorhabditis elegans*. *microPublication Biology*. <https://doi.org/10.17912/micropub.biology.000167>