

Short report

Open Access

## Activation of Erk in the anterior cingulate cortex during the induction and expression of chronic pain

Feng Wei\*<sup>1</sup> and Min Zhuo\*<sup>2</sup>

Address: <sup>1</sup>Department of Biomedical Sciences, University of Maryland Dental School, Baltimore, Maryland 20201, USA and <sup>2</sup>Department of Physiology, University of Toronto, Medical Sciences Bldg., Rm 3342, 1 King's College Circle, Toronto, Ontario, M5S 1A8, Canada

Email: Feng Wei\* - [fwei@umaryland.edu](mailto:fwei@umaryland.edu); Min Zhuo\* - [min.zhuo@utoronto.ca](mailto:min.zhuo@utoronto.ca)

\* Corresponding authors

Published: 23 July 2008

Received: 12 June 2008

*Molecular Pain* 2008, **4**:28 doi:10.1186/1744-8069-4-28

Accepted: 23 July 2008

This article is available from: <http://www.molecularpain.com/content/4/1/28>

© 2008 Wei and Zhuo; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

### Abstract

The extracellular signal-regulated kinase (Erk) activity contributes to synaptic plasticity, a key mechanism for learning, memory and chronic pain. Although the anterior cingulate cortex (ACC) has been reported as an important cortical region for neuronal mechanisms underlying the induction and expression of chronic pain, it has yet to be investigated whether or not Erk activity in the ACC may be affected by peripheral injury or in chronic pain state. In the present study, we use adult rat animal models of inflammatory and neuropathic pain and demonstrate that Erk signaling pathway in the ACC is potently activated after peripheral tissue or nerve injury. Furthermore, we demonstrate that mechanical allodynia significantly activated Erk activity at synaptic sites at two weeks after the injury. We propose a synaptic model for explaining the roles of Erk activity during different phases of chronic pain. Our findings suggest that cortical activation of Erk may contribute to both induction and expression of chronic pain.

### Short report

Recent studies have consistently indicated that Erk signaling cascade plays an important role in activity-dependent plasticity in the central nervous system (CNS) and may contribute to the molecular mechanisms underlying learning, memory and persistent pain. Previous studies have found that tissue and nerve injury transiently activates Erk pathway in the spinal dorsal horn neurons, and the activation of Erk is required for the central sensitization during the development of hyperalgesia and allodynia [1-5]. In supraspinal structures, it has been reported that activation of Erk in the amygdala is induced by peripheral injury [6], and the increased Erk activity in this region is acquired for behavioral sensitization to mechanical stimulation after injury [7]. The ACC has been found to be an important site for cortical regulation of nociception and persistent pain after amputation [8-11]. Long-

term potentiation (LTP) in ACC neurons is the likely synaptic model for persistent pain [12-16]. Recent studies using pharmacological inhibitors has showed that the activity of Erk contributes to synaptic potentiation caused by LTP induction protocols [17], and that such inhibitors are relatively selective and do not affect basic synaptic transmission. However, little is known about the possible involvement of Erk in the ACC after tissue or nerve injury in adult animals.


To determine the possible activation of Erk in the ACC during acute or chronic pain after peripheral injury, we carried out experiments in adult male rats using two different injury models. Activation of Erk was monitored by immunostaining with an antibody that detects the activated form of Erk (phosphorylated on both thr-202 and Try-204, P-Erk). For the first pain model, 5% formalin was

injected into the dorsal part of the unilateral hindpaw as previously reported [18]. At three different time points between 15 min to 90 min after the formalin injection, rats were killed by rapidly anesthetized and perfused through the heart with fixative. We found that formalin injection induced a rapid increase in P-Erk expression in some of layer II neurons in the bilateral ACC at 15 min after tissue injury (Fig. 1A). The expression of P-Erk was mainly restricted in the cell bodies but weak in their dendrites (Fig. 1A). The activated Erk level in the ACC was reduced at 45 min and declined at 90 min after formalin injection. There was no obvious Erk activation in the deep layer neurons in the ACC. Thus, the P-Erk expression pattern in the region is different from that of immediate early genes, such as c-Fos, which is widely expressed in ACC neurons located in all layers after formalin injection as previously reported [18]. These findings suggest that Erk activation caused by tissue injury is likely occurring at sub-population of the layer II neurons. It has been known that neurons in the Layer II/III receive ascending noxious inputs from the thalamus and communicate with other cortical areas [19]. Thus, the Erk activation in ACC neurons may play a role in Erk-dependent neuronal plasticity in the ACC during the induction or development of inflammatory pain.


Phantom pain is chronic pain occurring after losing or amputation of a part of limb or organ. Although the animal model for studying central mechanisms of phantom pain is rare, investigations of changes in central synapses in the ACC after amputation in animals may reveal molecular mechanism for plastic changes that are caused by amputation. We have previously demonstrated that amputation of a single digit resulted in a loss of synaptic cortical depression [20,9,11] in brain slice preparations, and caused LTP of synaptic responses in the ACC to peripheral sensory stimulation or local synaptic stimulation [11,21], demonstrating that central plasticity takes place in the synapses of ACC neurons after the amputation. Because the Erk activity is shown to be required for cingulate LTP [17], we decided to test if tissue and nerve injury after the amputation would activate the Erk pathway in the ACC neurons. As shown in Fig. 1B, we found that larger number of the layer II pyramidal-like neurons in the ACC expressed P-Erk immunoreactivity at two weeks after single digit amputation. Activation of Erk activity in the ACC is bilateral. Interestingly, the P-Erk expression in these neurons was present in cytoplasm of both cell body and dendrites (Fig. 1B). There was some P-Erk in neurons of the deeper layers, suggesting the possible interaction between neurons in different layers of the ACC. This data further indicates that there is a prolonged activation of Erk in the layer II neurons after amputation.

In addition to spontaneous pain and hyperalgesia, allodynia is the most common feature of pathological pain and is a painful response to a usually innocuous stimulus. Touch-evoked allodynia occurs often in patients with phantom pain after amputation. Most of previous studies focus on the activation of Erk at early time points after the injury, there is few studies for the possible involvement of Erk activity during allodynic stimulation. Therefore, we applied non-noxious mechanical stimuli (brushing) on amputated hindpaw in rats to identify if Erk signaling in the ACC may be activated. To our surprise, there was significantly enhanced activation of Erk in the ACC neurons after brushing the amputated hindpaw. In addition to the increased number of P-Erk labeled neurons in the layer II, we also observed that there was stronger expression of P-Erk in the dendrites distributed in the layer I (Fig. 1C), as compared with that in rats with amputation alone. Most of P-Erk labeled pyramidal neurons in the layer II exhibited strong immunoreactivity in the main and distal apical dendrites, which branched upward to the superficial layer I of the ACC (see Fig. 1C). We also observed that there were many P-Erk labeled granular-like neurons in the deeper layers of the ACC. Again, we found similar activation pattern at bilateral sides of the ACC. By contrast, non-noxious brushing alone in normal rats did not cause any detectable Erk activation in the ACC. The enhanced expression of P-Erk in ACC neurons and their dendrites after brushing normal skin of amputated hindpaw suggests that Erk activity are recruited at distal synapses in ACC neurons. It may contribute to local synaptic plasticity and/or neuronal modulation during allodynia after amputation.

Our results provide the first evidence for the activation of Erk activity in the ACC neurons after tissue or nerve injury. More importantly, we show that the enhanced activation of Erk activity in synaptic sites in the layer II/III neurons during the touch-evoked allodynia after the amputation; suggesting the likely contribution of Erk activity to the central mechanisms underlying pathological pain. Different activation pattern of Erk in the inflammatory and amputation pain model suggest that Erk may contribute to different types of chronic pain in different manners. This is further supported by our observation of distal dendrite Erk activation during mechanical allodynia. Fig. 2 is a proposed synaptic model showing different subcellular location of P-Erk during two different phases of chronic pain: induction and expression of mechanical allodynia. The expression of p-Erk in the cytoplasm near the nucleus during the induction of chronic pain may play roles in the activation of a series of plasticity-related immediate early genes such as cAMP response element binding protein (CREB) [16]. In this case, P-Erk may trigger a series of plasticity-related signaling molecules that are important for inducing plastic changes in the ACC. In the expression

**Figure 1**

**Enhanced Erk activation in the ACC after tissue and nerve injury.** **A.** Immunohistochemical staining for phosphorylation of Erk illustrated time course-dependent activation of Erk in layer II neurons of the contralateral ACC after unilateral hindpaw injection of formalin (5%, 50 µl, n = 4–5 rats for each time point). **B.** The P-Erk expression in the layer II ACC neurons and their main apical dendrites (arrows) was increased at 2 weeks after the amputation of the unilateral hindpaw third digit (n = 5), compared to sham animals (n = 3). **C.** Mechanical stimulation by brushing hindpaw of digit amputation induced P-Erk expression in more number of layer II ACC neurons and the more distinctive apical dendrites at 2 weeks after the amputation (n = 5), compared to that in rats with amputation alone. There was not P-Erk activation in the ACC in normal animals after the brushing (n = 3). Left and middle columns: low power of the coronal ACC sections. Scale bar= 50 µm; Right column: enlarged layer II regions corresponding to the small rectangle areas in the middle column, respectively. Scale bar = 25 µm.

**Figure 2**

**A model for activation of Erk in the induction and expression phases of chronic pain. A.** In the induction phase of chronic pain, peripheral injury triggers glutamate release in the ACC synapses. Activation of NMDA receptors leads to an increase in postsynaptic  $\text{Ca}^{2+}$ , and  $\text{Ca}^{2+}$  binds to CaM and leads to activation of  $\text{Ca}^{2+}$ -stimulated AC1. cAMP, a key second messenger, leads to activation of Erk, and PKA-dependent CREB. New protein synthesis is likely triggered as a consequence of CREB activation. **B.** In the expression phase of chronic pain (allodynia), a non-noxious stimuli triggers glutamate release in sensitized ACC synapses. Due to synaptic enhancement caused during the induction phase [13], glutamate triggers greater postsynaptic activation because AMPA and NMDA receptor mediated responses are likely enhanced after the injury [13,23]. Such postsynaptic sensitization makes activation of Erk at distal synaptic sites possible. Activated Erk at synaptic sites may contribute to AMPA receptor modulation, ion channel modulation and other synaptic modifications.

phase of chronic pain (or mechanical allodynia here), in part due to prolonged plastic changes triggered in the induction phase, a normally non-noxious stimulus (i.e., non-noxious brushing of the skin) triggers a wide spread activation of Erk in ACC neurons. In addition to the cytoplasm, activation of Erk at synaptic sites is found. These activated Erk may contribute to rapid synaptic potentiation [17], the regulation of neuronal excitability [5], and other local modulations of synaptic transmission and neuronal excitability (see Fig. 2).

Although we cannot distinguish the contribution of Erk activities to pain perception, pain-related emotional responses or pain-related memory in the present study, we believe that such increased Erk activity may contribute to long-term plastic changes in the ACC caused by the injury or amputation. In fact, in brain slices, we demonstrate that Erk activity is required for synaptic potentiation in the ACC excitatory synapses [17]. One major function of Erk may contribute to cortical plasticity during the expression of behavioral allodynia, and serve as one of key signaling protein kinase in chronic pain [16].

## Materials and methods

### Animals

Adult male Sprague Dawley rats weighing 250–300 gram were used. Animals were kept in cages (2 animals per cage) at an ambient temperature of 20–25°C under a 12 hr light/dark cycle and had free access to food and water. We adhered to the ethical guidelines for investigation of experimental pain in conscious animals [22].

### Animal models of chronic pain

For inflammatory pain model, fifty microliters of a 5% formalin solution (dissolved in saline) were injected into the plantar surface of the left hindpaw. For rat model of amputation, under brief anesthesia with halothane, the third digit of the left hindpaw was amputated. In some cases, at 2 week after the amputation, a brush stimulus (with number 4 camel's hair artist's brush) was applied for 3 min by stroking vertically at the dorsal part of amputated hindpaw. At 15 min, 45 min, 90 min and 120 min after formalin injection, 2 week after the amputation, or 15 min after brush stimulus, rats were deeply anesthetized with halothane and perfused transcardially with 100 ml of saline, followed by 500 ml of cold 0.1 M phosphate buffer containing 4% paraformaldehyde. The brain was removed, post-fixed for 4 hr, and then cryoprotected. Coronal sections (25 µm) through the ACC were cut using a cryostat. Sham groups without formalin injection or amputation were performed as controls. Sections from sham and experimental animals were processed simultaneously for immunostaining.

### Immunohistochemistry

Immunostaining was performed using free-floating sections [4]. Briefly, the ACC sections were first treated with 0.75% Triton X-100 and 1% H<sub>2</sub>O<sub>2</sub> in PBS for 1 hr, and then processed for 30 min in 3% normal goat serum, followed by incubation with anti-phospho-p44/42 Erk (Thr202/Tyr204) monoclonal antibody (diluted 1:500; Cell Signaling, Beverly, MA) overnight at room temperature. Secondary reactions with biotinylated goat anti-mouse immunoglobulin (1:400; Vector Laboratories, Burlingame, CA) for 1 h were followed by avidin-biotin-peroxidase complexes (1:100; Vector Laboratories) for 1 h. A nickel-intensified diaminobenzidine with glucose oxidase was used as the final chromogen. Sections were washed several times, mounted on gelatinized slides, dehydrated through a series of ethanol solutions, cleared in xylene, and covered with glass coverslips. Controls, performed by replacing primary antibody with 1% NGS in the protocol, exhibited no staining.

### Authors' contributions

The authors declare that they have no competing interests. FW and MZ participated in the conception, design, and interpretation of the study. FW carried out the experiments, FW and MZ wrote the manuscript.

### Acknowledgements

This work was supported by grants from the EJLB-CIHR Michael Smith Chair in Neurosciences and Mental Health, Canada Research Chair, CIHR84256, CIHR66975 and NIH NINDS NS42722 to Dr. Min Zhuo.

### References

1. Ji RR, Baba H, Brenner GJ, Woolf CJ: **Nociceptive-specific activation of ERK in spinal neurons contributes to pain hypersensitivity.** *Nat Neurosci* 1999, **2**:1114-1119.
2. Karim F, Wang CC, Gereau IV: **Metabotropic glutamate receptor subtypes 1 and 5 are activators of extracellular signal-regulated kinase signaling required for inflammatory pain in mice.** *J Neurosci* 2001, **21**:3771-3779.
3. Hu HJ, Gereau IV: **ERK integrates PKA and PKC signaling in superficial dorsal horn neurons. II. Modulation of neuronal excitability.** *J Neurophysiol* 2003, **90**:1680-1688.
4. Wei F, Vadakkan KI, Toyoda H, Wu LJ, Zhao MG, Xu H, Shum FW, Jia YH, Zhuo M: **Calcium calmodulin-stimulated adenylyl cyclases contribute to activation of extracellular signal-regulated kinase in spinal dorsal horn neurons in adult rats and mice.** *J Neurosci* 2006, **26**:851-861.
5. Hu HJ, Alter BJ, Carrasquillo Y, Qiu CS, Gereau IV: **Metabotropic glutamate receptor 5 modulates nociceptive plasticity via extracellular signal-regulated kinase-Kv4.2 signaling in spinal cord dorsal horn neurons.** *J Neurosci* 2007, **27**:13181-13191.
6. Carrasquillo Y, Gereau IV: **Activation of the extracellular signal-regulated kinase in the amygdala modulates pain perception.** *J Neurosci* 2007, **27**:1543-1551.
7. Carrasquillo Y, Gereau IV: **Hemispheric lateralization of a molecular signal for pain modulation in the amygdala.** *Mol Pain* 2008, **4**:24.
8. Calejesan AA, Kim SJ, Zhuo M: **Descending facilitatory modulation of a behavioral nociceptive response by stimulation in the adult rat anterior cingulate cortex.** *Eur J Pain* 2000, **4**:83-96.
9. Wei F, Li P, Zhuo M: **Loss of synaptic depression in mammalian anterior cingulate cortex after amputation.** *J Neurosci* 1999, **19**:9346-9354.
10. Wei F, Wang GD, Zhang C, Shokat KM, Wang H, Tsien JZ, Liauw J, Zhuo M: **Forebrain overexpression of CaMKII abolishes cin-**

- gulate long term depression and reduces mechanical allodynia and thermal hyperalgesia. *Mol Pain* 2006, **2**:21.
11. Wei F, Zhuo M: **Potentiation of sensory responses in the anterior cingulate cortex following digit amputation in the anaesthetised rat.** *J Physiol* 2001, **532**:823-833.
  12. Tang J, Ko S, Ding HK, Qiu CS, Calejesan AA, Zhuo M: **Pavlovian fear memory induced by activation in the anterior cingulate cortex.** *Mol Pain* 2005, **1**:6.
  13. Zhao MG, Ko SW, Wu LJ, Toyoda H, Xu H, Quan J, Li J, Jia Y, Ren M, Xu ZC, Zhuo M: **Enhanced presynaptic neurotransmitter release in the anterior cingulate cortex of mice with chronic pain.** *J Neurosci* 2006, **26**:8923-8930.
  14. Zhao MG, Toyoda H, Lee YS, Wu LJ, Ko SW, Zhang XH, Jia Y, Shum F, Xu H, Li BM, Kaang BK, Zhuo M: **Roles of NMDA NR2B subtype receptor in prefrontal long-term potentiation and contextual fear memory.** *Neuron* 2005, **47**:859-872.
  15. Zhuo M: **A synaptic model for pain: long-term potentiation in the anterior cingulate cortex.** *Mol Cells* 2007, **23**:259-271.
  16. Zhuo M: **Cortical excitation and chronic pain.** *Trends Neurosci* 2008, **31**:199-207.
  17. Toyoda H, Zhao MG, Xu H, Wu LJ, Ren M, Zhuo M: **Requirement of extracellular signal-regulated kinase/mitogen-activated protein kinase for long-term potentiation in adult mouse anterior cingulate cortex.** *Mol Pain* 2007, **3**:36.
  18. Wei F, Wang GD, Kerchner GA, Kim SJ, Xu HM, Chen ZF, Zhuo M: **Genetic enhancement of inflammatory pain by forebrain NR2B overexpression.** *Nat Neurosci* 2001, **4**:164-169.
  19. Wang CC, Shyu BC: **Differential projections from the medio-dorsal and centrolateral thalamic nuclei to the frontal cortex in rats.** *Brain Res* 2004, **995**:226-235.
  20. Zhuo M: **NMDA receptor-dependent long term hyperalgesia after tail amputation in mice.** *Eur J Pharmacol* 1998, **349**:211-220.
  21. Wu MF, Pang ZP, Zhuo M, Xu ZC: **Prolonged membrane potential depolarization in cingulate pyramidal cells after digit amputation in adult rats.** *Mol Pain* 2005, **1**:23.
  22. Zimmermann M: **Ethical guidelines for investigations of experimental pain in conscious animals.** *Pain* 1983, **16**:109-110.
  23. Wu LJ, Toyoda H, Zhao MG, Lee YS, Tang J, Ko SW, Jia YH, Shum FW, Zerbiniatti CV, Bu G, Wei F, Xu TL, Muglia LJ, Chen ZF, Auber-son YP, Kaang BK, Zhuo M: **Upregulation of forebrain NMDA NR2B receptors contributes to behavioral sensitization after inflammation.** *J Neurosci* 2005, **25**:11107-11116.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:  
[http://www.biomedcentral.com/info/publishing\\_adv.asp](http://www.biomedcentral.com/info/publishing_adv.asp)

