

Projected estimates of cancer in Canada in 2024

Darren R. Brenner PhD, Jennifer Gillis PhD, Alain A. Demers MSc PhD, Larry F. Ellison MSc, Jean-Michel Billette PhD, Shary Xinyu Zhang MSc, JiaQi Leon Liu MPH, Ryan R. Woods PhD, Christian Finley MD MPH, Natalie Fitzgerald MA, Nathalie Saint-Jacques PhD, Lorraine Shack PhD, Donna Turner PhD; for the Canadian Cancer Statistics Advisory Committee

■ Cite as: *CMAJ* 2024 May 13;196:E615-23. doi: 10.1503/cmaj.240095

See related editorial at www.cmaj.ca/lookup/doi/10.1503/cmaj.240612

Abstract

Background: Cancer surveillance data are essential to help understand where gaps exist and progress is being made in cancer control. We sought to summarize the expected impact of cancer in Canada in 2024, with projections of new cancer cases and deaths from cancer by sex and province or territory for all ages combined.

Methods: We obtained data on new cancer cases (i.e., incidence, 1984–2019) and deaths from cancer (i.e., mortality, 1984–2020) from the Canadian Cancer Registry and Canadian Vital Statistics Death Database, respectively. We projected cancer incidence and mortality counts and rates to 2024 for 23 types of cancer, overall, by sex, and by prov-

ince or territory. We calculated age-standardized rates using data from the 2011 Canadian standard population.

Results: In 2024, the number of new cancer cases and deaths from cancer are expected to reach 247 100 and 88 100, respectively. The age-standardized incidence rate (ASIR) and mortality rate (ASMR) are projected to decrease slightly from previous years for both males and females, with higher rates among males (ASIR 562.2 per 100 000 and ASMR 209.6 per 100 000 among males; ASIR 495.9 per 100 000 and ASMR 152.8 per 100 000 among females). The ASIRs and ASMRs of several common cancers are projected to continue to decrease (i.e.,

lung, colorectal, and prostate cancer), while those of several others are projected to increase (i.e., liver and intrahepatic bile duct cancer, kidney cancer, melanoma, and non-Hodgkin lymphoma).

Interpretation: Although the overall incidence of cancer and associated mortality are declining, new cases and deaths in Canada are expected to increase in 2024, largely because of the growing and aging population. Efforts in prevention, screening, and treatment have reduced the impact of some cancers, but these short-term projections highlight the potential effect of cancer on people and health care systems in Canada.

Cancer has a substantial impact on the lives of people living in Canada and on Canadian health care systems.^{1,2} Previous estimates suggest that 45% of all people in Canada will receive a cancer diagnosis in their lifetime.^{3,4} As the population grows and ages, new cancer cases and deaths from cancer in Canada are also increasing.^{5,6} In addition, cancer has a major economic impact. The most recent estimates indicated that the economic burden of cancer in Canada was \$26.2 billion in 2021, with 30% of costs being carried by patients and their families.⁷

Because of the time required for collecting and verifying data, cancer data necessarily lag several years. Short-term projections are estimated by extrapolating past trends into the future using statistical models. These projections of incidence and mortality provide a timely picture of the impact of cancer in Canada that is necessary for resource planning, research, and informing cancer control programs. Given the overall impact of cancer on the lives and economic livelihood of people in Canada, such estimates are

essential to highlight where progress has been made in cancer control and where more resources and effort are required.

The *Canadian Cancer Statistics 2023* included detailed estimates of cancer metrics, including incidence, mortality, survival by sex assigned at birth, age, and province or territory, as well as time trends, for 23 cancer types, from 1984 to 2023.³ We sought to update these analyses and determine projections of the counts and age-standardized rates of new cancer cases and deaths expected in 2024, by sex and by province or territory for all ages combined.

Methods

Study design

We conducted an analysis of population-level cancer data with updated projected estimates from the Canadian Cancer Statistics Advisory Committee, a collaboration between Statistics Canada,

the Public Health Agency of Canada, and the Canadian Cancer Society. The advisory committee includes representation from each of these organizations, as well as representatives from the Canadian Partnership Against Cancer, provincial cancer registries, and academic collaborators. Analytical and methodological decisions are made collaboratively and agreed upon by the committee based on documented approaches.³ Unless otherwise noted, the data sources and methodical decisions used in this study are as described in detail in *Canadian Cancer Statistics 2023*.³

Sources of data

We obtained data on cancer incidence from the National Cancer Incidence Reporting System (1984–1991) and the Canadian Cancer Registry Tabulation Master File (1992–2019), released May 16, 2023.⁸ We obtained data on deaths from the Canadian Vital Statistics Death Database (1984–2020), released Jan. 24, 2022.⁹ These national population-based databases are populated by data submissions from provincial and territorial cancer registries to Statistics Canada, where these databases are maintained. We also obtained actual and projected population estimates from Statistics Canada.^{10–12}

Statistical analysis

We projected counts and rates from 2020 to 2024 using the CANPROJ projection package to generate estimates of cancer incidence and mortality.¹³ The CANPROJ package uses trends in actual (i.e., historical) data to select the best-fit model for subsequent years based on a decision algorithm that compares 6 models, including the Nordpred (age–period–cohort drift), age–cohort, hybrid age-specific trend, hybrid age-common trend, hybrid age-only, and 5-year average models. Model selection is based on statistical tests for each model, starting with the most complex (Nordpred). Based on specific rules, a simpler model may be selected. Additional details on the CANPROJ modelling approach and model selection are provided in Appendix 1, available at www.cmaj.ca/lookup/doi/10.1503/cmaj.240095/tab-related-content. Validation of the CANPROJ software has been completed using data on cancer incidence in Canada (excluding Quebec).¹⁴

The Canadian Cancer Statistics Advisory Committee reviewed each model selected by the CANPROJ software, alongside the 5 other models produced, to suggest the most appropriate model for each jurisdiction. The committee is guided by a set of principles, along with use of a decision tree, in making its recommendation (Appendix 1). Decisions on the final models for each province or territory were made by consensus in consultation with that jurisdiction's cancer registry. For these analyses, we used the same models selected for *Canadian Cancer Statistics 2023*.³

For Quebec, we calculated incidence projections from 2018 to 2024 as cancer cases diagnosed in this province after 2017 had yet to be submitted to the Canadian Cancer Registry. In addition, Quebec transitioned from a registry exclusively based on hospital discharge data to a more integrated central registry, resulting in an artifactual increase for some type-specific cancer counts during the early years of the transition (2011–2012). Consequently, for these 2 diagnosis years, for some age- and sex-

specific rates in Quebec, we used estimates derived from locally estimated scatterplot smoothing to the 2006–2017 reference period (Appendix 1). We generated incidence projections for Nova Scotia from 2019 to 2024, because the data for 2019 (i.e., submission of cases from the province to the Canadian Cancer Registry) required further verification.

We generated projected estimates for 23 cancer types by sex assigned at birth (hereafter referred to as sex) and geographic region (province or territory). Definitions of cancer types are in Appendix 1, Supplementary Table 1.^{15–17} We computed projected estimates for Canada as sums of the projections for each individual province and territory. All incidence and mortality rates were age-standardized to the 2011 Canadian standard population using the direct method.¹⁸

Ethics approval

Because this study involved the analysis of administrative data publicly available in Statistics Canada's Research Data Centres and did not involve contacting individuals, consideration and approval by an ethics review board were not required.

Results

Incidence in 2024

In 2024, 247 100 new cancer cases are expected to be diagnosed (Table 1). Lung cancer is projected to remain the most diagnosed cancer in Canada, with 32 100 new cases expected in 2024. Breast (30 800 cases), prostate (27 900 cases) and colorectal (25 200 cases) cancers are expected to be the next most common cancers. Combined, these 4 cancer types are 47% of all new cancer cases expected in 2024.

Among males, the most commonly diagnosed cancers are expected to be prostate (22%), lung (12%), colorectal (11%), and bladder (7%) cancers. Among females, breast (25%), lung (14%), colorectal (9%), and uterine (7%) cancers are expected to be the most commonly diagnosed cancers. Apart from sex-specific cancers, cancer is more commonly diagnosed among males than females, except for breast, thyroid, and lung cancers, which are more common among females (Table 1). Overall, the age-standardized incidence rate (ASIR) for all cancers combined is expected to be 13% higher among males than females (562.2 v. 495.9 per 100 000) in 2024.

Mortality in 2024

In 2024, 88 100 people in Canada are expected to die of cancer (Table 2). Lung cancer is expected to remain the most common cause of cancer death, with an expected 20 700 deaths in 2024. These deaths will account for almost one-quarter of all cancer deaths in Canada. Colorectal (9400 deaths), pancreas (6100 deaths), breast (5500 deaths), and prostate (5000 deaths) are projected to be the next most common causes of cancer death. Combined, these 5 cancers are expected to account for more than half (53%) of all cancer deaths in Canada in 2024.

For both males and females, lung cancer accounts for the highest number and proportion of cancer deaths at 10 900 (23%) and 9800 (24%) projected deaths, respectively. Colorectal (11%),

Table 1: Projected estimates of new cases and age-standardized incidence rates (ASIRs) for cancers in Canada, by sex, 2024

Type of cancer*	No. of new cases			ASIR per 100 000†		
	Total‡	Males	Females	Both sexes	Males	Females
All cancers§	247 100	127 100	120 000	523.9	562.2	495.9
Lung and bronchus	32 100	14 800	17 300	63.8	63.4	64.9
Breast	30 800	290	30 500	69.5	1.3	133.1
Prostate	27 900	27 900	NA	NA	119.7	NA
Colorectal	25 200	14 100	11 100	53.1	63.0	44.2
Bladder	12 300	9300	3000	24.6	40.4	11.4
Non-Hodgkin lymphoma	11 700	6600	5100	24.8	29.8	20.3
Melanoma	11 300	6500	4800	24.5	29.4	20.6
Kidney and renal pelvis	9000	5900	3100	19.5	26.9	12.7
Uterus (body, NOS)	8600	NA	8600	NA	NA	36.7
Head and neck	8100	5800	2300	17.5	26.4	9.4
Pancreas	7100	3800	3300	14.5	16.8	12.4
Thyroid	6600	2000	4700	16.2	9.6	22.7
Leukemia	6600	4000	2700	14.1	17.9	10.7
Liver and intrahepatic bile duct	4700	3300	1450	9.6	14.2	5.5
Multiple myeloma	4100	2400	1750	8.4	10.5	6.6
Stomach	4000	2600	1400	8.3	11.3	5.6
Brain or CNS	3300	1850	1400	7.3	8.8	6.0
Ovary	3000	NA	3000	NA	NA	12.9
Esophagus	2700	2100	630	5.6	9.3	2.4
Soft tissue (including heart)	1700	960	730	3.8	4.5	3.3
Cervix	1600	NA	1600	NA	NA	7.8
Testis	1300	1300	NA	NA	6.7	NA
Hodgkin lymphoma	1200	650	530	2.9	3.2	2.6
All other cancers	22 100	11 000	11 200	46.0	49.3	43.9

Note: CNS = central nervous system, NA = not applicable, NOS = not otherwise specified.

*The complete definition of the specific cancers included here can be found in Appendix 1, Supplementary Table 1.

†Rates are age-standardized to the 2011 Canadian standard population.

‡Total may not be sum of males and females because of rounding.

§Includes in situ bladder cancer and excludes non-melanoma skin cancer (neoplasms, NOS; epithelial neoplasms, NOS; and basal and squamous skin cancer).

prostate (11%), pancreas (7%), and liver and intrahepatic bile duct (5%) cancers are the next most common causes of cancer death among males, while among females, the next leading causes of cancer death are breast (13%), colorectal (10%), pancreas (7%), and ovarian (5%) cancers.

The age-standardized mortality rate (ASMR) is expected to be markedly higher (37% higher) among males than females (209.6 v. 152.8 per 100 000, respectively) in 2024. Apart from sex-specific cancers, males are expected to have higher mortality for all cancer types except for breast cancer (Table 2).

Incidence over time

The trends in ASIR for the most common cancer types for both males and females are presented in Figure 1. Over the study period, ASIRs for all cancers combined and the most common cancer types have been higher among males than females.

Notably, however, incidence rates of lung cancer have converged for males and females, with females expected to have slightly higher incidence of lung cancer in 2024. Incidence rates of colorectal cancer have been declining in recent years for both males and females. In 2024, the incidence of colorectal cancer is expected to be 26% and 31% lower than in 1984 among males and females, respectively.

Mortality over time

The trends in ASMR for the most common causes of cancer death for both males and females are presented in Figure 2. For all cancers combined and the most common causes of cancer death, ASMRs have been higher among males than females over time. Mortality from lung cancer has been declining among males since its peak in 1988, with steady decline thereafter. For females, this peak occurred in 2006 and the rate has since declined.

Table 2: Projected estimates of deaths and age-standardized mortality rates (ASMRs) for cancers by sex, Canada, 2024

Type of cancer*	No. of deaths			ASMR per 100 000†		
	Total‡	Males	Females	Both sexes	Males	Females
All cancers	88 100	47 300	40 800	177.5	209.6	152.8
Lung and bronchus	20 700	10 900	9800	40.7	46.8	35.9
Colorectal	9400	5300	4200	19.2	23.7	15.3
Pancreas	6100	3200	2900	12.2	13.9	10.7
Breast	5500	60	5500	11.8	0.3	21.8
Prostate	5000	5000	NA	NA	22.7	NA
Liver and intrahepatic bile duct§	3700	2300	1350	7.4	10.1	5.0
Leukemia	3200	1850	1350	6.4	8.2	4.9
Non-Hodgkin lymphoma	3100	1800	1300	6.2	8.1	4.7
Bladder	2600	1900	740	5.1	8.5	2.6
Brain or CNS	2600	1500	1100	5.6	6.8	4.5
Esophagus	2500	1900	560	5.0	8.3	2.1
Head and neck	2100	1550	590	4.4	6.9	2.2
Stomach	2000	1250	760	4.1	5.6	2.8
Ovary	2000	NA	2000	NA	NA	7.8
Kidney and renal pelvis	1950	1250	670	3.9	5.7	2.4
Multiple myeloma	1750	1000	720	3.5	4.5	2.6
Uterus (body, NOS)	1600	NA	1600	NA	NA	6.0
Melanoma	1300	850	440	2.7	3.8	1.7
Soft tissue including heart	660	360	300	1.4	1.7	1.2
Cervix	400	NA	400	NA	NA	1.8
Thyroid	280	130	150	0.6	0.6	0.5
Hodgkin lymphoma	110	75	40	0.2	0.3	0.1
Testis	30	30	NA	NA	0.2	NA
All other cancers	9600	5200	4400	19.1	23.2	15.8

Note: CNS = central nervous system, NA = not applicable, NOS = not otherwise specified.
 *The complete definition of the specific cancers included here can be found in Appendix 1, Supplementary Table 1.
 †Rates are age-standardized to the 2011 Canadian standard population.
 ‡Total may not be sum of males and females because of rounding.
 §Liver cancer mortality was underestimated because deaths from liver cancer, unspecified (code C22.9 in the *International Classification of Diseases, 10th edition*), were excluded.

Notable trends outside of most common cancers

Over time, incidence and mortality for liver and intrahepatic bile duct cancers have been increasing among both males and females (Figure 1B and Figure 2B). Incidence of kidney and renal pelvis cancer has also been increasing, particularly among males, although mortality has decreased in recent years. Incidence of melanoma continues to increase for both males and females, although mortality has remained largely stable. Although relatively stable among females, incidence of head and neck cancer among males decreased considerably from 1984 to 2005, and has been rising since. Mortality from head and neck cancer has decreased among males and females.

Non-Hodgkin lymphoma is expected to be the fifth most diagnosed cancer, accounting for 5% of new cancers among males and 4% among females. Incidence of non-Hodgkin lymphoma

increased until the mid-2010s, more among males than females (Figure 1B). Conversely, mortality has been declining for this cancer type over the last 2 decades for both sexes (Figure 2B).

Incidence and mortality across Canada

Projected ASIRs and counts of new cancer cases for 2024 are provided by cancer type and region in Appendix 2, Supplementary Table 2 and Table 3, respectively, available at www.cmaj.ca/lookup/doi/10.1503/cmaj.240095/tab-related-content; ASMRs and cancer deaths in 2024 are provided in Appendix 2, Supplementary Table 4 and Table 5, respectively. Both ASIRs and ASMRs showed considerable variability across geographic regions. For all cancers combined, ASIRs are generally higher in eastern Canada (Appendix 3, Supplementary Figure 1A, available at www.cmaj.ca/lookup/doi/10.1503/cmaj.240095/tab-related-content).




Figure 1: Age-standardized incidence rates for (A) the most common cancers (lung, colorectal, prostate, breast) and (B) additional cancers (liver and hepatic bile duct, melanoma, kidney and renal pelvis, non-Hodgkin lymphoma, head and neck) in Canada, by sex, 1984–2024. Data from 1984 to 1991 were from the National Cancer Incidence Reporting System and data from 1992 to 2019 were from the Canadian Cancer Registry. Analyses were conducted by Statistics Canada at the Centre for Population Health Data. Projected data for 2020–2024 are shown in shaded areas.




Figure 2: Age-standardized mortality rates for (A) the most common cancers (lung, colorectal, prostate, breast, pancreas) and (B) additional cancers (liver and hepatic bile duct, melanoma, kidney and renal pelvis, non-Hodgkin lymphoma, head and neck), in Canada, by sex, 1984–2024. Data from 1984 to 2020 were from the Canadian Vital Statistics Death Database. Analyses were conducted by Statistics Canada at the Centre for Population Health Data. Projected data for 2020–2024 are shown in shaded areas.

Projected ASMRs are generally lower in western Canada (Appendix 3, Supplementary Figure 1B). We also observed notable differences in projected ASIRs and ASMRs for specific cancer types across provinces and territories, as shown in Appendix 2, Supplementary Table 2 and Table 4. For the most part, overall ASIRs are higher in the Atlantic and eastern provinces than in western provinces, driven by differences in common cancers such as lung, colorectal, and breast cancer.

Interpretation

Although the number of cancer cases in 2024 is expected to rise compared with previous years, overall cancer incidence and mortality continue to decline. These declines are likely owing to continued efforts across the cancer control spectrum and improvements in prevention, screening, and treatment, particularly for certain cancers where concerted efforts have occurred.¹⁹

The overall survival among people with cancer continues to improve,^{20–22} contributing to a growing number of people living with and beyond cancer. Our estimates in 2022 suggested that more than 1.5 million people in Canada were living with and beyond cancer up to 25 years after diagnosis.²³ This growing population of cancer survivors, combined with the projected annual increase in the total number of new primary cancers in 2024, will have a substantial impact on Canadian health care systems, given ongoing needs of people with a history of cancer. The increase in the number of cancer survivors will require both additional investment and innovation to provide necessary supports.²⁴

Cancer remains the leading cause of death in Canada. Estimates from the Centre for Population Health Data at Statistics Canada, released in 2023, suggest that cancer accounted for nearly 25% of all deaths in Canada in 2022.²⁵ In terms of potential years of life lost before age 75 years, the impact of cancer is even more profound. Between 2018 and 2020, more than 1 322 000 potential years of life were lost from cancer. These data suggest that cancer is also by far the leading cause of premature death in Canada.³

Notable improvements continue in terms of lung and colorectal cancer incidence and mortality. The impact of tobacco cessation and avoidance on incidence of lung cancer continues to be observed.^{26,27} However, among females, reductions in lung cancer incidence have not been as pronounced, and more cases are expected to be diagnosed among females than males in 2024 in Canada. In the United States, higher rates of lung cancer have been observed among younger women, which may not be fully explained by tobacco consumption patterns.²⁸ For colorectal cancer incidence, organized screening programs have led to large reductions in ASIRs in recent years.^{29,30} For both lung and colorectal cancers, improvements in diagnostic pathways and clinical management have led to reductions in mortality, with improved survival reported for most stages at diagnosis.^{20,31–33}

In our analyses, we highlighted several other cancers (i.e., liver and intrahepatic bile duct cancer, kidney cancer, melanoma and head and neck cancers) with concerning trends. Over the past 4 decades, some of the fastest increases in cancer rates in Canada were seen for these cancers. Increases in incidence of liver and intrahepatic bile duct cancers have been tied to factors such as

alcohol consumption,³⁴ hepatitis B and C infection,³⁵ excess body size,³⁶ and diabetes.³⁷ Kidney cancer has emerged as one of the cancers most strongly associated with excess body size,³⁶ diabetes,³⁸ and high blood pressure.³⁹ The rise in melanoma is a continued cause for concern, given the known relationship with exposure to ultraviolet radiation; continued investment is required for preventive personal and policy efforts.⁴⁰ Also included in the mid-incidence group are cancers of the head and neck. Diverse risk factors and epidemiological trends in tobacco and alcohol consumption,⁴¹ along with human papillomavirus (HPV) infection,⁴² are affecting these rates and merit continued attention.

Although substantial progress has been made in reducing incidence of cervical cancer over the past decades, recent trends suggest that rates have plateaued and have even started to increase. The rate remains relatively low (7.8 cases per 100 000), but the observed recent increase is concerning given that cervical cancer is largely preventable. Similar trends have been observed in the US, with increases in late-stage diagnoses⁴³ and overall rates among younger age groups (30–40 yr).⁴⁴ A study in British Columbia from 1971 to 2017 found similar attenuation of declining rates among younger age groups (< 45 yr).⁴⁵ These projections suggest that continued intervention is needed to reach the goal of eliminating cervical cancer in Canada.²⁴ The World Health Organization suggests that countries should strive for fewer than 4 cases per 100 000 by 2030 through the widespread uptake (> 90%) of HPV vaccination programs and HPV testing as the primary screening modality.⁴⁶

A major update to our analyses is the inclusion of data from Quebec, enabled by the submission of 2011–2017 incidence data to the Canadian Cancer Registry from the Quebec cancer registry.³ Compared with previous efforts to model national data, the projected estimates from this analysis are likely more accurate for Quebec and Canada, given the inclusion of the previously missing Quebec data.

Limitations

A major limitation of currently available population-level cancer data in Canada is the lack of individual-level data on social determinants of health such as income, education, and race and ethnicity. Although provincial cancer registries are mandated to collect data to capture pathologically confirmed diagnoses of cancer, measures of social determinants are not captured by provincial or regional health authorities for inclusion in registries. Despite these limitations, data from area-level analyses of socioeconomic status and regional-specific analyses of individual-level data in Canada show the strong effects of social factors on cancer incidence and mortality.^{47–49} In particular, large effects have been observed for cancers for which there are organized screening programs⁵⁰ or strong exposures related to social gradient such as smoking and lung cancer.⁵¹

The estimates here are projections based on the most up-to-date national cancer data and may be subject to variability over time. Because of the intensive nature of registering cancer diagnoses and deaths, data availability is delayed. Although we included 2020 mortality data from the Canadian Vital Statistics Death database, we included incidence data up to 2019 in the

projections to avoid biasing projections because of pandemic-related effects. Reports have suggested that decreases in diagnoses of cancer observed in 2020 were related to pandemic responses.^{52,53} These changes are more a reflection of differences in practice patterns during the pandemic than actual changes in the underlying incidence, which may bias short-term projections away from the actual trends in cancer incidence. Subsequent iterations of our analyses will inform projections of trends less affected by service interruptions or altered diagnostic patterns from the pandemic response.

We did not include data from Quebec (2018, 2019) and Nova Scotia (2019) because of reporting delays. These differences are likely to have minimal impact on the overall projections of national data. Furthermore, although the CANPROJ approach has been validated using Canadian data, specific validations for each province have not been performed. In addition, changes to cancer site definitions have occurred during the data years included in the projections (Appendix 1). These are likely to have minimal impact on the reported trends since the current definitions were retroactively applied to all diagnosis years used for projection purposes and are in alignment with changes in definitions for other national cancer registries (e.g., the North American Association of Central Cancer Registries).

Conclusion

These estimates highlight the substantial impact of cancer on people and health systems in Canada. Although the continued reduction in ASIRs and ASMRs that we project reflect increased survivorship and lower cancer incidence related to considerable efforts in prevention, screening, and treatment, continued investment across all areas of cancer control and care are needed to reduce the impact on all people affected by cancer in Canada.

References

- Brenner DR, Weir HK, Demers AA, et al. Projected estimates of cancer in Canada in 2020. *CMAJ* 2020;192:E199-205.
- Cancer survival statistics, 2020 update. Ottawa: Statistics Canada; 2020. Available: <https://www150.statcan.gc.ca/n1/daily-quotidien/201127/dq201127b-eng.htm> (accessed 2021 Apr. 15).
- Canadian Cancer Statistics Advisory Committee. Canadian Cancer Statistics 2023. Toronto: Canadian Cancer Society; 2023. Available: <http://cancer.ca/Canadian-Cancer-Statistics-2023-EN> (accessed 2024 Jan. 19).
- Liu JL, Zhang SX, Billette JM, et al. Lifetime probability of developing cancer and dying from cancer in Canada, 1997 to 2020. *Health Rep* 2023;34:14-21.
- Table 13-10-0111-01. Number and rates of new cases of primary cancer, by cancer type, age group and sex. Ottawa: Statistics Canada; modified 2024 Mar. 28. Available: <https://www150.statcan.gc.ca/t1/tbl1/en/tv.action?pid=1310011101> (accessed 2024 Jan. 19).
- Table 13-10-0142-01. Deaths, by cause, chapter II: neoplasms (C00-D48). Ottawa: Statistics Canada; 2023. Available: <https://www150.statcan.gc.ca/t1/tbl1/en/tv.action?pid=1310014201> (accessed 2024 Jan. 19).
- Garaszczuk R, Yong JHE, Sun Z, et al. The Economic burden of cancer in Canada from a societal perspective. *Curr Oncol* 2022;29:2735-48.
- Canadian Cancer Registry. Ottawa: Statistics Canada; 2023. Available: <http://www23.statcan.gc.ca/imdb/p2SV.pl?Function=getSurvey&SDDS=3207> (accessed 2023 Apr. 20).
- Canadian Vital Statistics – Death Database (CVS: D). Ottawa: Statistics Canada; 2023. Available: <http://www23.statcan.gc.ca/imdb/p2SV.pl?Function=getSurvey&lang=en&db=imdb&adm=8&dis=2&SDDS=3233> (accessed 2023 Apr. 20).
- Annual demographic estimates: Canada, provinces and territories. Ottawa: Statistics Canada; 2021. Cat no 91-215-X. Available: <https://www150.statcan.gc.ca/n1/pub/91-215-x/91-215-x2021001-eng.htm> (accessed 2022 Sept. 28).
- Annual demographic estimates: Canada, provinces and territories. Ottawa: Statistics Canada; 2022. Available: <https://www150.statcan.gc.ca/n1/pub/91-215-x/91-215-x2022001-eng.htm> (accessed 2024 Jan. 19).
- Population projections for Canada (2018 to 2068), provinces and territories (2018 to 2043). Ottawa: Statistics Canada; 2019. Cat. no. 91-520-X. Available: <https://www150.statcan.gc.ca/n1/en/pub/91-520-x/91-520-x2019001-eng.pdf?st=AtO08q7u> (accessed 2021 Apr. 15).
- Qiu Z, Hatcher J; Cancer Projection Analytical Network Working Team. CAN-PROJ: the R package of cancer projection methods based on generalized linear models for age, period, and/or cohort: version 1. Technique Report for Cancer Projections Network (C-Proj) Alberta. Edmonton: Alberta Health Services; 2013.
- Demers A, Qiu Z, Dewar R, et al. Validation of Canproj for projecting Canadian cancer incidence data. *Health Promot Chronic Dis Prev Can* 2020;40:267-80.
- Steliarova-Foucher E, Stiller C, Lacour B, et al. International classification of childhood cancer, third edition. *Cancer* 2005;103:1457-67.
- International classification of diseases, ninth revision. Volumes 1 and 2. Geneva: World Health Organization; 1977.
- International statistical classification of diseases and related health problems, tenth revision. Volumes 1 to 3. Geneva: World Health Organization; 1992.
- Canadian Cancer Registry – age-standardization: incidence. Ottawa: Statistics Canada; 2023. Available: https://www.statcan.gc.ca/en/statistical-programs/document/3207_D12_V4 (accessed 2024 Jan. 19).
- Warkentin MT, Ruan Y, Ellison LF, et al. Progress in site-specific cancer mortality in Canada over the last 70 years. *Sci Rep* 2024;14:5688.
- Ellison LF, Saint-Jacques N. Five-year cancer survival by stage at diagnosis in Canada. *Health Rep* 2023;34:3-15.
- Ellison LF. The cancer survival index: measuring progress in cancer survival to help evaluate cancer control efforts in Canada. *Health Rep* 2021;32:14-26.
- Ellison LF. Measuring progress in cancer survival across Canadian provinces: Extending the cancer survival index to further evaluate cancer control efforts. *Health Rep* 2022;33:17-29.
- Canadian Cancer Statistics Advisory Committee. Canadian cancer statistics: a special report on cancer prevalence. Toronto: Canadian Cancer Society; 2022. Available: <http://cancer.ca/Canadian-Cancer-Statistics-2022-EN> (accessed 2024 Jan. 19).
- Canadian Partnership Against Cancer. 2019-2029 Canadian strategy for cancer control. Available: <https://s22457.pcdn.co/wp-content/uploads/2019/06/Canadian-Strategy-Cancer-Control-2019-2029-EN.pdf> (accessed 2024 Jan. 19).
- Table 13-10-0394-01. Leading causes of death, total population, by age group. Ottawa: Statistics Canada; 2023. Available: <https://www150.statcan.gc.ca/t1/tbl1/en/tv.action?pid=1310039401> (accessed 2024 Jan. 19).
- Smoking and lung cancer in Canada. Toronto: Canadian Partnership Against Cancer; 2010.
- Poirier AE, Ruan Y, Grevers X, et al. Estimates of the current and future burden of cancer attributable to active and passive tobacco smoking in Canada. *Prev Med* 2019;122:9-19.
- Jemal A, Miller KD, Ma J, et al. Higher lung cancer incidence in young women than young men in the United States. *N Engl J Med* 2018;378:1999-2009.
- Sewitch MJ, Fournier C, Ciampi A, et al. Colorectal cancer screening in Canada: results of a national survey. *Chronic Dis Can* 2008;29:9-21.
- Brenner DR, Heer E, Sutherland RL, et al. National trends in colorectal cancer incidence among older and younger adults in Canada. *JAMA Netw Open* 2019;2:e198090.
- Brenner DR, O'Sullivan DE, Jarada TN, et al. The impact of population-based EGFR testing in non-squamous metastatic non-small cell lung cancer in Alberta, Canada. *Lung Cancer* 2023;175:60-7.
- Siegel RL, Miller KD, Wagle NS, et al. Cancer statistics, 2023. *CA Cancer J Clin* 2023;73:17-48.
- Akhtar-Danesh N, Akhtar-Danesh GG, Seow HY, et al. Trends in survival based on treatment modality in non-small cell lung cancer patients: a population-based study. *Cancer Invest* 2019;37:355-66.
- Chuang SC, Lee YC, Wu GJ, et al. Alcohol consumption and liver cancer risk: a meta-analysis. *Cancer Causes Control* 2015;26:1205-31.
- Song C, Lv J, Liu Y, et al. Associations between hepatitis B virus infection and risk of all cancer types. *JAMA Netw Open* 2019;2:e195718.
- Lauby-Secretan B, Scoccianti C, Loomis D, et al. Body Fatness and Cancer-Viewpoint of the IARC Working Group. *N Engl J Med* 2016;375:794-8.
- Davila JA, Morgan RO, Shaib Y, et al. Diabetes increases the risk of hepatocellular carcinoma in the United States: a population-based case control study. *Gut* 2005;54:533-9.
- Bonilla-Sanchez A, Rojas-Munoz J, Garcia-Perdomo HA. Association between diabetes and the risk of kidney cancer: systematic review and meta-analysis. *Clin Diabetes* 2022;40:270-82.
- Chow WH, Gridley G, Fraumeni JF Jr, et al. Obesity, hypertension, and the risk of kidney cancer in men. *N Engl J Med* 2000;343:1305-11.

40. O'Sullivan DE, Brenner DR, Villeneuve PJ, et al. Estimates of the current and future burden of melanoma attributable to ultraviolet radiation in Canada. *Prev Med* 2019;122:81-90.
41. Zhang Y, Wang R, Miao L, et al. Different levels in alcohol and tobacco consumption in head and neck cancer patients from 1957 to 2013. *PLoS One* 2015;10:e0124045.
42. Sabatini ME, Chioocca S. Human papillomavirus as a driver of head and neck cancers. *Br J Cancer* 2020;122:306-14.
43. Francoeur AA, Liao CI, Caesar MA, et al. The increasing incidence of stage IV cervical cancer in the USA: What factors are related? *Int J Gynecol Cancer* 2022 Aug 18;ijgc-2022-003728. doi: 10.1136/ijgc-2022-003728. [Epub ahead of print].
44. Shahmoradi Z, Damgacioglu H, Clarke MA, et al. Cervical cancer incidence among US women, 2001–2019. *JAMA* 2022;328:2267-9.
45. Raveinthiranathan N, Simkin J, Donken R, et al. Age-specific trends of invasive cervical cancer incidence in British Columbia, Canada, 1971–2017. *Curr Oncol* 2023;30:7692-705.
46. Global strategy to accelerate the elimination of cervical cancer as a public health problem. Geneva: World Health Organization; 2020. Available: <https://www.who.int/publications/i/item/9789240014107> (accessed 2024 Jan. 19).
47. Hwang SW, Wilkins R, Tjepkema M, et al. Mortality among residents of shelters, rooming houses, and hotels in Canada: 11-year follow-up study. *BMJ* 2009;339:b4036.
48. Tjepkema M, Wilkins R, Long A. Cause-specific mortality by occupational skill level in Canada: a 16-year follow-up study. *Chronic Dis Inj Can* 2013;33:195-203.
49. Ruan Y, Heer E, Warkentin MT, et al. The association between neighborhood-level income and cancer stage at diagnosis and survival in Alberta. *Cancer* 2024;130:563-75.
50. Booth CM, Li G, Zhang-Salomons J, et al. The impact of socioeconomic status on stage of cancer at diagnosis and survival: a population-based study in Ontario, Canada. *Cancer* 2010;116:4160-7.
51. Lung cancer and equity: a focus on income and geography. Toronto: Canadian Partnership Against Cancer; 2020. Available: <https://www.partnershipagainstcancer.ca/lung-equity> (accessed 2021 Apr. 15).
52. Yong JH, Mainprize JG, Yaffe MJ, et al. The impact of episodic screening interruption: COVID-19 and population-based cancer screening in Canada. *J Med Screen* 2021;28:100-7.
53. Walker MJ, Meggetto O, Gao J, et al. Measuring the impact of the COVID-19 pandemic on organized cancer screening and diagnostic follow-up care in Ontario, Canada: a provincial, population-based study. *Prev Med* 2021;151:106586.

Competing interests: Darren Brenner reports support from the Canadian Cancer Society and the Canadian Partnership Against Cancer. Ryan Woods reports funding from the BC Cancer Foundation, the Canadian Partnership Against Cancer, and Statistics Canada. Donna Turner reports support to attend meetings from the Canadian Cancer Society. No other competing interests were declared.

This article has been peer reviewed.

Affiliations: Departments of Oncology and Community Health Sciences (Brenner), Cumming School of Medicine, University of Calgary, Calgary, Alta.; Surveillance, Canadian Cancer Society (Gillis), Vancouver, BC; Centre for Surveillance and Applied Research (Demers), Public Health Agency of Canada; Centre for Population Health Data (Ellison, Billette, Zhang, Liu), Statistics Canada, Ottawa, Ont.; Population Oncology, BC Cancer (Woods), Vancouver, BC; Department of Surgery (Finley), McMaster University and St. Joseph's Health Care Centre, Hamilton, Ont.; System Performance & Analytics

(Fitzgerald), Canadian Partnership Against Cancer, Toronto, Ont.; Nova Scotia Health Cancer Care Program (Saint-Jacques), Halifax, NS; Cancer Advanced Analytics (Shack), Cancer Care Alberta, Calgary, Alta.; Department of Community Health Sciences (Turner), Max Rady College of Medicine, University of Manitoba; Paul Albrechtsen Research Institute (Turner), CancerCare Manitoba, Winnipeg, Man.

Contributors: All of the authors contributed to the conception and design of the work, and the acquisition, analysis, and interpretation of data. All of the authors drafted the manuscript, revised it critically for important intellectual content, gave final approval of the version to be published, and agreed to be accountable for all aspects of the work.

Content licence: This is an Open Access article distributed in accordance with the terms of the Creative Commons Attribution (CC BY-NC-ND 4.0) licence, which permits use, distribution and reproduction in any medium, provided that the original publication is properly cited, the use is noncommercial (i.e., research or educational

use), and no modifications or adaptations are made. See: <https://creativecommons.org/licenses/by-nc-nd/4.0/>

Funding: This study was developed by the Canadian Cancer Statistics Advisory Committee in collaboration with the Canadian Cancer Society, Statistics Canada, and the Public Health Agency of Canada and was therefore supported by public and donor funding. No external sources of funding were obtained for this study.

Data sharing: The data included in these analyses are available to the public and researchers through application and access from the Research Data Centres of Statistics Canada.

Disclaimer: The findings and conclusions in this report are those of the authors and do not necessarily represent the official position of the Canadian Cancer Society.

Accepted: Mar. 28, 2024

Correspondence to: Darren Brenner, Darren.Brenner@ucalgary.ca