

IMA Genome-F 10

Nine draft genome sequences of *Claviceps purpurea* s.lat., including *C. arundinis*, *C. humidiphila*, and *C. cf. spartinae*, pseudomolecules for the pitch canker pathogen *Fusarium circinatum*, draft genome of *Davidsoniella eucalypti*, *Grosmanniagaleiformis*, *Quambalaria eucalypti*, and *Teratosphaeria destructans*

Brenda D. Wingfield⁴, Miao Liu¹, Hai D.T. Nguyen¹, Frances A. Lane⁴, Seamus W. Morgan⁴, Lieschen De Vos⁴, P. Markus Wilken⁴, Tuan A. Duong⁴, Janneke Aylward^{4,5}, Martin P.A. Coetzee⁴, Kasia Dadej¹, Z. Wilhelm De Beer⁴, Wendy Findlay¹, Minette Havenga^{4,5}, Miroslav Kolařík³, Jim G. Menzies², Kershney Naidoo⁴, Olivia Pochopski¹, Parivash Shoukouhi¹, Quentin C. Santana⁴, Keith A. Seifert¹, Nicole Soal⁴, Emma T. Steenkamp⁴, Catherine T. Tatham⁴, Margriet A. van der Nest⁴, and Michael J. Wingfield⁴

¹Ottawa Research & Development Centre, Agriculture and Agri-Food Canada, 960 Carling Ave. Ottawa, Ontario K1A 0C6, Canada

²Morden Research and Development Centre, Agriculture and Agri-Food Canada, 101 Route 100, Morden, Manitoba R6M 1Y5, Canada

³Laboratory of Fungal Genetics and Metabolism, Institute of Microbiology, Academy of Sciences of Czech Republic, Videnska 1083, 142 20 Prague 4, Czech Republic

⁴Department of Biochemistry, Genetics and Microbiology (BGM), Forestry and Agricultural Biotechnology Institute (FABI), University of Pretoria, Private Bag x20, Hatfield, Pretoria, 0028, South Africa; corresponding author e-mail: Brenda.Wingfield@FABI.up.ac.za

⁵Department of Conservation Ecology and Entomology, Stellenbosch University, Private Bag X1, Matieland 7602, South Africa

Abstract: This genome announcement includes draft genomes from *Claviceps purpurea* s.lat., including *C. arundinis*, *C. humidiphila* and *C. cf. spartinae*. The draft genomes of *Davidsoniella eucalypti*, *Quambalaria eucalypti* and *Teratosphaeria destructans*, all three important eucalyptus pathogens, are presented. The insect associate *Grosmannia galeiformis* is also described. The pine pathogen genome of *Fusarium circinatum* has been assembled into pseudomolecules, based on additional sequence data and by harnessing the known synteny within the *Fusarium fujikuroi* species complex. This new assembly of the *F. circinatum* genome provides 12 pseudomolecules that correspond to the haploid chromosome number of *F. circinatum*. These are comparable to other chromosomal assemblies within the FFSC and will enable more robust genomic comparisons within this species complex.

Key words:

chromosome numbers
ergotism
Eucalyptus
pine pitch canker
Poaceae

Article info: Submitted: 21 November 2018; Accepted: 26 November 2018; Published: 14 December 2018.

IMA Genome-F 10A

Nine draft genome sequences of *Claviceps purpurea* s.lat., including *C. arundinis*, *C. humidiphila* and *C. cf. spartinae*

INTRODUCTION

Claviceps purpurea (*Clavicipitaceae*, *Hypocreales*) is a plant pathogen that infects the flowers of cereal crops and grasses (*Poaceae*) causing ergot disease. After floral infection by the pathogen, the seeds of grass hosts are replaced with

hard, dark fungal resting bodies called sclerotia or ergots. Consumption of grains contaminated with ergots is harmful to human and animal health, causing ergotism, the result of a spectrum of potent mycotoxins known as ergot alkaloids (Lyons *et al.* 1986, Miles *et al.* 1996, Scott 2009). These alkaloids have caused significant health, social and economic concerns at different times in history (Fuller 1968, Caporael 1976, Miles *et al.* 1996, De Groot *et al.* 1998, Alm 2003), but are also powerful pharmaceuticals for treating various medical conditions (De Groot *et al.* 1998, Crosignani 2006, Micale *et al.* 2006). Understanding the genetic diversity of species, their correlated toxin profiles and molecular backgrounds is important for the agricultural and pharmaceutical sectors, and regulatory agencies.

Intraspecific variations in morphology, alkaloid chemistry, genetics, and ecological niches have revealed the existence

© 2018 International Mycological Association

You are free to share - to copy, distribute and transmit the work, under the following conditions:

Attribution: You must attribute the work in the manner specified by the author or licensor (but not in any way that suggests that they endorse you or your use of the work).

Non-commercial: You may not use this work for commercial purposes.

No derivative works: You may not alter, transform, or build upon this work.

For any reuse or distribution, you must make clear to others the license terms of this work, which can be found at <http://creativecommons.org/licenses/by-nc-nd/3.0/legalcode>. Any of the above conditions can be waived if you get permission from the copyright holder. Nothing in this license impairs or restricts the author's moral rights.

of several subgroups (Pazoutová *et al.* 2000) in *C. purpurea* s. *lat.* These groups were later identified as cryptic species based on multi-gene phylogenetic and population genetic analyses, i.e. *C. arundinis*, *C. humidiphila*, and *C. spartinae* (Douhan *et al.* 2008, Pažoutová *et al.* 2015) and recently a few more species from South Africa were described as a part of the species complex (Van der Linde *et al.* 2016). In Canada, the incidence of ergot diseases in Alberta, Saskatchewan and Manitoba has been increasing since 2002 (Menzies & Turkington 2014). During a recent investigation of ergot fungi in agriculture areas in Canada, we discovered a few more new phylogenetic lineages: two closely related but different from *C. spartinae* (G3), one close to *C. humidiphila*, and another two located as basal branches in the *C. purpurea* s. *lat.* complex (Fig. 1). Here, we selected representatives of these new and previously designated lineages, sequenced and assembled their genomes. A complete multigene phylogenetic analysis, including a much greater sampling of strains, will be presented in a separate publication.

SEQUENCED STRAINS

Claviceps purpurea s.str.

Canada: Saskatchewan: Estavan SK1 49.14 N 102.99 W, isolated from *Triticum aestivum*, 2000, R. Clear [identified by J. G. Menzies] (LM28 = DAOMC 250647). **Czech Republic:** Bezdědice, 49.83 N 14.03 E, isolated from *Secale cereale*, 2003 [identified by S. Pažoutová] (LM582 = DAOMC 251723 = CCC771 ex-neotype).

Claviceps purpurea s.lat.

Canada: Alberta: North Star 58.53 N 118.12 W, isolated from *Bromus inermis*, 7 Sep. 1956 [identified by W. P. Campbell] (LM78 = DAOMC 250578); Metiskow, 52.41 N 110.63 W, isolated from *Elymus albicans*, 7 Sep. 1956 [identified by W. P. Campbell] (LM81 = DAOMC 250581). Quebec: Cote Nord, MRC Minganie, Pointe-Parent, 50.13 N, 61.08 W, isolated from *Ammophila* sp., 8 Sep. 2015, J. Cayouette & Y. Dalpé (LM458 = DAOMC 251898).

Claviceps cf. *spartinae*

Canada: Manitoba: Grant's Field Snowflake 49.05 N 98.66 W, isolated from *Phalaris arudinacea*, 2014, J. Menzies [identified by M. Liu] (LM218 = DAOMC 251843). Quebec: Maria-Chapdelaine, parc national de la Pointe-Taillon, river du Lac Saint-Jean, 48.67 N 71.87 W, isolated from *Ammophila breviligulata*, 31 Aug. 2014, J. Cayouette [identified by M. Liu] (LM454 = DAOMC 251845 = DAOM 550246b – specimen).

Claviceps humidiphila

Germany: Bavaria: near Phillipsreuth, 48.86 N 13.68 E, isolated from *Dactylis* sp., 1998 and S. Pažoutová (LM576 = DAOMC 251717 = CCC434 ex-epitype).

Claviceps arundinis

Czech Republic: Haklovy Dvory, Stary Vrbensky pond, 49.01 N 14.43 E, isolated from *Phragmites australis*, 11 Jan. 2008, M. Kolařík (LM583 = DAOMC 251724 = CCC933 ex-type).

NUCLEOTIDE SEQUENCE ACCESSION NUMBERS

The genome data for the nine strains were deposited at DDBJ/EMBL/GenBank under BioProject PRJNA449361. The versions described in this paper are version QERD01000000, QEQW01000000, QEQX01000000, QEQY01000000, QEQZ01000000, QERA01000000, QERB01000000, QERC01000000, QERE01000000. Raw reads were deposited in NCBI SRA (<http://www.ncbi.nlm.nih.gov/sra>) under accession number SRP139610.

MATERIALS AND METHODS

Fungal mycelia and spores were inoculated onto Potato Dextrose Agar (PDA; BD Difco™) by streaking the medium surface in Petri dishes using inoculation loops, to allow the fungus to cover the dish in a short time and incubated for 10 d at 20 °C in the dark. Mycelia of 1–2 dishes were harvested and ground using liquid N₂, followed by genomic DNA extraction using a modified cetyltrimethyl ammonium bromide (CTAB) protocol (Doyle & Doyle 1987). The protocol was modified to remove RNA by adding extra RNase and incubating for longer time as follows: (1) after homogenization of ground fungal tissues with 700 µl 2X CTAB, 140 µl RNase Cocktail™ Enzyme Mix (4x recommended volume ratio: 2.5 µl / 50 µl sample; Invitrogen by Thermo Fisher Scientific), and 28 µl RNASEA (4x recommended volume ratio: 1 µl / 100 µl sample) was added, and digested for 6–7 h at room temperature. (2) Afterwards, 10 µl Proteinase K (50 µg/µL stock; Fisher Scientific by Thermo Fisher Scientific) was added to the suspension and incubated for 2 h at 55 °C. (3) Before DNA precipitation, another 35 µl RNase Cocktail (recommended volume ratio without extra) was added into the supernatant, incubated for 45 min – 1 h at room temperature, followed by adding 650 µL CHCl₄ and centrifuging at 13 000 rpm for 15 min. Potential RNA contamination in the DNA samples was checked by running the samples on 1 % agarose gels, and determining the 260/280 ratio using NanoDrop 1000 Spectrophotometer v3.8 (Thermo Fisher Scientific) to ensure no RNA was present. gDNA was quantified using Qubit® 2.0 Fluorometer (Invitrogen by Life Technologies).

The extracted gDNAs were normalized to 400 ng and sheared to a 300 bp insert using a Covaris LE220 instrument. The fragmented inserts were used as a template to construct PCR free Libraries with a NxSeq AmpFREE Low DNA Library kit (Lucigen) following the manufacturer's instructions. Indexed libraries were pooled into two individual pools and sequencing runs were carried on a NextSeq (Illumina; Molecular Technologies Laboratory, Ottawa Research & Development Centre, Agriculture and Agri-Food Canada) using 2x150 bp NextSeq Mid Output Reagent Kits (Illumina).

The FastQC software v0.11.5 (<https://www.bioinformatics.babraham.ac.uk/projects/fastqc/>) was used to assess the raw read quality. Poor quality data was removed with Trimmomatic v0.36 (Bolger *et al.* 2014) using the following parameters: HEADCROP:20 SLIDINGWINDOW:4:20 MINLEN:36. The trimmed reads were error corrected with BayesHammer (Nikolenko *et al.* 2013). *De novo* genome assembly was




Fig. 1. One of the two MP trees showing nine strains (in bold) in relation to *Claviceps* lineages based on EF1- α partial region, 99 informative characters, length = 302, CI = 0.606, RI = 0.807, RC = 0.489, HI = 0.394, G-fit = -75.089. Values on branches are MP bootstrapping/BI posterior probability.

performed using SPAdes v3.10.1 (Bankevich *et al.* 2012) with the mismatch correction step enabled. Contigs shorter than

1000 bp were discarded. QFAST v4.5 program (Gurevich *et al.* 2013) was used to evaluate assemblies. Corrected reads

Table 1. Statistics of *Claviceps* genomes sequenced in this study.

Species	Strain	Voucher #	Version #	number of contigs	Number of protein coding gene models	Total length (bp)	N50 (bp)	L50	GC (%)	Coverage (x)	Complete BUSCO's (%)
<i>Claviceps purpurea</i> s.str.	LM28	DAOMC 250647	QERD01000000	1930	8977	30251759	32494	269	51,7	39	97,9
<i>Claviceps purpurea</i> s.str.	LM582	DAOMC 251723 = CCC771 ^a	QERA01000000	2207	8777	30199509	27733	327	51,7	72	98,6
<i>Claviceps purpurea</i> s.lat.	LM78	DAOMC 250578	QEQY01000000	2321	8410	28571566	21589	416	51,4	52	97,3
<i>Claviceps purpurea</i> s.lat.	LM81	DAOMC 250581	QEQX01000000	1423	9008	30694913	46550	190	51,5	305	96,9
<i>Claviceps purpurea</i> s.lat.	LM458	DAOMC 251898	QEQW01000000	1698	9230	35875594	41916	189	51,9	66	98
<i>Claviceps</i> cf. <i>spartinae</i>	LM218	DAOMC 251843	QERE01000000	1630	8960	30598250	39984	229	51,4	214	97,6
<i>Claviceps</i> cf. <i>spartinae</i>	LM454	DAOMC 251845	QERC01000000	2108	9038	30692584	28795	307	51,4	126	97,9
<i>Claviceps humidiphila</i>	LM576	DAOMC 251717 = CCC434 ^a	QERB01000000	1831	9040	30488243	35139	257	51,5	63	97,9
<i>Claviceps arundinis</i>	LM583	DAOMC 251724 = CCC933 ^a	QEQZ01000000	1613	8855	30055381	39989	219	51,4	56	96,9
<i>Claviceps purpurea</i> s.str. ^b	20,1			1442	8 823	30901872	46498	197	51,6	39	97,6

^a Ex-type cultures: CCC771 ex-neotype, CCC434 ex-epitype, CCC933 ex-isotype.^b This genome was sequenced by Schardl *et al.* (2013).

were mapped back onto the contigs using Bowtie2 v2.0.0 (Langmead & Salzberg 2012). Alignments produced by Bowtie2 in SAM format were converted to sorted BAM format by SAMtools v0.1.19 (Li *et al.* 2009) and statistics for nucleotide coverage were generated with Qualimap v2.2 (García-Alcalde *et al.* 2012). To evaluate the completeness of our genome assemblies Universal Single-Copy Orthologs, BUSCO v2 (Simão *et al.* 2015) was run on the contigs using the fungal database (obd9). Genome annotation was carried out using GeneMark-ES v4.38 (Lomsadze *et al.* 2005) with the “fungus” option enabled (Ter-Hovhannisyann *et al.* 2008). Annotations were validated using Genome Annotation Generator (Hall *et al.* 2014) and tbl2asn (<http://www.ncbi.nlm.nih.gov/genbank/tbl2asn2/>). All statistics are summarized in Table 1.

To confirm the identities of the strains, DNA sequences of partial elongation factor 1- α gene were extracted from each assembly of nine genomes using Geneious 10.0.9 (Kearse *et al.* 2012); the fragments were aligned with representative sequences developed by Pažoutová *et al.* (2015) and Van der Linde *et al.* (2016) using MAFFT version 7 (<https://mafft.cbrc.jp/alignment/server/>) Katoh & Standley 2013. Maximum Parsimony (MP) analyses were performed using PAUP* 4.0b10 (Swofford 2002) with heuristic search of 200 replicates random stepwise addition, bootstrapping of 1000 replicates. Bayesian Inference (BI) was conducted using Mr Bayes 3.2 (Ronquist *et al.* 2012), with two independent runs, each sets four chains of 100 000 000 MCMC generation, and 25 % burn-in.

RESULTS AND DISCUSSION

The strains sequenced include: two strains of *Claviceps purpurea* s.str. and *C. cf. spartinae*, one strain each of *C. humidiphila* and *C. arundinis*, and three strains of *C. purpurea* s.lat. (Table 1, Fig. 1). The genomes were assembled into 1423 to 2321 contigs with a mean assembly size 30.8 Mb ranging from 28.6 Mb to 35.9 Mb. The average GC content was 51.5 %. The N50 ranged from 21.6 kb to 46.6 kb. The assemblies had a BUSCO completeness score ranging from 96.9 % to 98.6 %; the number of gene models ranged from 8410 to 9230.

The genome of *Claviceps purpurea* strain 20.1 (NCBI accession no. CAGA00000000.1) was previously published by (Schardl *et al.* 2013). They obtained a similar number of contigs (1442), number of gene models (8823), total assembly size (30.9 Mb), N50 (46.5 kb) and GC content (51.6 %), compared to our assemblies, suggesting we obtained reasonable assemblies. However, they additionally ordered contigs into only 191 scaffolds and performed genome annotation. We ran BUSCO on their assembly and obtained a completeness score of 97.6 %. Our aim is to use these sequenced *Claviceps* genomes to further understand the species diversity, genetic variation, and establish correlations with ergot alkaloid chemical profiles.

Authors: H.D.T. Nguyen, W. Findlay, K. Dadej, O. Pochopski, P. Shoukouhi, J.G. Menzies, K.A. Seifert, M. Kolařík, and M. Liu
*Contact: Miao Liu, miao.liu@canada.ca

IMA Genome-F 10B

Assembling pseudomolecules for the pitch canker pathogen, *Fusarium circinatum*, utilising additional genome sequence data and synteny within the *Fusarium fujikuroi* complex

INTRODUCTION

Fusarium includes a diverse group of filamentous ascomycetes (Geiser *et al.* 2013). Many of these fungi cause diseases on economically important plants, with an estimated 80 % of cultivated crops having at least one associated *Fusarium* disease (Leslie & Summerell 2006). Within the *Fusarium fujikuroi* species complex (FFSC), more than 50 phylogenetically distinct species have been grouped into three biogeographical clades (O'Donnell *et al.* 1998, 2000). *Fusarium circinatum*, residing in the so-called “American clade”, is the causal agent of the disease known as pine pitch canker that damages susceptible *Pinus* spp. and *Pseudotsuga menziesii* (Douglas Fir). It has a cosmopolitan distribution and is associated with significant economic losses due to widespread seedling mortality in nurseries as well as the reduction of growth in mature trees due to dieback of infected branches (Gordon *et al.* 2015).

The importance of this pathogen has justified sequencing its genome (GenBank accession AYJV00000000, version AYJV01000000) (Wingfield *et al.* 2012). The isolate sequenced, FSP34 (Gordon *et al.* 1996), has a genetic linkage map available (De Vos *et al.* 2007) which has been anchored to the genome (De Vos *et al.* 2014) enabling localization of quantitative trait loci (QTLs) to the genomic sequence data (De Vos *et al.* 2011, Van Wyk *et al.* 2018). The draft assembly was 94.8 % complete (Waterhouse *et al.* 2017), but it included an exorbitant number of contigs (4145) (Wingfield *et al.* 2012, De Vos *et al.* 2014) and this limits its use in comparative genomic studies.

The whole genome sequences of other members of the FFSC are available and their genome complement is present in chromosomes. These include *Fusarium verticillioides* for which the sequences for only 11 chromosomes are available (Ma *et al.* 2010). This *F. verticillioides* assembly excludes that for the twelfth and smallest chromosome known to exist in members of the FFSC. This is due to the dispensable nature of this chromosome, with it being strain-specific within the FFSC (Xu *et al.* 1995, Ma *et al.* 2010, Wiemann *et al.* 2013, Van der Nest *et al.* 2014). In contrast, the whole molecule sequences have been determined for the full complement of the twelve chromosomes for *F. fujikuroi* (Wiemann *et al.* 2013). These two species represent two of the three biogeographical clades of the FFSC. Comparisons among them and *F. temperatum* have shown a significant level of macrosynteny at the genomic sequence level (Wiemann *et al.* 2013, De Vos *et al.* 2014). This highlights the fact that the genomic content on chromosomes is highly conserved

between various species in the FFSC.

The aim of this study was to improve the available draft assembly of *F. circinatum*, and assemble it into pseudomolecules that are comparable with the chromosomes of other members of the FFSC. For this purpose, we utilized additional genome sequence information (i.e. mate-pair sequence data) to allow for the scaffolding of contigs. We then exploited the macrosynteny that characterizes the genomes of species within FFSC (Wiemann *et al.* 2013, De Vos *et al.* 2014), to orientate and order these scaffolds into twelve pseudomolecules. In this study we present the pseudomolecule assemblies for the full chromosomal complement of *F. circinatum*. This improved genome assembly will aid in genome-sequence based studies, particularly those involving chromosomal comparisons. Addition of the *F. circinatum* pseudomolecule complement will furthermore enable comparative studies focusing on genomic synteny and architecture between the three biogeographic clades within the FFSC, as well as more broadly in the genus *Fusarium*.

SEQUENCED STRAIN

USA: *California:* isol. Monterey pine (*Pinus radiata*), 1996, T.R. Gordon, A.J. Storer & D. Okamoto (FSP34, MRC7870, CMW51752, PREM 62197–dried culture).

NUCLEOTIDE SEQUENCE ACCESSION NUMBER

The improved genome assembly for *Fusarium circinatum*, generated in this study, has been deposited at DDBJ/EMBL/GenBank under the accession AYJV00000000, version AYJV02000000.

MATERIALS AND METHODS

Fusarium circinatum was grown on half strength potato dextrose broth (20 % potato dextrose broth w/v) and incubated at 25 °C in the dark on an orbital shaker at 120 rpm. After 7 d, DNA was extracted following the protocol outlined (Möller *et al.* 1992) and the DNA quality was assessed using a NanoDrop™ Spectrophotometer.

Additional genomic sequence data from *F. circinatum* isolate FSP34 and a second isolate, KS17, were utilised for scaffolding the original 4145 contig assembly (Wingfield *et al.* 2012). Isolate KS17 was cultured from infected root tissue of *P. radiata* nursery seedlings collected from the Western Cape, South Africa in 2005 (Steenkamp *et al.* 2014). The genomes of *F. circinatum* isolates FSP34 and KS17 were sequenced using mate-pair libraries (1 kb insert size) by making use of the SOLiD™ V4 technology (Applied Biosystems) at Seqomics (Hungary). In total, 82.45 and 153.95 million mate-pair reads were obtained for the respective isolates. Poor quality reads (below Q20), reads smaller than 36 bp and duplicate reads were removed in CLC Genomics Workbench v.5.1 (CLCbio, Aarhus).




Fig. 2. Maximum likelihood tree based on partial gene sequences of β -tubulin and translation elongation factor 1- α (Scaufflaire *et al.* 2011, De Vos *et al.* 2014). Sequence alignments were assembled with MAFFT version 7 (Kato & Standley 2013). The program jModelTest v 2.1.7 (Guindon & Gascuel 2003, Darriba *et al.* 2012) was used to determine the best-fit substitution model (TIM2+I+G substitution model) with gamma correction (Tavare 1986). A maximum likelihood (ML) phylogenetic analysis was performed using PhyML v 3.1 (Guindon *et al.* 2010). Values at branch nodes are the bootstrapping confidence values with those $\geq 85\%$ shown. The *F. circinatum* FSP34 isolate used in this study is indicated in **bold**.

SSPACE v. 2.0 (Boetzer *et al.* 2011) was utilized to scaffold the pre-assembled contigs of the FSP34 assembly (GenBank accession no. AYJV00000000) (Wingfield *et al.* 2012), using the trimmed mate-pair data. Default parameters were used, but the minimum number of paired reads linking contigs to form a scaffold was set to 200. The average genome coverage was calculated using the Lander/Waterman equation (number of reads \times read length/genome size).

The resulting scaffolds were then assembled into 11 contiguous pseudomolecules (representing the first chromosome 1-11) using *F. verticillioides* as a reference genome. The scaffolds were ordered and orientated based on BLAST searches (Altschul *et al.* 1990) against a local database of the *F. verticillioides* genome (DDBJ/EMBL/GenBank accession number AAIM00000000.2) using CLC Genomics Workbench. To assemble pseudomolecule 12, scaffolds were ordered and orientated to chromosome 12 of *F. fujikuroi* (Wiemann *et al.* 2013) and *F. temperatum* (Wingfield *et al.* 2015b), as described above. To indicate a break between the various scaffolds comprising a pseudomolecule, 100 N's were inserted. Synteny maps were generated between the chromosomes of *F. verticillioides* and *F. fujikuroi* and the pseudomolecules of *F. circinatum* using the program MUMmer v. 3.22 (Kurtz *et al.* 2004).

The assembled genome was annotated using the MAKER annotation pipeline (Cantarel *et al.* 2008) utilizing Genemark ES (Ter-Hovhannisyanyan *et al.* 2008), Augustus (Stanke & Morgenstern 2005), and SNAP (Korf 2004). Manual curation of the predicted annotations was also performed (Wingfield *et al.* 2012). As additional evidence, genome data from *F. verticillioides*, *Fusarium oxysporum* f. sp. *lycopersici* and *F. graminearum* (Ma *et al.* 2010), as well as expressed sequence tag (EST) evidence for *F. circinatum* (Wingfield *et al.* 2012) were included.

RESULTS AND DISCUSSION

The improved genome assembly for *Fusarium circinatum*, generated in this study, has been deposited at DDBJ/EMBL/GenBank under accession no. AYJV00000000, version AYJV02000000. The *F. circinatum* genome was assembled into 585 scaffolds that cumulatively comprised 43 932 912 bases of DNA and had a N50 of 363 633bp. The genome coverage was 273.82x. A GC content of 47.41% was obtained, which is comparable to other sequenced *Fusarium* species within the FFSC (Ma *et al.* 2010, Wingfield *et al.* 2012, Jeong *et al.* 2013, Wiemann *et al.* 2013, Van der Nest *et al.* 2014, Chiara *et al.* 2015, Wingfield *et al.* 2015a, b, Niehaus *et al.* 2017a, b, Wingfield *et al.* 2017, Gardiner 2018, Srivastava *et al.* 2018, Van Wyk *et al.* 2018, Wingfield *et al.* 2018). A total of 14 923 genes were predicted to be protein-coding, yielding a gene density of 339.68 open reading frames (orfs) per million base pairs. Phylogenetic analysis of the sequenced genome confirmed the taxonomic identity as *F. circinatum* (Fig. 2).

Pseudomolecules, corresponding to each of the 11 chromosomes of *F. verticillioides*, were constructed in this study. Pseudomolecule 12 was assembled according to synteny observed with chromosome 12 of *F. fujikuroi*. We managed to genetically anchor 96.97% (ca. 42.60 Mb) of the *F. circinatum* scaffolds to these 12 pseudomolecules. These pseudomolecules harbour 99.09% of the 15 060 genes originally predicted for *F. circinatum* (Wingfield *et al.* 2012). Genomic alignments of these 12 pseudomolecules to the corresponding *F. verticillioides* and *F. fujikuroi* chromosomes are shown in Fig. 3. These dot-plots are indicative of the observable macrosynteny of *Fusarium* species within the FFSC.




Fig. 3. Whole genome comparisons of: **A.** *Fusarium verticillioides* chromosomes to *F. circinatum* pseudomolecules. **B.** *F. fujikuroi* chromosomes to *F. circinatum* pseudomolecules. In the dotplot alignments forward matches are indicated by purple dots, reverse matches with blue dots.

CONCLUSIONS

Resequencing has provided a vastly improved assembly for *Fusarium circinatum* characterized by fewer and significantly larger scaffolds. By making use of the macrosynteny known to characterize the genomes of FFSC species (Wiemann *et al.* 2013, De Vos *et al.* 2014), we were able to join these scaffolds into 12 pseudomolecules that represent the haploid chromosome number of this fungus.

This new assembly of the *F. circinatum* genome provides a considerably more robust and complete representation of the whole genome sequence of the pathogen, including pseudomolecules that correspond to the twelve chromosomes of *F. circinatum*. The availability of the sequence for these pseudomolecules will enable future comparative studies at the chromosomal level between/within the three biogeographic clades of the FFSC. In addition, comparisons of chromosomal architecture will expand our knowledge regarding the genomes of species in the FFSC and highlight inter- and intraspecific similarities and differences between them. This would broaden available knowledge regarding the evolution of an important group of plant pathogens.

Authors: L. De Vos*, Q.C. Santana, M.J. Wingfield, B.D. Wingfield, E.T. Steenkamp, and M.P.A. Coetzee

*Contact: lieschen.bahlmann@fabi.up.ac.za

IMA Genome-F 10C

Draft genome sequence of *Quambalaria eucalypti*

INTRODUCTION

The genus *Quambalaria* (*Quambalariaceae*, *Microstromatales*, *Exobasidiomycetes*, *Basidiomycotina*) includes mainly leaf and shoot pathogens of trees belonging to *Myrtaceae* (De Beer *et al.* 2006, Pegg *et al.* 2009). The exceptions to this are *Q. coyrecup* which is a canker pathogen of marri (*Corymbia calophylla*) in Western Australia (Paap *et al.* 2008), and *Q. cyanescens* which is an opportunistic pathogen of primarily immunocompromised or debilitated humans (Kuan *et al.* 2015). The latter species is also frequently isolated from galleries of bark beetles infesting hardwoods, although its ecological role in these ecosystems remains enigmatic (Kolarík *et al.* 2006). Of the leaf and shoot pathogens in the genus, *Q. pitereka* and *Q. eucalypti* are most important. *Quambalaria pitereka* affects *Corymbia* species in Australia (Pegg *et al.* 2009) and China (Zhou *et al.* 2008), while *Q. eucalypti* causes disease on *Eucalyptus* spp. in South Africa (Wingfield *et al.* 1993), Brazil (Alfenas *et al.* 2001), Uruguay (Perez *et al.* 2008), Australia (Pegg *et al.* 2008), and Portugal (Bragança *et al.* 2015).

Research on species of *Quambalariaceae* has mostly focussed on their classification and taxonomy (De Beer *et al.* 2006, Kijpornyongpan and Aime 2017, Paap *et al.* 2008), as well as their pathogenicity and impact on tree health (Pegg *et al.* 2009, 2011). However, little is understood about the life-cycle and general biology of these fungi, that are related to the smut fungi and human pathogenic members of *Malasseziales* (De Beer *et al.* 2006, Wang *et al.* 2015b). For




Fig. 4. Phylogram resulting from a ML analyses using RaxML, based on ITS sequences of selected reference sequences representing all species of *Quambalaria*. The isolate from which the genomic DNA was extracted is indicated in bold type. T = ex-type isolates; NT = Northern Territory; NSW = New South Wales; WA = Western Australia; and QLD = Queensland. Support values at branches resulted from 1000 bootstraps and only values above 75 % are indicated.

example, it is not yet known whether *Quambalariaceae* have the ability to reproduce sexually, and if so, what the mating system of these fungi encompasses.

Recently the first genome-based studies started exploring pathogenicity factors in *Ustilaginomycotina*, and although two species in *Microstromatales* were included in the comparative analyses, no representative of *Quambalariaceae* was incorporated (Kijpornyongpan *et al.* 2018). Whole genome sequences have also been shown to be extremely valuable to study mating systems in smut fungi (Que *et al.* 2014), the *Malasseziales* (Xu *et al.* 2007), and the more distantly related *Polyporales* (James *et al.* 2013).

The aim of this study was to produce a draft genome sequence of *Q. eucalypti*. This genome sequence will allow for the exploration and comparative analyses of genes involved in pathogenicity and mating for this pathogen. Here we report the draft genome sequence of isolate CMW1101, an isolate representing the holotype (PREM 51089) of *Q. eucalypti*.

SEQUENCED STRAIN

South Africa: KwaZulu Natal: Kwambonambi, *Eucalyptus grandis* clone TAG12, May 1987, M.J. Wingfield (CMW1101=CBS118844 ex-holotype isolate; PREM 51089 – holotype).

NUCLEOTIDE ACCESSION NUMBER

The draft genome sequence of *Quambalaria eucalypti* has been deposited at GenBank under accession no. PIRRYC00000000. The version presented here is RRYC01000000.

MATERIALS AND METHODS

Genomic DNA was extracted from cultures grown on Malt Yeast Agar (2 % Malt extract, 0.5 % yeast extract; 2 % agar,

Biolab, Midrand, South Africa) using the method described by Duong *et al.* (2013). The genomic DNA was sent to Macrogen (South Korea), where one pair-end library with 500 bp insert size was prepared and sequenced on Illumina Hiseq 2500 to get 250 bp pair-end reads, aiming for 100 X coverage.

The raw sequencing reads were imported into CLC Genomics Workbench v. 7.5.1 (CLCbio, Aarhus), and default settings were used to both trim the reads for quality and to produce a *de novo* genome assembly using the trimmed reads. The completeness of the assembly was evaluated using the Benchmarking Universal Single-Copy Orthologs (BUSCO v. 1.1b1) tool using the *Basidiomycota* dataset (Simao *et al.* 2015). The number of protein coding genes was determined using Augustus v. 3.3.2 (Stanke *et al.* 2008) using pre-optimised species models for *Ustilago maydis*.

RESULTS AND DISCUSSION

The paired end sequencing yielded just over 31 million reads. Assembly of the trimmed reads resulted in 966 contigs, with the largest contig being 225 583 bp, the smallest contig being 449 bp, with an average contig size of 24 384 bp and the N50 value was 62 600 bp. The genome size is estimated at around 23.5 Mb, estimated through the sum of the contig sizes, with a GC content of 60 %. This estimated size is in the larger size range of that reported in *Exobasidiomycetes*, which typically range from 17 Mb to 19 Mb with the exception of *Tilletia caries* with a genome size of 29.5 Mb (Konishi 2013, Saika *et al.* 2014, Toome *et al.* 2014, Wang *et al.* 2015a, Kijpornyongpan *et al.* 2018). BUSCO analysis indicated an assembly completeness of 84.5 %. The assembly contained 1128 complete BUSCOs (1093 complete single-copy BUSCOs, 35 complete and duplicated BUSCOs), 129 fragmented BUSCOs and 78 missing BUSCOs out of a total 1335 BUSCO groups searched. AUGUSTUS predicted 7241 putative protein coding regions. Phylogenetic analysis of sequences from the sequenced genome confirmed the taxonomic identity as *Q. eucalypti* (Fig. 4). The availability of the *Q. eucalypti* genome will enable the inclusion of this species as representative for the family *Quambalariaceae* in comparative studies with other members of the class *Exobasidiomycetes*. Such studies could focus on topics like the factors involved in pathogenicity, mating, evolution and more.

Authors: S.W. Morgan, T. A. Duong, M. Coetzee, M.J. Wingfield, and Z.W. De Beer*

*Contact: wilhelm.debeer@fabi.up.ac.za

IMA Genome-F 10D

Draft genome sequence of the *Eucalyptus* pathogen *Teratosphaeria destructans*

INTRODUCTION

The genus *Teratosphaeria* includes numerous economically

important tree pathogens of plantation eucalypts in tropical and subtropical areas (Park *et al.* 2000, Crous *et al.* 2009, Hunter *et al.* 2011). The aggressive pathogen *T. destructans* was initially reported from Indonesia in 1996 (Wingfield *et al.* 1996), followed by reports from Thailand, Vietnam, East Timor, Laos, China, and, most recently, South Africa (Old *et al.* 2003, Burgess *et al.* 2006, Barber *et al.* 2012, Greyling *et al.* 2016). It causes leaf, bud and shoot blight disease in one- to three-year-old trees of *Eucalyptus camaldulensis*, *E. grandis* and *E. urophylla* as well as on hybrids of these species (Wingfield *et al.* 1996, Old *et al.* 2003, Barber 2004). The rapid spread of *T. destructans* over large distances has been attributed to the anthropogenic movement of germplasm to establish clonal *Eucalyptus* nurseries (Andjic *et al.* 2011).

The discovery of *T. destructans* on clonal *E. grandis* × *E. urophylla* plantations in South Africa, coupled with its reported rapid spread to new areas, makes this pathogen of major concern to the forestry industries of southern African countries (Andjic *et al.* 2011, Greyling *et al.* 2016). Similarly, the ability of this pathogen to rapidly invade new areas are also a concern to Australia, where the prospect of *T. destructans* spreading to native *Eucalyptus* species could prove catastrophic to Australia's commercial and natural vegetation (Old *et al.* 2003). In this study, the genome sequence of a South African isolate of *T. destructans* is reported. The availability of a complete genome sequence for *T. destructans* will prove beneficial to studies on the genes and pathways involved in virulence, pathogenicity and sexual reproduction. Such studies will increase our understanding of the biology of this fungus, which is crucial for the development of preventative and control measures for this tree pathogen.

SEQUENCED STRAIN

South Africa: *Kwa-Zulu Natal:* Kwambonambi, isol. *Eucalyptus grandis* × *E. urophylla*, Apr. 2015, I. Greyling (CMW 44962, PREM 62207 – dried specimen).

NUCLEOTIDE ACCESSION NUMBER

The *Teratosphaeria destructans* Whole Genome Shotgun project has been deposited at DDBJ/ENA/GenBank under accession no. RIBY00000000. The version described in this paper is version RIBY01000000.

MATERIALS AND METHODS

Teratosphaeria destructans isolate CMW 44962 was obtained from the culture collection (CMW) of the Forestry and Agricultural Biotechnology Institute (FABI) at the University of Pretoria. A single-spore culture was generated (Greyling *et al.* 2016) and grown on sterile cellophane sheets placed onto MEA+Y plates (malt extract agar; Biolab, South Africa; amended with 0.3 % yeast extract; Oxoid, Basingstoke). After incubation for four weeks at 25 °C, the mycelia were collected and freeze-dried for DNA extraction. Approximately 30 mg of freeze-dried material, three 5 mm




Fig. 5. A Maximum Likelihood phylogeny showing *Teratosphaeria* species including the genome sequences of *Teratosphaeria destructans* reported here. The β -tubulin and EF1- α gene regions were used and were obtained from previous studies (Quaedvlieg *et al.* 2014, Aylward *et al.* 2018).

glass beads and 13 mg polyvinylpyrrolidone (PVP, Sigma Aldrich, Steinheim) were mixed and powdered in a FastPrep FP120 tissue lyzer (Qbiogene, Carlsbad, CA). Subsequently, 650 μ l of CTAB extraction buffer (100 mM Tris-HCl, pH8; 25 mM EDTA; 2 M NaCl; 3.5 % CTAB; 2 % SDS), 2 μ l of 500 mg/L spermidine (Sigma Aldrich, Steinheim) and 4.5 % (v/v) β -mercaptoethanol were added and the sample was incubated at 60 °C for 20 min. After cooling to room temperature, 1.5 volumes of chloroform:isoamylalcohol (24:1) was added, the sample was mixed and centrifuged for 15 min at 8100 *g*. The supernatant was re-extracted with 1.5 volumes of chloroform:isoamylalcohol (24:1) and centrifuged for 15 min at 16 300 *g*. The supernatant was combined with potassium acetate to a final concentration of 1.5 M. After incubation for 30 min at -20 °C, 1.5 volumes of cold isopropanol were added, followed by a 30 min incubation at 24 °C. Thereafter, the sample was centrifuged for 20 min at 16 300 *g* to collect the DNA which was washed twice with 70 % ethanol. The dried DNA pellet was dissolved in 50 μ l low TE (Tris-EDTA) buffer (Thermo Fisher Scientific, Wilmington, NC).

The genomic DNA sample was submitted to the Central Analytical Facilities (CAF) of Stellenbosch University (Stellenbosch, South Africa) for whole genome sequencing using the Ion GeneStudio S5 Next-Generation Sequencer. An Ion 530 chip with a capacity to generate 12 million reads of 600 bp was prepared as per the manufacturer's instructions. The resulting single reads were trimmed and assembled using SPAdes v. 3.12.0 (Nurk *et al.* 2013) with k-values of 21, 33, 55, 77, 99 and 127. The completeness of the genome

assembly was evaluated using the Benchmarking Universal Single-Copy Orthologs (BUSCO v. 2.0.1) tool in conjunction with the fungal data set (Simão *et al.* 2015). Bowtie2 v. 1.1.2 and SAMtools v. 1.5 were used to calculate the average base coverage by mapping the reads back to the genome assembly (Li *et al.* 2009, Langmead & Salzberg 2012). QUAST v. 5.0.1 (Mikheenko *et al.* 2018) was used to calculate general genome statistics.

The β -tubulin and elongation factor 1-alpha (EF1- α) gene regions, commonly used for species delineation in *Teratosphaeria* (Quaedvlieg *et al.* 2014) were extracted from the *T. destructans* genome sequence. These sequences, together with previously published sequences (Quaedvlieg *et al.* 2014, Aylward *et al.* 2018) sourced from the NCBI database (<https://www.ncbi.nlm.nih.gov/>) were subjected to a "one click" phylogeny analysis using the Phylogeny.fr online tool (Dereeper *et al.* 2008, 2010). This analysis included a MUSCLE alignment (Edgar 2004) and a Gblocks (Castresana 2000) curation step before phylogenetic analysis was conducted using PhyML (Guindon & Gascuel 2003, Anisimova & Gascuel 2006).

RESULTS AND DISCUSSION

The assembly yielded a genome of 32 316 120 bp assembled into 4132 contigs. Of these, 1837 were 500 bp or larger and contained 31.61 Mb of the genome. The genome had a GC content of 51.83 %, a N50 value of 103 119 bases (with

the largest contig 398 757 bp in size), and a L50 value of 94 contigs. The average coverage of this assembly was 170x. The genome completeness, as assessed by BUSCO analysis, was 95.86 %, with 278 complete, five fragmented and seven missing BUSCO terms out of the 290 searched.

The genomic sequence of *Teratosphaeria destructans* reported here is the first published genome sequence for a member of this genus (Fig. 5). Considering its status as an emerging pathogen (Andjic *et al.* 2011, Greyling *et al.* 2016, Burgess & Wingfield 2017) the availability of a genome sequence holds various benefits. These include its use in comparative genomic studies between different *Teratosphaeria* species, many of which are a concern to the global *Eucalyptus* industry (Hunter *et al.* 2011). Similar studies have already yielded insight into the factors that determine host range, while also elucidating an arsenal of pathogenic effectors and virulence factors important for the pathogenicity of fungal species (Klosterman *et al.* 2011, Condon *et al.* 2013, Zhao *et al.* 2013, Deng *et al.* 2017). The availability of a genome sequence also provides the opportunity to develop species-specific microsatellite markers (Gnocato *et al.* 2017, Rafiei *et al.* 2018) that would be important for studying invasive populations of this aggressive, emerging tree pathogen.

Authors: F.A. Lane*, C.T. Tatham, J. Aylward, M. Havenga, P.M. Wilken, T.A. Duong, M.A. van der Nest, and B.D. Wingfield

*Contact Frances.Lane@FABI.up.ac.za

IMA Genome-F 10E

Draft nuclear genome assembly for *Davidsoniella eucalypti*

INTRODUCTION

The family *Ceratocystidaceae* includes an ecologically diverse assemblage of fungi currently classified into 11 distinct genera (De Beer *et al.* 2014, 2017, Mayers *et al.* 2015, Nel *et al.* 2018). The best-known of these is *Ceratocystis*, a genus that includes economically important plant pathogens such as the mango and *Acacia mangium* pathogen *C. manginecans* (Al Adawi *et al.* 2013, Tarigan *et al.* 2011) and the causal agent of cacao wilt *C. cacaofunesta* (Baker Engelbrecht & Harrington 2005). Species of the genera *Endoconidiophora* and *Thielaviopsis* are well known as pathogens of conifers and monocot plants, respectively (Mbenoun *et al.* 2014, Wingfield *et al.* 2013). The genera *Bretziella* and *Berkeleyomyces* were recently erected to accommodate the casual-agent of oak wilt *Br. fagacearum* (previously *Ceratocystis fagacearum*; De Beer *et al.* 2017) and the multi-host root pathogen *Be. basiola* (formerly *Thielaviopsis basicola*; Nel *et al.* 2018). Many *Ceratocystidaceae* species rely on flies, picnic beetles, and ambrosia beetles for spread, forming casual associations with these insects (Van Wyk *et al.* 2013). In contrast, species in the genera *Ambrosiella*, *Meredithiella*, and *Phialophoropsis*

form obligate mutualisms with ambrosia beetles of the tribes *Xyleborini*, *Corthylini* and *Xyloterini*, respectively (Mayers *et al.* 2015). Three asexual species associated with woody substrates are present in *Chalaropsis*, while members of the genus *Huntia* are considered saprobes or weak pathogens, with only a handful of species known to cause sapstain of timber (De Beer *et al.* 2014). The genus *Davidsoniella* consists of four species, three of which (*D. eucalypti*, *D. neocaledoniae* and *D. australis*) are present in Australasia, with *D. virescens* being described from North America (De Beer *et al.* 2014).

In this study we report a draft nuclear genome assembly for *Davidsoniella eucalypti*, first isolated from stem wounds on living *Eucalyptus* species in Australia (Kile *et al.* 1996). Although able to colonize wounds made in these trees, the fungus causes limited damage and is not considered pathogenic (Kile *et al.* 1996). In contrast, *D. virescens*, *D. australis*, and *D. neocaledoniae* are all considered plant-pathogens, causing disease on maple trees (*Acer* spp.), *Nothofagus cunninghamii* trees (Kile 1993), and *Coffea robusta* plants (Dadant 1950), respectively.

Davidsoniella eucalypti and *D. virescens* are the only two species in the genus for which a sexual morph is known (De Beer *et al.* 2014), although the reproductive strategy between these species differ dramatically – homothallism in *D. virescens* and heterothallism in *D. eucalypti* (Harrington *et al.* 1998). With the genome sequence of *D. virescens* already published (Wingfield *et al.* 2015b), the addition of the *D. eucalypti* genome brings the total number of *Davidsoniella* sequences to two. This raises the possibility for interesting genomic studies on these two biologically diverse species.

SEQUENCED STRAIN

Australia: Victoria: Cabbage Tree Creek, isolated from *Eucalyptus sieberi*, Aug. 1989, M.J. Dudzinski (CMW 3254, C 639, DAR 70205 – dried culture).

NUCLEOTIDE SEQUENCE ACCESSION NUMBER

This Whole Genome Shotgun project for *Davidsoniella eucalypti* isolate CMW 3254 has been deposited at DDBJ/ENA/GenBank under the accession RMBW00000000. The version described in this paper is version RMBW01000000.

MATERIALS AND METHODS

Davidsoniella eucalypti isolate CMW 3254 was obtained from the culture collection of the Forestry and Agricultural Biotechnology Institute (FABI) and grown on 2 % malt extract agar (MEA: 2 % w/v, Biolab, South Africa) at 25 °C. A 14-d old culture was used for genomic DNA extraction using a previously described phenol-chloroform protocol (Roux *et al.* 2004). The isolated DNA was submitted to Macrogen (Seoul, Korea) to generate long-read sequences from three cells of the Pacific BioSciences Single-molecule real time (SMRT or

PacBio) sequencing platform. This was complemented by a second round of sequencing on the Illumina HiSeq 2500 instrument at Macrogen. For this, a single library with 550 bp insert size was generated and used to produce pair-end reads of 250 bp target length. The DNA for the Illumina sequencing was extracted as described by Duong *et al.* (2013).

The reads obtained from the three cells of the SMRT sequencing run were concatenated into a single fastq file which was used for read-correction, trimming and assembly using Canu v1.4 and default settings (Koren *et al.* 2017). The resulting assembly was scaffolded using SSPACE-LongRead with the default settings (Boetzer *et al.* 2011, Boetzer & Pirovano 2014), using the corrected long-read sequences produced by Canu. The paired-end Illumina reads were imported into the CLC Genomics Workbench v11.0.1 (Qiagen, South Africa) and trimmed using default settings. These trimmed reads were indexed and aligned to the scaffolded, long-read assembly using BWA (Li & Durbin 2009) and SAMtools (Li *et al.* 2009). The alignment files were used in three rounds of Pilon corrections (Walker *et al.* 2014) to improve the long-read assembly by correcting single base differences, small insertions/deletions and other mis-assemblies identified in the draft genome assembly. To produce the best version of the assembly, the trimmed Illumina pair-end reads were used by GapFiller (Boetzer & Pirovano 2012) to fill gaps produced in the assembly during the scaffolding process. The draft genome was assessed for completeness using the Benchmarking Universal Single Copy Orthologs tool (BUSCO v 2.0.1) (Simão *et al.* 2015) and the *Ascomycota* database. An estimation of the number of protein coding genes in the genome was made by the *de novo* prediction software AUGUSTUS using the *Fusarium graminearum* gene models (Keller *et al.* 2011, Stanke *et al.* 2006b), while general genome statistics (genome length, GC content, N50, L50 and largest contig size) were calculated using QUAST v5.0.1 (Mikheenko *et al.* 2018).

The 60S, LSU and MCM7 gene regions were extracted from the genome and, together with these regions from *D. eucalypti*, *D. virescens*, *D. neocaledoniae*, *D. australis*, *Endoconidiophora polonica*, and *E. laricicola* (De Beer *et al.* 2014) were used for phylogenetic analysis. The datasets were subjected to a “one click” phylogeny analysis at the Phylogeny.fr online tool (Dereeper *et al.* 2008, 2010) that included a MUSCLE alignment (Edgar 2004) and a Gblocks (Castresana 2000) curation step before phylogenetic analysis was conducted using PhyML (Guindon & Gascuel 2003). Branch support was calculated using the approximate likelihood ratio test (Anisimova & Gascuel 2006).

RESULTS AND DISCUSSION

The 41 874 515 bp genome assembly of *Davidsoniella eucalypti* was present in 1219 contigs of 1000 bp or large, the largest of which was 1 065 836 bp. The genome had a G/C content of 45.93 %, an average coverage of 69x, a N50 value of 230 092 bp and a L50 value of 51. AUGUSTUS predicted 9029 protein coding genes, while BUSCO reported




Fig. 6. A maximum-likelihood phylogeny showing the position of the *Davidsoniella eucalypti* isolate used for this genome. Represented are the four known species of *Davidsoniella*, with two *Endoconidiophora* species used as outgroup. Approximate likelihood ratio test values for branch support are shown as percentages.

a completeness score of 96.1 %. This was based on the analysis of 1315 orthologs, with 1236 present as complete single copies, 27 as complete duplicated copies, 16 as fragmented copies, and 36 copies absent.

The genome assembly of *D. eucalypti* differed dramatically from that of the sister-species *D. virescens* (Fig. 6) (Wingfield *et al.* 2015b). At 33.6 Mb, the latter genome was 8.3 Mb smaller than that of *D. eucalypti*. The latter genome is also predicted to encode more proteins (9029 vs. 6953 for *D. virescens*), although the gene densities were comparable at 207 and 215 genes/Mb for *D. virescens* and *D. eucalypti* respectively. It is known that the genome sizes of plant-pathogenic filamentous fungi tend to be larger than those of non-pathogenic relatives, mostly due to the presence of high amounts of repetitive DNA and an expansion of the effector repertoire (Frantzeskakis *et al.* 2018, Möller & Stukenbrock 2017, Raffaele & Kamoun 2012). Therefore, the larger genome size of the non-pathogenic *D. eucalypti* as compared to the pathogenic species *D. virescens* was surprising and warrants further study.

The hybrid assembly of *D. eucalypti* presented here has a N-50 value (230092 bp) twice that of *D. virescens* (Wingfield *et al.* 2015b). This improvement in contig contiguity can be attributed to the inclusion of long-read PacBio sequences (English *et al.* 2012), a trend seen for many other genomes (Huddleston *et al.* 2014, Koren *et al.* 2013, Koren & Phillippy 2015). The availability of a highly contiguous genome sequence for one *Davidsoniella* species could provide the basis for genomic comparative studies. These should be of much interest as the higher gene number and larger genome size of *D. eucalypti* might point to an interesting evolutionary history for the genome of this species.

Authors: P.M. Wilken*, N. Soal, K. Naidoo, T.A. Duong, and B.D. Wingfield

*Contact: Markus.Wilken@fabu.up.ac.za

IMA Genome-F 10F

Draft genome sequence of *Grosmannia galeiformis*

INTRODUCTION

Grosmannia galeiformis was first described as *Ceratocystis galeiformes* from conifer infesting bark and ambrosia beetles in Scotland in 1951 (Bakshi 1951). It was later transferred to *Ophiostoma* (as *Ophiostoma galeiforme*; Mathiesen-Käärik 1953), and thereafter to *Grosmannia* (as *G. galeiformis*; Zipfel *et al.* 2006).

Grosmannia galeiformis is often found associated with conifer-infesting bark beetles and has been reported from Europe (Bakshi 1951, Linnakoski *et al.* 2012, Mathiesen-Käärik 1953, Zhou *et al.* 2004), South America (Zhou *et al.* 2004), and Africa (Zhou *et al.* 2004). Phylogenetic studies indicated that this species is part of a complex of several closely related species known as the *G. galeiformis* species complex, which is distinct from other species complexes in *Leptographium s. lat.* (Chang *et al.* 2019, De Beer *et al.* 2013, Linnakoski *et al.* 2012; Fig. 7). In this study, we sequenced and assembled the draft genome sequence for *G. galeiformis*, the key species representing the *G. galeiformis* species complex.

SEQUENCED STRAIN

United Kingdom: *Elgin*: on *Pinus sylvestris* (Scotch pine) infested with *Tomicus piniperda*, 29 Aug 1997, T. Kirisits & M.J. Wingfield (epitype isolate CMW 5290 = CBS 115711, PREM57491).

NUCLEOTIDE SEQUENCE ACCESSION NUMBER

The genomic sequence of *Grosmannia galeiformis* (CMW 5290, CBS 115711) has been deposited at DDBJ/EMBL/GenBank under accession no. RQWE00000000. The version described in this paper is version RQWE01000000.

METHODS

Genomic DNA was extracted from freeze-dried mycelium obtained from a single spore culture of *G. galeiformis* (CMW 5290). DNA extraction was done following a previously described method (Duong *et al.* 2013). Genome sequencing was carried out on the Illumina HiSeq 2000 platform (University of California Davis, CA). Two libraries (350 and 550 bp insert sizes) were prepared and sequenced to obtain 100 bp pair-end reads. Obtained pair-end reads were trimmed using Trimmomatic v. 0.36 (Bolger *et al.* 2014), and *de novo* assembled using SPAdes v. 3.9.0 (Bankevich *et al.* 2012). Scaffolds obtained from SPAdes was further placed into larger scaffolds using SSPACE-Standard v. 3.0 (Boetzer




Fig. 7. Phylogenetic tree generated from maximum likelihood analysis of a dataset consisted of partial beta-tubulin gene to authenticate the identity of *G. galeiformis* used in this study. Bootstrap values (≥ 70 ; 1000 replicates) are indicated at nodes. Beta-tubulin gene sequence for *G. galeiformis* was extracted directly from genome assembly. Other authenticated reference sequences were obtained from GenBank database.

et al. 2011), and assembly gaps were filled using GapFiller v. 1.10 (Boetzer & Pirovano 2012). The completeness of the resulting assembly was estimated using Benchmarking Universal Single Copy Orthologs (BUSCO) program v. 2.0.1 with the *Sordariomyceta* odb9 dataset (Simão *et al.* 2015). Protein coding gene models were predicted using the MAKER genome annotation pipeline (Cantarel *et al.* 2008) with the combination of GeneMark v. 4.32 (using self-training; Lomsadze *et al.* 2005) and AUGUSTUS v3.2.2 (using species models optimised for *Neurospora crassa*; Stanke *et al.* 2006a) as gene predictors.

RESULTS AND DISCUSSION

Over 25 million 100 bp pair-end reads were obtained after filtering and trimming. The final draft assembly consisted of 869 scaffolds that were over 500 bp in size. The assembly had a N50 of 67.79 Kb and a genome size of around 26.44 Mb. BUSCO reported the score for the assembly of 97 % [D:5.8 %, F:1.8 %, M:0.6 %, $n = 1348$ (C: complete; D: duplicated; F: fragmented; M: missing, $n =$ number of genes), which is comparable to that from other species of *Leptographium s. lat.* generated from previous studies (Wingfield *et al.* 2015a, 2016). Genome annotation using MAKER pipeline with GeneMark and AUGUSTUS as gene predictors resulted in 8527 protein-coding gene models. The genome of *G. galeiformis* generated in this study will add to the already growing genome resources for species of ophiostomatoid fungi (Wingfield *et al.* 2015a, b, 2016), which will facilitate future comparative genomic and evolutionary studies of these fungi (Fig. 7).

Authors: T.A. Duong*, M.J. Wingfield, Z.W. de Beer,
and B.D. Wingfield

*Contact: Tuan.Duong@fabi.up.ac.za

ACKNOWLEDGEMENTS

The authors of the *Claviceps purpurea s.lat.* study thank Zlatko Popovic, Rafik Assabgui, and the Molecular Technologies Laboratory (MTL) at the Ottawa Research & Development Centre specifically Julie T. Chapados, Wayne McCormick and Lisa Koziol for technical assistance. They also thank the Canadian Collection of Fungal Cultures (CCFC), Canadian National Mycological Herbarium (DAOM), and Jacques Cayouette for providing ergot samples or cultures. Genome sequencing was funded by Agriculture and Agri-Food Canada's Growing Forward 2 for a research network on Emerging Mycotoxins (EmTox, project # J-000048).

The authors of the *Davidsoniella eucalypti*, *Fusarium circinatum*, *Grosmannia galeiformis* and *Teratosphaeria destructans* study acknowledge funding received from the Department of Science and Technology (DST)/National Research Foundation (NRF) Centre of Excellence in Tree Health Biotechnology (CTHB), South Africa and the DST-NRF SARCHI chair in Fungal Genomics. The *F. circinatum* study was also supported by an NRF Scarce Skills Postdoctoral Fellowship awarded to the first author. P.M.W. received a scholarship from the Claude Leon foundation.

The authors of the *Quambalaria eucalypti* study thank the University of Pretoria, the Tree Protection Co-operative Programme (TPCP), the DST-NRF Centre of Excellence in Tree Health Biotechnology the National Research Foundation and the DST-NRF SARCHI chair in Fungal Genomics for their financial support.

REFERENCES

- Al Adawi AO, Barnes I, Khan IA, Al Subhi AM, Al Jahwari AA, *et al.* (2013) *Ceratocystis manginecans* associated with a serious wilt disease of two native legume trees in Oman and Pakistan. *Australasian Plant Pathology* **42**: 179–193.
- Alfenas AC, Zauza EAV, Rosa OPP, Assi TF (2001) Sporothrix eucalypti, um novo patogeno do eucalipto no Brasil. *Fitopatologia Brasileira* **26**: 221.
- Alm T (2003) The witch trials of Finnmark, Northern Norway, during the 17th century: evidence for ergotism as a contributing factor. *Economic Botany* **57**: 403–416.
- Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ (1990) Basic local alignment search tool. *Journal of Molecular Biology* **215**: 403–410.
- Andjic V, Dell B, Barber P, Hardy G, Wingfield M, *et al.* (2011) Plants for planting; indirect evidence for the movement of a serious forest pathogen, *Teratosphaeria destructans*, in Asia. *European Journal of Plant Pathology* **131**: 49–58.
- Anisimova M, Gascuel O (2006) Approximate likelihood-ratio test for branches: a fast, accurate, and powerful alternative. *Systematic Biology* **55**: 539–552.
- Aylward J, Roets F, Dreyer LL, Wingfield MJ (2018) *Teratosphaeria* stem canker of *Eucalyptus*: two pathogens, one devastating disease. *Molecular Plant Pathology*. Epub ahead of print. DOI:10.1111/mpp.12758
- Baker Engelbrecht CJ, Harrington TC (2005) Intersterility, morphology and taxonomy of *Ceratocystis fimbriata* on sweet potato, cacao and sycamore. *Mycologia* **97**: 57–69.
- Bakshi BK (1951) Studies on four species of *Ceratocystis*, with a discussion on fungi causing sapstain in Britain. *Mycological Paper* **35**: 1–16.
- Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, *et al.* (2012) SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *Journal of Computational Biology* **19**: 455–477.
- Barber PA (2004) Forest pathology: the threat of disease to plantation forests in Indonesia. *Plant Pathology Journal* **3**: 97–104.
- Barber PA, Thu PQ, Hardy GE, Dell B (2012) Emerging disease problems in eucalypt plantations in Lao PDR. In: *Proceedings of International Conference on the Impacts of Climate Change to Forest Pests and Diseases in the Tropics, 8–10 October* (Mohammed C, Beadle C, Roux J, *et al.*, eds): 79–84, Indonesia: Yogyakarta.
- Boetzer M, Henkel CV, Jansen HJ, Butler D, Pirovano W (2011) Scaffolding pre-assembled contigs using SSPACE. *Bioinformatics* **27**: 578–579.
- Boetzer M, Pirovano W (2012) Toward almost closed genomes with GapFiller. *Genome Biology* **13**: R56.
- Boetzer M, Pirovano W (2014) SSPACE-LongRead: scaffolding bacterial draft genomes using long read sequence information. *BMC Bioinformatics* **15**: 211.
- Bolger AM, Lohse M, Usadel B (2014) Trimmomatic: A flexible trimmer for Illumina sequence data. *Bioinformatics* **30**: 2114–2120.
- Bragança H, Diogo ELF, Neves L, Valente C, Araujo C, *et al.* (2015) *Quambalaria eucalypti* a pathogen of *Eucalyptus globulus* newly reported in Portugal and in Europe. *Forest Pathology* **46**: 67–75.
- Burgess TI, Andjic V, Hardy GE, Dell B, Xu D (2006) First report of *Phaeocephalospora destructans* in China. *Journal of Tropical Forest Science* **18**: 144–146.
- Burgess TI, Wingfield MJ (2017) Pathogens on the move: a 100-year global experiment with planted eucalypts. *BioScience* **67**: 14–25.
- Cantarel BL, Korf I, Robb SMC, Parra G, Ross E, *et al.* (2008) MAKER: An easy-to-use annotation pipeline designed for emerging model organism genomes. *Genome Research* **18**: 188–196.
- Caporael LR (1976) Ergotism: the satan loosed in Salem? *Science* **192**: 21
- Castresana J (2000) Selection of conserved blocks from multiple alignments for their use in phylogenetic analysis. *Molecular Biology and Evolution* **17**: 540–552.
- Chang R, Duong TA, Taerum SJ, Wingfield MJ, Zhou X, De Beer ZW (2019) Ophiostomatoid fungi associated with the spruce bark beetle *Ips typographus*, including 11 new species from China. *Persoonia* **42**: 50–74.
- Chiara M, Fanelli F, Mulè G, Logrieco AF, Pesole G, *et al.* (2015) Genome sequencing of multiple isolates highlights sub-telomeric genomic diversity within *Fusarium fujikuroi*. *Genome Biology and Evolution* **7**: 3062–3069.
- Condon BJ, Leng Y, Wu D, Bushley KE, Ohm RA, *et al.* (2013) Comparative genome structure, secondary metabolite, and effector coding capacity across *Cochliobolus* pathogens. *PLoS Genetics* **9**: e1003233.
- Crosignani PG (2006) Current treatment issues in female hyperprolactinaemia. *European Journal of Obstetrics & Gynecology and Reproductive Biology* **125**: 152–164.
- Crous PW, Groenewald JZ, Summerell BA, Wingfield BD, Wingfield MJ (2009) Co-occurring species of *Teratosphaeria* on *Eucalyptus*.

- Persoonia* **22**: 38–48.
- Dadant R (1950) On a new disease of *Coffea robusta* in New Caledonia. *Revue Generale de Botanique* **57**: 1–11.
- Daribo D, Taboada GL, Doallo R, Posada D (2012) jModelTest 2: More models, new heuristics and parallel computing. *Nature Methods* **9**: 772.
- De Beer ZW, Begerow D, Bauer R, Pegg GS, Crous PW, *et al.* (2006) Phylogeny of the *Quambalariaceae* fam. nov., including important *Eucalyptus* pathogens in South Africa and Australia. *Studies in Mycology* **55**: 289–298.
- De Beer ZW, Duong TA, Barnes I, Wingfield BD, Wingfield MJ (2014) Redefining *Ceratocystis* and allied genera. *Studies in Mycology* **79**: 187–219.
- De Beer ZW, Marincowitz S, Duong TA, Wingfield MJ (2017) *Bretziella*, a new genus to accommodate the oak wilt fungus, *Ceratocystis fagacearum* (*Microascales*, *Ascomycota*). *Mycology* **27**: 1–19.
- De Beer ZW, Wingfield MJ (2013) Emerging lineages in the *Ophiostomatales*. In: *The Ophiostomatoid Fungi: expanding frontiers* (Seifert KA, *et al.*, eds): 21–46. [CBS Biodiversity Series no. 12.] Utrecht: CBS-KNAW Fungal Biodiversity Institute.
- De Groot ANJA, Van Dongen PWJ, Vree TB, Hekster YA, Van Roosmalen J (1998) Ergot alkaloids: current status and review of clinical pharmacology and therapeutic use compared with other oxytocics in obstetrics and gynaecology. *Drugs* **56**: 523–535.
- De Vos L, Myburg AA, Wingfield MJ, Desjardins AE, Gordon TR, *et al.* (2007) Complete genetic linkage maps from an interspecific cross between *Fusarium circinatum* and *Fusarium subglutinans*. *Fungal Genetics and Biology* **44**: 701–714.
- De Vos L, Steenkamp ET, Martin SH, Santana QC, Fourie G, *et al.* (2014) Genome-wide macrosynteny among *Fusarium* species within the *Gibberella fujikuroi* complex revealed by amplified fragment length polymorphisms. *PLoS One* **9**: e114682.
- De Vos L, Van der Nest MA, Van der Merwe NA, *et al.* (2011) Genetic analysis of growth, morphology and pathogenicity in the F_1 progeny of an interspecific cross between *Fusarium circinatum* and *Fusarium subglutinans*. *Fungal Biology* **115**: 902–908.
- Deng CH, Plummer KM, Jones DAB, Mesarich CH, Shiller J, *et al.* (2017) Comparative analysis of the predicted secretomes of Rosaceae scab pathogens *Venturia inaequalis* and *V. pirina* reveals expanded effector families and putative determinants of host range. *BMC Genomics* **18**: 339.
- Dereeper A, Audic S, Claverie JM, Blanc G (2010) BLAST-EXPLORER helps you building datasets for phylogenetic analysis. *BMC Evolutionary Biology* **10**: 8.
- Dereeper A, Guignon V, Blanc G, Audic S, Buffet S, *et al.* (2008) Phylogeny.fr: robust phylogenetic analysis for the non-specialist. *Nucleic Acids Research* **36**: W465–W469.
- Douhan GW, Smith ME, Huyrn KL, Westbrook A, Beerli P, *et al.* (2008) Multigene analysis suggests ecological speciation in the fungal pathogen *Claviceps purpurea*. *Molecular Ecology* **17**: 2276–2286.
- Doyle JJ, Doyle JL (1987) A rapid DNA isolation procedure for small quantities of fresh leaf tissue. *Phytochemistry Bulletin* **190**: 11–15
- Duong TA, de Beer ZW, Wingfield BD, Wingfield MJ (2013) Characterization of the mating-type genes in *Leptographium procerum* and *Leptographium profanum*. *Fungal Biology* **117**: 411–421.
- Edgar RC (2004) MUSCLE: multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids Research* **32**: 1792–1797.
- English AC, Richards S, Han Y, Wang M, Vee V, *et al.* (2012) Mind the gap: upgrading genomes with Pacific Biosciences RS long-read sequencing technology. *PLoS ONE* **7**: e47768.
- Frantzeskakis L, Kracher B, Kusch S, Yoshikawa-Maekawa M, Bauer S, *et al.* (2018) Signatures of host specialization and a recent transposable element burst in the dynamic one-speed genome of the fungal barley powdery mildew pathogen. *BMC Genomics* **19**: 381.
- Fuller JG (1968) *The Day of St. Anthony's Fire: the suspenseful, true account of a medieval plague in modern times, and of the scientific detective work that traced it to a surprising cause*. New York: Macmillan.
- Garc'a-Alcalde F, Okonechnikov K, Carbonell J, Cruz LM, G'ztz S, *et al.* (2012) Qualimap: evaluating next-generation sequencing alignment data. *Bioinformatics* **28**: 2678–2679
- Gardiner DM (2018) Genome sequences of three isolates of *Fusarium verticillioides*. *Microbiology Resource Announcements* **7**: e00918-18.
- Geiser DM, Aoki T, Bacon CW, Baker SE, Bhattacharyya MK, *et al.* (2013) One fungus, one name: Defining the genus *Fusarium* in a scientifically robust way that preserves longstanding use. *Phytopathology* **103**: 400–408.
- Gnocato FS, Dracatos PM, Karaoglu H, Zhang P, Berlin A, *et al.* (2017) Development, characterization and application of genomic SSR markers for the oat stem rust pathogen *Puccinia graminis* f. sp. *avenae*. *Plant Pathology* **67**: 457–466.
- Gordon TR, Storer AJ & Okamoto D (1996) Population structure of the pitch canker pathogen, *Fusarium subglutinans* f. sp. *pini*, in California. *Mycological Research* **100**: 850–854.
- Gordon TR, Swett CL & Wingfield MJ (2015) Management of *Fusarium* diseases affecting conifers. *Crop Protection* **73**: 28–39.
- Greyling I, Wingfield MJ, Coetzee MPA, Marincowitz S, Roux J (2016) The *Eucalyptus* shoot and leaf pathogen *Teratosphaeria destructans* recorded in South Africa. *Southern Forests* **78**: 123–129.
- Guindon S, Dufayard J-F, Lefort V, Anisimova M, Hordijk W, *et al.* (2010) New algorithms and methods to estimate maximum-likelihood phylogenies: Assessing the performance of PhyML 3.0. *Systematic Biology* **59**: 307–321.
- Guindon S, Gascuel O (2003) A simple, fast, and accurate algorithm to estimate large phylogenies by maximum likelihood. *Systematic Biology* **52**: 696–704.
- Gurevich A, Saveliev V, Vyahhi N, Tesler G (2013) QUASt: Quality assessment tool for genome assemblies. *Bioinformatics* **29**: 1072–1075.
- Hall B, DeRego T, Geib S (2014) GAG: the genome annotation generator (version 1.0): <http://genomeannotation.github.io/GAG>.
- Harrington TC, Steimel J, Kile G (1998) Genetic variation in three *Ceratocystis* species with outcrossing, selfing and asexual reproductive strategies. *European Journal of Forest Pathology* **28**: 217–226.
- Huddleston J, Ranade S, Malig M, Antonacci F, Chaisson M, *et al.* (2014) Reconstructing complex regions of genomes using long-read sequencing technology. *Genome Research* **24**: 688–696.
- Hunter GC, Crous PW, Carnegie AJ, Burgess TI, Wingfield MJ (2011) *Mycosphaerella* and *Teratosphaeria* diseases of *Eucalyptus*; easily confused and with serious consequences. *Fungal Diversity* **50**: 145–166.
- James TY, Sun S, Li W, Heitmen J, Kuo HC, *et al.* (2013) *Polyporales* genomes reveal the genetic architecture underlying tetrapolar and bipolar mating systems. *Mycologia* **105**: 1374–1390.

- Jeong H, Lee S, Choi GJ, Lee S, Yun S-H, *et al.* (2013) Draft genome sequence of *Fusarium fujikuroi* B14, the causal agent of the bakanae disease of rice. *Genome Announcements* **1**: e00035-13.
- Katoh K, Standley DM (2013) MAFFT multiple sequence alignment software version 7: Improvements in performance and usability. *Molecular Biology and Evolution* **30**: 772–780.
- Kearse M, Moir R, Wilson A, Stones-Havas S, Cheung M, *et al.* (2012) Geneious Basic: An integrated and extendable desktop software platform for the organization and analysis of sequence data. *Bioinformatics* **28**: 1647–1649.
- Keller O, Kollmar M, Stanke M, Waack S (2011) A novel hybrid gene prediction method employing protein multiple sequence alignments. *Bioinformatics* **27**: 757–763.
- Kijpornyongpan T, Aime MC (2017) Taxonomic revisions in the Microstromatales: two new yeast species, two new genera, and validation of *Jaminaea* and two *Symptodiomyopsis* species. *Mycological Progress* **16**: 495–505.
- Kijpornyongpan T, Mondo SJ, Barry K, Sandor L, Lee L, *et al.* (2018) Broad genomic sampling reveals a smut pathogenic ancestry of the fungal clade *Ustilaginomycotina*. *Molecular Biology and Evolution* **35**: 1840–1854.
- Kile G (1993). Plant diseases caused by species of *Ceratocystis sensu stricto* and *Chalara*. In: *Ceratocystis and Ophiostoma: taxonomy, ecology and pathogenicity* (Wingfield MJ, Seifert KA, Webber JF, eds): 173–184. S. Paul, MN: American Phytopathological Society Press.
- Kile GA, Harrington TC, Yuan ZQ, Dudzinski MJ, Old KM (1996) *Ceratocystis eucalypti* sp. nov., a vascular stain fungus from eucalypts in Australia. *Mycological Research* **100**: 571–579.
- Klosterman SJ, Subbarao K V, Kang S, Veronese P, Gold SE, *et al.* (2011) Comparative genomics yields insights into niche adaptation of plant vascular wilt pathogens. *PLoS Pathogens* **7**: e1002137.
- Kolarik M, Sláviková E, Pažoutová S (2006) The taxonomic and ecological characterisation of the clinically important heterobasidiomycete *Fugomyces cyanescens* and its association with bark beetles. *Czech Mycology* **58**: 81–98.
- Konishi M, Hatada Y, Horiuchi J (2013) Draft genome sequence of the basidiomycetous yeast-like fungus *Pseudozyma hubeiensis* SY62, which produces an abundant amount of the biosurfactant mannosylerythritol lipids. *Genome Announcements* **1**: e00409-13–e00409-13.
- Koren S, Harhay GP, Smith TP, Bono JL, Harhay DM, *et al.* (2013) Reducing assembly complexity of microbial genomes with single-molecule sequencing. *Genome Biology* **14**: R101.
- Koren S, Phillippy AM (2015) One chromosome, one contig: complete microbial genomes from long-read sequencing and assembly. *Current Opinion in Microbiology* **23**: 110–120.
- Koren S, Walenz BP, Berlin K, Miller JR, Bergman NH, *et al.* (2017) Canu: scalable and accurate long-read assembly via adaptive *k*-mer weighting and repeat separation. *Genome Research* **27**: 722–736.
- Korf I (2004) Gene finding in novel genomes. *BMC Bioinformatics* **5**: 59.
- Kuan CS, Yew SM, Toh YF, Chan CL, Lim SK, *et al.* (2015) Identification and characterization of a rare fungus, *Quambalaria cyanescens*, isolated from the peritoneal fluid of a patient after nocturnal intermittent peritoneal dialysis. *PLoS ONE* **10**: 1–15.
- Kurtz S, Phillippy A, Delcher AL, Smoot M, Shumway M, *et al.* (2004) Versatile and open software for comparing large genomes. *Genome Biology* **5**: R12.
- Langmead B, Salzberg SL (2012) Fast gapped-read alignment with Bowtie 2. *Nature Methods* **9**: 357–359.
- Leslie JF, Summerell BA (2006) *The Fusarium Laboratory Manual*. Iowa: Wiley-Blackwell.
- Li H, Durbin R (2009) Fast and accurate short read alignment with Burrows-Wheeler transform. *Bioinformatics* **25**: 1754–1760.
- Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, *et al.* (2009) The sequence alignment/map format and SAMtools. *Bioinformatics* **25**: 2078–2079.
- Linnakoski R, De Beer ZW, Duong TA, Niemelä P, Pappinen A, Wingfield MJ (2012) *Grossmannia* and *Leptographium* spp. associated with conifer-infesting bark beetles in Finland and Russia, including *L. taigense* sp. nov. *Antonie van Leeuwenhoek* **102**: 375–399.
- Lomsadze A, Burns PD, Borodovsky M (2014) Integration of mapped RNA-Seq reads into automatic training of eukaryotic gene finding algorithm. *Nucleic Acids Research* **42**: e119.
- Lomsadze A, Ter-Hovhannisyán V, Chernoff Y, Borodovsky M (2005) Gene identification in novel eukaryotic genomes by self-training algorithm. *Nucleic Acids Research* **33**: 6494–6506.
- Lyons PC, Plattner RD, Bacon CW (1986) Occurrence of peptide and clavine ergot alkaloids in tall fescue grass. *Science* **232**: 487–489.
- Ma L-J, Van der Does HC, Borkovich KA, Coleman JJ, Daboussi MJ, *et al.* (2010) Comparative genomics reveals mobile pathogenicity chromosomes in *Fusarium*. *Nature* **464**: 367–373.
- Mathiesen-Käärik A (1953) Eine Übersicht über die gewöhnlichsten mit Borkenkäfern assoziierten Bläuepilze in Schweden und einige für Schweden neue Bläuepilze. *Meddelanden från Statens Skogsforskningsinstitut* **43**: 1–74.
- Mayers CG, McNew DL, Harrington TC, Roeper RA, Fraedrich SW, *et al.* (2015) Three genera in the *Ceratocystidaceae* are the respective symbionts of three independent lineages of ambrosia beetles with large, complex mycangia. *Fungal Biology* **119**: 1075–1092.
- Mbenoun M, De Beer ZW, Wingfield BD, Roux J (2014) Reconsidering species boundaries in the *Ceratocystis paradoxa* complex, including a new species from oil palm and cacao in Cameroon. *Mycologia* **106**: 757–784.
- Menzies JG, Turkington TK (2015) An overview of the ergot (*Claviceps purpurea*) issue in western Canada: Challenges and solutions. *Canadian Journal of Plant Pathology* **37**: 40–51.
- Micale V, Incognito T, Ignoto A, Rampello L, Spart M, *et al.* (2006) Dopaminergic drugs may counteract behavioral and biochemical changes induced by models of brain injury. *European Neuropsychopharmacology* **16**: 195–203.
- Mikheenko A, Prjibelski A, Saveliev V, Antipov D, Gurevich A (2018) Versatile genome assembly evaluation with QUAST-LG. *Bioinformatics* **34**: i142–i150.
- Miles C, Lane G, E. di Menna M, Garthwaite I, L. Piper E, *et al.* (1996) High levels of ergonovine and lysergic acid amide in toxic *Achnatherum inebrians* accompany infection by an *Acremonium*-like endophytic fungus. *Journal of Agricultural and Food Chemistry* **44**: 1285–1290.
- Möller EM, Bahnweg G, Sandermann H, Geiger HH, (1992) A simple and efficient protocol for isolation of high molecular weight DNA from filamentous fungi, fruit bodies, and infected plant tissues. *Nucleic Acids Research* **20**: 6115–6116.

- Möller M, Stukenbrock EH (2017) Evolution and genome architecture in fungal plant pathogens. *Nature Reviews Microbiology* **15**: 756–771.
- Nel WJ, Duong TA, Wingfield BD, Wingfield MJ, De Beer ZW (2018) A new genus and species for the globally important, multihost root pathogen *Thielaviopsis basicola*. *Plant Pathology* **67**: 871–882.
- Niehaus E-M, Kim H-K, Münsterkötter M, Janevskis S, Arndt B, *et al.* (2017a) Comparative genomics of geographically distant *Fusarium fujikuroi* isolates revealed two distinct pathotypes correlating with secondary metabolite profiles. *PLoS Pathogens* **13**: e1006670.
- Niehaus E-M, Münsterkötter M, Proctor RH, Brown DW, Sharon A, *et al.* (2017b) Comparative “omics” of the *Fusarium fujikuroi* species complex highlights differences in genetic potential and metabolite synthesis. *Genome Biology and Evolution* **8**: 3574–3599.
- Nikolenko SI, Korobeynikov AI, Alekseyev MA (2013) BayesHammer: Bayesian clustering for error correction in single-cell sequencing. *BMC Genomics* **14** (suppl. 1) :S7.
- Nurk S, Bankevich A, Antipov D, Gurevich A, Korobeynikov A, *et al.* (2013) Assembling genomes and mini-metagenomes from highly chimeric reads. *Research in Computational Molecular Biology* **7821**: 158–170.
- O'Donnell K, Cigelnik E & Nirenberg HI (1998) Molecular systematics and phylogeography of the *Gibberella fujikuroi* species complex. *Mycologia* **90**: 465–493.
- O'Donnell K, Nirenberg HI, Aoki T, Cigelnik E (2000) A multigene phylogeny of the *Gibberella fujikuroi* species complex: detection of additional phylogenetically distinct species. *Mycoscience* **41**: 61–78.
- Old KM, Pongpanich K, Thu PQ, Wingfield MJ, Yuan ZQ (2003) *Phaeophleospora destructans* causing leafblight epidemics in South East Asia. *Feb. Vol. 2 – Offered papers* ICPP: 165, Christchurch, New Zealand
- Paap T, Burgess TI, McComb JA, Shearer, BL, Hardy GE (2008) *Quambalaria* species, including *Q. coyrecup* sp. nov., implicated in canker and shoot blight diseases causing decline of *Corymbia* species in the southwest of Western Australia. *Mycological Research* **112**: 57–69.
- Park RF, Keane PJ, Wingfield MJ, Crous PW (2000) Fungal diseases of eucalypt foliage. In: *Diseases and Pathogens of Eucalypts* (Keane PJ, Kile GA, Podger FD, *et al.*, eds): 153–239. Collingwood: CSIRO Publishing.
- Pažoutová S, Pešicov K, Chud'čkov M, Šrůtka P, Kolařík M (2015) Delimitation of cryptic species inside *Claviceps purpurea*. *Fungal Biology* **119**: 7–26.
- Pažoutová S, Olsovská J, Linka M, Kolínská R, Flieger M (2000) Chemoraces and habitat specialization of *Claviceps purpurea* populations. *Applied and Environmental Microbiology* **66**: 5419–5425.
- Pegg GS, O'Dwyer C, Carnegie AJ, Burgess TI, Wingfield MJ, *et al.* (2008) *Quambalaria* species associated with plantation and native eucalypts in Australia. *Plant Pathology* **57**: 702–714.
- Pegg GS, Shuey LS, Carnegie AJ, Wingfield MJ, Drenth A (2011) Variability in aggressiveness of *Quambalaria pitereka* isolates. *Plant Pathology* **60**: 1107–1117.
- Pegg GS, Webb RI, Carnegie AJ, Wingfield MJ, Drenth A (2009) Infection and disease development of *Quambalaria* spp. on *Corymbia* and *Eucalyptus* species. *Plant Pathology* **58**: 642–654.
- Perez CA, De Beer ZW, Altier NA, Wingfield MJ, Blanchette RA (2008) Discovery of the eucalypt pathogen *Quambalaria eucalypti* infecting a non-Eucalyptus host in Uruguay. *Australasian Plant Pathology* **37**: 600–604.
- Quaedvlieg W, Binder M, Groenewald JZ, Summerell BA, Carnegie AJ, *et al.* (2014) Introducing the Consolidated Species Concept to resolve species in the *Teratosphaeriaceae*. *Persoonia* **33**: 1–40.
- Que Y, Xu L, Wu Q, Liu, Y, Ling H, *et al.* (2014) Genome sequencing of *Sporisorium scitamineum* provides insights into the pathogenic mechanisms of sugarcane smut. *BMC Genomics* **15**: 996.
- Raffaele S, Kamoun S (2012) Genome evolution in filamentous plant pathogens: why bigger can be better. *Nature Reviews Microbiology* **10**: 417–430.
- Rafiei V, Banihashemi Z, Bautista-Jalon LS, Del Mar Jiménez-Gasco M, Turgeon BG, *et al.* (2018) Population genetics of *Verticillium dahliae* in Iran based on microsatellite and single nucleotide polymorphism markers. *Phytopathology* **108**: 780–788.
- Ronquist F, Teslenko M, van der Mark P, Ayres DL, Darling A, *et al.* (2012) MrBayes 3.2: Efficient bayesian phylogenetic inference and model choice across a large model space. *Systematic Biology* **61**: 539–542.
- Roux J, van Wyk M, Hatting H, Wingfield MJ (2004) *Ceratocystis* species infecting stem wounds on *Eucalyptus grandis* in South Africa. *Plant Pathology* **53**: 414–421.
- Saika A, Koike H, Hori T, Fukuoak T (2014) Draft genome sequence of the yeast *Pseudozyma antarctica* type strain JCM10317, a producer of the glycolipid biosurfactants, mannosylerythritol lipids. *Genome Announcements* **2**: 4–5.
- Scauflaire J, Gourgue M, & Munaut F (2011) *Fusarium temperatum* sp. nov. from maize, an emergent species closely related to *Fusarium subglutinans*. *Mycologia* **103**: 586–597.
- Schardl CL, Young CA, Hesse U, Amyotte SG, Andreeva K, *et al.* (2013) Plant-symbiotic fungi as chemical engineers: multi-genome analysis of the clavicipitaceae reveals dynamics of alkaloid loci. *PLoS Genetics* **9**: e1003323.
- Scott P (2009) Ergot alkaloids: extent of human and animal exposure. *World Mycotoxin Journal* **2**: 141–149.
- Simão FA, Waterhouse RM, Ioannidis P, Kriventseva E V, Zdobnov EM (2015) BUSCO: assessing genome assembly and annotation completeness with single-copy orthologs. *Bioinformatics* **31**: 3210–3212.
- Srivastava, AK, Kashyap, PL, Chakdar, H, Kumar M, Srivastava AK, *et al.* (2018) First *de novo* draft genome sequence of the pathogenic fungus *Fusarium udum* F02845, associated with pigeonpea (*Cajanus cajan* L. Millspaugh) wilt. *Microbiology Resource Announcements* **7**: e01001-18.
- Stanke M, Morgenstern B (2005) AUGUSTUS: a web server for gene prediction in eukaryotes that allows user-defined constraints. *Nucleic Acids Research* **33**: W465–W467.
- Stanke M, Diekhans M, Baertsch R, Haussler D (2008) Using native and syntenically mapped cDNA alignments to improve de novo gene finding. *Bioinformatics* **24**: 637–644.
- Stanke M, Keller O, Gunduz I, Hayes A, Waack S, Morgenstern B (2006a) AUGUSTUS: *ab initio* prediction of alternative transcripts. *Nucleic Acids Research* **34**: W435–W439.
- Stanke M, Tzvetkova A, Morgenstern B (2006b) AUGUSTUS at EGASP: using EST, protein and genomic alignments for improved gene prediction in the human genome. *Genome Biology* **7** (suppl. 1): S11.11–S11.18.

- Steenkamp ET, Makhari OM, Coutinho TA, Wingfield BD, Wingfield MJ (2014) Evidence for a new introduction of the pitch canker fungus *Fusarium circinatum* in South Africa. *Plant Pathology* **63**: 530–538.
- Swofford DL (2002) *PAUP**. *Phylogenetic analysis using parsimony (*and other methods)*. Version 4.0 b10 (computer program). Sunderland MA: Sinauer Associates
- Tarigan M, Roux J, van Wyk M, Tjahjono B, Wingfield MJ (2011) A new wilt and die-back disease of *Acacia mangium* associated with *Ceratocystis manginecans* and *C. acaciivora* sp. nov. in Indonesia. *South African Journal of Botany* **77**: 292–304.
- Tavare S (1986) Some probabilistic and statistical problems in the analysis of DNA sequences. In: *Lectures on Mathematics in the Life Sciences* **17**: 57–86, American Mathematical Society, Rhode Island.
- Ter-Hovhannissyan V, Lomsadze A, Chernoff YO, Borodovsky M (2008) Gene prediction in novel fungal genomes using an *ab initio* algorithm with unsupervised training. *Genome Research* **18**: 1979–1990.
- Toome M, Kuo A, Henrissat B, Lipzen A, Tritt A, et al. (2014) Draft genome sequence of a rare smut relative, *Tilletiaria anomala* UBC 951. *Genome Announcements* **2**: 2–3.
- Van der Linde EJ, Pešicov K, Pažoutová S, Stodůlkov E, Flieger M, et al. (2016) Ergot species of the *Claviceps purpurea* group from South Africa. *Fungal Biology* **120**: 917–930.
- Van der Nest MA, Beirn LA, Crouch JA, Demers JE, De Beer ZW, et al. (2014) Draft genomes of *Amanita jacksonii*, *Ceratocystis albifundus*, *Fusarium circinatum*, *Huntia omanensis*, *Leptographium procerum*, *Rutstroemia sydowiana*, and *Sclerotinia echinophila*. *IMA Fungus* **5**: 473–486.
- Van Wyk M, Wingfield BD, Wingfield MJ (2013) *Ceratocystis* species in the *Ceratocystis fimbriata* complex. In: *The Ophiostomatoid Fungi: expanding frontiers* (Seifert KA, De Beer ZW, Wingfield MJ, eds): 65–73. [CBS Biodiversity Series no. 12.] Utrecht: CBS-KNAW Fungal Biodiversity Institute.
- Van Wyk S, Wingfield BD, De Vos L, Santana QC, Van der Merwe NA, et al. (2018) Multiple independent origins for a subtelomeric locus associated with growth rate in *Fusarium circinatum*. *IMA Fungus* **9**: 27–36.
- Walker BJ, Abeel T, Shea T, Priest M, Abouelliel A, et al. (2014) Pilon: an integrated tool for comprehensive microbial variant detection and genome assembly improvement. *PLoS ONE* **9**: e112963.
- Wang N, Ai P, Tang Y, Zhang J, Dai X, et al. (2015a) Draft genome sequence of the rice kernel smut *Tilletia horrida* Strain QB-1. *Genome Announcements* **3**: e00621-15.
- Wang QM, Begerow D, Groenewald M, Liu XZ, Theelen B, et al. (2015b) Multigene phylogeny and taxonomic revision of yeasts and related fungi in the Ustilaginomycotina. *Studies in Mycology* **81**: 55–83.
- Waterhouse RM, Seppey M, Simão FA, Manni M, Ioannidis P, et al. (2017) BUSCO applications from quality assessments to gene prediction and phylogenomics. *Molecular Biology and Evolution* **35**: 543–548.
- Wiemann P, Sieber CMK, Von Bargen K, Studt L, Niehaus EM, et al. (2013) Unleashing the cryptic genome: genome-wide analyses of the rice pathogen *Fusarium fujikuroi* reveal complex regulation of secondary metabolism and novel metabolites. *PLoS Pathogens* **9**: e1003475.
- Wingfield BD, Ades PK, Al-Naemi FA, Beir LA, Bihon W, et al. (2015a) IMA Genome-F 4. Draft genome sequences of *Chrysosporthe austroafricana*, *Diplodia scrobiculata*, *Fusarium nygamai*, *Leptographium lundbergii*, *Limonomyces culmigenus*, *Stagonosporopsis tanacetii*, and *Thielaviopsis punctulata*. *IMA Fungus* **6**: 233–248.
- Wingfield BD, Barnes I, De Beer ZW, de Vos L, Duong TA, et al. (2015b) IMA Genome-F 5. Draft genome sequences of *Ceratocystis eucalypticola*, *Chrysosporthe cubensis*, *C. deuterocubensis*, *Davidsoniella virescens*, *Fusarium temperatum*, *Graphilbum fragrans*, *Penicillium nordicum*, and *Thielaviopsis musarum*. *IMA Fungus* **6**: 493–506.
- Wingfield BD, Berger DK, Steenkamp ET, Lim HJ, Duong TA, et al. (2017) IMA Genome-F 6: draft genome of *Cercospora zeina*, *Fusarium pininemorale*, *Hawksworthiomyces lignivorus*, *Huntia decipiens* and *Ophiostoma ips*. *IMA Fungus* **8**: 385–396.
- Wingfield BD, Bills GF, Dong Y, Huang W, Nei WJ, et al. (2018) IMA Genome-F 9. Draft genome sequence of *Annulohyphoxylon stygium*, *Aspergillus mulundensis*, *Berkeleyomyces basicola* (syn. *Thielaviopsis basicola*), *Ceratocystis smalleyi*, two *Cercospora beticola* strains, *Coleophoma cylindrospora*, *Fusarium fracticaudum*, *Phialophora* cf. *hyalina*, and *Morchella septimelata*. *IMA Fungus* **9**: 199–223.
- Wingfield BD, Duong TA, Hammerbacher A, Van der Nest MA, Wilson A, et al. (2016) IMA Genome-F 7. Draft genome sequences for *Ceratocystis fagacearum*, *C. harringtonii*, *Grosmannia penicillata*, and *Huntia bhutanensis*. *IMA Fungus* **7**: 317–323.
- Wingfield BD, Steenkamp ET, Santana QC, Coetzee MPA, Bam S, et al. (2012) First fungal genome sequence from Africa: a preliminary analysis. *South African Journal of Science* **108**: 1–2.
- Wingfield BD, Van Wyk M, Roos H, Wingfield MJ (2013) *Ceratocystis*: emerging evidence for discrete generic boundaries. In: *The Ophiostomatoid Fungi: expanding frontiers* (Seifert KA, De Beer ZW, Wingfield MJ, eds): 57–64. [CBS Biodiversity Series no. 12.] Utrecht: CBS-KNAW Fungal Biodiversity Institute.
- Wingfield MJ, Crous PW, Boden D (1996) *Kiramyces destructans* sp. nov., a serious leaf pathogen of *Eucalyptus* in Indonesia. *South African Journal of Botany* **62**: 325–327.
- Wingfield MJ, Crous PW, Swart WJ (1993) *Sporothrix eucalypti* (sp. nov.), a shoot and leaf pathogen of *Eucalyptus* in South Africa. *Mycopathologia* **123**: 159–164.
- Xu J, Saunders CW, Hu P, Grant RA, Noekhout T, et al. (2007) Dandruff-associated *Malassezia* genomes reveal convergent and divergent virulence traits shared with plant and human fungal pathogens. *Proceedings of the National Academy of Sciences, USA* **104**: 18730–18735.
- Xu J-R, Yan K, Dickman MB, Leslie JF, (1995) Electrophoretic karyotypes distinguish the biological species of *Gibberella fujikuroi* (*Fusarium* section *Liseola*). *Molecular Plant-Microbe Interactions* **8**: 74–84.
- Zhao Z, Liu H, Wang C, Xu JR (2013) Comparative analysis of fungal genomes reveals different plant cell wall degrading capacity in fungi. *BMC Genomics* **14**: 274.
- Zhou XD, De Beer ZW, Harrington TC, McNew D, Kirisits T, Wingfield MJ (2004) Epitypification of *Ophiostoma galeiformis* and phylogeny of species in the *O. galeiformis* complex. *Mycologia* **96**: 1306–1315.
- Zhou ZD, Xie YJ, Chen SF, Wingfield MJ (2008) Diseases of eucalypt plantations in China: challenges and opportunities. *Fungal Diversity* **32**: 1–8.
- Zipfel RD, De Beer ZW, Jacobs K, Wingfield BD, Wingfield MJ (2006) Multigene phylogenies define *Ceratocystiopsis* and *Grosmannia* distinct from *Ophiostoma*. *Studies in Mycology* **55**: 75–97.