

Images in medicine

Atypical erythema multiforme

Fatima Zahra Elfatoiki^{1, &}, Soumia Chiheb¹

¹Department of Dermatology, Ibn Rochd UHC of Casablanca, Morocco

[&]Corresponding author: Fatima Zahra Elfatoiki, Department of Dermatology, Ibn Rochd UHC of Casablanca, Morocco

Key words: Genital erythematous, target-like lesions, apoptotic individual keratinocytes

Received: 04/12/2014 - Accepted: 16/12/2014 - Published: 30/04/2015

Pan African Medical Journal. 2015; 20:436 doi:10.11604/pamj.2015.20.436.5882

This article is available online at: <http://www.panafrican-med-journal.com/content/article/20/436/full/>

© Fatima Zahra Elfatoiki et al. The Pan African Medical Journal - ISSN 1937-8688. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Image in medicine

4-year-old child presented oral and genital eruptions for 2 days, he had history of herpetic stomatitis 3 days before. Examination revealed oral painful erosive lesions with genital erythematous purpuric patch mimicking atypical target-like lesions . No others cutaneous lesions were observed. Skin biopsy showed apoptotic individual keratinocytes, degeneration of basal keratinocytes and intercellular oedema, it concluded to diagnosis of erythema multiforme. The patient was treated by Aciclovir 5mg/kg/day for 1 week with oral antihistamines, analgesics and local skin care. A complete resolution was noted after 2 weeks.



Figure 1: Génital atypical target-like lesions