

Poster presentation

## Clues to remind leukemia in patients with arthritis: case presentations

B Makay\*, Y Topçu, S Yilmaz and E Unsal

Address: Dokuz Eylul University Hospital, Izmir, Turkey

\* Corresponding author

from 15<sup>th</sup> Paediatric Rheumatology European Society (PreS) Congress  
London, UK. 14–17 September 2008

Published: 15 September 2008

*Pediatric Rheumatology* 2008, **6**(Suppl 1):P172 doi:10.1186/1546-0096-6-S1-P172

This abstract is available from: <http://www.ped-rheum.com/content/6/S1/P172>

© 2008 Makay et al; licensee BioMed Central Ltd.

Acute lymphocytic leukemia (ALL) may initially present with musculoskeletal symptoms such as pain or swelling, even before appearance of blasts in the peripheral blood. Fifteen to 30% of all ALL patients have musculoskeletal complaints at disease onset. Such presentation may lead to misdiagnosis of the patients causing a delay in proper management. Here, we present 5 patients who were referred to Pediatric Rheumatology Department for the evaluation of arthritis and were finally diagnosed as ALL (table 1).

The peripheral blood changes were absent or subtle in these patients. Some clinical, laboratory, and radiological clues helped to establish the diagnosis of ALL. We aimed to discuss the usefulness of history, physical findings, and simple laboratory and radiographic tests in decision-making.

**Table 1: Five patients who were referred to Pediatric Rheumatology Department for the evaluation of arthritis and were finally diagnosed as ALL**

Patient	Presentation	Provisional diagnosis	Final diagnosis
10 y, ♂	Migratory arthritis, fatigue	Acute rheumatic fever	ALL
3 y, ♂	Fever, arthritis of the left hip	Septic arthritis	ALL
2 y, ♀	Arthritis of the right hip	Transient synovitis	ALL
3 y, ♂	Pain in legs, inability to walk	JIA	ALL
6 y, ♀	Arthritis of the left elbow, fatigue	Postviral arthritis	ALL