



Draft Genome Sequences of Two Acidophilic, Mesophilic Verrucomicrobial Methanotrophs Contain Only One *pmoCAB* Operon

Geert Cremers,^a Arjan Pol,^a  Mike S. M. Jetten,^a  Huub J. M. Op den Camp^a

^aDepartment of Microbiology, IWWR, Radboud University, Nijmegen, The Netherlands

ABSTRACT *Methylacidimicrobium cyclopophantes* 3B and *Methylacidimicrobium tartarophylax* 4AC are Gram-negative rod-shaped mesophilic methanotrophs isolated from soil samples with low pH at the Solfatara Crater, near Naples, Italy. The genomes of these extremophilic verrucomicrobia were sequenced using Illumina technology, and both species possess one *pmoCAB* operon and two *soxB* genes.

Methane is a potent greenhouse gas, and methanotrophs play a significant role in mitigating methane emissions to the atmosphere. Methane-oxidizing bacteria have been detected in a variety of environments, including the hot and acidic soils of volcanic regions (1). The majority of the methanotrophs in these hostile environments belong to the verrucomicrobial methanotrophs. *Methylacidiphilum* species growing at 40°C to 60°C and pH 1 to 6 were described in 2007 (2). Later, mesophilic, highly acid-tolerant isolates belonging to a second genus, *Methylacidimicrobium*, were discovered (3, 4). Here, we report the draft genome sequences of two representatives of this genus (3). *Methylacidimicrobium cyclopophantes* 3B and *Methylacidimicrobium tartarophylax* 4AC were isolated from soil samples taken at the Solfatara Crater, at the center of the Campi Flegrei caldera, near Naples, Italy, and were phylogenetically characterized (3). The strains were grown in a medium containing lanthanides (5) based on local soil concentrations and with a headspace of 5% (vol/vol) carbon dioxide and 10% (vol/vol) methane at 29°C with shaking at 350 rpm.

DNA was isolated using the DNeasy Powersoil kit (Qiagen, Venlo, The Netherlands) with 5-min sonication at 15-s intervals (Bioruptor Next Gen; Diagenode AS, Ougrée, Belgium), and a library was prepared according to the manufacturer's protocol (Nextera DNA sample preparation kit; Illumina, San Diego, CA, USA). The library was single-end sequenced using the cBot single-read cluster generation system (catalog number GD-300-1001) and a 36-cycle sequencing kit V4 (catalog number FC-104-4002) on the Genome Analyzer II system (Illumina). The quality of the sequence reads was checked with CLC workbench 4 (Qiagen Aarhus A/S, Denmark) and trimmed using a quality limit of 0.001 and minimum number of nucleotides in reads of 30. Assemblies were performed using CLC workbench 4 (Qiagen Aarhus A/S) with default settings and a minimum contig length of 500 bp. Binning was done on G+C content (no threshold) and coverage (>250×) by plotting G+C content against coverage per contig in a scatterplot. For *Methylacidimicrobium tartarophylax* 4AC, the trimmed reads (27,600,863) resulted in 137 contigs ranging from 502 to 140,500 bp (N_{50} , 31,024 bp). For *Methylacidimicrobium cyclopophantes* 3B, the trimmed reads (29,990,414) resulted in 273 contigs ranging from 513 to 63,014 bp (N_{50} , 14,841 bp). Completeness (strain 3B, 98.7%; strain, 4AC, 98.0%) and contamination (strain 3B, 0.7%; strain 4AC, 1.4%) were determined using CheckM (6). The genomes were annotated by the NCBI Prokaryotic Genome Annotation Pipeline (PGAP) (7) and analyzed with

Citation Cremers G, Pol A, Jetten MSM, Op den Camp HJM. 2020. Draft genome sequences of two acidophilic, mesophilic verrucomicrobial methanotrophs contain only one *pmoCAB* operon. *Microbiol Resour Announc* 9:e00315-20. <https://doi.org/10.1128/MRA.00315-20>.

Editor Christina A. Cuomo, Broad Institute

Copyright © 2020 Cremers et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Huub J. M. Op den Camp, h.opdenkamp@science.ru.nl.

Received 25 March 2020

Accepted 31 March 2020

Published 16 April 2020

TABLE 1 Characteristics of the *Methylocidimicrobium* genomes

Isolate	Genome size (bp)	G+C content (%)	No. of contigs	N_{50} (bp)	Coverage (×)	No. of genes by type				No. of CRISPR arrays
						Total	CDS ^a	RNA	Pseudogenes	
<i>M. tartarophylax</i> 4AC	2,327,085	61.2	137	31,024	665	2,241	2,188	53	121	1
<i>M. cyclopophantes</i> 3B	2,276,790	61.2	273	14,841	598	2,178	2,121	57	141	3

^aCDS, coding DNA sequences.

emphasis on the major methanotrophic pathways. Additional information on the genome sequences is compiled in Table 1.

Unlike the *Methylocidiphilum* species (8–10) which contain three *pmoCAB* operons, strains 3B and 4AC possess only one *pmoCAB* operon encoding the membrane-bound methane monooxygenase. Furthermore, they also lack genes encoding the soluble methane monooxygenase (*mmoXYZ*). Both genomes contain two *xoxF* genes encoding lanthanide-dependent pyrroloquinoline quinone (PQQ)-methanol dehydrogenases (5, 11), the accompanying *xoxG* (encoding the electron acceptor *cytochrome c*) and *xoxJ* genes, and all genes for cofactor PQQ synthesis. All genes required for complete oxidation of formaldehyde/formate were identified. In addition, the genomes encode a complete Calvin cycle for CO₂ fixation, including the genes for two subunits of the key enzyme RuBisCO (*cbbL* and *cbbS*) (3, 12). The presence of hydrogenase gene clusters in both genomes supports potential growth as chemolithotrophic Knallgas bacteria (13, 14). Both strains contain *nifHDK* genes, indicating that, similar to *Methylocidiphilum fumariolicum* SolV, nitrogen fixation might be possible (15).

Data availability. This whole-genome shotgun project has been deposited in ENA under project number [PRJEB32513](https://www.ebi.ac.uk/ena/record/PRJEB32513). The assembled genomes are deposited under accession numbers [GCA_902143385](https://www.ncbi.nlm.nih.gov/nuccore/GCA_902143385) and [GCA_902143375](https://www.ncbi.nlm.nih.gov/nuccore/GCA_902143375). The versions described in this paper are the second versions. Raw reads are available under SRA accession numbers [ERR3674889](https://www.ncbi.nlm.nih.gov/sra/ERR3674889) and [ERR3675335](https://www.ncbi.nlm.nih.gov/sra/ERR3675335).

ACKNOWLEDGMENTS

H.J.M.O.D.C. and M.S.M.J. were supported by European Research Council Advanced grants VOLCANO 669371 and EcoMoM 339880, respectively.

REFERENCES

- Knief C. 2015. Diversity and habitat preferences of cultivated and uncultivated aerobic methanotrophic bacteria evaluated based on *pmoA* as molecular marker. *Front Microbiol* 6:1346. <https://doi.org/10.3389/fmicb.2015.01346>.
- Op den Camp HJM, Islam T, Stott MB, Harhangi HR, Hynes A, Schouten S, Jetten M, Birkeland NK, Pol A, Dunfield PF. 2009. Environmental, genomic and taxonomic perspectives on methanotrophic Verrucomicrobia. *Environ Microbiol Rep* 1:293–306. <https://doi.org/10.1111/j.1758-2229.2009.00022.x>.
- van Teeseling MCF, Pol A, Harhangi HR, van der Zwart S, Jetten MSM, Op den Camp HJM, van Niftrik L. 2014. Expanding the verrucomicrobial methanotrophic world: description of three novel species of *Methylocidimicrobium* gen. nov. *Appl Environ Microbiol* 80:6782–6791. <https://doi.org/10.1128/AEM.01838-14>.
- Sharp CE, Smirnova AV, Graham JM, Stott MB, Khadka R, Moore TR, Grasby SE, Strack M, Dunfield PF. 2014. Distribution and diversity of Verrucomicrobia methanotrophs in geothermal and acidic environments. *Environ Microbiol* 16:1867–1878. <https://doi.org/10.1111/1462-2920.12454>.
- Pol A, Barends TRM, Dietl A, Khadem AF, Eygensteyn J, Jetten MSM, Op den Camp H. 2014. Rare earth metals are essential for methanotrophic life in volcanic mudpots. *Environ Microbiol* 16:255–264. <https://doi.org/10.1111/1462-2920.12249>.
- Parks DH, Imelfort M, Skennerton CT, Hugenholtz P, Tyson GW. 2015. CheckM: assessing the quality of microbial genomes recovered from isolates, single cells, and metagenomes. *Genome Res* 25:1043–1055. <https://doi.org/10.1101/gr.186072.114>.
- Tatusova T, DiCuccio M, Badretdin A, Chetvernin V, Nawrocki EP, Zaslavsky L, Lomsadze A, Pruitt KD, Borodovsky M, Ostell J. 2016. NCBI Prokaryotic Genome Annotation Pipeline. *Nucleic Acids Res* 44:6614–6624. <https://doi.org/10.1093/nar/gkw569>.
- Hou S, Makarova KS, Saw JHW, Senin P, Ly BV, Zhou Z, Ren Y, Wang J, Galperin MY, Omelchenko MV, Wolf YI, Yutin N, Koonin EV, Stott MB, Mountain BW, Crowe MA, Smirnova AV, Dunfield PF, Feng L, Wang L, Alam M. 2008. Complete genome sequence of the extremely acidophilic methanotroph isolate V4, *Methylocidiphilum infernorum*, a representative of the bacterial phylum Verrucomicrobia. *Biol Direct* 3:26. <https://doi.org/10.1186/1745-6150-3-26>.
- Anvar SY, Frank J, Pol A, Schmitz A, Kraaijeveld K, den Dunnen JT, Op den Camp H. 2014. The genomic landscape of the verrucomicrobial methanotroph *Methylocidiphilum fumariolicum* SolV. *BMC Genomics* 15:914. <https://doi.org/10.1186/1471-2164-15-914>.
- Kruse T, Ratnadevi CM, Erikstad HA, Birkeland NK. 2019. Complete genome sequence analysis of the thermoacidophilic verrucomicrobial methanotroph “*Candidatus* *Methylocidiphilum kamchatkense*” strain Kam1 and comparison with its closest relatives. *BMC Genomics* 20:642. <https://doi.org/10.1186/s12864-019-5995-4>.
- Keltjens JT, Pol A, Reimann J, Op den Camp H. 2014. PQQ-dependent methanol dehydrogenases: rare earth elements make a difference. *Appl Microbiol Biotechnol* 98:6163–6183. <https://doi.org/10.1007/s00253-014-5766-8>.
- Khadem AF, Pol A, Wiczorek A, Mohammadi SS, Francoijs KJ, Stunnenberg HG, Jetten MSM, Op den Camp H. 2011. Autotrophic methanotrophy in Verrucomicrobia: *Methylocidiphilum fumariolicum* SolV uses the Calvin-Benson-Bassham cycle for carbon dioxide fixation. *J Bacteriol* 193:4438–4446. <https://doi.org/10.1128/JB.00407-11>.

13. Mohammadi S, Pol A, van Alen TA, Jetten MS, Op den Camp H. 2017. *Methylacidiphilum fumariolicum* SoIV, a thermoacidophilic “Knallgas” methanotroph with both an oxygen-sensitive and -insensitive hydrogenase. ISME J 11:945–958. <https://doi.org/10.1038/ismej.2016.171>.
14. Mohammadi SS, Schmitz RA, Pol A, Berben T, Jetten MSM, Op den Camp HJM. 2019. The acidophilic methanotroph *Methylacidimicrobium tartarophilax* 4AC grows as autotroph on H₂ under microoxic conditions. Front Microbiol 10:2352. <https://doi.org/10.3389/fmicb.2019.02352>.
15. Khadem AF, Pol A, Jetten MSM, Op den Camp H. 2010. Nitrogen fixation by the verrucomicrobial methanotroph *Methylacidiphilum fumariolicum* SoIV. Microbiology 156:1052–1059. <https://doi.org/10.1099/mic.0.036061-0>.