



ORAL PRESENTATION

Open Access

Gastric emptying is rapid in adolescents with type 1 diabetes and relates to gastrointestinal symptoms

Shiree Perano^{1,2*}, Christopher Rayner^{2,3}, Stamatiki Kritas⁴, Christine Mpundu-Kaambwa⁵, Kim Donaghue⁶, Michael Horowitz^{2,7}, Jennifer Couper^{1,2}

From 8th APPEs Biennial Scientific Meeting Darwin, Australia. 29 October – 1 November 2014

Gastric emptying is a critical determinant in postprandial glycaemic control. This study aimed to assess whether gastric emptying in adolescents with type 1 diabetes (T1D) relates to gastrointestinal symptoms and to heart rate variability (HRV) as a measure of autonomic function.

We studied 30 adolescents (age 15 ± 2.5 years, BMI 22 ± 3.1 kg/m²) with T1D. Subjects consumed a ¹³C labelled pancake meal. Gastric emptying was measured by ¹³C breath test. Blood glucose was monitored frequently over 4 hours and gastrointestinal symptoms at 30-60 minute intervals, by a visual analogue questionnaire. Chronic gastrointestinal symptoms over the previous 3 months were assessed by a validated Diabetes Bowel Symptoms Questionnaire [1]. HRV was assessed by LabChart Pro [2]. 15 age and sex matched controls were also studied.

Gastric half emptying time was accelerated in adolescents with T1D compared to controls; 77.6 (61.4-99.3) minutes [median (IQR)] versus 109.1 (70.8-124.2), $P = 0.02$, independent of hyperglycaemia during the study, HbA1c, duration of diabetes, and BMI.

There was no difference in the prevalence of chronic symptoms or symptoms of a severity that affected lifestyle between the two groups.

The presence of nausea, vomiting, bloating and/or fullness during the study in T1D was associated with faster gastric emptying compared to asymptomatic T1D ($r = 0.55$; $p = 0.04$), and this was independent of peak glucose and glucose at 4 hours.

Rate of gastric emptying in T1D did not correlate with HRV.

Adolescents with T1D have rapid gastric emptying associated with acute gastrointestinal symptoms. Symptomatology could be used as a clinical tool to determine the need for further investigation.

Authors' details

¹Diabetes and Endocrinology Department, Women's and Children's Hospital, North Adelaide, SA, Australia. ²University of Adelaide, Adelaide, SA, Australia. ³Gastroenterology Department, Royal Adelaide Hospital, SA, Australia. ⁴Gastroenterology Department, Women's and Children's Hospital, North Adelaide, SA, Australia. ⁵Research and Evaluation Unit, Women's and Children's Hospital, North Adelaide, SA, Australia. ⁶Institute of Endocrinology, Children's Hospital Westmead, Sydney, NSW, Australia. ⁷Endocrinology Department, Royal Adelaide Hospital, North Terrace, SA, Australia.

Published: 28 April 2015

References

1. Quan C, Talley NJ, Cross S, Jones M, Hammer J, Giles N, et al: **Development and validation of the Diabetes Bowel Symptom Questionnaire.** *Aliment Pharmacol Ther* 2003, **17**(9):1179-1187.
2. Task Force of the European Society of Cardiology and the North American Society of Pacing and Electrophysiology.: **Heart rate variability: standards of measurement, physiological interpretation and clinical use.** *Circulation* 1996, **93**(5):1043-1065.

doi:10.1186/1687-9856-2015-S1-O46

Cite this article as: Perano et al.: Gastric emptying is rapid in adolescents with type 1 diabetes and relates to gastrointestinal symptoms. *International Journal of Pediatric Endocrinology* 2015 **2015** (Suppl 1):O46.

¹Diabetes and Endocrinology Department, Women's and Children's Hospital, North Adelaide, SA, Australia

Full list of author information is available at the end of the article