

Evaluation of tumor necrosis factor-alpha (TNF- α) and interleukin (IL)-1 β levels among subjects vaping e-cigarettes and nonsmokers

Dharmendra Kumar Sinha¹, Vishal², Ashish Kumar³, Mohsin Khan⁴,
Ritu Kumari¹, Mritunjay Kesari⁵

¹Department of Prosthodontics, Awadh Dental College and Hospital, Jamshedpur, Jharkhand, ²Tutor, Oral and Maxillofacial Surgery, Dental Institute, RIMS, Ranchi, ³Department of Oral and Maxillofacial Surgery, Awadh Dental College and Hospital, Jamshedpur, Jharkhand, ⁴Department of Oral and Maxillofacial Surgery, TMDCRC, Moradabad, Uttar Pradesh, ⁵Department of Prosthodontics, Hazaribag College of Dental Sciences and Hospital, Jharkhand, India

ABSTRACT

Aim: The present study aimed to evaluate peri-implant immunological parameters along with clinical and radiographic parameters amongst subjects vaping e-cigarettes and nonsmokers (NS). Two immunological parameters that were included were measurement of tumor necrosis factor-alpha (TNF- α) and interleukin (IL)-1 β . **Materials and Methods:** A statistically significant number of subjects were included in the study and further divided into two groups: Group-1: Vaping e-cigs ($n = 47$) Group-2: Nonsmokers ($n = 45$) A structured questionnaire was used at baseline for data collection related to demographic and peri-implant data of each individual. Baseline parameters included plaque index, bleeding on probing, probing pocket depth, and peri-implant bone loss (PI, BOP, PD, and PIBL, respectively) for those vaping e-cigs and nonsmokers with the aid of standardized digital radiographs. ELISA kit was used to assess immunologic parameters using sulcular fluid collected from peri-implant region (PISF). **Results:** Bleeding on probing (BOP) was found significantly high in nonsmokers compared to the vaping group. But probing depth greater than 4 mm and PIBL scores were significantly higher in the vaping group than nonsmokers. Mean concentrations of immunologic parameters were significantly higher in vaping group than in nonsmokers. TNF- α levels and IL-1 β levels were found to have a positive correlation with bone loss around implants (PIBL). Besides, TNF- α also had a positive correlation with bleeding on probing in vaping group compared to nonsmokers. **Conclusions:** Owing to the effect of nicotine on the peri-implant tissue, levels of inflammatory mediators as detected by ELISA tests were found to be higher showing a greater amount of localized inflammatory tissue destruction and a compromised peri-implant area in vaping subjects. PISF concentrations were also found relatively higher than nonsmokers.

Keywords: Electronic cigarettes, IL-1 β levels, interleukin, TNF- α

Introduction

Electronic cigarette (e-cig) is the latest trend amongst nicotine users. It is a battery-operated device that releases nicotine in vaporized form simulating the effect of an actual cigarette. It

Address for correspondence: Dr. Dharmendra Kumar Sinha,
Department of Prosthodontics, Awadh Dental College and
Hospital, Jamshedpur, Jharkhand, India.
E-mail: designmypaper@gmail.com

Received: 16-10-2019

Revised: 27-12-2019

Accepted: 09-01-2020

Published: 28-02-2020

has gained quite popularity among today's youth in the age group of 18 to 25 years. Nevertheless, it poses deleterious effect on systemic health of vaping individuals.^[1]

E-cigs are now available in various forms ranging from a simple pen-like device to a USB like portable device that operates on a battery, has a metallic element that heats up covered by stainless steel casing from outside. There is a cartridge inside, an atomizer that is directly connected to the battery. The metal element on

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: reprints@medknow.com

How to cite this article: Sinha DK, Vishal, Kumar A, Khan M, Kumari R, Kesari M. Evaluation of tumor necrosis factor-alpha (TNF- α) and interleukin (IL)-1 β levels among subjects vaping e-cigarettes and nonsmokers. J Family Med Prim Care 2020;9:1072-5.

Access this article online

Quick Response Code:



Website:
www.jfmpc.com

DOI:
10.4103/jfmpc.jfmpc_902_19

heating changes the solution filled in the cartridge that contains nicotine and flavoring agents mixed in a chemical solution. Other additives are propylene glycol, glycerin, and so on.^[2] Many people consider e-cigs as a more sophisticated version of smoking tobacco with less deleterious effects.^[3] However, this is contrary to the findings put forward by several researchers. They have drawn our attention to the harmful aerosols liberated during vaping these electronic cigarettes that increase oxidative stress, enhance release of inflammatory mediators, changes in cellular function of liver, and cause damage in DNA constitution.^[4,5]

From the perspective of oral health, vaping nicotine in any form has shown the precipitating factor for serious oral conditions like oral submucous fibrosis with a marked reduction in mouth opening and damage to the integrity of periodontal tissues thus resulting in periodontal tissue-breakdown and, in turn, periodontal diseases.^[6,7] This aims for the conduction of the study which would be helpful for general public in creating awareness so as to avoid debilitating diseases like cancer.

Sundar *et al.*^[8] in their laboratory study drew the conclusion that oral epithelial cells and gingival fibroblasts have been negatively affected by aerosols of e-cig via damage to DNA and enhanced release of inflammatory mediators like cyclooxygenase as well as prostaglandins (COX-2 and PGE2). The present study aimed to evaluate immunological parameters (TNF- α) and (IL)-1 β between vaping individuals and nonsmokers and draw a comparison between clinical and radiographic parameters in the peri-implant structure.

Materials and Methods

Ethical clearance

The present study was done in accordance with the 2013 revised Helsinki Declaration protocol and was approved by the institutional ethical review board (Taken on 16-08-2019). Participants have explained the study protocol thoroughly and written informed consent was taken from each subject participating in the study. Individuals had the discretion of leaving the study as deemed fit by them at any phase of the study.

Study groups

Study subjects were selected from March 2018 to March 2019. Around 100 subjects were initially enrolled out of which 8 refused to participate so the remaining 92 subjects were finalized for taking up the study. The following inclusion and exclusion criteria were devised for selection and further divided into two groups:

Group-1: Vaping e-cigs ($n = 47$)

Group-2: Nonsmokers ($n = 45$)

Inclusion and exclusion criteria

Inclusion criteria:

- Group-1: Vaping e-cigs for a minimum of 1 year

- Group-2: Nonsmokers who never consumed any form of tobacco ever in their lives.
- A minimum of one dental implant (functional for 3 years) in both of the groups.

Exclusion criteria:

- Current smokers/pipe smokers/smokeless tobacco chewers
- Systemic disorders such as diabetes, AIDS, heart and kidney dysfunction
- Subjects having edentulous ridges.
- Antibiotics/NSAIDs for the past 6 months
- Periodontal therapy for the past 6 months.

Study questionnaire

A structured questionnaire was used at baseline for data collection related to demographic and peri-implant data of each individual. They included

- Total number of subjects per group (n)
- Sex (all males)
- Mean age of each individual in years along with standard deviation
- Total number of implants placed in both arches
- Position of the implant (maxilla/mandible)
- Duration of implants in months (mean \pm SD)
- Mean duration of habit (in years)
- Vaping frequency per day
- Duration of the session (mean in min)
- Frequency of brushing either once or twice a day (%).

Clinical assessment

Baseline clinical parameters included plaque index, bleeding on probing, pocket depth (PI, BOP, and PD) and were recorded by the single examiner to avoid interexaminer bias. The kappa statistics for intraexaminer variation were 0.91 which was a very good score. Around 6 conventional periodontal sites were used for charting and probing pocket depth was calibrated to the nearest mm using UNC-15 periodontal probe.

Radiographic evaluation (PIBL)

A single masked evaluator recorded all radiographic parameters. The kappa statistics for intraexaminer variation was .86 and was considered as a decent agreement. Digital periapical radiographs were shot and assessed using an image analyzer on a standardized desktop screen. These RVGs were further scanned at 800 dots per inch using a cam-scanner. PIBL was assessed by measuring the distance from the crest of bone to the implant platform.^[9]

Collection of PISF

Supragingival oral prophylaxis was done using ultrasonic scalers from the crown surface judiciously and the peri-implant sites were isolated using dry and sterile cotton roles. Using GCF collecting periopaper/paper strips PISF was collected in two separate vials. The paper was held still into the sulcus for 30 s reaching 2 mm into the sulcus/pocket. Any contamination with blood or

saliva was checked by discarding PISF after every 15 min from a particular site. Peritron was used for measuring PISF volume.

Measurement of TNF- α and IL-1 β in PISF

ELISA kit was used to analyze and quantify the inflammatory chemical mediators TNF- α and IL-1 β . PISF was centrifuged for 15 min at 5 kg at 48 $^{\circ}$.^[10]

Statistical analyses

Statistical software version SPSS 23 was used for carrying out statistical analysis. Intergroup comparison was carried out using the Kruskal–Wallis test. In addition, Mann-Whitney U test was used for comparison of inflammatory mediators. Multiple comparisons were done using the Post-hoc Bonferroni Test. A P value of < 0.05 was regarded as statistically significant.

Demographic and implant-related characteristics of the study groups

Clinical and radiographic peri-implant parameters

Cytokine levels in PISF among study groups

Pearson’s correlation analysis among pro-inflammatory cytokines and peri-implant parameters

Results

Bleeding on probing (BOP) was found significantly high in nonsmokers compared to the vaping group. But probing depth greater than 4 mm and PIBL scores were significantly higher in a vaping group than nonsmokers [Table 1]. Mean concentrations of immunologic parameters were significantly higher in a vaping group than in nonsmokers [Table 2]. TNF- α levels and IL-1 β levels were found to have a positive correlation with bone loss around implants (PIBL) [Table 3]. Besides TNF- α also had a positive correlation with bleeding on probing in a vaping group compared to nonsmokers [Table 4].

Discussion

Cellular healing relatively is affected by smoking tobacco which is rich in nicotine content. Smokers show less bleeding tendency. In the present study too, it was observed that bleeding on probing was more in nonsmokers than in vapers. The potential threat thus posed by smoking tobacco remains clinically less evident as basic indicator of periodontal tissue breakdown i.e. bleeding is not significantly appreciable. The mechanism behind reduced bleeding is the vasoconstriction caused by nicotine on the suprapariosteal blood vessels. Nonetheless, the higher concentration of sulcular fluid and increased amount of cytokines are indicative of rapid tissue destruction caused by nicotine. Not just smokers but vapers too remain unaware of potential threat e-cigs pose to the health of oral epithelium and gingival fibroblasts. It remains a common misconception that e-cigs are less harmful as strongly opposed by the findings of present study which showed a significant hike in the chemical mediators at the peri-implant sites compared to those of nonsmokers.

The findings of our study are clearly indicating that the severity of periodontal tissue destruction is higher in individuals vaping electronic cigarettes.^[11,12] Both probing pocket depth and peri-implant bone loss were significantly higher in group 1 than in group 2. An increased amount of interleukins in the peri-implant area is a well-established indicator of advanced bone loss or osteoclastic activity and findings showed that IL-1 β was significantly higher in PISF sample of vapers than smokers. Pro-inflammatory cytokines TNF- α plays a pivotal and central role in destruction of peri-implant tissue. It is suggestive of gram-negative microbes which liberate lipopolysaccharides and further advance destruction. Vaping e-cigs result in increase in advanced glycated end products AGEs in periodontal tissues, the AGE-RAGE interaction i.e. augmented interface of AGEs with their receptors liberates reactive oxygen species (ROS) creating an oxidative stress, inducing metabolic changes, altered chemotaxis, neutrophilic dysfunction, reduction in antibodies production, increased bacterial cell to cell adhesion, and thus enhancing

Table 1: Demographic and implant-related characteristics of the study groups

Demographic Parameters	Vapers (group 1)	Non-smokers (group 2)
Total participants (n)	47	45
Sex (males)	47	45
Mean age \pm SD (in years)	34.6 \pm 6.1	44.8 \pm 2.5
Total number of implants Implant position (maxilla/mandible)	66	55
Duration of implants in months (mean \pm SD)	45/21	35/20
Mean duration of habit (in years)	53.2 \pm 13.7	51.2 \pm 19.5
Daily frequency of the habit	4.6 \pm 1.8	-
Mean duration of a session (in min)	6.8 \pm 0.9	-
Brushing frequency (%)	37.7 \pm 11.3	-
Once	54	58
Twice	46	42

Table 2: Mean clinical and radiographic peri-implant parameters among vaping individuals and NS

Peri-implant parameters	Vapers (group 1) n=47	Non-smokers (group 2) n=45
Plaque index (%)	54.6 \pm 10.9	48.46 \pm 9.76
Bleeding on probing (%)	23.7 \pm 5.3 ^a	37.82 \pm 17.81
Probing depth \geq 4 mm (%)		
Peri-implant bone loss (mm)	5.87 \pm 1.42 ^b	4.65 \pm 0.71
Mesial	1.56 \pm 0.7 ^b	0.98 \pm 0.2
Distal	2.09 \pm 1.0 ^b	1.12 \pm 0.5
Total	1.78 \pm 0.9 ^b	0.93 \pm 0.3

^aCompared with group 2 (P<0.01). ^bCompared with group 2 (P<0.05)

Table 3: Concentrations of pro-inflammatory cytokines (TNF-a and IL-1b) in PISF among vaping individuals and NS

Immunological parameters	Vaping individuals (group 1) n=47	Never smokers (group 2) n=45
PISF volume collected (in IL)	3.17 \pm 0.6 ^a	1.5 \pm 0.5
TNF- α (in pg/mL)	24.3 \pm 32.4 ^b	6.7 \pm 8.1
IL1b (in pg/mL)	205.2 \pm 230.7 ^a	19.7 \pm 22.3

^aCompared with group 2 (P<0.01). ^bCompared with group 2 (P<0.001)

Table 4: Pearson's correlation analysis among pro-inflammatory cytokines and peri-implant parameters

Parameters	Vaping individuals (group 1)	Never smokers (group 2)
TNF- α		
PI		
Correlation coefficient	0.9111	-0.5268
P	0.924	0.3871
BOP		
Correlation coefficient	0.1441	-0.1245
P	0.0214	0.7847
PD		
Correlation coefficient	0.6127	-0.9572
P	0.1728	0.4213
PIBL		
Correlation coefficient	0.3200	-0.4455
P	0.0167	0.8914
IL-1b		
PI		
Correlation coefficient	0.5067	-0.4276
P	0.7436	0.1983
BOP		
Correlation coefficient	0.8116	-0.3145
P	0.2867	0.4810
PD		
Correlation coefficient	0.3345	-0.7669
P	0.9723	0.0912
PIBL		
Correlation coefficient	0.4201	-0.3772
P	0.0181	0.8168

Significant at $P < 0.05$

production of localized and systemic inflammatory mediators thus their expression is higher in sulcular fluid.

Further, such investigations should be carried out to understand the mechanism behind increased peri-implant tissue destruction in vaping individuals.

Conclusions

Owing to the effect of nicotine on the peri-implant tissue, levels of inflammatory mediators as detected by ELISA tests were found to be higher showing a greater amount of localized inflammatory tissue destruction and a compromised peri-implant area in vaping subjects. PISF concentrations were also found relatively higher than nonsmokers.

Financial support and sponsorship

Nil.

Conflict of interest

There is no conflict of interest.

References

- Choi K, Forster JL. Beliefs and experimentation with electronic cigarettes: A prospective analysis among young adults. *Am J Prev Med* 2014;46:175-8.
- Morean ME, Lipshie N, Josephson M, Foster D. Predictors of adult e-cigarette users vaporizing cannabis using e-cigarettes and vape- pens. *Subst Use Misuse* 2017;52:974-81.
- Ayers JW, Leas EC, Allem JP, Benton A, Dredze M, Althouse BM, *et al.* Why do people use electronic nicotine delivery systems (electronic cigarettes)? A content analysis of Twitter, 2012-2015. *PLoS One* 2017;12:e0170702.
- Lerner CA, Rutagarama P, Ahmad T, Sundar IK, Elder A, Rahman I. Electronic cigarette aerosols and copper nanoparticles induce mito-chondrial stress and promote DNA fragmentation in lung fibroblasts. *BiochemBiophys Res Commun* 2016;477:620-5.
- Lerner CA, Sundar IK, Watson RM, Elder A, Jones R, Done D, *et al.* Environmental health hazards of e-cigarettes and their components: Oxidants and copper in e-cigarette aerosols. *Environ Pollut* 2015;198:100-7.
- Wadia R, Booth V, Yap HF, Moyes DL. A pilot study of the gingival response when smokers switch from smoking to vaping. *Br Dent J* 2016;221:722-6.
- Javed F, Kellesarian SV, Sundar IK, Romanos GE, Rahman I. Recent updates on electronic cigarette aerosol and inhaled nicotine effects on periodontal and pulmonary tissues. *Oral Dis* 2017;23:1052-7.
- Sundar IK, Javed F, Romanos GE, Rahman I. E-cigarettes and flavorings induce inflammatory and pro-senescence responses in oral epithelial cells and periodontal fibroblasts. *Oncotarget* 2016;7:77196-204.
- Vohra F, Alkhudhairy F, Al-Kheraif AA, Akram Z, Javed F. Peri-implant parameters and C-reactive protein levels among patients with different obesity levels. *Clin Implant Dent Relat Res* 2018;20:130-6.
- Abduljabbar T, Akram Z, Vohra F, Warnakulasuriya S, Javed F. Assessment of interleukin-1b, interleukin-6, and tumor necrosis factor-A levels in the peri-implant sulcular fluid among water pipe (narghile) smokers and never-smokers with peri-implantitis. *Clin Implant Dent Relat Res* 2018;20:144-50.
- Ling Chen, Ian Todd, Lucy C. Fairclough. Immunological and pathological effects of E-cigarettes. *Basic Clin Pharmacol Toxicol* 2019;125:1-16.
- Kaur G, Pinkston R, Mclemore B, Dorsey WC, Batra S. Immunological and toxicological risk assessment of e-cigarettes. *Eur Respir Rev* 2018;27.doi: 10.1183/16000617.0119-2017.