

Original Article

## Exposure to High Doses of Lipopolysaccharide during Ovalbumin Sensitization Prevents the Development of Allergic Th2 Responses to a Dietary Antigen

Ikuko Torii<sup>1\*</sup>, Shigeki Shimizu<sup>1</sup>, Takashi Daimon<sup>2</sup>, Yoshiyasu Shinohara<sup>1</sup>, Tomoo Kudo<sup>1</sup>, Ayuko Sato<sup>1</sup>, and Tohru Tsujimura<sup>1</sup>

<sup>1</sup> Department of Molecular Pathology, Hyogo College of Medicine, 1–1 Mukogawa-cho, Nishinomiya, Hyogo 663–8501, Japan

<sup>2</sup> Department of Biostatistics, Hyogo College of Medicine, 1–1 Mukogawa-cho, Nishinomiya, Hyogo 663–8501, Japan

**Abstract:** Food allergies are driven by aberrant T helper (Th) 2 cells. Lipopolysaccharide (LPS) influences the development of Th2-mediated diseases, but its role in food allergy and tolerance remains unclear. To address this issue, we established mouse models presenting allergic or tolerant responses to ovalbumin (OVA). Mice sensitized with crude OVA developed Th2 responses including acute diarrhea, increases in serum OVA-specific IgE, dominant production of serum OVA-specific IgG1, increases in Th2-type cytokines and proliferation of mast cells in duodenal and colonic tissues. Sensitization of mice with crude OVA and LPS abrogated Th2-type responses observed in allergic mice. The level of OVA-specific proliferation in mesenteric lymph node CD4<sup>+</sup> T cells was comparable in allergic and tolerant mice, indicating that the tolerance is not caused by anergy and apoptosis of antigen-primed T cells. Expression of Th1- and Th2-type cytokines was suppressed in whole spleen cells and/or purified spleen CD4<sup>+</sup> T cells of tolerant mice, indicating that the tolerance was not caused by the shift from Th2 to Th1. On the other hand, interleukin (IL)-10, a regulatory cytokine produced by regulatory T cells, was upregulated in whole spleen cells and purified spleen CD4<sup>+</sup> T cells of tolerant mice. Furthermore, spleen CD4<sup>+</sup> T cells from tolerant mice suppressed the growth of CD4<sup>+</sup> T cells from DO11.10 mice in co-culture. These results indicate that tolerance is induced in allergic mice by simultaneous exposure to LPS during sensitization with OVA and that a population of T cells producing IL-10 plays an important role in the tolerance induction. (DOI: 10.1293/tox.2014-0023; J Toxicol Pathol 2014; 27: 205–215)

**Keywords:** allergy, lipopolysaccharide, ovalbumin, tolerance

### Introduction

Food allergy is defined as an aberrant T helper (Th) 2-mediated immune response to a dietary antigen<sup>1,2</sup>. Cytokines produced by Th2 cells, such as interleukin (IL)-4, IL-5 and IL-13, participate not only in dietary antigen-specific IgE production but also in mucus secretion, muscle contraction, and eosinophil/mast cell proliferation in intestinal tissues<sup>1,3</sup>. Mast cells express FcεRI, which binds antigen-specific IgE. Upon reexposure to the dietary antigen, cross-linking of the surface IgE stimulates mast cells to release a variety of chemical mediators, which can induce food allergic symptoms, such as diarrhea and anaphylactic reactions<sup>4</sup>.

Components of environmental microbes are potent activators of the immune system with the capacity to shift active immune responses towards priming of Th1 or Th2<sup>5</sup>.

Lipopolysaccharide (LPS), a cell wall component of Gram-negative bacteria, has the capacity to induce Th1-polarized adaptive immune responses via toll-like receptor (TLR) 4 signaling<sup>6,7</sup> and exerts crucial effects on the development of asthma, which is a Th2-mediated allergic disease<sup>8–11</sup>. It has also been shown that LPS exposure drives the development of Th2 airway hypersensitivity through TLR4 signaling in murine models of pulmonary inflammation<sup>12–15</sup>. It is therefore likely that the condition of LPS determines the subsequent immune responses. In the digestive tract, to maintain tolerance to food antigens, the immune system is strictly controlled by several regulatory mechanisms, such as the enzymatic digestion of immunogenic epitopes in food, the constitutive stimulation of gut-associated lymphoid tissue by commensal flora and the action of unique cell populations with regulatory function. The stimulation of gut-associated lymphoid tissue with noninvasive commensal microbiota has a profound impact on the responsiveness of lymphocytes and the generation of Th1-biased memory effector cells<sup>16–18</sup>. It also induces tolerance to orally administered antigens<sup>19</sup>. Since LPS is contained in commensal flora as a basic component with biological activity, it may function as a crucial regulator for food allergy and produce tolerance to dietary antigens.

Received: 16 May 2014, Accepted: 4 July 2014

Published online in J-STAGE: 22 August 2014

\*Corresponding author: I Torii (e-mail: ikukot@hyo-med.ac.jp)

©2014 The Japanese Society of Toxicologic Pathology

This is an open-access article distributed under the terms of the Creative Commons Attribution Non-Commercial No Derivatives (by-nc-nd) License <<http://creativecommons.org/licenses/by-nc-nd/3.0/>>.

The involvement of regulatory T cells (Treg) in food allergy has been reported<sup>20, 21</sup>. CD4<sup>+</sup> CD25<sup>+</sup> Foxp3<sup>+</sup> Treg participate in the induction of oral tolerance to cow's milk and peanut extract allergy<sup>22, 23</sup>. Treg producing IL-10, Tr1, inhibit Th2-mediated allergic diarrhea and eosinophilic infiltration into the small intestine of Peyer's patch-lacking mice<sup>24, 25</sup>. Treg producing transforming growth factor (TGF)- $\beta$ , Th3, are crucial for intestinal homeostasis. Food allergy and anaphylaxis caused by impaired production of TGF- $\beta$  can be prevented by administration of TGF- $\beta$ <sup>26–28</sup>. Recently, it has been shown that systemic stimulation with LPS enhances the production of IL-10 and TGF- $\beta$  in mice, suggesting that LPS regulates immune responses to antigens through induction of Treg producing IL-10 or TGF- $\beta$ <sup>29, 30</sup>.

In this study, to assess the role of LPS in food allergy, we sensitized BALB/c mice with crude ovalbumin (OVA) (hereafter called cOVA) or with cOVA and LPS by intraperitoneal (i.p.) injection. Th2-mediated allergy was observed in the former case, and tolerance was observed in the latter case. The abrogation of the allergic reaction in tolerant mice was not due to induction of anergy and apoptosis in primed T cells or to the shift from Th2 to Th1. On the other hand, IL-10 expression was upregulated in spleen CD4<sup>+</sup> T cells of tolerant mice. These spleen CD4<sup>+</sup> T cells suppressed the growth of CD4<sup>+</sup> T cells expressing an OVA-specific T cell receptor (TCR) in co-culture. These results indicate that systemic exposure to LPS during OVA sensitization activates a population of CD4<sup>+</sup> T cells producing IL-10, resulting in suppression of Th2-mediated allergic reactions.

## Materials and Methods

### Mice

Female BALB/c ByJ and BALB/c Cr mice were purchased from Clea Japan Inc. (Tokyo, Japan) and Japan SLC (Hamamatsu, Japan), respectively. OVA<sub>323–339</sub>-specific TCR transgenic mice (DO11.10) on a BALB/c background were obtained from the Jackson Laboratory (Bar Harbor, ME, USA). All mice were housed under specific pathogen-free conditions and used for experiments at 6–8 weeks of age.

### Endotoxin activity in OVA

To evaluate the contamination of LPS in cOVA (Grade V, A-5503; Sigma-Aldrich, St. Louis, Mo, USA) and purified OVA (hereafter called pOVA) (Profos AG, Regensburg, Germany), endotoxin activities of cOVA and pOVA were examined by the *Limulus* amoebocyte lysate assay (Seikagaku Biobusiness Corp., Tokyo, Japan). LPS derived from *Escherichia coli* 0013:H10 was used as a standard for endotoxin activity. Endotoxin activity of cOVA was assayed after it was diluted so that the activity fell within the linear range of the standard. pOVA was assayed for endotoxin activity without dilution. The endotoxin activity of cOVA (5  $\mu$ g/ml) and that of pOVA (5  $\mu$ g/ml) were estimated to be 52 EU/mg and < 1 EU/mg, respectively, based on the value of 545 nm absorbance. For further experiments, cOVA was used as an antigen for *in vivo* sensitization, and pOVA was used as that

for *ex vivo* stimulation.

### Sensitization and challenge protocol

Based on a previous study about oral allergen-induced diarrhea, we generated the following protocol for the food allergic and tolerant models<sup>31</sup>. Mice (n=3–10) were sensitized with 100  $\mu$ g of cOVA by i.p. injection with or without various doses of LPS derived from *Escherichia coli* 055:B5 (LIST Biological Laboratories Inc., Campbell, CA, USA) in 200  $\mu$ l phosphate-buffered saline (PBS) on days 0 and 14. Mice (n=3–5) sham sensitized with PBS on days 0 and 14 were used as the control. One week after the second sensitization, mice were orally challenged with 100 mg of cOVA in 250  $\mu$ l of sterile saline at 2- or 3-day intervals. Before each intragastric challenge, mice were deprived of food for 3–4 h to minimize degradation of the antigen in the stomach. Mice were carefully monitored for 1 h after each challenge, and those defecating profusely liquid stool were marked as acute diarrhea positive. Diarrhea occurrence was assessed for 15–60 min after each challenge with cOVA. Mice were sacrificed within 1 h after the last challenge. Blood was taken from the femoral artery, and sera were prepared and stored at –20°C until use. Duodenum (approximately 10 cm distal to the stomach) and colons (distal part of the cecum) were promptly removed. A part of each of these tissues was embedded in OCT compound (Sakura Finetech Co., Ltd., Tokyo, Japan), and the remaining tissues were quickly frozen and stored at –80°C until use. To characterize spleen or mesenteric lymph node (MLN) cells, mice were sensitized with 100  $\mu$ g of cOVA by i.p. injection with or without 100  $\mu$ g of LPS in 200  $\mu$ l PBS on days 0 and 7. Mice sham sensitized with PBS on days 0 and 7 were used as the control. Four days or 1 week after the second sensitization, spleens and MLNs were removed. All experimental procedures were approved by the Animal Care Committee of Hyogo College of Medicine and performed in accordance with the criteria outlined in the “Guide for the Care and Use of Laboratory Animals” prepared by the National Academy of Sciences.

### Cells and culture

Spleens and MLNs were removed from sensitized and control (PBS-sensitized) BALB/c ByJ mice at the indicated days after the second sensitization and dispersed mechanically. Single-cell suspensions from spleens and MLNs were prepared from the dispersed tissues by passing through nylon mesh cell strainers (BD Biosciences, Bedford, MA, USA) and excluding RBCs by lysis. CD4<sup>+</sup> T cells were negatively selected from single-cell suspensions by magnetic separation with a CD4<sup>+</sup> T cell Isolation Kit (Miltenyi Biotec, Auburn, CA, USA) according to the manufacturer's instructions (> 95% CD4<sup>+</sup> T cells as determined by FACS analysis). Spleen CD4<sup>+</sup> T cells from naive DO11.10 mice were isolated in a similar manner.

For preparation of antigen-presenting cells (APCs), the spleens removed from naive BALB/c ByJ and DO11.10 mice were cut into small pieces, treated with 400 Mandl units/ml collagenase D (Roche, Mannheim, Germany) and 10  $\mu$ g/ml

DNase I (Roche) with continuous stirring at 37°C for 35 min and incubated at 37°C for 5 min after adding EDTA to adjust the final concentration to 10 mM. Single cells were prepared from the spleen by vigorously pipetting the suspension of small tissue pieces and passing the suspension through nylon mesh cell strainers. After lysing of RBCs, the CD4<sup>+</sup> T cell-depleted fraction was isolated from the cell suspensions with a CD4<sup>+</sup> T cell Isolation Kit (Miltenyi Biotec). The isolated cells were irradiated at 30 Gray and used as APCs.

All cells used in this study were cultured in RPMI 1640 medium containing 10% FCS at 37°C in a 5% CO<sub>2</sub> humidified atmosphere.

#### *OVA-specific proliferation assay*

Co-culture of  $3.5 \times 10^5$  MLN CD4<sup>+</sup> T cells from control (PBS-sensitized) and sensitized BALB/c ByJ mice was performed with  $1 \times 10^5$  APCs from BALB/c ByJ mice in the presence of the indicated concentration of pOVA in 96-well flat bottom microplates for 72 h. MLN CD4<sup>+</sup> T cells in each well were labeled with 0.25  $\mu$ Ci [<sup>3</sup>H]-thymidine (PerkinElmer, Waltham, MA, USA) for the last 12 h of incubation, and the incorporated radioactivity in MLN CD4<sup>+</sup> T cells was measured by a liquid scintillation counter (TopCount, Packard BioScience Company, Meriden, CT, USA).

#### *Ex vivo co-culture assay*

Co-culture of  $5 \times 10^5$  spleen CD4<sup>+</sup> T cells from naive DO11.10 mice was performed with  $5 \times 10^5$  nonirradiated or irradiated spleen CD4<sup>+</sup> T cells from sensitized BALB/c ByJ mice in the presence of pOVA (50  $\mu$ g/ml) and  $1 \times 10^5$  APCs from DO11.10 mice in 96-well round bottom microplates for 72 h. Cells in each well were labeled with 0.25  $\mu$ Ci [<sup>3</sup>H]-thymidine (PerkinElmer) for the last 12 h of incubation, and the incorporated radioactivity was measured by a liquid scintillation counter (TopCount).

#### *Enzyme-linked immunosorbent assay (ELISA) for OVA-specific IgE, IgG and IgG2a*

The level of serum OVA-specific IgE was measured by ELISA (Dainippon Sumitomo Pharma, Tokyo, Japan). For measurement of serum OVA-specific IgG1 and IgG2a, sera were diluted 1:1,000 for IgG1, 1:10 for IgG2a and serially diluted 1:4. Immunoplates (Nalge Nunc International, Naperville, IL, USA) were coated with 50  $\mu$ g/ml cOVA in borate saline containing 1% (W/V) bovine serum albumin (BSA) and incubated overnight at 4°C. After blocking with 1% BSA in PBS for 1 h at room temperature, serially diluted serum samples were added into the immunoplates, and the immunoplates were incubated for 2 h at room temperature. After washing with borate saline containing 1% BSA and 0.05% Tween-20, horseradish peroxidase-conjugated anti-mouse IgG1 diluted 1:1,000 (X56; BD Biosciences Pharmingen, Franklin Lakes, NJ, USA) or horseradish peroxidase-conjugated anti-mouse IgG2a diluted in 1:1,000 (R19-15; BD Biosciences Pharmingen) was added the immunoplates, and the immunoplates were incubated for 1 h at room temperature. The color reaction was developed with 3,3',5,5'-

tetramethylbenzidine blue (DAKO, Glostrup, Denmark) for 30 min and stopped by the addition with 6N HCl. Optical density (OD) was measured at 405 nm within 30 min. Data represent OD values at a serum dilution of 1:125,000 for IgG1 and 1:50 for IgG2a, which were in the middle of the linear part of the OD curve, respectively.

#### *Immunohistochemistry*

Frozen sections (6  $\mu$ m) were prepared from OCT-embedded tissues and fixed with 10% formalin. Sections were immunostained with rat anti-mouse c-kit mAb (ACK2), a gift from Dr. Shin-Ichi Nishikawa (Biohistory Research Hall, Osaka, Japan) and then with Histofine Simple Stain mouse MAX-PO (Rat) (Nichirei Corporation, Tokyo, Japan). Cells immunoreactive for c-kit were visualized with Simple Stain DAB Solution (Nichirei). The sections were lightly counterstained with hematoxylin. The average number of c-kit-positive cells in a high-power field (HPF) was calculated from the total number of c-kit-positive cells in 10 HPFs of 5 mice.

#### *Quantitative Real-time RT-PCR*

Frozen tissues were immersed in RNA later (Ambion, Austin, TX, USA) at 4°C overnight. Total RNA was extracted from the frozen tissues, whole spleen cells and CD4<sup>+</sup> T cells by using ISOGEN (Nippon Gene Co., Ltd., Tokyo, Japan) and reverse transcribed to produce cDNA with a PrimeScript RT kit (Takara Bio Inc., Otsu, Japan). The expression of Th1-, Th2- and Treg-type cytokine mRNA was examined by SYBR Green-based quantitative real-time RT-PCR with 40 cycles of 5 seconds of denaturation at 95°C, 30 seconds of annealing at 64°C and 15 seconds of dissociation at 95°C with a TP800 Thermal Cycler (Takara Bio Inc.). The amount of PCR products was analyzed by the standard curve method and calculated by software (Multiplate RQ) with  $\beta$ -actin cDNA as an internal control. The following oligonucleotides were used as primers for PCR: 5'-ACC-GTT-TAA-CCA-GAA-CGT-TGA-ATT-G-3' (sense) and 5'-TCC-ACG-AAT-TTG-GAC-AGG-TTT-ACT-C-3' (antisense) for the *IL-3* gene, 5'-GGA-TGT-GCC-AAA-CGT-CCT-CAC-AGC-3' (sense) and 5'-GCT-TAT-CGA-TGA-ATC-CAG-GC-ATC-G-3' (antisense) for the *IL-4* gene, 5'-AAT-TCC-TGT-AGC-GCA-GGC-TGG-3' (sense) and 5'-TAC-TTG-AGA-CCC-TGA-TGC-AAC-G-3' (antisense) for the *IL-5* gene, 5'-CTG-AAG-ACC-CTC-AGG-ATG-CGG-CT-3' (sense) and 5'-TCT-AGG-TCC-TGG-AGT-CCA-GCA-GAC-3' (antisense) for the *IL-10* gene, 5'-GCT-CAT-GGC-TGG-TGC-AAA-GA-3' (sense) and 5'-GAG-ACG-CCA-TTC-CAC-ATG-TCA-3' (antisense) for the *IL-12 $\beta$*  gene, 5'-CGG-CAC-AGT-CAT-TGA-AAG-CCT-A-3' (sense) and 5'-GTT-GCT-GAT-GGC-CTG-ATT-GTC-3' (antisense) for the *interferone (IFN)- $\gamma$*  gene, 5'-CAA-TTC-CTG-GCG-TTA-CCT-TGG-3' (sense) and 5'-TGG-AGT-TTG-TTA-TCT-TTG-CTG-TCA-C-3' (antisense) for the *TGF- $\beta$*  gene and 5'-CAT-CCG-TAA-AGA-CCT-CTA-TGC-CAA-C-3' (sense) and 5'-ATG-GAG-CCA-CCG-ATC-CAC-A-3' (antisense) for the  *$\beta$ -actin*



gene. The PCR products were confirmed by electrophoresis in 0.7% agarose gel and staining with ethidium bromide.

### Statistical analysis


Data are summarized as means  $\pm$  standard deviations (SDs). Comparisons between two groups and among three groups were made with the independent *t*-test and one-way analysis of variance, followed by the post hoc pairwise independent *t*-test, respectively. The multiplicity in pairwise comparisons was corrected by Bonferroni's procedure. All probability values are 2-sided, and values of  $P < 0.05$  were considered to indicate statistical significance. Statistical analysis was performed using the SPSS software (version 17.0, SPSS Inc., Chicago, IL, USA).

## Results


### Establishment of an allergic model

BALB/c ByJ or BALB/c Cr mice were sensitized twice by i.p. injection of cOVA (100  $\mu$ g) prior to repeated intragastric challenges with a high dose of cOVA (100 mg) (Fig. 1A). Acute diarrhea occurred in both strains, but BALB/c ByJ mice were found to be less sensitive to lethal anaphylactic shock than BALB/c Cr mice (data not shown). BALB/c ByJ mice were chosen for the following experiments.

BALB/c ByJ mice sensitized with cOVA developed acute diarrhea after the 4th or 5th challenge with cOVA (Fig. 1B). The diarrhea usually occurred within 30 min after each challenge and lasted for at least 1 h (Fig. 1C). Mice also showed systemic manifestations, such as hypothermia, less movement and hair ruffles. The ceca and colons removed from these mice were severely edematous and hyperemic and contained watery soft feces (Fig. 1C). In contrast, no control mice sham sensitized with PBS and challenged with cOVA developed diarrhea and systemic manifestations during the experimental period (Fig. 1B). The ceca and colons removed from control mice showed no edematous and hyperemic changes and contained solid feces (Fig. 1C). These results indicate that acute diarrhea resulted from immune reactions to cOVA sensitization but not from an osmotic load and/or nonspecific irritation in the digestive tract. In addition, the allergic reaction was induced, without an additional exogenous adjuvant, in mice sensitized with cOVA and challenged with cOVA. Mice sensitized with 100  $\mu$ g cOVA, hereafter called allergic mice, are simple to produce and useful for studying food allergic diseases.




**Fig. 1.** Establishment of an allergic model. (A) The experimental protocol for induction of allergy in mice. BALB/c ByJ or BALB/c Cr mice were sensitized twice by i.p. injection of cOVA (100  $\mu$ g) at a 2-week interval (black arrows) and then challenged by intragastric administration (ig) of cOVA (100 mg) on the indicated days (white arrows). Mice were sacrificed within 1 h after the last challenge (dotted arrow). Mice sensitized with PBS were used as a control. (B) Incidence of diarrhea. cOVA-sensitized (closed circle) ( $n=8$ ) and PBS control BALB/c ByJ mice (open circle) ( $n=3$ ) were monitored for 1 h, and diarrhea occurrence was assessed for 15 - 60 min after each challenge with cOVA. Similar results were obtained in 5 experiments carried out in this study, and the representative results are shown. (C) Allergic reactions of the alimentary tract. (a) Mice sensitized with cOVA showed severe mucous diarrhea within 30 min after challenge with cOVA. (b) The ceca and colons removed from cOVA-sensitized, cOVA-challenged mice showed severe edematous and hyperemic changes and contained watery soft feces after repeated oral challenges with cOVA, and (c) those from control mice showed no edematous and hyperemic changes and contained solid feces.



**Fig. 2.** Prevention of diarrhea by simultaneous sensitization with cOVA and LPS. (A) The experimental protocol for induction of LPS-mediated tolerance in allergic mice. Mice were sensitized by simultaneous i.p. injection of cOVA (100  $\mu$ g) and various doses of LPS (striped arrows) twice and repeatedly challenged by the intragastric administration (ig) of cOVA (100 mg) on the indicated days (white arrows). Mice were sacrificed within 1 h after the last challenge (dotted arrow). Allergic mice, sensitized with cOVA alone, were used as a positive control, and PBS-sensitized mice were used as a negative control. (B) Prevention of diarrhea. The incidence of diarrhea decreased with increasing doses of LPS ( $\blacktriangle$  0.1  $\mu$ g (n=4)  $\blacksquare$  1  $\mu$ g (n=4) and  $\blacklozenge$  100  $\mu$ g (n=4)). The presence of LPS (100  $\mu$ g) completely abrogated diarrhea. Mice sensitized with cOVA alone and those sensitized with PBS are indicated by closed ( $\bullet$  (n=4)) and open ( $\circ$  (n=3)) circles, respectively. Similar results were obtained in 5 experiments carried out in this study, and representative results are shown.

#### Establishment of a tolerant model

Mice were simultaneously sensitized with a constant dose of cOVA (100  $\mu$ g) and various doses of LPS prior to cOVA challenges (Fig. 2A). The incidence of diarrhea decreased with increasing doses of LPS, and the presence of 100  $\mu$ g LPS completely abrogated diarrhea (Fig. 2B). These results indicate that the presence of a high dose of LPS during cOVA sensitization converted the allergic reactions to tolerant ones against a dietary antigen even if mice were repeatedly challenged with cOVA subsequently. The mice sensitized with 100  $\mu$ g cOVA and 100  $\mu$ g LPS, hereafter called the tolerant mice/model, are useful for studying the mechanism by which tolerance to a food allergy is induced




**Fig. 3.** The expression of Th1-, Th2- and Treg-type cytokine mRNAs in duodenal and colonic tissues. Th1-, Th2- and Treg-type cytokine mRNAs were measured by quantitative real-time RT-PCR in duodenal (A) and colonic (B) tissues removed from control mice (white columns), allergic mice (black columns) and tolerant mice (dotted columns). Each value of mRNA expression was calibrated to that of  $\beta$ -actin expression. Each bar represents the mean  $\pm$  SD of 5 control mice, 5 tolerant mice and 6 allergic mice. Significant differences ( $*P < 0.05$ ) were found between columns by the independent *t*-test with Bonferroni's correction for pairwise comparisons. In control mice, IL-3 mRNA was below the level of detection. Similar results were obtained in 3 experiments, and representative results are shown.

in the digestive tract.

#### Cytokine profiles in the intestinal tissues

To ascertain the involvement of the Th2 immune response in the allergic model, we examined the mRNA expression of Th1-, Th2- and Treg-type cytokines in duodenal and colonic tissues of control, allergic and tolerant mice by quantitative real-time RT-PCR (Fig. 3A and B). The relative expression of IL-3, IL-4 and IL-10 in allergic mice was higher than that in control mice. On the other hand, the levels of IL-4 and IL-10 were almost the same in control and tolerant mice. The IL-5, IFN- $\gamma$  and TGF- $\beta$  expression levels were almost equal among these three types of mice. These




**Fig. 4.** Production of OVA-specific antibodies in allergic mice. Blood samples were obtained from PBS-sensitized control, allergic and tolerant mice within 1 h after the last challenge, and their sera were prepared. (A) Increases in OVA-specific IgE levels in allergic mice. Serum levels of OVA-specific IgE were determined by ELISA. Each dot indicates the OVA-specific IgE concentration in one mouse. We used control (n=5), allergic (n=10) and tolerant (n=8) mice. (B) Increases in OVA-specific IgG1, but not IgG2a, levels in allergic mice. The titers of OVA-specific IgG1 (a) and IgG2a (b) in sera were determined by ELISA. Each dot indicates the values of OVA-specific IgG1 and IgG2a in one mouse. We used control (n=5), allergic (n=10) and tolerant (n=5) mice. Lines depict mean values. ND, not detectable.

results indicate that enhanced expression of Th2 cytokines, IL-3 and IL-4 and IL-10, contributes to the induction of acute diarrhea in allergic mice and that downregulation of these cytokines prevents acute diarrhea in tolerant mice. In addition, it appears that IL-5, IFN- $\gamma$  and TGF- $\beta$  are not in-

involved in the induction of diarrhea in allergic mice and the prevention of diarrhea in tolerant mice.

#### *OVA-specific antibodies in cOVA-challenged allergic and tolerant mice*


We examined serum levels of OVA-specific IgE and serum titers of OVA-specific IgG1 and IgG2a in control, allergic and tolerant mice by ELISA. The concentrations of OVA-specific IgE were significantly higher in allergic mice (11.7 mg/ml) than in tolerant mice (1.5 mg/ml) (Fig. 4A). The titers of OVA-specific IgG1 were undetectable, 2.53 at an OD of 405 nm and 1.51 at an OD of 405 nm in control, allergic and tolerant mice, respectively (Fig. 4B(a)). The titers of OVA-specific IgG2a were undetectable, 0.25 at an OD of 405 nm and 2.92 at an OD of 405 nm in control, allergic and tolerant mice, respectively (Fig. 4B(b)). Thus, the levels of OVA-specific IgG1 were significantly higher in allergic mice than in control and tolerant mice, and the levels of OVA-specific IgG2a were markedly reduced in allergic mice as compared with tolerant mice. Additionally, the level of OVA-specific IgG1 was significantly higher than that of OVA-specific IgG2a, suggesting that OVA-specific



**Fig. 5.**

**Fig. 5.** Involvement of mast cells in allergic and tolerant mice. (A) Duodenal sections of allergic mice, tolerant mice and control mice were immunostained with anti-*c-kit* antibody (ACK2). Numerous mast cells infiltrated into the duodenal mucosa of allergic mice (a), but mast cells were hardly detectable in the duodenal mucosa of tolerant mice (b) and control mice (c). The secondary antibody does not show a nonspecific reaction (d). Scale bars represent 200  $\mu$ m. (B) Increase of mast cells in allergic mice. The numbers of mast cells in the duodenal mucosa of allergic (n=5), tolerant (n=5) and control mice (n=5) were counted under a light microscope. Each bar indicates the mean  $\pm$  SD of 5 mice. Significant differences (\* $P$  < 0.01; \*\* $P$  < 0.005) were found between columns by the independent *t*-test with Bonferroni's correction for pairwise comparisons.





**Fig. 6.** Proliferative ability for OVA stimulation of MLN CD4<sup>+</sup> T cells from allergic and tolerant mice. MLNs were removed from control mice (n=5) (white columns), allergic mice (n=5) (black columns) and tolerant mice (n=5) (dotted columns) 1 week after the second sensitization. MLN CD4<sup>+</sup> T cells were purified by magnetic separation and co-cultured with APCs from BALB/c ByJ mice in the presence of the indicated concentration of pOVA. Proliferative ability of MLN CD4<sup>+</sup> T cells was examined with a [<sup>3</sup>H]-thymidine incorporation assay (cpm). Each bar represents the mean ± SD of 5 mice. Significant differences (\*P < 0.05; \*\*P < 0.01) were found between columns by the independent *t*-test with Bonferroni's correction for pairwise comparisons. Similar results were obtained in 3 experiments, and representative results are shown.


IgG1 was predominantly produced in allergic mice. These data indicate that IgE-mediated Th2 immune responses participate in the induction of food allergy in mice sensitized with cOVA.

#### *Mast cells in cOVA-challenged allergic and tolerant mice*

Mast cells are known to be effectors for allergic reactions. Immunostaining with an antibody to c-kit, a marker of mast cells, showed that numerous c-kit-positive mast cells infiltrated into the duodenal mucosa of allergic mice (Fig. 5A). In contrast, c-kit-positive mast cells were hardly detectable in the duodenal mucosa of tolerant and control mice (Fig. 5A). The secondary antibody did not show a nonspecific reaction. The number of mast cells in the duodenal mucosa was significantly increased in allergic mice (Fig. 5B). Proliferation of mast cells in allergic mice was also observed in the colonic mucosa (data not shown). These results indicate that the acute diarrhea observed in allergic mice results from extensive proliferation of intestinal mast cells and that tolerance is induced by suppressing mast cell proliferation.

#### *Antigen-specific responses of MLN CD4<sup>+</sup> T cells in the allergic and tolerant models*

To examine whether the tolerant reaction against a dietary antigen is caused by anergy and/or apoptosis of primed T cells in tolerant mice, we compared the proliferative ability of MLN CD4<sup>+</sup> T cells in response to pOVA stimulation between allergic and tolerant mice. The level of [<sup>3</sup>H]-thymi-



**Fig. 7.** The expression of Th2-, Th1- and Treg-type cytokine mRNA in whole spleen cells and spleen CD4<sup>+</sup> T cells. Spleens were removed from control mice (white columns), allergic mice (black columns) and tolerant mice (dotted columns) 4 days after the second sensitization. Total RNA was extracted from whole spleen cells (A) and purified CD4<sup>+</sup> T cells (B) and converted to cDNA. Expression of cytokines at the mRNA level was examined by quantitative real-time RT-PCR. Each value of mRNA expression was calibrated to that of β-actin expression. Each bar represents the mean ± SD of 3 mice. Significant differences (\*P < 0.05; \*\*P < 0.01) were found between columns by the independent *t*-test with Bonferroni's correction for pairwise comparisons. Similar results were obtained in 3 experiments, and representative results are shown.

dine incorporation of CD4<sup>+</sup> T cells from tolerant mice was similar to that from allergic mice at a low dose of pOVA, albeit higher than that from allergic mice at a high dose of pOVA (Fig. 6). In addition, there was no apparent difference in the number of MLN lymphoid cells and their viability between allergic and tolerant mice (data not shown). These results indicate that abrogation of the allergic reaction in the tolerant mice is not due to the unresponsiveness of CD4<sup>+</sup> T cells to OVA restimulation and the decrease in MLN cells caused by the apoptotic mechanism.

#### *Cytokine profiles in the spleen*


Whole spleen cells were prepared from control, allergic and tolerant mice, and the expression of Th1-, Th2- and

Treg-type cytokines was examined by quantitative real-time RT-PCR (Fig. 7A). IL-4 expression in tolerant mice was similar to that in control mice and lower than that in allergic mice. The levels of IFN- $\gamma$  and TGF- $\beta$  expression were comparable among control, allergic and tolerant mice. IL-12 expression was markedly lower in tolerant mice than in control and allergic mice. On the other hand, IL-10 expression was upregulated in tolerant mice as compared with control and allergic mice. Next, CD4<sup>+</sup> T cells were isolated from spleen cells of control, allergic and tolerant mice, and the expression of Th1-, Th2- and Treg-type cytokines was examined by quantitative real-time RT-PCR (Fig. 7B). IL-4 expression was comparable among control, allergic and tolerant mice. IL-5 expression in tolerant mice was similar to that in control mice and lower than that in allergic mice. IL-13 expression was lower in tolerant mice than in control and allergic mice. IFN- $\gamma$  expression was markedly lower in tolerant mice than in control mice and slightly lower in tolerant mice than in allergic mice. IL-10 expression was markedly upregulated in tolerant mice as compared with control and allergic mice. In addition, there was no apparent difference in TGF- $\beta$  expression among control, allergic and tolerant mice (data not shown). These results showed that the cytokine profile in tolerant mice corresponded to neither a Th1- or Th2-mediated immune response, indicating that the unresponsiveness to cOVA challenge in tolerant mice is not caused by the shift from Th2 to Th1.

#### *Effect of CD4<sup>+</sup> T cells from tolerant mice on the proliferation of OVA-specific CD4<sup>+</sup> T cells from DO11.10 mice*

To examine whether CD4<sup>+</sup> T cells from tolerant mice have the capability to suppress proliferation of OVA-specific CD4<sup>+</sup> T cells, spleen CD4<sup>+</sup> T cells from naive DO11.10 mice, which contained a transgene that expressed OVA-specific TCR in T cells, were co-cultured with either spleen CD4<sup>+</sup> T cells from allergic or tolerant mice in the presence of pOVA and APCs, and a [<sup>3</sup>H]-thymidine uptake assay was performed in each co-culture. [<sup>3</sup>H]-thymidine incorporation in the co-culture of DO11.10 CD4<sup>+</sup> T cells with CD4<sup>+</sup> T cells from allergic mice was almost equal to that in the culture of DO11.10 CD4<sup>+</sup> T cells alone (Fig. 8A). On the other hand, [<sup>3</sup>H]-thymidine incorporation was remarkably decreased when DO11.10 CD4<sup>+</sup> T cells were co-cultured with CD4<sup>+</sup> T cells from tolerant mice. In addition, the reduction of [<sup>3</sup>H]-thymidine incorporation in the co-culture of DO11.10 CD4<sup>+</sup> T cells with CD4<sup>+</sup> T cells from tolerant mice was dependent on the number of CD4<sup>+</sup> T cells from tolerant mice (data not shown).

To estimate [<sup>3</sup>H]-thymidine uptake by growth of only DO11.10 CD4<sup>+</sup> T cells, DO11.10 CD4<sup>+</sup> T cells were co-cultured with X ray-irradiated CD4<sup>+</sup> T cells from allergic or tolerant mice, and [<sup>3</sup>H]-thymidine uptake was examined in each co-culture. The levels of [<sup>3</sup>H]-thymidine incorporation in the co-culture of DO11.10 CD4<sup>+</sup> T cells with X ray-irradiated CD4<sup>+</sup> T cells from allergic and tolerant mice were lower than that in the culture of DO11.10 CD4<sup>+</sup> T cells



**Fig. 8.** Inhibition of DO11.10 CD4<sup>+</sup> T cell proliferation by CD4<sup>+</sup> T cells from tolerant mice. Naive DO11.10 spleen CD4<sup>+</sup> T cells ( $5 \times 10^5$ ) were co-cultured with nonirradiated or irradiated spleen CD4<sup>+</sup> T cells ( $5 \times 10^5$ ) from allergic or tolerant mice in the presence of pOVA (50  $\mu$ g/ml) and APCs ( $1 \times 10^5$ ) from DO11.10 mice. Proliferative ability of cells was examined with a [<sup>3</sup>H]-thymidine incorporation assay (cpm). (A) [<sup>3</sup>H]-thymidine incorporation in the co-culture of naive DO11.10 CD4<sup>+</sup> T cells with CD4<sup>+</sup> T cells from allergic or tolerant mice. (B) [<sup>3</sup>H]-thymidine incorporation in the coculture of naive DO11.10 CD4<sup>+</sup> T cells with irradiated CD4<sup>+</sup> T cells from allergic or tolerant mice. The levels of [<sup>3</sup>H]-thymidine incorporation of each cell culture and co-culture as designated below the graph are shown. Each bar indicates the mean  $\pm$  SD of triplicate cultures. Significant differences (\* $P < 0.01$ ) were found between columns by the independent *t*-test with Bonferroni's correction for pairwise comparisons. Similar results were obtained in 3 experiments, and representative results are shown.

alone, but the levels of [<sup>3</sup>H]-thymidine incorporation were comparable in X ray-irradiated CD4<sup>+</sup> T cells from allergic and tolerant mice (Fig. 8B). In addition, when the levels of [<sup>3</sup>H]-thymidine incorporation in co-culture were compared between nonirradiated and X ray-irradiated CD4<sup>+</sup> T cells from tolerant mice, the level was higher in the co-culture



with X ray-irradiated CD4<sup>+</sup> T cells than in that with nonirradiated CD4<sup>+</sup> T cells. These results showed that viable CD4<sup>+</sup> T cells from tolerant mice are able to suppress the growth of DO11.10 CD4<sup>+</sup> T cells, suggesting that they produce a suppressor for the OVA-specific response of DO11.10 CD4<sup>+</sup> T cells.

## Discussion

Food allergy is considered to be an aberrant Th2-mediated immune response against dietary antigens<sup>1, 2</sup>. LPS, an outer wall component of Gram-negative bacteria, is a potent activator of the immune system, and it may play a crucial role in T cell priming<sup>5</sup>. Recent studies have demonstrated that LPS exposure primes Th2 cells in airway allergy, but the role of LPS in food allergy has not been well investigated<sup>12–15</sup>. In the present study, we established two distinct mouse models: one is an IgE-mediated, Th2-type allergic model induced by cOVA sensitization alone, and the other is a tolerant model induced by sensitization with both cOVA and LPS. In the allergic model, sensitization with cOVA contaminated with a little LPS could induce Th2-mediated reactions in mice. In the tolerant model, tolerance was increased with increasing doses of LPS during sensitization, and a high dose (100 µg) of LPS completely suppressed Th2-mediated allergic reactions. These results indicate that the development of food allergy may be induced by a low dose of LPS exposure and prevented by a high dose of LPS exposure during sensitization.

It is generally accepted that Th1 priming requires TLR4 signaling. It is also thought that the priming of Th2 occurs through a default pathway at a low level of TLR signaling, but the mechanism has been poorly defined<sup>6–11</sup>. Regarding airway allergy, Eisenbarth *et al.* reported that a high dose of LPS exposure during sensitization induces both IL-12 production and a Th1 response, whereas a low dose of LPS exposure is not sufficient to induce a Th1 response but is required to induce Th2 inflammation<sup>12</sup>. In our model, an IgE-mediated, Th2-type food allergy could be induced by cOVA sensitization. Because there is a possibility that a little LPS contaminated the cOVA and influenced the induction of allergic reactions, a low dose of LPS exposure during sensitization may play a crucial role in the priming of Th2 in both airway and food allergic reactions. On the other hand, unlike airway hypersensitivity, a high dose of LPS exposure failed to induce both IL-12 mRNA expression and Th1-mediated responses in our model. The reason for this discrepancy may lie in the environmental variation between the respiratory tract, where LPS is supplied mainly from exogenous sources, and the digestive tract, where an abundance of LPS is constitutively present.

Endotoxin tolerance is a phenomenon in which cells or organisms exposed to a low dose of endotoxin enter into a transient unresponsive state and are unable to respond to further challenges with the endotoxin<sup>32, 33</sup>. In endotoxin tolerance, dendritic cells (DCs) and monocytes/macrophages are unresponsive to subsequent antigen stimulation

by downregulating the expression of MHC class II and costimulatory molecules. It has been reported that endotoxin tolerance attenuates airway allergic inflammation through direct suppression of the T-cell stimulatory effect of DCs<sup>34</sup>. On the other hand, in our food allergic and tolerant models, MLN CD4<sup>+</sup> T cells from tolerant mice showed sufficient antigen responses after reexposure to the antigen. These results indicate that unresponsiveness in tolerant mice is not caused by endotoxin tolerance.

The production state of IL-12 is crucial to determine the subsequent acquired immunity<sup>13, 30, 35</sup>. Upregulation of IL-12 is generally related to the induction of Th1-mediated responses, and its downregulation is generally related to that of Th2-mediated responses. It is therefore thought that downregulation of IL-12 results in allergic reactions mediated by aberrant Th2 cells. However, IL-12 mRNA expression decreased markedly in spleen cells of our tolerant mice, showing that the development of Th2-mediated allergic reactions is prevented under IL-12 exhausted conditions. It is therefore likely that factors other than IL-12 participate in the suppression of Th2-mediated allergic reactions and the induction of tolerance. Our preliminary FACS analysis showed prominent decreases in CD11<sub>C</sub><sup>high</sup> DCs, which produce IL-12 and induce Th1-mediated responses, in the spleen of tolerant mice (data not shown). So, it is possible that downregulation of IL-12 mRNA expression is caused by a decrease in CD11<sub>C</sub><sup>high</sup> DCs in the spleen of tolerant mice.

Several studies have shown the crucial role of Treg in allergic diseases<sup>20, 21, 36</sup>. The involvement of Treg producing TGF-β, Th3, in the induction of oral tolerance and the maintenance of intestinal homeostasis has been reported<sup>27, 28</sup>. In our tolerant mice, upregulation of TGF-β mRNA expression was not observed. These results indicate that TGF-β does not participate in the induction of tolerance. On the other hand, the enhanced mRNA expression of IL-10, originally known as a Th2-derived cytokine synthesis inhibitory factor, was clearly identified in our tolerant mice. Groux *et al.* first reported Treg producing IL-10, Tr1, which suppress antigen-specific immune responses and actively downregulate a pathological immune response *in vivo*<sup>37</sup>. Subsequent studies have shown that Tr1 are able to inhibit the development of airway allergic inflammation in mouse models<sup>21, 38, 39</sup>. In a similar way, Tr1 may be induced by sensitization with cOVA and LPS and regulate the immune response to OVA in our tolerant mice. Wakkache *et al.* reported that CD11c<sup>low</sup>CD45RB<sup>high</sup> DCs that produce IL-10 are enriched after LPS stimulation, inducing the differentiation of Tr1<sup>40</sup>. In addition to the decrease in CD11<sub>C</sub><sup>high</sup> DCs, our preliminary FACS analysis showed that the proportion of CD11c<sup>low</sup>CD45RB<sup>high</sup> DCs remarkably increased in the spleen of tolerant mice (data not shown). There is a possibility that, in our tolerant mice, LPS exposure during sensitization functions as a trigger for activation of CD11c<sup>low</sup>CD45RB<sup>high</sup> DCs, promoting Tr1 development and IL-10 production by Tr1. Our tolerant mice may be useful in study of the function of CD11c<sup>low</sup>CD45RB<sup>high</sup> DCs and Tr1 as crucial regulators for

food allergy.

Thus, LPS is crucial for the regulation of Th2-mediated allergic reactions to dietary antigens. Understanding of the mechanisms involved in the induction of LPS-induced tolerogenic cells can potentially provide useful information for developing novel strategies to prevent food allergy or to treat allergic disease by therapeutic manipulation of the innate immune system.

**Declaration of Conflicting of Interests:** The authors declare that they have no conflicts of interest.

**Acknowledgements:** This study was supported in part by a grant from the Ministry of Education, Culture, Sports, Science and Technology of Japan, grants-in-aid for Scientific Research and the Hitec Research Center and grants-in-aid for Researchers from Hyogo College of Medicine.

## References

1. Sampson HA, and Burks AW. Mechanisms of food allergy. *Annu Rev Nutr.* **16**: 161–177. 1996. [Medline] [CrossRef]
2. Sampson HA, and Ho DG. Relationship between food-specific IgE concentrations and the risk of positive food challenges in children and adolescents. *J Allergy Clin Immunol.* **100**: 444–451. 1997. [Medline] [CrossRef]
3. Verdú EF, Bercik P, Bergonzelli GE, Huang XX, Blennerhasset P, Rochat F, Fiaux M, Mansourian R, Corthésy-Theulaz I, and Collins SM. *Lactobacillus paracasei* normalizes muscle hypercontractility in a murine model of postinfective gut dysfunction. *Gastroenterology.* **127**: 826–837. 2004. [Medline] [CrossRef]
4. Moon A, and Kleinman RE. Allergic gastroenteropathy in children. *Ann Allergy Asthma Immunol.* **74**: 5–12, quiz 12–16. 1995. [Medline]
5. Brix S, Kjaer TM, Barkholt V, and Frøkiaer H. Lipopolysaccharide contamination of beta-lactoglobulin affects the immune response against intraperitoneally and orally administered antigen. *Int Arch Allergy Immunol.* **135**: 216–220. 2004. [Medline] [CrossRef]
6. Poltorak A, He X, Smirnova I, Liu MY, Van Huffel C, Du X, Birdwell D, Alejos E, Silva M, Galanos C, Freudenberg M, Ricciardi-Castagnoli P, Layton B, and Beutler B. Defective LPS signaling in C3H/HeJ and C57BL/10ScCr mice: mutations in Tlr4 gene. *Science.* **282**: 2085–2088. 1998. [Medline] [CrossRef]
7. Qureshi ST, Larivière L, Leveque G, Clermont S, Moore KJ, Gros P, and Malo D. Endotoxin-tolerant mice have mutations in Toll-like receptor 4 (Tlr4). *J Exp Med.* **189**: 615–625. 1999. [Medline] [CrossRef]
8. Braun-Fahrlander C, Riedler J, Herz U, Eder W, Waser M, Grize L, Maisch S, Carr D, Gerlach F, Bufe A, Lauener RP, Schierl R, Renz H, Nowak D, von Mutius E. Allergy and Endotoxin Study Team. Environmental exposure to endotoxin and its relation to asthma in school-age children. *N Engl J Med.* **347**: 869–877. 2002. [Medline] [CrossRef]
9. Liu AH. Endotoxin exposure in allergy and asthma: reconciling a paradox. *J Allergy Clin Immunol.* **109**: 379–392. 2002. [Medline] [CrossRef]
10. Reed CE, and Milton DK. Endotoxin-stimulated innate immunity: A contributing factor for asthma. *J Allergy Clin Immunol.* **108**: 157–166. 2001. [Medline] [CrossRef]
11. Schwartz DA. Does inhalation of endotoxin cause asthma? *Am J Respir Crit Care Med.* **163**: 305–306. 2001. [Medline]
12. Eisenbarth SC, Piggott DA, Huleatt JW, Visintin I, Herrick CA, and Bottomly K. Lipopolysaccharide-enhanced, toll-like receptor 4-dependent T helper cell type 2 responses to inhaled antigen. *J Exp Med.* **196**: 1645–1651. 2002. [Medline] [CrossRef]
13. Kuipers H, Hijdra D, De Vries VC, Hammad H, Prins JB, Coyle AJ, Hoogsteden HC, and Lambrecht BN. Lipopolysaccharide-induced suppression of airway Th2 responses does not require IL-12 production by dendritic cells. *J Immunol.* **171**: 3645–3654. 2003. [Medline] [CrossRef]
14. Langenkamp A, Messi M, Lanzavecchia A, and Sallusto F. Kinetics of dendritic cell activation: impact on priming of TH1, TH2 and nonpolarized T cells. *Nat Immunol.* **1**: 311–316. 2000. [Medline] [CrossRef]
15. Tan AM, Chen HC, Pochard P, Eisenbarth SC, Herrick CA, and Bottomly HK. TLR4 signaling in stromal cells is critical for the initiation of allergic Th2 responses to inhaled antigen. *J Immunol.* **184**: 3535–3544. 2010. [Medline] [CrossRef]
16. Bashir ME, Louie S, Shi HN, and Nagler-Anderson C. Toll-like receptor 4 signaling by intestinal microbes influences susceptibility to food allergy. *J Immunol.* **172**: 6978–6987. 2004. [Medline] [CrossRef]
17. Berg RD. The indigenous gastrointestinal microflora. *Trends Microbiol.* **4**: 430–435. 1996. [Medline] [CrossRef]
18. Cebra JJ. Influences of microbiota on intestinal immune system development. *Am J Clin Nutr.* **69**: 1046S–1051S. 1999. [Medline]
19. Sudo N, Sawamura S, Tanaka K, Aiba Y, Kubo C, and Koga Y. The requirement of intestinal bacterial flora for the development of an IgE production system fully susceptible to oral tolerance induction. *J Immunol.* **159**: 1739–1745. 1997. [Medline]
20. Strobel S, and Mowat AM. Oral tolerance and allergic responses to food proteins. *Curr Opin Allergy Clin Immunol.* **6**: 207–213. 2006. [Medline] [CrossRef]
21. Umetsu DT, Akbari O, and Dekruyff RH. Regulatory T cells control the development of allergic disease and asthma. *J Allergy Clin Immunol.* **112**: 480–487, quiz 488. 2003. [Medline] [CrossRef]
22. van Wijk F, Wehrens EJ, Nierkens S, Boon L, Kasran A, Pieters R, and Knippels LM. CD4+CD25+ T cells regulate the intensity of hypersensitivity responses to peanut, but are not decisive in the induction of oral sensitization. *Clin Exp Allergy.* **37**: 572–581. 2007. [Medline] [CrossRef]
23. Karlsson MR, Rugtveit J, and Brandtzaeg P. Allergen-responsive CD4+CD25+ regulatory T cells in children who have outgrown cow's milk allergy. *J Exp Med.* **199**: 1679–1688. 2004. [Medline] [CrossRef]
24. Roncarolo MG, Bacchetta R, Bordignon C, Narula S, and Levings MK. Type 1 T regulatory cells. *Immunol Rev.* **182**: 68–79. 2001. [Medline] [CrossRef]
25. Takayama N, Igarashi O, Kweon MN, and Kiyono H. Regulatory role of Peyer's patches for the inhibition of OVA-induced allergic diarrhea. *Clin Immunol.* **123**: 199–208. 2007. [Medline] [CrossRef]

26. Ando T, Hatsushika K, Wako M, Ohba T, Koyama K, Ohnuma Y, Katoh R, Ogawa H, Okumura K, Luo J, Wyss-Coray T, and Nakao A. Orally administered TGF- $\beta$  is biologically active in the intestinal mucosa and enhances oral tolerance. *J Allergy Clin Immunol.* **120**: 916–923. 2007. [[Medline](#)] [[CrossRef](#)]
27. Chen Y, Kuchroo VK, Inobe J, Hafler DA, and Weiner HL. Regulatory T cell clones induced by oral tolerance: suppression of autoimmune encephalomyelitis. *Science.* **265**: 1237–1240. 1994. [[Medline](#)] [[CrossRef](#)]
28. Nakao A, Miike S, Hatano M, Okumura K, Tokuhisa T, Ra C, and Iwamoto I. Blockade of transforming growth factor  $\beta$ /Smad signaling in T cells by overexpression of Smad7 enhances antigen-induced airway inflammation and airway reactivity. *J Exp Med.* **192**: 151–158. 2000. [[Medline](#)] [[CrossRef](#)]
29. Lin AH, Luo J, Mondschein LH, ten Dijke P, Vivien D, Contag CH, and Wyss-Coray T. Global analysis of Smad2/3-dependent TGF- $\beta$  signaling in living mice reveals prominent tissue-specific responses to injury. *J Immunol.* **175**: 547–554. 2005. [[Medline](#)] [[CrossRef](#)]
30. Yanagawa Y, and Onoé K. Enhanced IL-10 production by TLR4- and TLR2-primed dendritic cells upon TLR restimulation. *J Immunol.* **178**: 6173–6180. 2007. [[Medline](#)] [[CrossRef](#)]
31. Brandt EB, Strait RT, Hershko D, Wang Q, Muntel EE, Scribner TA, Zimmermann N, Finkelman FD, and Rothenberg ME. Mast cells are required for experimental oral allergen-induced diarrhea. *J Clin Invest.* **112**: 1666–1677. 2003. [[Medline](#)] [[CrossRef](#)]
32. Biswas SK, and Lopez-Collazo E. Endotoxin tolerance: new mechanisms, molecules and clinical significance. *Trends Immunol.* **30**: 475–487. 2009. [[Medline](#)] [[CrossRef](#)]
33. Freudenberg MA, and Galanos C. Induction of tolerance to lipopolysaccharide (LPS)-D-galactosamine lethality by pretreatment with LPS is mediated by macrophages. *Infect Immun.* **56**: 1352–1357. 1988. [[Medline](#)]
34. Matsushita H, Ohta S, Shiraishi H, Suzuki S, Arima K, Toda S, Tanaka H, Nagai H, Kimoto M, Inokuchi A, and Izuhara K. Endotoxin tolerance attenuates airway allergic inflammation in model mice by suppression of the T-cell stimulatory effect of dendritic cells. *Int Immunol.* **22**: 739–747. 2010. [[Medline](#)] [[CrossRef](#)]
35. Akbari O, DeKruyff RH, and Umetsu DT. Pulmonary dendritic cells producing IL-10 mediate tolerance induced by respiratory exposure to antigen. *Nat Immunol.* **2**: 725–731. 2001. [[Medline](#)] [[CrossRef](#)]
36. Shreffler WG, Wanich N, Moloney M, Nowak-Wegrzyn A, and Sampson HA. Association of allergen-specific regulatory T cells with the onset of clinical tolerance to milk protein. *J Allergy Clin Immunol.* **123**: 43–52 e7. 2009.
37. Groux H, O'Garra A, Bigler M, Rouleau M, Antonenko S, de Vries JE, and Roncarolo MGA. A CD4<sup>+</sup> T-cell subset inhibits antigen-specific T-cell responses and prevents colitis. *Nature.* **389**: 737–742. 1997. [[Medline](#)] [[CrossRef](#)]
38. Bellinghausen I, Knop J, and Saloga J. The role of interleukin 10 in the regulation of allergic immune responses. *Int Arch Allergy Immunol.* **126**: 97–101. 2001. [[Medline](#)] [[CrossRef](#)]
39. Cottrez F, Hurst SD, Coffman RL, and Groux H. T regulatory cells 1 inhibit a Th2-specific response *in vivo*. *J Immunol.* **165**: 4848–4853. 2000. [[Medline](#)] [[CrossRef](#)]
40. Wakkach A, Fournier N, Brun V, Breitmayer JP, Cottrez F, and Groux H. Characterization of dendritic cells that induce tolerance and T regulatory 1 cell differentiation *in vivo*. *Immunity.* **18**: 605–617. 2003. [[Medline](#)] [[CrossRef](#)]