

In Silico Analysis of the *cadF* Gene and Development of a Duplex Polymerase Chain Reaction for Species-Specific Identification of *Campylobacter jejuni* and *Campylobacter coli*

Saeed Shams,¹ Bita Bakhshi,^{1,*} and Tahereh Tohidi Moghadam²

¹Department of Medical Bacteriology, Faculty of Medical Sciences, Tarbiat Modares University, Tehran, IR Iran

²Department of Nanobiotechnology, Faculty of Biological Sciences, Tarbiat Modares University, Tehran, IR Iran

*Corresponding author: Bita Bakhshi, Department of Medical Bacteriology, Faculty of Medical Sciences, Tarbiat Modares University, Tehran, IR Iran. Tel: +98-2182884558, Fax: +98-2182884555, E-mail: b.bakhshi@modares.ac.ir

Received 2015 May 4; Revised 2015 September 28; Accepted 2015 November 3.

Abstract

Background: Campylobacteriosis is a zoonotic infectious disease caused by *Campylobacter jejuni* and *C. coli*. The *cadF* gene is considered as a genus-specific gene while other genes are mainly used for discrimination at the species level.

Objectives: This study aimed to analyze the *cadF* gene and to develop a duplex PCR assay for simultaneous detection of *C. coli* and *C. jejuni*, the two commonly encountered species.

Materials and Methods: *In silico* analysis of the *cadF* gene was carried out by several software and available online tools. A duplex PCR optimized with specific primers was used for detection and differentiation of both species. To evaluate specificity and sensitivity of the test, a panel of different *Campylobacter* spp. together with several intestinal bacterial pathogens was tested. The limit of detection (LOD) of method was determined using serial dilutions of standard genomes.

Results: The analysis of the full size *cadF* gene indicated variations in this gene, which can be used to differentiate *C. jejuni* and *C. coli*. The duplex PCR designed in this study showed that it could simultaneously detect and differentiate both *C. jejuni* and *C. coli* with product sizes of 737 bp and 461 bp, respectively. This assay, with 100% specificity and sensitivity, had a limit of detection (LOD) of about 14 and 0.7 µg/mL for *C. jejuni* and *C. coli*, respectively.

Conclusions: *In silico* analysis of the *cadF* full-gene showed variations between the two species that can be used as a molecular target for differentiating *C. jejuni* and *C. coli* in a single-step duplex-PCR assay with high specificity and sensitivity.

Keywords: *In Silico*, Duplex PCR, *cadF*, *Campylobacter jejuni*, *C. coli*

1. Background

Campylobacter enteritis is one of the most frequent food-borne infections worldwide (1). Thermophilic *C. jejuni* and *C. coli* have been recognized as the most common causes of bacterial diarrhea in humans, especially among children less than five years of age and young adults (2). Although, poultry and poultry products are important sources of Campylobacteriosis, yet the organism can be transmitted to humans via contact with other warm-blooded animals such as cattle, pigs, sheep, ostriches, shellfish, and pets (3, 4).

The symptoms of campylobacteriosis can vary from mild to severe complications, including abdominal pain, fever, myalgia and watery or bloody diarrhea. Although, in most cases the illness is self-limited and rarely fatal yet post-infectious acute immune-mediated neurologic complications such as Guillain-Barre syndrome and Miller Fisher syndrome can occur, which are the consequence of molecular mimicry between lipooligosaccharides (LOS) of bacterial cell wall and gangliosides in periph-

eral nerves of humans (5, 6). These complications can be prevented or lowered with rapid and accurate detection of etiological agents of the disease. Diagnosis of campylobacteriosis is performed through microbiological, molecular and serological tests. Culture is the gold standard of diagnosis of *C. coli* and *C. jejuni*; however, the culture conditions for detection of these fastidious bacteria are complicated and time consuming, which in some cases make the recovery of bacteria unsuccessful. Moreover, the emergence of viable but non-culturable (VBNC) phenotypes should not be ignored.

Differentiation of the two species is only performed through hippurate hydrolysis biochemical test or molecular-based detections (7-10). In molecular methods different genetic targets have been used for the detection *Campylobacter* species (e.g. *asp*, *hipO*, *ceuE*, *cadF*, *16SrRNA*, *23S rRNA* and *cdt*, *fur*, *glyA*, *cdtABC*, *ceuB-E* and *fliY*) (9). Among them, the *cadF* gene encodes a fibronectin-binding protein that promotes bacteria-host cell interaction and has

been described as a conserve and genus-specific gene. In most studies a fragment from this gene with a length of 400 bp is used for identification of *Campylobacter* spp. at the genus level (1, 11-15). There is no documented bioinformatics study on the *cadF* gene full-sequence analysis in *C. jejuni* and *C. coli*.

2. Objectives

The aim of this study was to analyze the *cadF* gene and to develop and evaluate a single-step duplex polymerase chain reaction (PCR) assay for simultaneous detection of *C. coli* and *C. jejuni*, the two commonly encountered species in human Campylobacteriosis.

3. Materials and Methods

3.1. Alignment of *cadF* Sequences From GenBank

The *cadF* sequences from the complete genome of *C. jejuni* and *C. coli* were acquired from NCBI GenBank (<http://www.ncbi.nlm.nih.gov/>) (Table 1). Multiple alignments were performed using the CLC sequence viewer 7.6 software (CLC bio, Aarhus, Denmark).

3.2. In Silico Analysis of the *cadF* Gene

The conserved internal fragment (400 bp) of the *cadF* gene, reported by Konkel et al. (15) as a specific gene for detection of *Campylobacter* spp., was used as a reference sequence in this study. This fragment and other selected sequences from the full gene of *cadF* were subjected to *in silico* analysis with the online NEB cutter program (<http://tools.neb.com/NEBcutter>) to compare and select a proper

restriction endonuclease for discriminating between *C. jejuni* and *C. coli* using analysis of enzymatic digestion pattern.

3.3. Designing a Duplex Polymerase Chain Reaction Assay for Specific Detection of *Campylobacter jejuni* and *C. coli*

The entire *cadF* sequence obtained from GenBank was robustly examined for the presence of intra-species conserved regions, which could differentiate inter-species. Universal forward primer, FU, (position 101 - 120) and reverse primer, R1, (position 478 - 497) were selected for the *cadF* gene, and were previously described by Konkel et al. (15). Other reverse primers, R2 (position 542 - 561) and R3 (position 818 - 837), were designed in this study using the Genrunner and CLC sequence viewer software (Table 2). Analysis of the designed primers was performed by the Primer-BLAST on NCBI (<http://www.ncbi.nlm.nih.gov/>). Schematic representation of the PCR amplification of fragments related to *C. jejuni* and *C. coli* in duplex PCR is shown in Figure 1. Oligonucleotide primers were synthesized by TAG Copenhagen (Denmark).

The duplex-PCR was carried out in a 25- μ L reaction mixture, containing 10 ng of DNA template extracted by the boiling method, 2.5 μ L PCR buffer 10X, 200 μ M dNTP, 5 mM MgCl₂, 0.1 μ M of each primer, 1 unit of Taq DNA polymerase, and sterile deionized water (12, 14). Amplification conditions were 95°C for three minutes (one cycle), then denaturation at 94°C for 30 seconds, annealing at 43°C for 30 seconds and extension at 72°C for 30 seconds for 32 cycles in a thermocycler (Eppendorf, Hamburg, Germany). Finally, an additional extension step (five minutes, 72°C) was carried out.


Table 1. *Campylobacter cadF* Sequences Used in This Study

Definition	Accession No.
<i>Campylobacter jejuni</i> subsp. <i>jejuni</i> strain MTVDSJ20, complete genome	CP008787.1
<i>Campylobacter jejuni</i> subsp. <i>jejuni</i> 00-2538, complete genome	CP006707.2
<i>Campylobacter jejuni</i> subsp. <i>jejuni</i> 00-2544, complete genome	CP006709.2
<i>Campylobacter jejuni</i> subsp. <i>jejuni</i> PT14, complete genome	NC_018709.2
<i>Campylobacter jejuni</i> subsp. <i>jejuni</i> NCTC 11168 complete genome	AL111168.1
<i>Campylobacter jejuni</i> RM1221, complete genome	CP000025.1
<i>Campylobacter jejuni</i> subsp. <i>jejuni</i> 81116, complete genome	CP000814.1
<i>Campylobacter coli</i> RM2228 cont193, whole genome shotgun sequence	AAFL01000010.1
<i>Campylobacter coli</i> RM1875, complete genome	CP007183.1
<i>Campylobacter coli</i> 15-537360, complete genome	CP006702.1
<i>Campylobacter coli</i> RM5611, complete genome	CP007179.1
<i>Campylobacter coli</i> CVM N29710, complete genome	CP004066.1
<i>Campylobacter coli</i> RM4661, complete genome	CP007181.1
<i>Campylobacter coli</i> JV20 contig00034, whole genome shotgun sequence	AEER01000022.1
<i>Campylobacter coli</i> JV20 genomic scaffold SCAFFOLD1, whole genome shotgun sequence	GL405235.1

Table 2. Primer Sequences for The *cadF* Gene Used in the Duplex Polymerase Chain Reaction Assay^a

Primer	Sequence (5' to 3')	Size of Product, bp	Target	Reference
FU	TTGAAGGTAATTAGATATG	400	<i>Campylobacter</i> spp.	(15)
R1	CTAATACCTAAAGTTGAAAC	400	<i>Campylobacter</i> spp.	(15)
R2	TTTATTAACACTCTCTTTTG	461	<i>C. coli</i>	This study
R3	ATATTTTCAAGTTCATTAG	737	<i>C. jejuni</i>	This study

^aAnnealing temperature is 43°C for all the primers.

Figure 1. Position of Primers for Developing the Duplex Polymerase Chain Reaction Assay

Black thin lines of 999 and 960 bp are related to the *cadF* gene of *C. coli* and *C. jejuni*, respectively. The discontinuity is related to deletion in *C. jejuni*. A, produced a 400-bp fragment from both *C. jejuni* and *C. coli* (as the positive control of the assay); B, produced a 461-bp fragment from *C. coli*; C, produced a 737-bp fragment from *C. jejuni*.

3.4. Limit of Detection, Sensitivity and Specificity of the Duplex Polymerase Chain Reaction

Limit of detection (LOD) of the amplification assay was evaluated using serial 10-fold dilutions of genomes with initial concentrations of 140 (*C. jejuni*) and 7 (*C. coli*) µg/mL. A total of 20 clinical and envi-

ronmental *Campylobacter* isolates were examined for further evaluation of the sensitivity. Specificity of the test was evaluated using genomic DNA from standard and isolated clinical strains of other enteric non-*Campylobacter* bacterial pathogens (Table 3). Sensitivity and specificity were calculated according to the following Equations (16):

$$(1) \quad \text{Sensitivity} = \frac{(\text{number of positive isolates, as determined by duplex PCR})}{(\text{total number of positive isolates as determined by three genes } (cadF / hipO / asp) \text{ PCR})} \times 100$$

$$(2) \quad \text{Specificity} = \frac{(\text{number of negative isolates, as determined PCR})}{(\text{total number of negative isolates, as determined by three genes } (cadF / hipO / asp) \text{ PCR})} \times 100$$

Table 3. List of Bacteria Used for the Determination of Specificity and Sensitivity of *cadF* Targeted Species-Specific Duplex Polymerase Chain Reaction^a

Organism Name	Strain Name	Amplification with Newly Designed Primers
<i>Shigella sonnei</i>	ATCC 25931	negative
<i>Shigella flexneri</i>	ATCC 12022	negative
<i>Shigella boydii</i>	ATCC 8700	negative
<i>Shigella dysenteriae</i>	ATCC 13313	negative
<i>Aeromonas hydrophila</i>	ATCC 7966	negative
<i>Enterobacter aerogenes</i>	ATCC 13048	negative
<i>Vibrio cholerae</i>	ATCC 39315	negative
<i>Enteropathogenic Escherichia coli</i>	ATCC 43887	negative
<i>Escherichia coli</i> O157:H7	ATCC 35150	negative
<i>Enteroinvasive Escherichia coli</i>	ATCC 43893	negative
<i>Enteroaggregative Escherichia coli</i>	ATCC 33780	negative
<i>Enterotoxigenic Escherichia coli</i>	ATCC 35401	negative
<i>Salmonella typhimurium</i>	ATCC 29946	negative
<i>Salmonella typhi</i>	ATCC 19430	negative
<i>Campylobacter jejuni</i>	ATCC 29428	positive
<i>Campylobacter coli</i>	ATCC 43478	positive
<i>Campylobacter coli</i>	Isolate 1	positive
<i>Campylobacter coli</i>	Isolate 2	positive
<i>Campylobacter coli</i>	Isolate 3	positive
<i>Campylobacter coli</i>	Isolate 4	positive
<i>Campylobacter coli</i>	Isolate 5	positive
<i>Campylobacter coli</i>	Isolate 6	positive
<i>Campylobacter coli</i>	Isolate 7	positive
<i>Campylobacter coli</i>	Isolate 8	positive
<i>Campylobacter coli</i>	Isolate 9	positive
<i>Campylobacter jejuni</i>	Isolate 1	positive
<i>Campylobacter jejuni</i>	Isolate 2	positive
<i>Campylobacter jejuni</i>	Isolate 3	positive
<i>Campylobacter jejuni</i>	Isolate 4	positive
<i>Campylobacter jejuni</i>	Isolate 5	positive
<i>Campylobacter jejuni</i>	Isolate 6	positive
<i>Campylobacter jejuni</i>	Isolate 7	positive
<i>Campylobacter jejuni</i>	Isolate 8	positive
<i>Campylobacter jejuni</i>	Isolate 9	positive
<i>Campylobacter jejuni</i>	Isolate 10	positive
<i>Campylobacter jejuni</i>	Isolate 11	positive

^aSource of isolations is clinical.

4. Results


4.1. In Silico Analysis of the *cadF* Gene

The length of the *cadF* sequences extracted from complete genome of *C. jejuni* and *C. coli* was 960 (with C + G 31.8% and A + T 68.2%) and 999 bp (with C + G 34% and A + T 66%), respectively. The *cadF* gene in both species was located after the *rpsL* gene, which coded for a 30S ribosomal

protein. Although there were some nucleotide variations along the sequence of *cadF* between the two species, yet the main difference was related to the 39-bp deletion in the positions of 533 - 544 and 560 - 586 of *C. jejuni* (Figure 2). The results of the BLAST analysis of *cadF* gene showed an average sequence identity of 98.5% and 94% among *C. jejuni* and *C. coli* strains, respectively. The identity between the two species was also estimated as 88%, approximately (Table 4).

terial species. The assay showed limit of detection (LOD) of 14 and 0.7 µg/mL (approximately equal to 7×10^9 and 3×10^8 copy number) for *C. jejuni* and *C. coli*, respectively.

Figure 3. Agarose Gel Electrophoresis of the Duplex Polymerase Chain Reaction Assay with Specific Primers



Lanes 1 and 2, 400-bp fragment of the *cadF* gene of *Campylobacter jejuni* and *Campylobacter coli*, respectively as positive controls; lanes 4 - 6, 737 bp fragment of *C. jejuni*; lanes 9 - 11, 461 bp fragment of *C. coli*; lanes 3, 7, 8, 12 and 13, negative controls; lane M, 1 kb molecular weight marker.

5. Discussion

Campylobacter jejuni and *C. coli* are now recognized as important causes of acute bacterial diarrhea in most countries. The isolation and discrimination of *C. jejuni* and *C. coli* by biochemical tests at the species level is limited and laborious, thus there is a crucial need to develop a sensitive, validated and rapid DNA-based method for detection of *Campylobacter* at the species level (10, 17). In some studies, multiple genes have been used for distinguishing *C. jejuni* and *C. coli*. Al Amri et al. (2007) developed a multiplex PCR assay using the combination of a genus-specific virulence gene (*cadF*) together with hippuricase and aspartokinase genes (*asp*) for species-specific identification of *C. jejuni* and *C. coli*, respectively (11). In a study by Cloak and Fratamico (2002), a multiplex PCR was designed for differentiation of *C. jejuni* and *C. coli* by means of *cadF* and *ceuE* genes. In another work by Adzitey and Corry (2011), *lpxA*, *hipO* and *glyA* genes were used for differentiating *C. jejuni* and *C. coli* species. The study of Nayak et al. (2005) was also designed with *cadF*, *ceuE* and oxidoreductase subunit genes as fragments of 400-bp conserved region in *Campylobacter* spp. 894-bp specific for *C. coli* and 160-bp specific for *C. jejuni*, respectively (14, 18, 19).

Our duplex PCR method was developed only with the *cadF* gene and the specificity and sensitivity of novel reverse primers (R_2 and R_3) in association with a previously described forward primer (FU) was studied. The PCR assay designed in this work showed 100% sensitivity and specificity while no amplification product was seen for the genomic DNA from non-*Campylobacter* enteric bacteria. One applicable advantage of this newly designed duplex PCR assay is that the amplified products are of different sizes, which can be concurrently visualized on agarose gel without the need to duplicate the reaction or further electrophoresis and sequencing. In a similar study, Klena et al. used divergence and conservation

regions of *lpxA* to develop a robust PCR assay. They differentiated *C. coli*, *C. jejuni*, *C. lari* and *C. upsaliensis* using multiplex PCR with the lipid A gene *lpxA*, encoding a UDP-N-acetyl glucosamine acyl transferase. Another work similar to our research was the study of Gonzalez et al. They discriminated *C. jejuni* and *C. coli* by using *ceuE* gene diversity (approximately 13%) between two species (20, 21).

The lowest concentration of genomic DNA for detection of *C. jejuni* and *C. coli* was 14 and 0.7 µg/mL (approximately equal to 7×10^9 and 3×10^8 copy number), respectively. These LODs are almost comparable with the study conducted by Wisessombat et al., in which the sensitivity of the multiplex PCR for the detection of *Campylobacter* spp. was 2×10^5 CFU/PCR (22). Another study indicated that the colony multiplex PCR sensitivity range for *C. jejuni* and *C. coli* was 10^8 to 10^{13} and 10^6 to 10^{13} CFU/mL, respectively (23).

The bioinformatics data analysis of the 400-bp internal section of the *cadF* introduced by Konkel et al. which has been used by many investigators for genus-specific detection of *Campylobacter* spp. showed that this fragment is highly conserved among *C. jejuni* and *C. coli* strains and is significantly validated for the identification of both species. It seems that there is a concomitant general misjudged belief that the *cadF* full-gene is genus-specific. Our analysis of total *cadF* sequence revealed that other than single-nucleotide variations between two bacteria, an approximately 4% deletion has occurred in the *cadF* sequence of *C. jejuni* compared with *C. coli*, which could be useful for our work. The intra-species identity level among *C. jejuni* and *C. coli* strains was about 98.5% and 94%, respectively. The identity level was approximately 88% between the two species. These results were similar to the report of Konkel et al. with 87% identity between *C. jejuni* and *C. coli* and 98.6% among *C. jejuni* strains, individually (15).

There are several articles about the PCR-RFLP method for the differentiation of *Campylobacter* spp. using genes other than *cadF* (24, 25). Although the restriction pattern of enzymatic digestion of the 400-bp fragment introduced by Konkel et al. is not suitable for separation of the two species, yet the enzymatic digestion of the full-length gene may be useful for differentiation and clinical diagnosis of *C. jejuni* and *C. coli* (15) The *cadF* full-gene has some variations in its sequence and length between species, which can be beneficial for developing a duplex PCR. The designed PCR assay in this study is highly sensitive and specific and provides an accurate, inexpensive, sensitive and specific tool for rapid and simultaneous detection and differentiation of *C. coli* and *C. jejuni* in clinical settings.

Footnotes

Authors' Contribution: Study concept, design analysis and interpretation of data: Bitra Bakhshi, Tahereh Tohidi Moghadam and Saeed Shams; performance and drafting

of the manuscript: Saeed Shams; critical revision of the manuscript: Bitra Bakhshi.

Funding/Support: This work was supported by a grant from the research council of Tarbiat Modares university and it was part of a PhD thesis by Saeed Shams in medical bacteriology.

References

- Randall L, Lemma F, Rodgers J, Vidal A, Clifton-Hadley F. Development and evaluation of internal amplification controls for use in a real-time duplex PCR assay for detection of *Campylobacter coli* and *Campylobacter jejuni*. *J Med Microbiol*. 2010;**59**(Pt 2):172-8. doi: 10.1099/jmm.0.014415-0. [PubMed: 19833779]
- Galanis E. *Campylobacter* and bacterial gastroenteritis. *CMAJ*. 2007;**177**(6):570-1. doi: 10.1503/cmaj.070660. [PubMed: 17846438]
- Sarkar SR, Hossain MA, Paul SK, Ray NC, Sultana S, Rahman MM, et al. *Campylobacteriosis* - an overview. *Mymensingh Med J*. 2014;**23**(1):173-80. [PubMed: 24584395]
- World Health Organization. *Campylobacter Fact Sheet N255*. Geneva: WHO; October 2011. Available from: <http://www.who.int/mediacentre/factsheets/fs255/en/>.
- Haag LM, Fischer A, Otto B, Grundmann U, Kuhl AA, Gobel UB, et al. *Campylobacter jejuni* infection of infant mice: acute enterocolitis is followed by asymptomatic intestinal and extra-intestinal immune responses. *Eur J Microbiol Immunol (Bp)*. 2012;**2**(1):2-11. doi: 10.1556/EuJMI.2.2012.1.2. [PubMed: 24611115]
- Godschalk PC, Kuijf ML, Li J, St Michael F, Ang CW, Jacobs BC, et al. Structural characterization of *Campylobacter jejuni* lipooligosaccharide outer cores associated with Guillain-Barre and Miller Fisher syndromes. *Infect Immun*. 2007;**75**(3):1245-54. doi: 10.1128/IAI.00872-06. [PubMed: 17261613]
- Williams LK, McMeechan A, Baalham T, Ward L, Humphrey TJ, Jorgensen F. Detection of *Campylobacter jejuni* and *Campylobacter coli* from broiler chicken-related samples using BAX PCR and conventional International Organization for Standardization culture. *J Food Prot*. 2008;**71**(4):835-8. [PubMed: 18468043]
- Jackson DN, Davis B, Tirado SM, Duggal M, van Frankenhuyzen JK, Deaville D, et al. Survival mechanisms and culturability of *Campylobacter jejuni* under stress conditions. *Antonie Van Leeuwenhoek*. 2009;**96**(4):377-94. doi: 10.1007/s10482-009-9378-8. [PubMed: 19774483]
- Persson S, Olsen KE. Multiplex PCR for identification of *Campylobacter coli* and *Campylobacter jejuni* from pure cultures and directly on stool samples. *J Med Microbiol*. 2005;**54**(Pt 11):1043-7. doi: 10.1099/jmm.0.46203-0. [PubMed: 16192435]
- Kulkarni SP, Lever S, Logan JM, Lawson AJ, Stanley J, Shafi MS. Detection of *Campylobacter* species: a comparison of culture and polymerase chain reaction based methods. *J Clin Pathol*. 2002;**55**(10):749-53. [PubMed: 12354800]
- Al Amri A, Senok AC, Ismael AY, Al-Mahmeed AE, Botta GA. Multiplex PCR for direct identification of *Campylobacter* spp. in human and chicken stools. *J Med Microbiol*. 2007;**56**(Pt 10):1350-5. doi: 10.1099/jmm.0.47220-0. [PubMed: 17893173]
- Ghorbanalizadgan M, Bakhshi B, Kazemnejad Lili A, Najar-Peerayeh S, Nikmanesh B. A molecular survey of *Campylobacter jejuni* and *Campylobacter coli* virulence and diversity. *Iran Biomed J*. 2014;**18**(3):158-64. [PubMed: 24842142]
- Patrone V, Campana R, Vallorani L, Dominici S, Federici S, Casadei L, et al. CadF expression in *Campylobacter jejuni* strains incubated under low-temperature water microcosm conditions which induce the viable but non-culturable (VBNC) state. *Antonie Van Leeuwenhoek*. 2013;**103**(5):979-88. doi: 10.1007/s10482-013-9877-5. [PubMed: 23314927]
- Nayak R, Stewart TM, Nawaz MS. PCR identification of *Campylobacter coli* and *Campylobacter jejuni* by partial sequencing of virulence genes. *Mol Cell Probes*. 2005;**19**(3):187-93. doi: 10.1016/j.mcp.2004.11.005. [PubMed: 15797819]
- Konkel ME, Gray SA, Kim BJ, Garvis SG, Yoon J. Identification of the enteropathogens *Campylobacter jejuni* and *Campylobacter coli* based on the cadF virulence gene and its product. *J Clin Microbiol*. 1999;**37**(3):510-7. [PubMed: 9986804]
- Roozbehani AD, Bakhshi B, Pourshafie MR, Katouli M. A rapid and reliable species-specific identification of clinical and environmental isolates of *Vibrio cholerae* using a three-test procedure and recA polymerase chain reaction. *Indian J Med Microbiol*. 2012;**30**(1):39-43. doi: 10.4103/0255-0857.93027. [PubMed: 22361759]
- Keramas G, Bang DD, Lund M, Madsen M, Bunkenborg H, Telleman P, et al. Use of culture, PCR analysis, and DNA microarrays for detection of *Campylobacter jejuni* and *Campylobacter coli* from chicken feces. *J Clin Microbiol*. 2004;**42**(9):3985-91. doi: 10.1128/JCM.42.9.3985-3991.2004. [PubMed: 15364980]
- Cloak OM, Fratamico PM. A multiplex polymerase chain reaction for the differentiation of *Campylobacter jejuni* and *Campylobacter coli* from a swine processing facility and characterization of isolates by pulsed-field gel electrophoresis and antibiotic resistance profiles. *J Food Prot*. 2002;**65**(2):266-73. [PubMed: 11848556]
- Adzitey F, Corry J. A Comparison between Hippurate Hydrolysis and Multiplex PCR for Differentiating *Campylobacter coli* and *Campylobacter jejuni*. *Trop Life Sci Res*. 2011;**22**(1):91-8. [PubMed: 24575212]
- Klena JD, Parker CT, Knibb K, Ibbitt JC, Devane PM, Horn ST, et al. Differentiation of *Campylobacter coli*, *Campylobacter jejuni*, *Campylobacter lari*, and *Campylobacter upsaliensis* by a multiplex PCR developed from the nucleotide sequence of the lipid A gene *lpxA*. *J Clin Microbiol*. 2004;**42**(12):5549-57. doi: 10.1128/JCM.42.12.5549-5557.2004. [PubMed: 15583280]
- Gonzalez I, Grant KA, Richardson PT, Park SF, Collins MD. Specific identification of the enteropathogens *Campylobacter jejuni* and *Campylobacter coli* by using a PCR test based on the *ceuE* gene encoding a putative virulence determinant. *J Clin Microbiol*. 1997;**35**(3):759-63. [PubMed: 9041429]
- Wisessombat S, Inthagard J, Kittiniyom K, Srimanote P, Wonglumsom W, Voravuthikunchai SP. Multiplex PCR for direct identification of thermophilic *Campylobacter*, *C. jejuni*, *C. coli*, *C. lari* and *C. upsaliensis* and simultaneous detection of CDTB gene. *J Rapid Methods Autom Microbiol*. 2009;**17**(4):438-54.
- Wang G, Clark CG, Taylor TM, Pucknell C, Barton C, Price L, et al. Colony multiplex PCR assay for identification and differentiation of *Campylobacter jejuni*, *C. coli*, *C. lari*, *C. upsaliensis*, and *C. fetus* subsp. *fetus*. *J Clin Microbiol*. 2002;**40**(12):4744-7. [PubMed: 12454184]
- Kashoma IP, Kumar A, Sanad YM, Gebreyes W, Kazwala RR, Garabed R, et al. Phenotypic and genotypic diversity of thermophilic *Campylobacter* spp. in commercial turkey flocks: a longitudinal study. *Foodborne Pathog Dis*. 2014;**11**(11):850-60. doi: 10.1089/fpd.2014.1794. [PubMed: 25184688]
- Kamei K, Asakura M, Somroop S, Hatanaka N, Hinenoya A, Nagita A, et al. A PCR-RFLP assay for the detection and differentiation of *Campylobacter jejuni*, *C. coli*, *C. fetus*, *C. hyointestinalis*, *C. lari*, *C. helveticus* and *C. upsaliensis*. *J Med Microbiol*. 2014;**63**(Pt 5):659-66. doi: 10.1099/jmm.0.071498-0. [PubMed: 24568882]