

# Description of a *Zostera marina* catalase gene involved in responses to temperature stress

Yu Zang<sup>1</sup>, Jiao Liu<sup>3</sup>, Xue Xi Tang<sup>1,2</sup> and Bin Zhou<sup>1,2</sup>

<sup>1</sup> College of Marine Life Science, Ocean University of China, Qingdao, China

<sup>2</sup> Laboratory for Marine Ecology and Environmental Science, Qingdao National Laboratory for Marine Science and Technology, Qingdao, China

<sup>3</sup> Institute of Oceanology, Chinese Academy of Sciences, Qingdao, China

## ABSTRACT

Catalase (CAT) is an antioxidant enzyme that plays a significant role in cellular protection against oxidative damage by degradation of hydrogen peroxide to oxygen and water. In the present study, the complete CAT cDNA sequence of *Zostera marina* was identified through expressed sequence tags (EST) analysis and the rapid amplification of cDNA ends (RACE) technique. The nucleotide sequence of *ZmCAT* cDNA consisted of 1,816 bp with a 1,434 bp open reading frame (ORF), encoding a polypeptide of 477 amino acid residues, which possessed significant homology to other known plant CATs. The molecular mass of the predicted protein was 55.3 kDa with an estimated isoelectric point of 6.40. Phylogenetic analysis showed that *ZmCAT* was closely related to CAT from gramineous species. In response to temperature stress, H<sub>2</sub>O<sub>2</sub> and MDA contents in *Z. marina* increased significantly with cold stress (<10 °C) and heat stress (>25 °C). *ZmCAT* expression was significantly upregulated at temperatures from 5 to 10 °C and then gradually downregulated, reaching its lowest expression at 30 °C. Recombinant *ZmCAT* protein exhibited strong antioxidant activity over a wide temperature range, with the highest r*ZmCAT* activity observed at 25 °C and a higher relative activity retained even with heat stress. All these results indicated that *ZmCAT* was a member of the plant CAT family and involved in minimizing oxidative damage effects in *Z. marina* under temperature stress.

**Subjects** Ecology, Marine Biology, Molecular Biology

**Keywords** CAT, Temperature stress, mRNA expression, Enzyme activity

## INTRODUCTION

Seagrass meadows, covering ~0.1–0.2% of the global ocean floor, are highly productive ecosystems that provide key ecological services to the marine environment (*Erftemeijer & Robin Lewis III, 2006; Orth et al., 2006*). *Zostera marina* (or eelgrass), one of the most widespread seagrass species throughout the temperate northern hemisphere, plays an important ecological engineering role by influencing sediment stabilization, carbon cycling, and food web structure (*Fourqurean et al., 2012; Hemminga & Duarte, 2000*). In recent decades, *Z. marina*, as well as other species of seagrass is declining as a result of aquatic environmental changes, which will become more noticeable with sustained global climate change (*Orth et al., 2010; Short et al., 2011*). Temperature has long been recognized

Submitted 27 October 2017

Accepted 4 March 2018

Published 26 March 2018

Corresponding author

Bin Zhou, zhoubin@ouc.edu.cn

Academic editor

Bruno Marino

Additional Information and  
Declarations can be found on  
page 15

DOI 10.7717/peerj.4532

© Copyright  
2018 Zang et al.

Distributed under  
Creative Commons CC-BY 4.0

**OPEN ACCESS**

as a major environmental factor affecting the biogeographical distribution and health of seagrass meadows (Bulthuis, 1987; Ralph, 1998). Furthermore, recent large-scale seagrass populations have been precipitously decimated worldwide from historical abundances, caused by extreme summer heat waves (Moore, Shields & Parrish, 2014; Thomson et al., 2015), suggesting the question regarding how global climate change might aggravate this decline.

Temperature stress is known to accelerate the generation of reactive oxygen species (ROS), such as singlet oxygen ( $^1\text{O}_2$ ), hydrogen peroxide ( $\text{H}_2\text{O}_2$ ), superoxide radical ( $\text{O}_2\cdot^-$ ), and hydroxyl radicals ( $\cdot\text{OH}$ ), thereby leading to oxidative damage (Potters et al., 2007; Jena et al., 2013; Mittler, 2002). ROS production is a necessary process that can occur naturally and is related to both respiratory and photosynthetic metabolism, representing normal aerobic metabolism of plants. However, under abiotic stressful situations, homeostasis between the production and scavenging of ROS is altered and causes ROS accumulation (Wang et al., 2010). Excessive accumulation of ROS can cause oxidative damage through increasing lipid peroxidation (LPO), protein oxidation and degradation, and double-strand DNA breakage (Matés, 2000). To maintain homeostasis and protect against oxidative stress, plant cells have evolved antioxidant defense mechanisms, such as antioxidant enzymes, including catalase (CAT), superoxide dismutase, ascorbate peroxidase, glutathione peroxidase, glutathione reductase, and glutathione peroxidase (Hasanuzzaman, Nahar & Fujita, 2013). Among these enzymes, CAT plays a critical role in antioxidant defense pathways by efficiently catalyzing, in peroxisomes and glyoxysomes, the conversion of two molecules of  $\text{H}_2\text{O}_2$  to two molecules of  $\text{H}_2\text{O}$  and an  $\text{O}_2$  ( $2\text{H}_2\text{O}_2 \rightarrow 2\text{H}_2\text{O} + \text{O}_2$ ) through the transfer of two electrons, thus counteracting  $\text{H}_2\text{O}_2$  toxicity (Kashiwagi et al., 1997; Srivastava et al., 2012). Regulation mechanisms of CAT have been investigated at the transcriptional and enzymatic levels, which demonstrated that mechanisms of plant adaptations to heating and chilling stress might be positively correlated with CAT responses in maize (Scandalios, Acevedo & Ruzsa, 2000) cucumber (Gao, Qin & Yu, 2009) olive (Cansev, Gulen & Eris, 2011), broccoli (Lin, Huang & Lin, 2010) and banana (Figuerola-Yáñez et al., 2012).

The recent development of molecular resources for different seagrass species has provided important new insights into the knowledge of the transcriptional control of temperature stress responses (Franssen et al., 2014; Gu et al., 2012; Marin-Guirao et al., 2017). Our first attempts have already been carried out in the MnSOD gene of *Z. marina* and found that it was beneficial in minimizing oxidative damage effects of temperature stress (Liu et al., 2016). For seagrass, CAT activity has been detected in many species, such as *Zostera japonica* (Lin et al., 2016) and *Posidonia oceanica* (Sureda et al., 2008), but there have been no reports regarding detailed analysis and characterization of genes encoding CAT from any species of seagrass except for a partial CDS deposited in GenBank from *Cymodocea nodosa*. Study of the structural characteristics and antioxidant responses of *ZmCAT* will help us to determine the functional roles of *ZmCAT* in response to temperature stress. Thus, the purposes of this study were to: (1) clone the full-length cDNA sequence of *ZmCAT* and analyze the evolution of *ZmCAT* among other species (2) investigate the effects of temperature stress on  $\text{H}_2\text{O}_2$  and malondialdehyde (MDA) contents in *Z. marina*

(3) validate mRNA expression and recombinant protein activity of *ZmCAT* in response to temperature treatments.

## MATERIALS AND METHODS

### Plant material

Eelgrass vegetative shoots with attached roots were collected from the subtidal zone in Huiquan Bay (36°05'19.4"N, 120°34'46.2"E; Qingdao City, China) The research permit was issued to the College of Marine Life Science, Ocean University of China (Permit number: 2013OUCOLS0618). Plants were maintained in plastic boxes and transported within 2 h to the laboratory. Plant surface epiphytes and the older leaves were removed and individual shoots planted in plastic hydroponic planting baskets (100 mm inner diameter), which were filled with washed and sterilized silt (sediment particle size <63  $\mu\text{m}$ ) from a collection site in Huiquan Bay. The baskets were placed at aquaria bottoms and provided aerated filtered natural seawater at a constant temperature of 15 °C and allowed to acclimatize for 7 d. Fluorescent lamps provided an overhead light source with an applied light intensity of 150  $\mu\text{mol m}^{-2} \text{s}^{-1}$  and light/dark regime of 12/12 h. Healthy shoots of *Z.marina* were used for the following experiments.

### Temperature treatments

A total of 30 eelgrass plants were employed for the temperature stress stimulation treatment, with plants were randomly divided into 6 groups. Five eelgrass plants were evenly spaced and kept upright in each of 6 glass aquariums (195 L). Temperature treatments were at 5, 10, 15, 20, 25, and 30 °C for 96 h, maintained by an automatic water temperature regulator (RESUN, 800W/S2TS2500) in each aquarium. During the experiment, aquaria seawater was adequately aerated and circulated using multifunctional submersible pumps to ensure mixing and homogeneous temperature within each aquarium. All experimental seawater was completely changed daily to avoid nutrient limitations, and the salinity and water temperature was checked daily. After treatment, leaf samples ( $n = 5$ ) were cut into 100 mg (fresh weight, FW) sections after 5 cm segment from the leaf base, flash frozen in liquid nitrogen and rapidly stored at  $-80$  °C until further use.

### Total RNA extraction and cDNA synthesis of *ZmCAT*

Total RNA was extracted from fresh leaf samples (50 mg) of eelgrass with RNA Extract Kit (Aidlab Biotechnologies Ltd., Beijing, China) according to the manufacturer's instructions. The quality and integrity of total RNA were assessed by a NanoDrop (Thermo Fisher Scientific, Delaware, USA) and 1% agarose gel electrophoresis. The first-strand synthesis was carried out using the RQ1 RNase-free DNase I (Promega Corp., Madison, WI, USA) treated raw RNA (4  $\mu\text{g}$ ) as template and adaptor primer-oligo (dT) as primer (Table S1). The reaction were performed at 55 °C for 1 h and terminated by heating at 70 °C for 5 min. Products from this reaction were subsequently stored at  $-80$  °C.

## Expressed sequence tags analysis and cloning of full-length *ZmCAT* cDNA

BLAST analysis of all ESTs sequences from the *Z. marina* cDNA library in National Center for Biotechnology Information (NCBI) showed that an EST of 487 bps sequence (Accession No. [AM768877](#)) showed high similarity to identified CATs from *Musa acuminata subsp. Malaccensis* (accession number [XM009402721](#)); thus, this EST sequence was selected for further cloning the full-length cDNA sequence of *ZmCAT*. Two gene specific primers, *ZmCAT*-Race-F<sub>1</sub> and *ZmCAT*- Race-F<sub>2</sub> ([Table S1](#)), were designed to amplify the sequence of *ZmCAT* cDNA by the rapid amplification of cDNA ends (RACE) technique. PCR products were gel-purified and cloned into the T-A cloning vector pMD19-T simple (Takara Bio Inc. Shiga, Japan). After transformation into competent cells of *E. coli* DH5  $\alpha$ 0, positive recombinants were recognized by blue-white selection on ampicillin-containing Luria-Bertani plates and white colonies subsequently screened by PCR with M13-47 and RV-M primers ([Table S1](#)). Positive clones were sequenced on an ABI Prism<sup>TM</sup> 3730 automated DNA sequencer (Thermo Fisher Scientific Inc., Pittsburgh, PA, USA).

## Bioinformatical analysis of nucleotide and amino acid sequences

The nucleotide and deduced amino acid sequences of *ZmCAT* cDNA were analyzed using the BLAST software of NCBI and signal peptides predicted with the SignalP 4.1 Server. Besides, functional domains and signature motifs were identified using the normal mode of the Simple Modular Architecture Research Tool ([Letunic, Doerks & Bork, 2012](#)) and PROSITE database. Cellular localization of the inferred amino acid sequence was carried out with the PSORT Prediction tool of the PSORT WWW Server. A multiple sequence alignment was generated using the Clustal Version 2.0 program ([Larkin et al., 2007](#)). A phylogenetic tree was then produced by the neighbor-joining (NJ) approach of the MEGA 6.0 program ([Tamura et al., 2013](#)), using 5,000 bootstrap replications.

## Analysis of the H<sub>2</sub>O<sub>2</sub> and MDA contents

Leaf samples were ground under liquid nitrogen and homogenized in 0.9 mL of precooled extraction solution, containing 0.05 mol/L phosphate buffer solution (pH 7.8) and 1% polyvinylpyrrolidone. Then, tissue homogenates were chilled to 4 °C until used for further processes. H<sub>2</sub>O<sub>2</sub> concentrations were measured using the Hydrogen Peroxide Assay Kit (Beyotime Biotechnology Inc, Shanghai, China) following manufacturer's instructions. Absorbance was read at 560 nm using a microplate reader (PerkinElmer Inc., Waltham, USA). H<sub>2</sub>O<sub>2</sub> concentrations were calculated according to a standard concentration curve and expressed as nmol/ml. MDA contents were measured using a Plant Malondialdehyde assay kit (Nanjing Jiancheng Bioengineering Inst., Nanjing, China), which was based on the reaction of thiobarbituric acid (TBA) with MDA. The absorbance of red TBA-MDA complex was read at 532 nm using a microplate reader and the MDA content (nmol/g) was calculated according to manufacturer's instructions.

### Analysis of *ZmCAT* mRNA expression by quantitative real-time PCR (qRT-PCR)

Total RNA was extracted from leaf samples and assessed as described above. Raw RNA (0.5  $\mu\text{g}$ ) was digested using RNase-free DNase I and then reverse-transcribed to cDNA using an AMV First Strand cDNA Synthesis Kit (Sangon Biotech Inc., Shanghai, China). The resulting cDNA samples were analyzed and quantitative PCR was performed in a Bio-Rad iQ5<sup>TM</sup> Multicolor Real-time PCR Detection System (Bio-Rad Laboratories Inc., Hercules, CA, USA), according to manufacturer's instructions. The qPCR analysis was performed in a total liquid volume of 25.0  $\mu\text{L}$ , containing 1  $\times$  SYBR Green Master Mix, 4.0  $\mu\text{L}$  of the diluted cDNA templates ( $n = 5$ ), 0.4 mM of each primer and 7.5  $\mu\text{L}$  of DEPC-treated RNA-free water. A 140 bps product was amplified with *Zm-CAT*-qRT F and *ZmCAT*-qRT-R (Table S1) and sequenced to verify PCR specificity. The housekeeping gene eukaryotic initiation factor 4A (eIF4A) was used as an internal control to check the RT-qPCR reaction (Ransbotyn & Reusch, 2006). Primers of the internal control gene used for RT-qPCR were listed in Table S1 and produced a fragment of 125 bps. Conditions for the reactions were 95 °C for 3 min, followed by 40 cycles of 95 °C for 15 s, and then 60 °C for 40 s. A melting curve was generated for each sample at the end of the RT-qPCR reaction to confirm the purity of the amplified products. Data from RT-qPCR were analyzed using the  $2^{-\Delta\Delta C_t}$  method of the ABI 7500 software 2.0.6 (Applied Biosystems, Inc., Foster City, USA).

### Recombinant overexpression and purification of *ZmCAT*

The pEASY-E1 vector was used for high level, inducible expression of *ZmCAT* recombinant proteins, as has been reported by Liu *et al.* (2016). In brief, the cDNA fragment encoding the mature peptide of *ZmCAT* was obtained after PCR amplification with the primers, *ZmCAT*-recombinant-F and *ZmCAT*-recombinant-R (Table S1). The plasmid pEASY-E1/*ZmCAT* was isolated by E.Z.N.A.<sup>®</sup> Plasmid Mini Kit (Omega Bio-Tek Inc., Norcross, GA, USA) and then transformed into *E. coli* strain BL21 (DE3) to produce recombinant protein. The resulting strain was grown at 37 °C in liquid SOB medium coupled with ampicillin (100 mg mL<sup>-1</sup>) until the OD<sub>600</sub> reached 0.4–0.6. The bacteria were induced for protein expression by IPTG addition to 1 mM. After incubation for 4 h at 30 °C, the bacteria were harvested by centrifugation and the recombinant protein purified by MagExtractor<sup>TM</sup> His Tag Kit (Toyobo Co Ltd., Osaka, Japan) under denaturing conditions (8 mol L<sup>-1</sup> urea). The purified protein was next refolded in a gradient urea-TBS glycerol buffer and the resultant protein mixed 1/1 (wt/wt) with 2 $\times$  Coomassie blue loading dye. Each sample was loaded on a 12% SDS-PAGE gel and Precision Plus Protein<sup>TM</sup> standards (Bio-Rad Laboratories Inc., Hercules, CA, USA) added to provide molecular mass markers.

### Analysis of enzymatic activity of recombinant *ZmCAT*

Enzymatic activities of recombinant *ZmCAT* (r*ZmCAT*) were examined using a Catalase Assay Kit (Beyotime Biotechnology Inc., Jiangsu, China) as previously developed methods (Liu *et al.*, 2016). The purified r*ZmCAT* protein was quantified using an Enhanced BCA Protein Assay Kit (Beyotime Biotechnology Inc, Shanghai, China). The enzymatic activities

of rZmCAT protein was characterized by treating protein samples at 5 °C intervals between 0 and 40 °C for 24 h.

### Statistical analysis

The data of mRNA expression and enzymatic activities were normalized to 20 °C. All data were presented as the mean  $\pm$ SD ( $n = 5$ ). One-way analysis of variance (ANOVA) followed by a Student-Newman-Keuls (S-N-K) multiple comparison method was used to analyze the significance of differences in the SPSS 22.0 software and  $p$ -values of  $<0.05$  considered statistically significant differences.

## RESULTS

### Cloning and sequence analysis of full-length ZmCAT cDNA

The full-length cDNA of ZmCAT was obtained by the RACE technique and deposited into GenBank (accession number [KJ766310](#)). The complete cDNA of ZmCAT consisted 1,816 bp, including a 1,434 bp open reading frame (ORF), 73 bp 5'-untranslated region (UTR), and a 309 bp 3'-UTR ([Fig. 1](#)). The 3'-UTR contained a 16 bp poly(A)-tail but no putative polyadenylation signal (AATAAA). The ORF encoded a polypeptide of 477 amino acid residues with a calculated molecular mass of 55.3 kDa, and a theoretical pI point of 6.40. No signal peptide in the deduced amino acid sequence was revealed by SignalP program analysis. Localization prediction software suggested that ZmCAT would be localized to the chloroplast. PROSITE analysis showed that the ZmCAT amino acid sequence contained highly conserved motifs, which included the proximal active site signature (<sup>54</sup>FDRERIPERVVHARGAS<sup>70</sup>) and proximal heme-ligand signature sequences (<sup>344</sup>RIFSYADTQ<sup>352</sup>). In addition, the His<sup>65</sup> catalytic residue was found in the active site responsible for proper binding and reduction of peroxide ([Fig. 1](#)). The putative ZmCAT contained two potential N-glycosylation sites at N-28 (NNSS) and N-247 (NHSH). A CAT core domain at 18–398 (Thr<sup>18</sup> to Rrg<sup>401</sup>) and CAT immune-responsive domain (Pro<sup>420</sup> to Lys<sup>475</sup>) were also detected.

### Multiple sequence alignment and phylogenetic analysis of ZmCAT

A multiple alignment of ZmCAT was performed by selecting CAT sequences of nine representative plant species ([Fig. 2](#)). The deduced amino acid sequence of ZmCAT exhibited high similarity with other previously identified CAT, such as an 88% identity with CAT from *Ananas comosus*, 85% identity with *Zea mays*, 84% identity with CATs from *Zantedeschia aethiopica*, *Bruguiera gymnorhiza*, *Oryza sativa*, *Triticum aestivum*, *Saccharum arundinaceum* and *Festuca arundinacea* and 83% identity with *Arabidopsis thaliana*. The molecular evolutionary relationship of ZmCAT was examined by aligning sequences of known CAT members from different taxa and constructing a phylogenetic tree from the conserved regions by the NJ method. The NJ phylogenetic tree was positioned separately into three main branches and ZmCAT found to be clustered with CAT from *Zostera muelleri* and located in the angiospermous cluster sub-branch ([Fig. 3](#)).




```

1 aggggaagaagaacggagggttcgacgacgacagaagaacggatgaatcaagatagttggtgc
1      M D P T K F R P S S D F N S P Y
61 gatcgattcaactATGGATCCCACAAAATCCGTCGTCAGTATTCAACTCCTCTTA
17 W T T N S G A P V W N N N S S L T V G S
121 CTGGACTACAAATCTGGAGCTCCTGTTTGAACAACAACCTTCATTGACCGTTGGTTC
37 R G P I L L E D Y H L I E K I A Q F D R
181 TCGAGGACCAATACTCCTTGAGGACTATCATTGATTGAAAAGATTGCTCAGTTGATAG
57 E R I P E R V V H A R G A S A K G F F E
241 GGAGCGTATCCAGAGCGTGTGTTTCATGCAAGAGGAGCTAGTGCAAAAGGCTTTTTGA
77 V T H D I S H L T C A D F L R A P A V Q
301 GGCTACTCATGATATATCACATCTAACATGTGCTGATTCTCCGTCACCAGCAGTTCA
97 T P V I V R F S T V I H E R G S P E T L
361 AACTCCGGTTATTGTCGTTTTTCAACTGTCAATTCATGAGAGAGGAAGCCCTGAAACCTT
117 R D P R G F A V K F Y T R E G N F D L V
421 GAGAGATCCTCGAGGCTTGTGCTGAAGTTCTACACTCGAGAGGGAATTTTGATCTTGT
137 G N N F P V F F T R D G M K F P E K H H
481 GGGGAACAATTTCCAGTTTTTTTCCACCGTGACGGAATGAAGTTTCTGAAAAGCATCA
157 A K K T N P K T H I Q D N W R I M D F F
541 CGCCAAGAAAACAACCCCAAGACTCACATCCAAGATAATTGGAGAATTATGGATTCTT
177 S H H P E S L N M F T F L F D D L G N P
601 CTCACACCATCCAGAGAGTTGAACATGTTCACTTTTCTATTGATGATCTTGGAAATCC
197 L N Y R H M D G S G V N T Y T L I N K A
661 ACTGAATTATAGACACATGGATGGTTCTGGTGTCAATACTTATACCTAATTAACAAGC
217 G K A H Y V K F H W K P T C G V K C L L
721 TGGAAAAGCTCATTATGGAAGTCCATTGGAAGCCTACATGTGGAGTGAATGCTGTT
237 D D E A V T V G G S N H S H A T K D L Y
781 GGATGATGAAGCTGTTACAGTGGAGGTAGTAACCACAGCCATGCAACTAAGGATCTTTA
257 D S I D A G N Y P E W T L Y I Q T I D P
841 TGATTCCATAGATGCTGAAAATATCCAGAATGGACACTTACATTCAGACCATTGATCC
277 D F E D T F D F D P L D V T K T W P E D
901 TGATTTGAAGATACATTGATTTTCGATCCACTTGATGTAACAAAGACATGGCCGGAAGA
297 I F P L Q P V G R L V L N R N V D N F F
961 CATCTTCTCTCAACCTGTTGGACGCTGGTCTTGAACAGGAATGTAGATAATTTCTT
317 A E N E Q L A F C P A I I V P G I Y Y S
1021 TGCAGAGAATGAGCAGCTCGCTTCTGTCCAGCAATTATTGTGCCTGGGATCTATTACT
337 N D K L L Q T R I F S Y A D T Q R H R L
1081 AAATGATAAACTTCTCAGACTAGGATTTTCTCTTATGCCGACTCAACGGCATCGTCT
357 G P N Y L M L P V N A P K C A H H N N H
1141 TGGGCCAAACTATCTGATGCTCCAGTGAATGCTCCAAAATGTGCACATCACAACAATCA
377 Y D G S M N F M H R N E E V D Y F P S R
1201 TTATGATGGGTCCATGAATTTTATGCATAGAAATGAAGAGGTGGACTATTTCCCTTCAAG
397 F D P A R H A E Q V P I P N P V L S G K
1261 ATTTGATCCTGCTCGACATGCCGAACAAGTCCCGATACCAATCTGTTCTTTCAGGGAA
417 R E K P I I V K E N N F K Q P G E R Y R
1321 GCGGAAAAGCCATAATTGTGAAAGAGAACAACCTCAAGCAGCCAGGAGAAAGTACAG
437 S W E S D R Q E R Y I C R W I D A L S D
1381 GTCCTGGGAATCTGACAGACAAGAGCGTTACATCTGCCGCTGGATTGATGCACTATCTGA
457 S R V T P E I R S I W I S Y W S Q V K C
1441 TTCCAGAGTCACTCCTGAAATCCGTTTCGATATGGATCTCTTATGGTCTCAGGTGAAATG
477 F *
1501 CTTTTAActc caatgccagtgcaactctttategettaatetg gatctcttactgctattt
1561 ttttgtgtgtgttgctgcaggetgacatgtccctgggcccagaaacttaccagtcgtc
1621 tcaatacgaacccaatatgtaaaaaagatgaatctcattttgtggaatcaacttggtg
1681 aaagagaagcagcactatttattattcttaaatctattgctgtgcttcatagggtt
1741 gataaacataatgataaacattgatgaatcaataatagcaataaacattgctttcc
1801 aaaaaaaaaaaaaaa

```

**Figure 1** Nucleotide sequence and deduced amino acid sequences of *ZmCAT* catalase. The number of nucleotides and deduced amino acid residues are shown in the left margin. A catalase domain (Thr<sup>18</sup> to Rrg<sup>401</sup>) and a catalase-rel domain (Pro<sup>420</sup> to Lys<sup>475</sup>) are shaded. The asterisk indicated the stop codon. The lowercase letters represent nucleotide sequences, and the capital letters indicate amino acid sequences.

Full-size  DOI: [10.7717/peerj.4532/fig-1](https://doi.org/10.7717/peerj.4532/fig-1)




**Figure 2** Multiple alignment of deduced amino acid sequences of *ZmCAT* with selected sequences from other plant species. *Ananas comosus* (OAY65449), *Zea mays* (NP\_001241808), *Zantedeschia aethiopica* (AAF19965), *Bruguiera gymnorhiza* (ADC95629), *Oryza sativa* (BAA34205) *Triticum aestivum* (AIZ77475), *Saccharum arundinaceum* (AIU99479), *Festuca arundinacea* (CAG23920), *Arabidopsis thaliana* (AAK96854). The black shaded regions represent identical amino acids, while the gray regions represent similar amino acids.

Full-size DOI: 10.7717/peerj.4532/fig-2

## Effects of temperature stress on the content of H<sub>2</sub>O<sub>2</sub> and MDA

The ROS levels and membrane injury in *Z. marina* under different temperature treatments (5, 10, 15, 20, 25, and 30 °C) after 96 h was investigated in an experiment with different H<sub>2</sub>O<sub>2</sub> and MDA contents. When the temperature decreased from 15 to 5 °C and increased from 20 to 30 °C, the H<sub>2</sub>O<sub>2</sub> content increased significantly (Fig. 4A). The greatest H<sub>2</sub>O<sub>2</sub> content increases occurred in the 5 treatment (8.20 ± 1.54 nmol/mL) with 3.1-fold higher than 20 °C (*p* < 0.05), while the H<sub>2</sub>O<sub>2</sub> content exhibited no significant change and kept






**Figure 3** Phylogenetic analysis based on amino acid sequences of *ZmCAT* and other known CATs. Numbers above or below branches at tree nodes indicate percent bootstrap values after 5,000 replicates. The GenBank accession numbers are displayed next to each sequence. Accession NO.Z: The protein sequences of CAT in *Zosteramuelleri* was download from [http://appliedbioinformatics.com.au/index.php/Seagrass\\_Zmu\\_Genome](http://appliedbioinformatics.com.au/index.php/Seagrass_Zmu_Genome).

Full-size DOI: 10.7717/peerj.4532/fig-3

at the temperature ranged from 15 to 20 °C. In response to temperature treatments, there were significantly increased MDA contents at the temperature decreased from 15 to 5 °C and increased from 15 °C to 30 °C (Fig. 4B). The highest MDA content occurred in the 30 °C treatments ( $69.74 \pm 10.32$  nmol/g) with 2.2-fold higher than 15 °C ( $p < 0.05$ ).


### Effects of temperature stress on *ZmCAT* mRNA expression

RNA levels of CAT gene were quantified using RT-qPCR technique of reverse transcripts of RNA from *Z. marina* that had been subjected to various temperature treatments for 96 h (Fig. 5). The data were then standardized relative to the RNA expression levels of eIF4A, a housekeeping gene consistently expressed in plants. Although relative mRNA expression levels were detected in all treatments, the CAT gene was significantly upregulated from 5 to 15 °C, with the highest expression at 10 °C. In contrast, this gene revealed rather low expression in heat-stressed conditions (25 and 30 °C); and *ZmCAT* expression at 10 °C was 13-fold higher than at 30 °C ( $p < 0.05$ ). Hence, temperature stress suppressed *ZmCAT* expression, but *ZmCAT* was expressed at lower values with elevated heat stress than with cold stress.



**Figure 4** The effects of temperature stress on H<sub>2</sub>O<sub>2</sub> and MDA contents in *Z. marina* during a 96 h experiment. (A) H<sub>2</sub>O<sub>2</sub> content. (B) MDA content. Values shown as means  $\pm$  SD ( $n = 5$ ), bars marked with dissimilar letters represent significant differences ( $p < 0.05$ ) from each other according to S-N-K multiple comparisons tests.

Full-size  DOI: [10.7717/peerj.4532/fig-4](https://doi.org/10.7717/peerj.4532/fig-4)




**Figure 5** The effects of temperature stress on the relative mRNA expression levels of *ZmCAT*. Amplification of *eIF4A* gene was used as an internal control. The *ZmCAT* expression levels at 20 °C was set to 100%. Values shown as means  $\pm$  SD ( $n = 5$ ), bars marked with dissimilar letters represent significant differences ( $p < 0.05$ ) from each other according to S-N-K multiple comparisons tests.

Full-size DOI: 10.7717/peerj.4532/fig-5


### Effects of temperature on recombinant *ZmCAT* activity

The recombinant protein activity of *ZmCAT* was determined to examine whether it was functionally active in the antioxidant response under temperature stress. Highly purified *ZmCAT* proteins for biochemical characterization under temperature stress were obtained using *E. coli* BL21 (DE3) as a heterologous expression system to express *ZmCAT* protein and demonstrate *ZmCAT*'s antioxidant activity, based on  $H_2O_2$  consumption. Recombinant *ZmCAT* (*rZmCAT*) protein was analyzed by SDS-PAGE, appearing as a distinct band at a molecular weight of  $\sim 55$  kDa, which was close to the calculated molecular mass of *ZmCAT* (55.3 kDa, Fig. 6). The enzymatic activity of recombinant protein was  $19,310.5 \pm 83.5$  U  $mg^{-1}$  at 20 °C. The stability of *rZmCAT* was investigated by measuring the enzymatic activities of its recombinant protein under different temperature treatments (0–40 °C). The enzymatic activity increased rapidly from 0 to 15 °C and remained at a high level from 15 to 25 °C, with  $>95\%$  of the relative enzymatic activity retained in the temperature range of 15–25 °C. After reaching peak activity at 25 °C the *rZmCAT* gradually lost enzymatic activity and almost devitalized at 40 °C. It was noteworthy that, although the relative activity was clearly reduced from 25 to 40 °C,  $>20\%$  of its activity was retained after incubation at 35 °C for 96 h (Fig. 7).



**Figure 6** SDS-PAGE analysis of recombinant *Zm. CAT* protein expressed in *E. coli* BL21 (DE3). Samples analyzed included low molecular weight protein markers (M, lane 1), un-induced expression of pEASY-E1/*ZmCAT* in total BL21 (DE3) cell lysates (U, lane 2), 1 mM IPTG induced expression of pEASY-E1/*ZmCAT* in BL21 (DE3) (I, lane 3), and the purified recombinant protein of *ZmCAT* (P, lane 4).

Full-size  DOI: [10.7717/peerj.4532/fig-6](https://doi.org/10.7717/peerj.4532/fig-6)



**Figure 7** The effects of temperature stress on the enzymatic activity of recombinant *ZmCAT* protein. The enzymatic activity at 20 °C was set to 100%. Each values is shown as mean  $\pm$  S.D. ( $n = 5$ ), and bars with different characters were significantly different ( $p < 0.05$ ).

Full-size DOI: 10.7717/peerj.4532/fig-7

## DISCUSSION

Catalase plays an important role in the antioxidant defense system by degrading the  $H_2O_2$  to  $H_2O$  and  $O_2$  (Unwin, 1975). In recent years, several CAT genes have been identified in different studies (Du, Zhou & Huang, 2013; Lin, Huang & Lin, 2010), but the present study was the first to deal with CAT cloned and characterized from eelgrass. *ZmCAT* was found to possess most of the main characteristic amino acid residues, motifs, and elements of the CAT protein family. The sequence and multiple alignment results indicated that the active site signature ( $^{54}FDRERIPERVVHARGAS^{70}$ ) and heme-ligand signature motifs ( $^{344}RIFSYADTQ^{352}$ ) were conserved in *ZmCAT* (Fig. 1). Multiple alignments revealed high sequence identity with CATs from other known plant species, ranging from 83 to 88% (Fig. 2). Phylogenetic relationship collectively suggested that *ZmCAT* was grouped within the angiosperms and might have similar functions to CATs from other plant species (Fig. 3). All present results showed that *ZmCAT* was a representative member of the plant CAT family.

Temperature stress usually causes injury such as excessive accumulation of ROS in plant cells, which seriously damage cell membranes, protein, and nucleic acids (Gill & Tuteja, 2010; Vighi et al., 2016). Low temperature stress can induce destructive oxidative processes leading to an increase of MDA content in tissue, which could reflect the degree



of LPO and structural integrity of plant membranes (Li et al., 2013). Therefore, redox states and membrane integrity responses of *Z. marina* exposed to temperature stress were identified by investigating the effects of such stress on H<sub>2</sub>O<sub>2</sub> and MDA contents. In response to temperature stress, H<sub>2</sub>O<sub>2</sub> and MDA contents in *Z. marina* significantly increased at cold stress (<10 °C) and heat stress (>25 °C) and the highest H<sub>2</sub>O<sub>2</sub> and MDA content respectively occurred at 5 and 30 °C (Fig. 4). The observed increased H<sub>2</sub>O<sub>2</sub> and MDA contents were indications that *Z. marina* was unable to suppress oxidative stress under extreme temperature stress, which might influenced its physiological functions and survival.

The expression profile of CAT was a key part of the plant defense system against oxidative stress (Bi et al., 2017). In the present study, *ZmCAT* appeared to be regulated at the transcriptional level for detoxifying increased H<sub>2</sub>O<sub>2</sub> during cold stress conditions. The optimal temperature for *Z. marina* growth has been reported to be 15.3 ± 1.6 °C (Lee, Sang & Kim, 2007), while the expression concentration increased rapidly from 5 to 10 °C and the highest *ZmCAT* gene expression observed at 10 °C (Fig. 4). Cold stress can lead to excess H<sub>2</sub>O<sub>2</sub> accumulation and increased H<sub>2</sub>O<sub>2</sub> content might act as signals for inducing antioxidant gene expressions. Therefore, it was deduced that the *ZmCAT* gene was induced and upregulated to reduce oxidative damage under lower temperature stress, but sustained extreme low temperature stress (5 °C) damages tissue and leads to decreased *ZmCAT* expression and increased H<sub>2</sub>O<sub>2</sub> content and membrane injury. Similar results have been obtained by Baek & Skinner (2003), who found that CAT gene expression was significantly higher in winter wheat after cold-stress exposure. However, *ZmCAT* gene expression was suppressed when the temperature increased from 25 to 30 °C, which showed the inhibiting effects of heat stress conditions on *ZmCAT* gene expression that, in turn, might have resulted in intracellular H<sub>2</sub>O<sub>2</sub> accumulation.

Information on the recombinant enzyme activity of the *ZmCAT* after temperature treatments was helpful for a better understanding of its physiological function. The relative enzymatic activity retained with >95% by the r*ZmCAT* at 15–25 °C. Field surveys and experiments have confirmed that high water temperatures >25 °C in summer heat waves increasingly threatens seagrass performance and >30 °C can be lethal (Ehlers, Worm & Reusch, 2008; Kaldy, 2014; Nejrup & Pedersen, 2008). However, the highest H<sub>2</sub>O<sub>2</sub> removal activity by r*ZmCAT* was observed at 25 °C. Moreover, a relative activity of 55% was retained by r*ZmCAT* at 30 °C (Fig. 6). On the basis of its strong antioxidant activity at high temperatures, *ZmCAT* was suggested here to play a role against heat stress, as a potent ROS-detoxifying enzyme in eelgrass. Several CATs have been demonstrated to have high thermal stability, including *Pyropia yezoensis* CAT (*Py* CAT), *Oryza sativa* CAT-A (*Os* CatA), and CAT-C (*Os* CatC), with reported optimal temperatures of ~30 °C (Li et al., 2012; Vighi et al., 2016). In addition, r*ZmCAT* exhibited higher heat sensitivity than that reported for *Py* CAT. The r*ZmCAT* activity declined rapidly as temperature exceeded 25 °C and <10% activity was retained at 40 °C, similar to that reported for *Os*CatC. A potential reason for this difference might have been because CAT in *Z. marina* and *P. yezoensis* must cope with different living environments, with *P. yezoensis* living in the upper intertidal zone with high and variable temperature, while *Z. marina* is a subtidal population and

in a more stable environment. Thus, a more stable enzyme is generated in *P. yezoensis* as an adaptation to harsh living conditions. It should be noted that the increased H<sub>2</sub>O<sub>2</sub> and MDA content at extreme temperature stress (5 °C and 30 °C) might have resulted from both *ZmCAT* gene expression and enzymatic activity, which were significantly inhibited. Interestingly, *ZmCAT* expression characterization was different from enzyme activity under heat stress. *ZmCAT* mRNA expression was downregulated after 25 and 30 °C temperature treatments for 96 h, while the enzymatic activity first increased and then declined. *ZmCAT* appeared to respond more sensitively at the transcriptional level than at the enzymatic level. These results indicated that *ZmCAT* in eelgrass survived under heat stress through elevated enzymatic activity rather than upregulated gene transcription to protect cellular components against ROS effects produced as a consequence of oxidative stress. This information might be valuable for further analysis of the profile and functional diversity of CAT.

## CONCLUSIONS

In this study, a novel CAT gene, named as *ZmCAT*, was identified in *Z. marina* and analysis of its deduced amino acid sequence showed typical CAT functional domains, including the CAT proximal heme-ligand signature and CAT proximal activity site. Multiple sequence alignment as well as phylogenetic analysis provided evidence for the evolutionary conservation of *ZmCAT*, which was a typical member of the plant CAT family. Temperature stress increased oxidative stress and membrane injury in *Z. marina*. To counter this stress, *ZmCAT* presented an antioxidant defense and protection by high enzymatic activity under heat stress and by upregulated gene transcription under cold stress. However, both *ZmCAT* gene expression and enzymatic activity were significantly suppressed at extreme temperatures (5 and 30 °C) which indicated that the ability of *ZmCAT* to scavenge H<sub>2</sub>O<sub>2</sub> was weakened and led to changes in the cellular redox state and influenced its physiological function in eelgrass. Gene expression and enzymatic activity of *ZmCAT* varied under different temperature stresses, which suggested that *ZmCAT* might have played a central role in mitigating oxidative damage under temperature stress.

## ACKNOWLEDGEMENTS

We are greatly appreciative of the laboratory members who prepared experimental materials and provided technical assistance. We also thank the three expert reviewers for their enlightening suggestions, which substantially improved the final version of this article.

## ADDITIONAL INFORMATION AND DECLARATIONS

### Funding

This work is financially supported by the National Natural Science Foundation of China (No. 41476091) and Key Special Program on “Marine Environmental Security Protection” of National Research and Development Project of China (No. 2016YFC1402305). The

fundere had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

### **Grant Disclosures**

The following grant information was disclosed by the authors:

National Natural Science Foundation of China: 41476091.

National Research and Development Project of China: 2016YFC1402305.

### **Competing Interests**

The authors declare there are no competing interests.

### **Author Contributions**

- Yu Zang conceived and designed the experiments, performed the experiments, analyzed the data, contributed reagents/materials/analysis tools, prepared figures and/or tables, authored or reviewed drafts of the paper, approved the final draft.
- Jiao Liu performed the experiments, analyzed the data, contributed reagents/materials/analysis tools, prepared figures and/or tables, authored or reviewed drafts of the paper, approved the final draft.
- Xue Xi Tang analyzed the data, contributed reagents/materials/analysis tools, authored or reviewed drafts of the paper, approved the final draft.
- Bin Zhou conceived and designed the experiments, analyzed the data, contributed reagents/materials/analysis tools, authored or reviewed drafts of the paper, approved the final draft.

### **Field Study Permissions**

The following information was supplied relating to field study approvals (i.e., approving body and any reference numbers):

The research permit was issued to the College of Marine Life Science, Ocean University of China.

### **DNA Deposition**

The following information was supplied regarding the deposition of DNA sequences.

The catalase gene sequences of *Zostera marina* described here are accessible via GenBank accession number [KJ766310](#).

### **Data Availability**

The following information was supplied regarding data availability:

NCBI databases.

GenBank: [KJ766310](#).

### **Supplemental Information**

Supplemental information for this article can be found online at <http://dx.doi.org/10.7717/peerj.4532#supplemental-information>.

## REFERENCES

- Baek KH, Skinner DZ. 2003. Alteration of antioxidant enzyme gene expression during cold acclimation of near-isogenic wheat lines. *Plant Science* 165:1221–1227 DOI 10.1016/S0168-9452(03)00329-7.
- Bi C, Ma Y, Wu Z, Yu YT, Liang S, Lu K, Wang XF. 2017. Arabidopsis ABI5 plays a role in regulating ROS homeostasis by activating CATALASE 1 transcription in seed germination. *Plant Molecular Biology* 94:197–213 DOI 10.1007/s11103-017-0603-y.
- Bulthuis DA. 1987. Effects of temperature on photosynthesis and growth of seagrasses. *Aquatic Botany* 27:27–40 DOI 10.1016/0304-3770(87)90084-2.
- Cansev A, Gulen H, Eris A. 2011. The activities of catalase and ascorbate peroxidase in olive (*Olea europaea* L. cv. Gemlik) under low temperature stress. *Horticulture Environment & Biotechnology* 52:113–120 DOI 10.1007/s13580-011-0126-4.
- Du H, Zhou P, Huang B. 2013. Antioxidant enzymatic activities and gene expression associated with heat tolerance in a cool-season perennial grass species. *Environmental & Experimental Botany* 87:159–166 DOI 10.1016/j.envexpbot.2012.09.009.
- Ehlers A, Worm B, Reusch TBH. 2008. Importance of genetic diversity in eelgrass *Zostera marina* for its resilience to global warming. *Marine Ecology Progress* 355:1–7 DOI 10.3354/meps07369.
- Erfteimeijer PLA, Robin Lewis III RR. 2006. Environmental impacts of dredging on seagrasses: a review. *Marine Pollution Bulletin* 52:1553–1572 DOI 10.1016/j.marpolbul.2006.09.006.
- Figuroa-Yáñez L, Cano-Sosa J, Castaño E, Arroyo-Herrera AL, Caamal-Velazquez JH, Sanchez-Teyer F, López-Gómez R, Santos-Briones CDL, Rodríguez-Zapata L. 2012. Phylogenetic relationships and expression in response to low temperature of a catalase gene in banana (*Musa acuminata* cv. “Grand Nain”) fruit. *Plant Cell Tissue & Organ Culture* 109:429–438 DOI 10.1007/s11240-011-0107-4.
- Fourqurean JW, Duarte CM, Kennedy H, Marbá N, Agrave N, Holmer M, Mateo MA, Apostolaki ET, Kendrick GA, Krause-Jensen D, Mcglathery KJ, Serrano O. 2012. Seagrass ecosystems as a globally significant carbon stock. *Nature Geoscience* 1:297–315 DOI 10.1038/ngeo1477.
- Franssen SU, Gu J, Winters G, Huylmans AK, Wienpahl I, Sparwel M, Coyer JA, Olsen JL, Reusch TB, Bornberg-Bauer E. 2014. Genome-wide transcriptional responses of the seagrasses *Zostera marina* and *Nanozostera noltii* under a simulated heatwave confirm functional types. *Marine Genomics* 15:65–73 DOI 10.1016/j.margen.2014.03.004.
- Gao JJ, Qin AG, Yu XC. 2009. Effects of grafting on cucumber leaf SOD and CAT gene expression and activities under low temperature stress. *Chinese Journal of Applied Ecology* 20:213–217.
- Gill SS, Tuteja N. 2010. Reactive oxygen species and antioxidant machinery in abiotic stress tolerance in crop plants. *Plant Physiology & Biochemistry* 48:909–930 DOI 10.1016/j.plaphy.2010.08.016.

- Gu J, Weber K, Klemp E, Winters G, Franssen SU, Wienpahl I, Huylmans AK, Zecher K, Reusch TB, Bornberg-Bauer E. 2012. Identifying core features of adaptive metabolic mechanisms for chronic heat stress attenuation contributing to systems robustness. *Integrative Biology Quantitative Biosciences from Nano To Macro* 4:480 DOI 10.1039/c2ib00109h.
- Hasanuzzaman M, Nahar K, Fujita M. 2013. Extreme temperature responses, oxidative stress and antioxidant defense in plants. *Journal of the National Cancer Institute* 65:81–93 DOI 10.5772/54833.
- Hemminga M, Duarte C. 2000. *Seagrass ecology*. Cambridge: Cambridge University Press.
- Jena K, Kar PK, Kausar Z, Babu CS. 2013. Effects of temperature on modulation of oxidative stress and antioxidant defenses in testes of tropical tasar silkworm *Antheraea mylitta*. *Journal of Thermal Biology* 38:199–204 DOI 10.1016/j.jtherbio.2013.02.008.
- Kaldy JE. 2014. Effect of temperature and nutrient manipulations on eelgrass *Zostera marina* L. from the Pacific Northwest, USA. *Journal of Experimental Marine Biology & Ecology* 453:108–115 DOI 10.1016/j.jembe.2013.12.020.
- Kashiwagi A, Kashiwagi K, Takase M, Hanada H, Nakamura M. 1997. Comparison of catalase in diploid and haploid *Rana rugosa* using heat and chemical inactivation techniques. *Comparative Biochemistry & Physiology Part B Biochemistry & Molecular Biology* 118:499–503 DOI 10.1016/S0305-0491(97)00216-2.
- Larkin MA, Blackshields G, Brown NP, Chenna R, Mcgettigan PA, Mcwilliam H, Valentin F, Wallace IM, Wilm A, Lopez R, Thompson JD, Gibson TJ, Higgins DG. 2007. Clustal W and Clustal X version 2.0. *Bioinformatics* 23:2947–2948 DOI 10.1093/bioinformatics/btm404.
- Lee KS, Sang RP, Kim YK. 2007. Effects of irradiance, temperature, and nutrients on growth dynamics of seagrasses: a review. *Journal of Experimental Marine Biology & Ecology* 350:144–175 DOI 10.1016/j.jembe.2007.06.016.
- Letunic I, Doerks T, Bork P. 2012. SMART 7: recent updates to the protein domain annotation resource. *Nucleic Acids Research* 40:302–305 DOI 10.1093/nar/gkr931.
- Li HX, Xiao Y, Cao LL, Yan X, Li C, Shi HY, Wang JW, Ye YH. 2013. Cerebroside C increases tolerance to chilling injury and alters lipid composition in wheat roots. *PLOS ONE* 8:e73380 DOI 10.1371/journal.pone.0073380.
- Li XC, Xing YZ, Jiang X, Qiao J, Tan HL, Tian Y, Zhou B. 2012. identification and characterization of the catalase gene pycat from the red alga *pyropia yezoensis* (Bangiales, Rhodophyta)1. *Journal of Phycology* 48:664–669 DOI 10.1111/j.1529-8817.2012.01152.x.
- Lin KH, Huang HC, Lin CY. 2010. Cloning, expression and physiological analysis of broccoli catalase gene and Chinese cabbage ascorbate peroxidase gene under heat stress. *Plant Cell Reports* 29:575–593 DOI 10.1007/s00299-010-0846-4.
- Liu J, Tang X, Wang Y, Zang Y, Zhou B. 2016. A *Zostera marina* manganese superoxide dismutase gene involved in the responses to temperature stress. *Gene* 575:718–724 DOI 10.1016/j.gene.2015.09.050.



- Lin H, Sun T, Zhou Y, Zhang X. 2016.** Anti-oxidative feedback and biomarkers in the intertidal seagrass *Zostera japonica* induced by exposure to copper, lead and cadmium. *Marine pollution bulletin* **109**:325–333 DOI [10.1016/j.marpolbul.2016.05.062](https://doi.org/10.1016/j.marpolbul.2016.05.062).
- Marin-Guirao L, Entrambasaguas L, Dattolo E, Ruiz JM, Procaccini G. 2017.** Molecular mechanisms behind the physiological resistance to intense transient warming in an iconic marine plant. *Frontiers in Plant Science* **8**:Article 1142 DOI [10.3389/fpls.2017.01142](https://doi.org/10.3389/fpls.2017.01142).
- Matés JM. 2000.** Effects of antioxidant enzymes in the molecular control of reactive oxygen species toxicology. *Toxicology* **153**:83–104 DOI [10.1016/S0300-483X\(00\)00306-1](https://doi.org/10.1016/S0300-483X(00)00306-1).
- Mittler R. 2002.** Oxidative stress, antioxidants and stress tolerance. *Trends in Plant Science* **7**:405–410 DOI [10.1016/S1360-1385\(02\)02312-9](https://doi.org/10.1016/S1360-1385(02)02312-9).
- Moore KA, Shields EC, Parrish DB. 2014.** Impacts of varying estuarine temperature and light conditions on *Zostera marina* (Eelgrass) and its interactions with *Ruppia maritima* (Widgeongrass). *Estuaries & Coasts* **37**:20–30 DOI [10.1007/s12237-013-9667-3](https://doi.org/10.1007/s12237-013-9667-3).
- Nejrup LB, Pedersen MF. 2008.** Effects of salinity and water temperature on the ecological performance of *Zostera marina*. *Aquatic Botany* **88**:239–246 DOI [10.1016/j.aquabot.2007.10.006](https://doi.org/10.1016/j.aquabot.2007.10.006).
- Orth RJ, Carruthers TJB, Dennison WC, Duarte CM, Fourqurean JW, Heck KL, Hughes AR, Kendrick GA, Kenworthy WJ, Olyarnik S, Short FT, Waycott M, Williams SL. 2006.** A global crisis for seagrass ecosystems. *Bioscience* **56**:987–996 DOI [10.1641/0006-3568\(2006\)56\[987:AGCFSE\]2.0.CO;2](https://doi.org/10.1641/0006-3568(2006)56[987:AGCFSE]2.0.CO;2).
- Orth RJ, Marion SR, Moore KA, Wilcox DJ. 2010.** Eelgrass (*Zostera marina* L.) in the Chesapeake Bay Region of Mid-Atlantic Coast of the USA: challenges in conservation and restoration. *Estuaries & Coasts* **33**:139–150 DOI [10.1007/s12237-009-9234-0](https://doi.org/10.1007/s12237-009-9234-0).
- Potters G, Pasternak TP, Guisez Y, Palme KJ, Jansen MA. 2007.** Stress-induced morphogenic responses: growing out of trouble? *Trends in Plant Science* **12**:98–105 DOI [10.1016/j.tplants.2007.01.004](https://doi.org/10.1016/j.tplants.2007.01.004).
- Ralph PJ. 1998.** Photosynthetic response of laboratory-cultured *Halophila ovalis* to thermal stress. *Marine Ecology Progress* **171**:123–130 DOI [10.3354/meps171123](https://doi.org/10.3354/meps171123).
- Ransbotyn V, Reusch TBH. 2006.** Housekeeping gene selection for quantitative real-time PCR assays in the seagrass *Zostera marina* subjected to heat stress. *Limnology & Oceanography Methods* **4**:367–373 DOI [10.4319/lom.2006.4.367](https://doi.org/10.4319/lom.2006.4.367).
- Scandalios JG, Acevedo A, Ruzsa S. 2000.** Catalase gene expression in response to chronic high temperature stress in maize. *Plant Science* **156**:103–110 DOI [10.1016/S0168-9452\(00\)00235-1](https://doi.org/10.1016/S0168-9452(00)00235-1).
- Short FT, Polidoro B, Livingstone SR, Carpenter KE, Bandeira S, Bujang JS, Calumpong HP, Carruthers TJB, Coles RG, Dennison WC, Erftemeijer PLA, Fortes MD, Freeman AS, Japtap TG, Kamal AHM, Kendrick GA, Kenworthy WJ, La Nafie YA, Nasution IM, Orth RJ, Prathep A, Sanciangco JC, Van Tussenbroek B, Vergara SG, Waycott M, Zieman JC. 2011.** Extinction risk assessment of the world's seagrass species. *Biological Conservation* **144**:1961–1971 DOI [10.1016/j.biocon.2011.04.010](https://doi.org/10.1016/j.biocon.2011.04.010).

- Srivastava S, Pathak AD, Gupta PS, Shrivastava AK, Srivastava AK. 2012.** Hydrogen peroxide-scavenging enzymes impart tolerance to high temperature induced oxidative stress in sugarcane. *Journal of Environmental Biology* **33**:657–661.
- Sureda A, Box A, Terrados J, Deudero S, Pons A. 2008.** Antioxidant response of the seagrass *Posidonia oceanica* when epiphytized by the invasive macroalgae *Lophocladia lallemandii*. *Marine Environmental Research* **66**:359–363  
DOI [10.1016/j.marenvres.2008.05.009](https://doi.org/10.1016/j.marenvres.2008.05.009).
- Tamura K, Stecher G, Peterson D, Filipski A, Kumar S. 2013.** MEGA6: molecular evolutionary genetics analysis version 6.0. *Molecular biology and evolution* **30**:2725–2729  
DOI [10.1093/molbev/mst197](https://doi.org/10.1093/molbev/mst197).
- Thomson JA, Burkholder DA, Heithaus MR, Fourqurean JW, Fraser MW, Statton J, Kendrick GA. 2015.** Extreme temperatures, foundation species, and abrupt ecosystem change: an example from an iconic seagrass ecosystem. *Global Change Biology* **21**:1463–1474 DOI [10.1111/gcb.12694](https://doi.org/10.1111/gcb.12694).
- Unwin PN. 1975.** Beef liver catalase structure: interpretation of electron micrographs. *Journal of Molecular Biology* **98**:235–242 DOI [10.1016/S0022-2836\(75\)80111-2](https://doi.org/10.1016/S0022-2836(75)80111-2).
- Vighi IL, Benitez LC, do Amaral MN, Auler PA, Moraes GP, Rodrigues GS, Da ML, Pinto LS, Braga EJ. 2016.** Changes in gene expression and catalase activity in *Oryza sativa* L. under abiotic stress. *Genetics & Molecular Research* **15**  
DOI [10.4238/gmr15048977](https://doi.org/10.4238/gmr15048977).
- Wang YC, Qu GZ, Li HY, Wu YJ, Chao W, Liu GF, Yang CP. 2010.** Enhanced salt tolerance of transgenic poplar plants expressing a manganese superoxide dismutase from *Tamarix androssowii*. *Molecular Biology Reports* **37**:1119–1124  
DOI [10.1007/s11033-009-9884-9](https://doi.org/10.1007/s11033-009-9884-9).