



A 9 cm robotic thymectomy and pericardial repair case report

Karishma Kodia, Dao M. Nguyen, Nestor R. Villamizar

Section of Thoracic Surgery, Department of Surgery, Miller School of Medicine, University of Miami, Miami, FL, USA

Correspondence to: Nestor Villamizar, MD. 1295 NW 14th Street Suite J 20, Miami, FL 33125 21, USA. Email: nvillamizar@med.miami.edu.

Abstract: In the modern surgical era, improved technology has allowed for the increasing use of the robotic platform for thymoma resection. Historically, tumors >5 cm were deemed inappropriate for minimally invasive approaches; thoracic surgeons, however, have become adept with performing increasingly complex thymectomies using minimally invasive techniques. Excision of large thymomas using the robotic platform is no longer considered a rare event, however few publications have described the use of minimally invasive surgery for *en bloc* excision of the pericardium with mesh reconstruction. We present a case of an asymptomatic, incidentally discovered 9 cm thymoma involving the pericardium and right lung upper lobe that was resected via bilateral robotic-assisted thymectomy *en bloc* with wedge resection and pericardial resection with mesh reconstruction. The case highlights the use of the robotic platform to avoid a conversion to open thymectomy. The patient was discharged home on postoperative day 3 with minimal pain and narcotic requirement. We aim to contribute to the existing literature supporting the use of the robotic platform during complex thymectomy. The associated video presentation serves as a visual instructional guide for the thymoma resection, *en bloc* with the right upper lobe and pericardium and the pericardial reconstruction. This minimally invasive technique has been associated with shorter hospital stay, reduced pain and faster recovery.

Keywords: Robotic thymectomy; pericardial resection; pericardial reconstruction; thymoma; case report

Received: 22 July 2020; Accepted: 09 September 2020; Published: 30 December 2020.

doi: 10.21037/med-20-52

View this article at: <http://dx.doi.org/10.21037/med-20-52>

Introduction

Over the past fifteen years, there has been an increasing use of video-assisted thoracoscopic surgery (VATS) and robotic-assisted thoracoscopic surgery (RATS) in the surgical treatment of thymomas (1). Some surgeons consider size over 5 cm as a contraindication for VATS thymectomy (2). However, some literature suggests that adequate oncologic resection can be achieved with minimally invasive surgery for tumor sizes larger than 3–5 cm without increasing complications (intraoperative capsular disruption) in experienced centers (3–5). Several previous publications have reported the excision of large thymomas using the robotic platform, however only a few publications have described the use of minimally invasive surgery for *en bloc* excision of the pericardium with mesh reconstruction. We present here a case report and associated video presentation of a robotic approach to a nearly 9 cm thymoma involving

the pericardium and the right lung upper lobe, in accordance with the CARE reporting checklist (6) (available at <http://dx.doi.org/10.21037/med-20-52>).

Case presentation

The patient is a 58-year-old female with an incidental finding of an anterior mediastinal mass during a calcium score chest computed tomography (CT) scan. She did not have any symptoms associated with myasthenia gravis preoperatively, however she had positive acetylcholine receptor (AR) antibodies [AR modulating: 44 (elevated); AR blocking: <15 (normal); AR binding 1.91 (elevated)]. Her chest CT scan reported an anterior mediastinal mass measuring 3.9 cm × 5.2 cm × 8.2 cm, in the anteroposterior, transverse, and cranio-caudal dimensions respectively (Figures 1–3).




Figure 1 CT Chest (axial) of patient's presenting CT Chest scan prompting initial clinic visit.




Figure 2 CT Chest (coronal) of patient's initial CT Chest scan.



Figure 3 CT Chest (sagittal) of patient's original CT Chest scan.

The patient underwent a bilateral robotic-assisted thoracoscopic thymectomy, right lung wedge resection and pericardial resection (*en bloc* with tumor), and reconstruction with pericardial membrane (Gor-Tex Preclude Pericardial Membrane, Gore Medical, Newark, Delaware) in March 2019.

Robotic trocar port sites in the left pleural cavity were placed as follows: 8 mm camera port in the 5th intercostal space (ICS) of the anterior axillary line; 8 mm port in the 3rd ICS between the midclavicular and anterior axillary line; 8 mm port in the 6th ICS between the midclavicular and anterior axillary line; and 15 mm assistant port in the 8th ICS at the anterior axillary line superior to the diaphragm. The left phrenic nerve was identified and protected. Thymic and mediastinal adipose tissue were dissected off of the aortopulmonary window, innominate vein and pericardium; thymic veins were divided with the vessel sealer. The left superior horn of the thymus was dissected and a mediastinal lymph node station 6 was excised. A left chest tube was placed and the rest of the case was performed via a right-side approach.

The right lung was attached to the anterior aspect of the mediastinal mass and was separated with green loads of the robotic stapler. A bipolar dissector was used to incise the anterior mediastinal pleura medial to the right internal mammary vessels and division was extended to reach the left internal mammary vessels. Thymic tissue was subsequently dissected off the innominate vein and the draining thymic veins were divided with the vessel sealer. The tumor was then resected *en bloc* with a segment of pericardium, which was firmly adherent to the tumor. The specimen was removed using an Anchor bag (Anchor Products Company, Addison, Illinois) with the capsule intact. Upon further exploration, there appeared to be no further thymic tissue remaining. The pericardial defect was reconstructed using a Gortex preclude pericardial membrane sutured with non-absorbable 0 V-loc suture; the membrane was fenestrated with robotic scissors to prevent tamponade in case of unrecognized bleeding. A right chest tube was placed. *Video 1* highlights key portions of the case including the thymic and pericardial resection and pericardial reconstruction.

The pathology report indicated a WHO type B2 9.0 cm thymoma with negative resection margins; all 6 mediastinal lymph nodes removed were negative for malignancy. The patient did well post operatively and had an uneventful course. The patient was discharged on post-op day 3 in stable condition. Her CT scan 6- and 12-month post-operatively did not demonstrate any evidence of recurrence. The patient developed moderate diplopia 5 months postop and was treated with pyridostigmine (30–60 mg every 4 hours) for myasthenia gravis by a neurologist at our institution; she is currently asymptomatic, and she has

weaned off and stopped her pyridostigmine. All procedures performed in studies involving human participants were in accordance with the ethical standards of the institutional and/or national research committee(s) and with the Helsinki Declaration (as revised in 2013). Written informed consent was obtained from the patient.

Discussion

Technological advances in thoracic surgery have enabled the increasing use of video-assisted and robotic-assisted surgery to perform thymectomy. Frequently cited benefits of robotic surgery for thymectomy include improved visualization of the gland and nearby structures, tremor reduction, decreased hospital length of stay (LOS), and better patient pain control (7).

The earliest report of a robotic thymectomy was in 2003 for a 28-year-old male with myasthenia gravis (8). Since then, a significant increase in utilization of robotic thymectomy has been reported. Burt *et al.* in their query of the National Cancer Database (NCDB) between 2010 and 2014 noted a doubling in the percentage of patients treated by minimally invasive thymectomy (MIT) (21.0% in 2010 to 40.2% in 2014, $P < 0.001$). They reported in their MIT group, a robotic approach accounted for 59.7% of cases (9).

In an attempt to avoid compromising oncologic outcomes with the increasing use of minimally invasive surgery, the International Thymic Malignancy Interest Group (ITMIG) established a precedent of nine principles for MIT. These include avoiding rib spreading or sternal cutting; ensuring a complete resection including the thymoma, thymus, and mediastinal fat; dissection and visualization of bilateral phrenic nerves and innominate vein; conversion to open if oncologic principles are at risk of being violated; an appropriate access incision size to prevent disruption of specimen; appropriate exploration of pleura if there is pleural involvement; retrieval of specimen in bag and evaluation of removed mass for completeness of resection; and communication with pathologist for suspicious areas, well demarcated tissue orientation, and areas of tissue disruption (10).

Critically, comparisons of R0 resections have been performed for open *vs.* MIT. Burt *et al.* analyzed an international thymectomy registry in 2017: the open (OT) approach (sternotomy or thoracotomy) was utilized in 2,053 (82%) patients while MIT (VATS or RATS) was used in 416 patients (18%). The rate of R0 resection was 86% and

94% ($P < 0.001$), respectively. This discrepancy, however, disappeared after propensity matching, which included variables such as geographic continent, time period, age, sex, paraneoplastic syndrome, WHO histologic type, Masaoka stage, tumor size, extent of thymectomy, and use of chemotherapy and radiotherapy; completeness of resection was not included (7). Notably, there was a higher rate of R1 and R2 resection in North America as compared to Asia; in a separate NCDB analysis by Burt *et al.* they noted a similar trend and attributed the continental differences to the relatively early experience of MIT in North America at that time. Rates of R0 resection increased over time (9). In their 2019 analysis of the NCDB, Burt and colleagues again found R0 resection was independent of surgical approach ($P = 0.17$) and MIT was associated with significantly shorter LOS (-1.03 days, 95% confidence interval (CI): -1.68 to -0.38) (9). More recently, Yang *et al.* performed a propensity-matched analysis of open versus MIT for stages I to III thymoma using the NCDB. Their propensity match included age, sex, race, education, facility type, insurance, Masaoka stage, tumor size and histology, induction therapy, and year of diagnosis; they reported shorter hospital LOS for MIT (3 *vs.* 4 days, $P < 0.01$) and no significant differences in margin positivity ($P = 0.84$), 30-day readmission ($P = 0.28$), 30-day mortality ($P = 0.60$), and 5-year survival (89.4% MIT *vs.* 81.6% open, $P = 0.20$). Their subgroup analysis of tumors < 4 cm and tumors > 4 cm also demonstrated no significant differences in margin positivity or overall survival. When robotic *vs.* VATS thymectomies were compared in the propensity-score analysis, there were no differences in LOS, perioperative outcomes, or overall survival (11).

Long term survival outcomes and risk of recurrence are difficult to assess as patients have long disease-free survivals (~ 10 years) after thymectomy for thymoma. As such, MIT studies have relatively short follow-up intervals. The Japanese Association for Research on the Thymus (JART) has performed the longest follow-up after MIT. After propensity matching patients with Masaoka stage I and II thymomas, median sternotomy (OT) and VATS thymectomy resulted in similar 5-year overall survival (OT = 97.1% *vs.* VATS = 97.0%) and recurrence free survival (OT = 95% *vs.* VATS = 93.9%) (12).

A systematic review comparing robotic, VATS, and open thymectomy was published by O'Sullivan and colleagues in 2019 (13). The authors noted significantly lower post-operative complication rates when comparing robotic *vs.* open thymectomy (OR: 0.37; 95% CI: 0.22–0.60, $P < 0.0001$;

heterogeneity $P=0.17$, $I^2=27\%$). There was no significant difference in overall mortality [risk difference (RD): -0.00 ; 95% CI: -0.02 to 0.02 , $P=0.86$] (13). Suda and colleagues compared single port VATS thymectomy (SPT) vs. multi-port robotic thymectomy (TRT) and concluded that both techniques are “equally minimally invasive” (14). They reported similar outcomes in terms of blood loss (5.9 ± 16.8 vs. 5.4 ± 4.6 mL; $P=0.48$), LOS (4.0 ± 2.0 vs. 4.3 ± 3.6 days; $P=0.21$), analgesic use (0.7 ± 5.4 vs. 10.1 ± 3.4 days; $P=0.89$), and complications (no intraoperative complications in either group; the SPT group had one case of left phrenic nerve paralysis and one case of transient paroxysmal atrial fibrillation). The VATS group had shorter operative times (35 ± 48 vs. 20 ± 40 min; $P=0.0004$).

There has been a recent focus on larger thymomas that were historically contraindicated for robotic thymectomy. Wilshire and colleagues performed a single-center retrospective review of patients who underwent a robotic vs. sternotomy approach for thymomas greater than 3 cm between 2004 and 2014 and found R0 resection was statistically equivalent (91% robotic and 88% open, $P=0.932$) and robotic surgery associated with shorter chest tube duration (1 vs. 3 days, $P=0.001$, respectively), intensive care unit (0 vs. 1 day, respectively $P=0.024$) and hospital LOS (2 vs. 5 days, $P<0.001$) (5). Kneuert and coworkers similarly conducted a retrospective, propensity-matched analysis based on age, sex, tumor size, and Masaoka stage for large (>4 cm) thymomas via robotic and transsternal approaches. They reported no difference in R0 resection rates between robotic (90%) and transsternal (85%, $P=0.62$) techniques (4).

A few cases of robotic thymectomy requiring pericardial reconstruction have been reported in the literature. The thoracic group at the University of Pittsburgh presented a similar case report for a 7 cm thymoma; they report *en bloc* resection of the mass along with pericardium and lung tissue for thymic squamous cell carcinoma. They employed TilePro software in order to enhance visualization of the phrenic nerve. Similarly, a 10 cm \times 10 cm pericardial defect was reconstructed with polytetrafluorethylene (Gore-Tex, Newark, DE) membrane patch and small cuts were made in the pericardial patch to allow for fluid drainage (15). The robotic platform allows for a facile reconstruction of pericardium (16). Various reports included use of a vented Gore-Tex patch for pericardial reconstruction, interrupted vs. running sutures, and suture type (proline vs. V-Loc) varied by institution (15,17). The thoracic department at University of North Carolina reported

routinely reconstructing pericardial defects greater than 3 cm \times 3 cm (18).

Our patient kindly shared the following thoughts on her pre-, peri-, and post-operative experience: “*My thymoma was found incidentally on a CT Calcium score of my heart ordered by my cardiologist. Once the thymoma was identified I was referred to a thoracic surgeon for removal. When I met the thoracic surgeon, Dr. Villamizar, for the first time, he explained that most surgeons would not attempt to remove a thymoma as large as mine robotically, but that he had experience with this approach and felt confident that he would be successful. The robotic removal would be less invasive with less downtime, easier recovery, less painful and less scarring were all factors that I considered when choosing the robotic route. My surgery was more difficult and more complicated than originally thought, as the thymoma was larger and more involved than was anticipated. Nonetheless, because of the surgeon’s tenacity and experience, he was successful in removing the tumor robotically. I stayed a total of three days in the hospital. I stopped taking pain medication, other than ibuprofen, the day I came home... [and continued to use ice packs and rest thereafter]. Despite still having some discomfort [...] after surgery, I believe that the robotic approach was the right choice for me. [I wonder] how much more pain might I have if the traditional thoracotomy had been necessary?*”

Conclusions

Robotic thymectomy continues to gain international acceptability across North America, Europe, and Asia. MIT has been associated with equivalent oncologic R0 resection rates while decreasing post-operative complications and pain requirements. Survival outcomes at 10 years are still under investigation. Our group and others have demonstrated that in experienced hands, the robotic approach is safe and has oncologic equivalence to median sternotomy for tumors larger than 5 cm in size, even if associated with invasion of surrounding structures. The type of approach should not compromise oncologic principles of achieving R0 resection; surgeons with inadequate experience should not attempt a complex thymectomy robotically if there is a significant risk of disrupting the capsule or performing an incomplete resection. There should be no hesitation for conversion to open if deemed necessary to preserve oncologic principles.

Acknowledgments

The authors would like to thank the University of Miami for support in writing this manuscript.

Funding: None.

Footnote

Reporting Checklist: The authors have completed the CARE reporting checklist. Available at <http://dx.doi.org/10.21037/med-20-52>

Peer Review File: Available at <http://dx.doi.org/10.21037/med-20-52>

Conflicts of Interest: All authors have completed the ICMJE uniform disclosure form (available at <http://dx.doi.org/10.21037/med-20-52>). The authors have no conflicts of interest to declare.

Ethical Statement: The authors are accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved. All procedures performed in studies involving human participants were in accordance with the ethical standards of the institutional and/or national research committee(s) and with the Helsinki Declaration (as revised in 2013). Written informed consent was obtained from the patient.

Open Access Statement: This is an Open Access article distributed in accordance with the Creative Commons Attribution-NonCommercial-NoDerivs 4.0 International License (CC BY-NC-ND 4.0), which permits the non-commercial replication and distribution of the article with the strict proviso that no changes or edits are made and the original work is properly cited (including links to both the formal publication through the relevant DOI and the license). See: <https://creativecommons.org/licenses/by-nc-nd/4.0/>.

References

- Hess NR, Sarkaria IS, Pennathur A, et al. Minimally invasive versus open thymectomy: a systematic review of surgical techniques, patient demographics, and perioperative outcomes. *Ann Cardiothorac Surg* 2016;5:1-9.
- Fiorelli A, Mazzella A, Cascone R, et al. Bilateral thoracoscopic extended thymectomy versus sternotomy. *Asian Cardiovasc Thorac Ann* 2016;24:555-61.
- Odaka M, Shibasaki T, Kato D, et al. Comparison of oncological results for early- and advanced-stage thymomas: thoracoscopic thymectomy versus open thymectomy. *Surg Endosc* 2017;31:734-42.
- Kneuert PJ, Kamel MK, Stiles BM, et al. Robotic Thymectomy Is Feasible for Large Thymomas: A Propensity-Matched Comparison. *Ann Thorac Surg* 2017;104:1673-8.
- Wilshire CL, Vallières E, Shultz D, et al. Robotic Resection of 3 cm and Larger Thymomas Is Associated With Low Perioperative Morbidity and Mortality. *Innovations (Phila)* 2016;11:321-6.
- Gagnier JJ, Kienle G, Altman DG, et al. The CARE guidelines: consensus-based clinical case reporting guideline development. *J Med Case Rep* 2013;7:223.
- Burt BM, Yao X, Shrager J, et al. Determinants of Complete Resection of Thymoma by Minimally Invasive and Open Thymectomy: Analysis of an International Registry. *J Thorac Oncol* 2017;12:129-36.
- Ashton RC Jr, McGinnis KM, Connery CP, et al. Totally endoscopic robotic thymectomy for myasthenia gravis. *Ann Thorac Surg* 2003;75:569-71.
- Burt BM, Nguyen D, Groth SS, et al. Utilization of Minimally Invasive Thymectomy and Margin-Negative Resection for Early-Stage Thymoma. *Ann Thorac Surg* 2019;108:405-11.
- Toker A, Sonett J, Zielinski M, et al. Standard terms, definitions, and policies for minimally invasive resection of thymoma. *J Thorac Oncol* 2011;6:S1739-42.
- Yang CFJ, Hurd J, Shah SA, et al. A National Analysis of Open vs Minimally Invasive Thymectomy for Stage I-III Thymoma. *J Thorac Cardiovasc Surg* 2020;160:555-567. e15.
- Agatsuma H, Yoshida K, Yoshino I, et al. Video-Assisted Thoracic Surgery Thymectomy Versus Sternotomy Thymectomy in Patients With Thymoma. *Ann Thorac Surg* 2017;104:1047-53.
- O'Sullivan KE, Kreaden US, Hebert AE, et al. A systematic review of robotic versus open and video assisted thoracoscopic surgery (VATS) approaches for thymectomy. *Ann Cardiothorac Surg* 2019;8:174-93.
- Suda T, Kaneda S, Hachimaru A, et al. Thymectomy via a subxiphoid approach: single-port and robot-assisted. *J Thorac Dis* 2016;8:S265-71.
- Kang CH, Na KJ, Song JW, et al. The robotic thymectomy via the subxiphoid approach: technique and early outcomes. *Eur J Cardiothorac Surg* 2020;58:i39-i43.
- Schneiter D, Tomaszek S, Kestenholz P, et al. Minimally invasive resection of thymomas with the da Vinci® Surgical System. *Eur J Cardiothorac Surg* 2013;43:288-92.

17. Yang HC, Coyan G, Vercauteren M, et al. Robot-assisted en bloc anterior mediastinal mass excision with pericardium and adjacent lung for locally advanced thymic carcinoma. *J Vis Surg* 2018;4:115.
18. Espey J, Acosta S, Kolarczyk L, et al. Case report: cardiac herniation following robotic-assisted thymectomy. *J Cardiothorac Surg* 2020;15:54.

doi: 10.21037/med-20-52

Cite this article as: Kodia K, Nguyen DM, Villamizar NR. A 9 cm robotic thymectomy and pericardial repair case report. *Mediastinum* 2020;4:38.